

## Natural Philosophy of Protoplanetary and Planetary Discs: On the Origin and Evolution of Life

XIAOMING LI

**Abstract:** This paper examines the four pivotal and most contentious issues within the field of abiogenesis by analyzing the life-originating processes centred on protoplanetary and planetary discs. These include: 1. Geological environment; 2. Source of nutrients/initial molecules; 3. Source of energy; 4. RNA world versus metabolic world: which came first? To date, scientists have been unable to reach a consensus on these questions. Regarding geological environment, the distinct conditions within the inner, middle, snow line, and outer regions of the protoplanetary disc—including solar, cosmic, and isotopic radiation—along with the exchange of dust and gases between these zones, provided an exceptional reaction platform for prebiotic chemistry. Concerning the origin of life's food/precursor molecules, the protoplanetary disc—the stellar gas and dust disc formed during star formation—served as a crucial transitional stage linking interstellar matter to life's chemical potential. Transition metals, short- and long-lived radioactive isotopes, polycyclic aromatic hydrocarbons (PAHs), fullerenes, and the metal-organic compounds formed from these substances play a crucial role in life's composition. Analysis of meteorites reinforces the credibility of the protoplanetary and planetary discs as potential sites for the origin of life. Quantum mechanics may significantly influence the possibility of life emerging within protoplanetary discs. Regarding energy sources, the energy required for the emergence of life is quantifiable. The energy driving life's birth and evolution within protoplanetary and planetary discs can be summarized by the equations  $F = ma$  and  $\Delta E = \Delta m \times c^2$ . These energy sources derive not only from the kinetic energy of the protoplanetary disc and interactions between dust particles, expressed by  $F=ma$ , but this kinetic energy can also be converted into chemical reactions, ultimately yielding prebiotic effects. Energy generated by radiation from the decay of radioactive isotopes within the protoplanetary and planetary discs, alongside energy produced by solar nuclear fusion, can be expressed by the formula  $\Delta E = \Delta m \times c^2$ . This energy not only generates heat but, crucially, produces radiolysis effects through solar radiation, cosmic rays, and isotope decay. These effects facilitate prebiotic chemical reactions and provide sustenance and nutrients for life within the planetary disc. Concerning the RNA-world versus metabolism-first debate, the high concentrations of PAHs and fullerenes in protoplanetary discs and planetary discs, coupled with their influence on the formation of aromatic amino acids and aromatic hydrophobic proteins affecting nucleic acid molecular chirality, underscore the metabolism-first theory. The discovery of Hemolithin in meteorites and its implications for RNA generation reinforce the "metabolism precedes genetics" theoretical framework. Starting from Hemolithin and primitive metabolic

systems found in meteorites, lipid membrane vesicles encapsulated metabolic small molecules and informational polymers. Sustained by external energy sources, these gradually evolved into primitive systems possessing the tripartite functions of "replication–metabolism–boundary". It subsequently led to the integration of energy, protein metabolism, and membrane–genetic systems within protoplanetary and planetary discs. Ultimately, a self-maintaining and self-replicating cycle system was established within the planetary disc. The core emphasis of this paper lies in the continuity and consistency, spanning hundreds of millions of years, of the gas and molecular dust composition within the protoplanetary disc and the gases and molecules that could be generated within planetary disc planetesimals. It provided a stable energy source and an excellent sanctuary for the birth of life within the protoplanetary disc and the evolution of life within planetary disc planetesimals. The philosophical coherence that life should emerge wherever gases and molecules exist elucidates the availability of protoplanetary disc gases and molecules (such as H<sub>2</sub>, CO, and electrons) alongside the bacteria consuming these substances within the disc's planetesimals. These gases and molecules are generated within planetary disc planetesimals by radioactive isotopes (such as <sup>26</sup>Al, <sup>60</sup>F, <sup>238</sup>U, <sup>40</sup>K), simultaneously producing oxygen radicals and hydrogen peroxide (e.g., H<sub>2</sub>O<sub>2</sub>, •OH, O<sub>2</sub>•<sup>-</sup>, NO<sub>3</sub><sup>-</sup>). Thus, isotopes provided both the energy for bacterial survival and the nutrients required for bacterial metabolism, while simultaneously generating oxygen-free radicals and hydrogen peroxide that constituted a hazard to these bacteria. Consequently, bacteria acquired the capacity to resist oxygen-free radicals and hydrogen peroxide while obtaining the nutrients essential for their metabolism. The core of this process lies in the "radiation-metabolism coupling" and "radiation-metabolism co-evolution" hypotheses concerning the origin of life in protoplanetary and planetary disks. Within this theoretical framework, radiation-generated gases, molecules, and oxygen free radicals coexisted long-term with bacterial metabolism and antioxidant functions, forming the "radiation-metabolism co-evolution" model for life's origin. Only this "radiation environment-metabolism co-evolution" hypothesis can explain the survival characteristics exhibited by all bacteria and viruses we observe today. The fundamental reason why bacteria, archaea, and viruses possess exponentially higher radiation resistance than mammals lies in the direct consequence of these microorganisms' prolonged coexistence with radiation within planetary disks. Affected by the reducing atmosphere of the protoplanetary disk, the internal environment of planetesimals in the protoplanetary disk is mainly reduced, while the internal environment of planetesimals in the planetary disk is mainly oxidized due to the influence of short- and long-lived isotopes. Evidence from the significantly higher abundances of short-lived radioactive isotopes (SLRs) and long-lived radioactive isotopes (LLRs) in Murchison meteorite and other meteorites compared to the Earth's crust has highlighted the central theme of 'radiation-

metabolism co-evolution'. Research on the Great Oxygenation Event (GOE) demonstrates that this event, dominated by cyanobacteria, represents a consistent progression from 'radiation-metabolism coupling' and 'radiation-metabolism co-evolution' to radiation-metabolism-atmosphere environmental change. Comparing sulphur isotope mass-independent fractionation (S-MIF,  $\Delta^{33}\text{S}$ ) before and after the Great Oxygenation Event (GOE) with planetary disc material reveals that the GOE effectively restored Earth's S-MIF state to the oxidized conditions present in the planetary disc. It further demonstrates that bacteria originating from the planetary disc transformed Earth's atmosphere into the oxidized environment they inhabited within the disc – the very environment that sustains our existence today. Finally, it is discussed that the highly efficient aerobic metabolism of eukaryotes is also directly descended from the antioxidant bacteria present in protoplanetary disc planetesimals. Migratory birds, which traverse Earth's magnetic field lines, share connections with bacterial antioxidant mechanisms and primordial genetic material. Finally, constructing a novel Tree of Life reveals that the evolutionary lineage of life differs entirely from our currently accepted model. The roots and trunk of this evolutionary tree do not originate on Earth, but rather on the planetary disc. Two branches diverge from the trunk: bacteria and archaea. The archaea branch further gives rise to a branch of eukaryotes, constituting the primary life forms on our planet today. Meanwhile, bacteria, archaea, and viruses continue to evolve both on the planetary disc and on Earth, exerting influence upon the life forms evolving on our planet. Given that fullerene ( $\text{C}_{60}$ ) itself exhibits quantum wave-particle duality, akin to the quantum effects observed in light, electrons, protons, and neutrons, the associated electro-nutritive bacteria—which convert electrons into nutrients—also possess quantum properties. It constitutes the pivotal link in this paper's integration of quantum mechanics into the origins of life. This analytical discussion further suggests that the essence of life and consciousness may fundamentally be quantum in nature. Under this premise that the essence of life and consciousness is quantum, the boundaries between physics, life, and consciousness may dissolve. At a deeper philosophical level, electro-nutritive life embodies the biological manifestation of quantum order. The electron resonance, coherence, and wave-particle duality exhibited by fullerene do not vanish with the emergence of life but gain structural continuity at the cellular level. It indicates that life's metabolism and consciousness are not entities 'transcending' the physical, but rather self-organized stages of the physical quantum order. From the cosmos to life, electron flow constitutes the sole truly continuous language of energy. As 'cosmic quantum energy nodes', fullerenes become, in a metaphysical sense, the mediating symbols between matter and life. Life's existence depends on the flow of energy. From stellar radiation to molecular chemistry, through neural activity to conscious experience, energy remains uninterrupted, merely organized and encoded at different levels. At the cosmos's inception, energy existed as quantum field oscillations. Within protoplanetary discs, it

underwent physical transformation through electron flow via fullerenes, PAHs, and Fe-S systems; within living organisms, this electron flow becomes the carrier of information, driving neural activity and metabolic equilibrium. Thus, spanning from the physical to the biological, from molecules to consciousness, there exists a continuous "quantum electron flow continuum"—one that not only conveys energy but also carries information, forming the quantum unity of life and consciousness. Life is the process by which cosmic energy achieves self-organization and self-awareness through the flow of quantum electrons. From this perspective, consciousness does not "emerge within life"; rather, it is the manifestation of the universe's own quantum order attaining a perceptible state within living organisms. Quantum conductor molecules such as fullerenes represent the earliest form of this order—already "thinking" quantum-mechanically in the lifeless cosmos, storing energy, transmitting information, and creating structures. Fullerenes are the quantum expression of energy; electro-nutritive microorganisms are the metabolic expression of energy flow; neural networks are the cognitive expression of energy-information. There is no discontinuity between them, only transitions in complexity. Thus, the universality of life lies in its being not an exception to the cosmos, but a form of cosmic energy becoming self-aware. The overarching trend of the universe becoming life—its evolution from quantum fields, stars, and molecules to organisms—can be viewed as a progressive increase in the density of energy and information. Within this framework, life is not a localized event, but a natural stage in the cosmic complexity of information. In other words, the universe progressively 'becomes life' during its evolution.

Ultimately, this paper establishes three laws governing the origins of life at distinct levels: the Solar System Law, the Stellar System Law, and the Cosmic Life Law. The Solar System Law elucidates the unity of life discovered across planets and asteroids within our solar system, demonstrating that all life forms observed here are unified by the "Central Law". The Stellar System Law states that on any planetary body within a stellar system, even if primitive life exists, without a "Great Oxidation Event" akin to the one facilitated by cyanobacteria, life would remain confined to prokaryotic forms and never evolve. Such life would vanish alongside the stellar system's demise. The Cosmic Life Law elucidates the protoplanetary disc as a decisive factor in the formation of amino acid chirality. It further posits that quantum life and quantum consciousness constitute the very essence of cosmic life and consciousness.

**Keywords:** Origin of Life, Evolution of Life, Natural Philosophy, Philosophy of Biology, Dust in the Universe, Protoplanetary Disc, Planetary Disc, Planetesimal, PAHs, SLRs, LLRs, S-MIF, Radiation-Metabolism Coupling, Radiation-Metabolism Co-evolution, Great Oxidation

Event, Radiation-Metabolism-Atmospheric Change- Emergence of Eukaryotes, Fullerenes and Quantum Mechanics, Quantum Chemistry, Quantum Biology, Quantum–Biological Coupling, Quantum Life, Quantum Organic Conscious Entity, Quantum Information Dynamics, Tree of Life, Planetary Disc Panspermia

University of Toronto, Department of Molecular Genetics

### **Introduction:**

Three fundamental questions of origin exist: the origin of the universe, the origin of life (OoL), and the origin of mind and consciousness. All these questions have been debated for thousands of years and are now subtopics of serious scientific research<sup>1</sup>. The perspective that Earth's life originated from Earth itself, also known as "endogenic origin" or "abiogenesis," is one of the mainstream theories of life's origin in the life sciences. This theory posits that life on Earth evolved gradually from inorganic matter through natural processes, without reliance on extraterrestrial life or imported organic material. However, whether it be the "primordial soup, Oparin–Hordern hypothesis"<sup>2</sup>; the "Miller–Urey experiment"<sup>3</sup>; the "RNA world hypothesis"<sup>4</sup>; or the "hydrothermal vent theory",<sup>5</sup> these represent the mainstream theories for the origin of life on Earth from within the planet itself.

An alternative theory proposed by a minority of scientists is Panspermia, which posits that life did not originate solely on Earth but is ubiquitous throughout the cosmos. Such life forms could be microorganisms or simpler organic entities<sup>6</sup>. Panspermia suggests that microbes or seeds of life could attach to cosmic dust, comets, meteorites, or other celestial bodies, propagating through space.<sup>7</sup> These celestial bodies may have transported life from one planet in the cosmos to another, including Earth.

Both theories of life's origin, in my view, have made significant contributions to understanding its emergence. Scientists have elucidated the potential causes of life's origin through diverse approaches and perspectives. In 1859, Darwin published "On the Origin of

---

<sup>1</sup> JACK W. SZOSTAK, ON THE ORIGIN OF LIFE, MEDICINA (Buenos Aires) 2016; 76: 199-203

<sup>2</sup> Haldane, J. B. S. (1929). "The Origin of Life." Rationalist Annual, 148, 3-10.

<sup>3</sup> Miller, S. L. (1953). "A Production of Amino Acids Under Possible Primitive Earth Conditions." Science, 117(3046), 528-529.

<sup>4</sup> Joyce, G. F. (2002). "The Antiquity of RNA-Based Evolution." Nature, 418(6894), 214-221.

<sup>5</sup> Russell, M. J., Hall, A. J., & Martin, W. (2010). "Serpentinisation as a Source of Energy at the Origin of Life." Geobiology, 8(5), 355-371.

<sup>6</sup> Wickramasinghe, N. C. (2011). Theories of Cosmic Dust. Springer Science & Business Media.

<sup>7</sup> Hoyle, F., & Wickramasinghe, N. C. (1979). Lifecloud: The Origin of Life in the Universe. J.M. Dent & Sons.

Species”<sup>8</sup>. Darwin's evolutionary tree features a single trunk, representing the scientific consensus that all life forms originated from a common ancestor—a theory termed monogenesis or the single-origin theory. This theory posits that all modern organisms can be traced back to a single-celled organism that emerged on Earth approximately 3.5 billion years ago. Thus, the single trunk of the evolutionary tree symbolizes this common ancestor, while the branches and leaves represent subsequent differentiation and evolution.

Scientists are increasingly converging on a consensus that Earth's life emerged through a combination of "exogenous delivery" of organic matter and "endogenous synthesis" via chemical evolution. Exogenous delivery refers to the process by which organic compounds and prebiotic molecules from space enter Earth via celestial bodies, such as meteorites and comets. Endogenous synthesis on Earth refers to the direct generation of organic molecules and prebiotic molecules through physical or chemical processes. These include diverse energy sources such as shock waves, ultraviolet radiation, electrical discharges, and hydrothermal vents, which may have catalyzed the synthesis of more complex organic compounds from simpler molecules. These two processes complemented each other, collectively accomplishing the origin of life. However, scientists have yet to piece together the whole puzzle of life's origins. The most contentious debates among contemporary scientists regarding life's origins centre on four key points: 1. Geological environment; 2. Source of nutrients/initial molecules; 3. Source of energy; 4. The distinction between the RNA world and metabolism-first scenarios<sup>9</sup>. To date, no consensus has been reached on any of these issues.

The preceding article (Chemistry Analysis of Origin of Life in Protoplanetary Disks) primarily discusses the possibility of life originating in protoplanetary disks from a chemical perspective. It highlights the environmental and water paradoxes inherent in the theory of life originating on Earth<sup>10</sup>. It contends that life is highly improbable to have arisen on Earth or any terrestrial planet, but rather is highly probable to have emerged within protoplanetary discs. Having established the possibility of life originating in protoplanetary discs, it becomes imperative to elucidate the associated prebiotic chemical mechanisms. It would further clarify the various potential pathways for life's origin within these discs. My reasoning and hypotheses are derived from analyses of existing scientific literature.

---

<sup>8</sup> Darwin, C. (1859). *On the Origin of Species by Means of Natural Selection, or the Preservation of Favoured Races in the Struggle for Life*. John Murray.

<sup>9</sup> Patel, Bhavesh H., Claudia Percivalle, Dougal J. Ritson, Colm D. Duffy, and John D. Sutherland. 2020. "The Future of Origin of Life Research: Bridging Decades-Old Divisions." *Life (Basel)* 10 (3): 20. <https://doi.org/10.3390/life10030020>.

<sup>10</sup> Li, XM. (2024). *Chemistry Analysis for the Origin of Life in Protoplanetary Disks*. Geoinfor Geostat An Overview Vol: 12 Issue: 5

Previous articles have discussed the origin of life within protoplanetary discs, encompassing viruses, archaea, and bacteria. Given that bacteria may have emerged within protoplanetary discs. The mechanisms related to cellular metabolism, cell membrane formation, and the establishment of cellular genetic mechanisms should be considered. The synthesis of amino acids, the generation of cell membranes, the production of RNA and DNA, and the mechanisms for translating genetic codes into proteins are all critical issues concerning the origin of life that require discussion.

This article primarily analyses the origin of life within protoplanetary discs and planetary discs from a prebiotic chemistry perspective, along with its evolution within planetary disc planetesimals. It also examines processes and principles related to quantum chemistry, quantum biology, cosmic environmental science, astrobiology, cosmic mechanics, and radiobiology as they pertain to protoplanetary and planetary discs. Concurrently, it discusses key issues concerning the environmental characteristics of the solar system's protoplanetary and planetary discs, as well as their relevance to the metabolism of life within the context of prebiotic chemistry.

This paper concentrates on four interrelated and most contentious themes concerning life's origin: (i) the environment in which life originated; (ii) the source of food or initial organic molecules; (iii) the source of energy; (iv) the divergence between the RNA world hypothesis and the metabolic primacy hypothesis. Concurrently, two further questions are addressed, which it considers more significant: (v) the establishment of life's metabolic networks, encompassing the central dogma of protein synthesis and translation, the formation of cell membranes, and the emergence of genetic systems; (vi) the origin of life within protoplanetary discs and the mechanisms governing its evolution within planetary disc planetesimals.

Because fullerene ( $C_{60}$ ) itself exhibits quantum wave-particle duality, akin to the quantum effects observed in light, electrons, protons, and neutrons, the associated electro-consuming bacteria—which convert electrons into nutrients—also possess quantum properties. The very essence of life and consciousness may well be quantum in nature. Under this premise that the essence of life and consciousness is quantum, the boundaries between physics, life, and consciousness may dissolve. The universality of life lies in its being not an exception within the cosmos, but rather a form of cosmic energy becoming self-aware. The overarching trend of life-forming within the universe—its evolution from quantum fields, through stars and molecules, to biological organisms—can be viewed as a progressive increase in the density of energy-information. Within this framework, life is not a localized event, but a natural stage in the cosmic progression of information complexity. In other words, the universe progressively 'becomes alive' during its evolution. The emergence

of these organic compounds aligns with the principles governing the origin of life within Earth's protoplanetary disc, as understood through physics, quantum mechanics, quantum chemistry, chemistry, and prebiotic chemistry.

This paper begins with the atomic constituents present in the early universe, progresses through the evolution of atomic composition within the cosmos, and then examines how supernova explosions gave rise to the protoplanetary disc of the Solar System, ultimately fostering the conditions for life. The concluding section focuses on panspermia theories grounded in protoplanetary and planetary discs, specifically "planetary disc panspermia." It explores the potential structure of the tree of life based on the Solar System's protoplanetary and planetary discs.

### **Discussion:**

#### **(i) The Environment of Life's Origin in the Protoplanetary Disk:**

This article primarily explores the environment in which life originated, one of the core questions in origin-of-life theories. Essentially, it addresses the fundamental question of where life actually emerged. In the preceding article, I concluded by discussing the impossibility of life originating on Earth and the various paradoxes inherent in Earth-centred theories of life's origin. That is, life did not originate on Earth or any Earth-like planet, but rather within protoplanetary disks. Guided by this conclusion, I must now examine the conditions necessary for life's emergence within the protoplanetary disc. These conditions encompass: (i) the environmental parameters of the origin site; (ii) the source of food or initial organic molecules; and (iii) the origin of energy. These represent the core problems requiring resolution in the origin of life. This article primarily addresses the environmental aspects of life's origin, which also constitutes the foundational question. Suppose we presuppose that life originated exclusively on Earth. In that case, we remain forever bound by the constraints of our own thinking, perpetually confined to the narrow and limited spatial and geological conditions of Earth's environment.

This section constitutes one of the most crucial parts of this article. It primarily explores how the protoplanetary disc provides the necessary conditions for the origin of life. It examines the entire environmental chain: from the initial growth of life-forming substances within the protoplanetary disc, through the emergence of life itself, to the eventual arrival of life (such as bacteria) on Earth via meteorites originating from the planetary disc. It necessitates an investigation into astrophysics, astrochemistry, protoplanetary disc physics, protoplanetary disc chemistry, and protoplanetary disc biology. If we distil this to a single critical point, it is the science of protoplanetary disc environments. Protoplanetary disc environmental



science investigates explicitly the environmental conditions that facilitate the origin of life within protoplanetary discs. It stands in contrast to the various environmental hypotheses concerning the origin of life on Earth, which are currently the focus of widespread research among scientists studying the origins of life.

This chapter aims to construct all conceivable environmental systems within the protoplanetary disc that could potentially influence the origin of life there. This endeavour seeks to establish a comprehensive, logically coherent framework for protoplanetary disc biogenesis that reflects scientific reality. It lays the groundwork for future research into protoplanetary disc biogenesis by creating a self-contained scientific and theoretical system.

This paper will explore the astrophysical environment of the Solar System, particularly the decisive influence of supernova explosions on the emergence of life within protoplanetary discs; the chemistry of the Solar System's protoplanetary disc, especially the chemical conditions relevant to the origin of life; the impact of physical factors within protoplanetary discs on the emergence of life; and the potential influence of astrobiology on the origin of life within protoplanetary discs.

## **1. The Origin of Protoplanetary Discs**

The origins of stars and planets are inextricably linked, both emerging from dense molecular clouds within the interstellar medium. Under the influence of gravitational forces, these molecular cloud clusters undergo collapse due to localized density increases, ultimately forming young stars. During this process, governed by the conservation of angular momentum, the collapsing material gradually spreads into a flattened, rotating structure within the equatorial plane—the protoplanetary disk<sup>11</sup>. This structure not only marks a crucial stage in stellar formation but also serves as the primary site for the formation of planets, comets, minor bodies, and precursor molecules for life.

The protoplanetary disc primarily consists of gas (predominantly hydrogen and helium) and dust particles, typically extending to several hundred astronomical units (AU), with a lifespan of approximately one to ten million years. During this period, dust particles within the disc undergo continuous collisions and aggregation, gradually forming embryonic celestial bodies ranging from micrometre to kilometre scale, which subsequently evolve into complete planetary systems. However, beyond its role in planetary formation, the protoplanetary disc also serves as a natural laboratory brimming with potential for complex

---

<sup>11</sup> Shu, F. H., Adams, F. C., & Lizano, S. (1987). Star formation in molecular clouds: Observation and theory. *Annual Review of Astronomy and Astrophysics*, 25, 23–81.  
<https://doi.org/10.1146/annurev.aa.25.090187.000323>

chemical evolution. Observations and meteorite sample studies reveal that these discs are not only rich in organic molecules (such as formaldehyde, HCN, polycyclic aromatic hydrocarbons PAHs, etc.)<sup>12</sup> but also contain significant quantities of transition metal elements (e.g., Fe, Ni, Co), short-lived radioactive isotopes (e.g., <sup>26</sup>Al, <sup>60</sup>Fe), and long-lived radioactive isotopes (such as potassium-40 (<sup>40</sup>K), uranium-238 (<sup>238</sup>U), uranium-235 (<sup>235</sup>U), and thorium-232 (<sup>232</sup>Th) provide catalysis and energy sources for diverse chemical reactions<sup>13</sup>.

Moreover, the surfaces of dust particles within protoplanetary discs function as 'catalytic carriers', facilitating the adsorption, directed alignment, and reactive polymerization of simple small molecules. The thermal energy released by high-energy radiation (X-rays, ultraviolet, gamma rays) and radioactive decay serves as a crucial energy injection mechanism driving these reactions.

Taking the Solar System as an example, approximately 4.6 billion years ago, shockwaves from one or more supernova explosions triggered the local collapse of the molecular cloud within which the Solar System formed, subsequently giving rise to the young Sun and its surrounding protoplanetary disc<sup>14</sup>. During the early stages of protoplanetary disc formation, as high temperatures gradually cooled and organic compounds accumulated and differentiated, ideal conditions were created for the transition of inorganic molecules into higher-order organic molecules (such as amino acids, sugars, and nucleosides). Some models even propose that early life components, such as lipid vesicles and RNA precursors, may have formed within the protoplanetary disc itself, subsequently being 'seeded' onto the young Earth via later planetary transport or cometary impacts<sup>15</sup>.

In summary, protoplanetary discs are not merely by-products of stellar system formation but may well serve as "chemical incubators" for life's origins. However, to comprehend "where life came from," a crucial scientific question remains unanswered: How precisely did the protoplanetary disc environment provide conditions for the precursors of life chemistry? This study will systematically explore the role and potential of protoplanetary discs in prebiotic chemical evolution by examining the mechanisms of organic molecule synthesis,

---

<sup>12</sup> Herbst, E., & van Dishoeck, E. F. (2009). Complex Organic Interstellar Molecules. *Annual Review of Astronomy and Astrophysics*, 47, 427–480. <https://doi.org/10.1146/annurev-astro-082708-101654>

<sup>13</sup> Boss, A. P. (1995). Collapse and fragmentation of molecular cloud cores. *Astrophysical Journal*, 439, 224. <https://doi.org/10.1086/175167>

<sup>14</sup> Cameron, A. G. W., & Truran, J. W. (1977). The Supernova Trigger for Formation of the Solar System. *Icarus*, 30(3), 447–461.

<sup>15</sup> Ehrenfreund, P., & Charnley, S. B. (2000). Organic Molecules in the Interstellar Medium, Comets, and Meteorites: A Voyage from Dark Clouds to the Early Earth. *Annual Review of Astronomy and Astrophysics*, 38, 427–483. <https://doi.org/10.1146/annurev.astro.38.1.427>

metal-catalyzed reactions, the effects of high-energy radiation, and mineral surface selectivity.

## **2. Structure and Evolutionary Characteristics of Protoplanetary Discs**

Protoplanetary discs constitute an indispensable component of stellar formation, exhibiting not only complex physical structures but also pronounced temporal evolution and spatial differentiation. Understanding the physical mechanisms governing protoplanetary discs holds fundamental significance for elucidating the origins and evolutionary trajectories of prebiotic chemical reactions.

### **2.1 Formation of the Flat Structure: An Inevitable Consequence of Angular Momentum Conservation**

The flattened morphology of protoplanetary discs originates from the physical effect of angular momentum conservation during the collapse of molecular clouds. As gravity causes a molecular cloud to collapse gradually, its initial weak rotation rapidly intensifies. Material within the cloud converges towards the centre, while centrifugal forces constrain motion along the rotational axis. Consequently, matter accumulates predominantly within the equatorial plane, forming a structure that thins vertically and expands horizontally<sup>16</sup>.

This disc, formed by the redistribution of angular momentum, is commonly referred to as an accretion disc. During the early stages of star formation, this disc continuously supplies material to the central star. Through mechanisms such as viscous forces, angular momentum is progressively transferred outward, thereby permitting the inward collapse of the material<sup>17</sup>.

### **2.2 Accretion Processes and Temperature Gradient Distribution**

Accretion serves as the primary mechanism for stellar mass growth while also driving the formation of complex structures and chemical differentiation within protoplanetary discs<sup>18</sup>. During accretion, material flows inward, releasing potential energy that is converted into heat, thereby generating a temperature gradient that decreases progressively from the center outward<sup>19</sup>.

---

<sup>16</sup> Williams, J. P., & Cieza, L. A. (2011). Protoplanetary Disks and Their Evolution. *Annual Review of Astronomy and Astrophysics*, 49, 67–117. <https://doi.org/10.1146/annurev-astro-081710-102548>

<sup>17</sup> Armitage, P. J. (2011). Dynamics of Protoplanetary Disks. *Annual Review of Astronomy and Astrophysics*, 49, 195–236. <https://doi.org/10.1146/annurev-astro-081710-102521>

<sup>18</sup> D'Alessio, P., Canto, J., Calvet, N., & Lizano, S. (1998). Accretion disks around young objects: I. Grain growth. *Astrophysical Journal*, 500, 411. <https://doi.org/10.1086/305698>

<sup>19</sup> Chiang, E. I., & Goldreich, P. (1997). Spectral energy distributions of T Tauri stars with passive circumstellar disks. *Astrophysical Journal*, 490, 368. <https://doi.org/10.1086/304869>

- Inner disc regions (<1 AU) may reach temperatures of several thousand Kelvin, where gas exists in atomic or ionized states, favouring the formation of high-temperature refractories such as calcium-aluminum inclusions (CAIs)<sup>20</sup> ;
- Intermediate zone (1–10 AU) Moderate temperatures facilitate gas molecular reactions, organic synthesis, and surface catalysis on minerals;
- Outer Disk Region (>10 AU) Temperatures drop to tens of kelvins, rich in ice, volatiles, and carbon-chain molecules, favourable for the aggregation of complex organic molecules (e.g., methanol, ethanol, PAHs).

This temperature gradient induces strong spatial dependence in material distribution, phase transitions, and chemical reaction pathways, forming a crucial backdrop for molecular complexity and the prebiotic selective formation of life.

### 2.3 Lifecycle and Physical Segmentation

The lifecycle of a protoplanetary disc typically spans approximately 1 to 10 million years (Myr), influenced by the combined effects of stellar radiation pressure, disc winds, photoevaporation, and the formation of planetary embryos<sup>21</sup> . Observations indicate that substantial quantities of dust and gas persist within the disc for one to two million years after star formation. By 5–10 million years, the gaseous discs in most systems gradually dissipate, transitioning into debris discs or vanishing entirely<sup>22</sup> . Structurally, protoplanetary discs can be divided into several characteristic regions<sup>23</sup> :

- Inner disk (<1 AU): High-temperature, close to the star, forming metal oxides, silicate crystals, and evaporating volatiles;
- Middle disc (1–10 AU): Reaction conditions most favourable for complex organic chemistry and the aggregation of protoplanetary embryos;
- Near the ice line: Critical zone where water, methane, ammonia and other icy molecules condense, forming a chemical enrichment belt (<sup>24</sup>);

---

<sup>20</sup> Dullemond, C. P., & Monnier, J. D. (2010). The inner regions of protoplanetary disks. *Annual Review of Astronomy and Astrophysics*, 48, 205–239.

<sup>21</sup> Williams, J. P., & Cieza, L. A. (2011). Protoplanetary disks and their evolution. *Annual Review of Astronomy and Astrophysics*, 49, 67–117. <https://doi.org/10.1146/annurev-astro-081710-102548>

<sup>22</sup> Haisch, K. E., Lada, E. A., & Lada, C. J. (2001). Disk frequencies and lifetimes in young clusters. *Astrophysical Journal Letters*, 553, L153–L156. <https://doi.org/10.1086/320685>

<sup>23</sup> Andrews, S. M., et al. (2009). Discovery of a large cavity in the disk around LkHa 330. *Astrophysical Journal*, 700, 1502. <https://doi.org/10.1088/0004-637X/700/2/1502>

<sup>24</sup> Öberg, K. I., Murray-Clay, R., & Bergin, E. A. (2011). The Effects of Snowlines on C/O in Planetary Atmospheres. *Astrophysical Journal Letters*, 743(1), L16. <https://doi.org/10.1088/2041-8205/743/1/L16>

- Outer Disk (>10 AU): Low temperatures and high densities favour gas adsorption onto icy dust particles and promote surface chemical reactions.

## 2.4 Supernova-injected short-lived radionuclides and long-half-life isotopes

During the early formation of the Solar System, several short-lived radionuclides (SLRs) were detected within the protoplanetary disc, such as  $^{26}\text{Al}$  (half-life  $\sim 0.7$  Myr) and  $^{60}\text{Fe}$  (half-life  $\sim 2.6$  Myr). These radionuclides cannot be synthesized within the Sun and most likely originated from one or more nearby supernova explosions that injected material into the molecular cloud or directly into the protoplanetary disc<sup>25</sup>.  $^{26}\text{Al}$  provided an internal heat source for planetesimals and protoplanets, inducing the formation of hydrated minerals and organic rearrangement reactions.  $^{60}\text{Fe}$  is recognized as a distinct supernova product; its presence suggests that the solar system formed in proximity to high-energy cosmic events<sup>26</sup>. Their decay not only delivers sustained thermal energy release over millions of years (approximately tens of kJ/mol) but also accompanies  $\beta$ -ray and gamma-ray radiation, potentially providing excitation mechanisms for the synthesis, cleavage, and repolymerization of organic molecules<sup>27</sup>. This exogenous injection mechanism enhances the chemical reactivity of the protoplanetary disc, establishing its energy foundation as a "prebiotic chemical reactor"<sup>28</sup>.

During the early formation of protoplanetary discs, long-lived radionuclides such as potassium-40 ( $^{40}\text{K}$ ), Uranium-238 ( $^{238}\text{U}$ ), Uranium-235 ( $^{235}\text{U}$ ), and Thorium-232 ( $^{232}\text{Th}$ ) were present. These long-lived radionuclides, inherited from the nucleosynthesis of progenitor stars, became incorporated into primordial dust and mineral grains, subsequently accumulating within protoplanetary bodies as nebular material coalesced. The continuous radioactive decay of these isotopes generates significant radiogenic heating within small planetesimals that lack effective mechanisms for heat dissipation<sup>29</sup>. It creates locally elevated temperature zones, driving phase transitions, mineral recrystallization, and the

---

<sup>25</sup> Cameron, A. G. W., & Truran, J. W. (1977). The supernova trigger for formation of the solar system. *Icarus*, 30, 447–461. [https://doi.org/10.1016/0019-1035\(77\)90101-6](https://doi.org/10.1016/0019-1035(77)90101-6)

<sup>26</sup> Gounelle, M., & Meynet, G. (2012). Solar system genealogy revealed by extinct short-lived radionuclides in meteorites. *Astronomy & Astrophysics*, 545, A4. <https://doi.org/10.1051/0004-6361/201219031>

<sup>27</sup> Lugaro, M., et al. (2018). Radioactive nuclei from cosmochronology to habitability. *Progress in Particle and Nuclear Physics*, 102, 1–47. <https://doi.org/10.1016/j.ppnp.2018.05.002>

<sup>28</sup> Trinquier, A., et al. (2009).  $^{26}\text{Al}$  and  $^{60}\text{Fe}$  in the early solar system: Solar wind implantation versus supernova injection. *Science*, 324, 374–376. <https://doi.org/10.1126/science.1168221>

<sup>29</sup> Travis et al. (2018). "Radioactive Heating and Its Influence on the Habitability of Icy Worlds." *Astrobiology*, 18(7). DOI: 10.1089/ast.2017.1766

emergence of a liquid water environment<sup>30</sup>. This process provides the necessary thermal energy and microenvironments for a series of prebiotic chemical reactions<sup>31</sup>.

### **3. Distribution of metallic elements in protoplanetary discs and the prebiotic significance of transition metals**

As a high-temperature, high-energy environment that persists during young star formation, the protoplanetary disc concentrates the fundamental elements CHNOPS (carbon, hydrogen, nitrogen, oxygen, phosphorus, sulphur), which constitute the essential organic molecules for life, alongside various metallic elements. These metals primarily originate from the interstellar medium, which is infused by metal-rich celestial phenomena such as supernova explosions, AGB stellar winds, and binary neutron star mergers, and subsequently become enriched within the disc through dust condensation. Evidence of extraterrestrial metallic content in meteorite samples, such as the substantial quantities of Fe, Ni, Co, Mg, and Zn detected in carbonaceous chondrites like the Allende and Murchison meteorites, further corroborates the widespread presence of metallic elements within protoplanetary discs.<sup>32</sup> Among all metals, transition metals played a pivotal catalytic role in prebiotic processes due to their electronic structure and chemical versatility. Firstly, transition metals exhibit multivalent properties, which enable electron transfer and facilitate redox reactions within catalytic cycles. Secondly, they form numerous stable coordination compounds, constructing catalytic microenvironments on mineral or dust surfaces. These functions remain widely conserved in modern life systems through metalloproteins, such as iron-containing nitrogenase<sup>33</sup>, copper-containing respiratory chain complexes, and zinc-containing DNA polymerases, suggesting their origins may be traced back to metal-assisted reactions in early non-living environments.

Moreover, in recent years, transition metals (such as Fe, Ni, and Co) have demonstrated highly efficient catalytic capabilities towards hydrocarbons, ammonia, and simple nitrogen-containing organic compounds in simulated experiments, including Fischer-Tropsch-type synthesis (FTT) and Haber synthesis simulations<sup>34</sup>. In these experiments, metal catalysts

---

<sup>30</sup> Zolotov & Shock (2004) "A Model for Low-Temperature Alteration of CI Chondrites." *Meteoritics & Planetary Science*, 39(3).

<sup>31</sup> Russell et al. (2010) "The Drive to Life on Wet and Icy Worlds." *Astrobiology*, 14(4). DOI: 10.1089/ast.2013.1110

<sup>32</sup> Beinert, H., Holm, R. H., & Münck, E. (1997). Iron-sulfur clusters: Nature's modular, multipurpose structures. *Science*, 277(5326), 653–659. <https://doi.org/10.1126/science.277.5326.653>

<sup>33</sup> Rees, D. C., et al. (2005). Structural basis of biological nitrogen fixation. *Philosophical Transactions of the Royal Society B*, 360(1458), 1221–1234. <https://doi.org/10.1098/rstb.2005.1658>

<sup>34</sup> Fioroni, M., & DeYonker, N. J. (2016). Iron and ruthenium centers grafted onto silica surfaces as models for heterogeneous Fischer-Tropsch and Haber-Bosch catalysis. *Journal of Physical Chemistry C*, 120(50), 28744–28752. <https://doi.org/10.1021/acs.jpcc.6b10696>

frequently deposit upon carbonaceous or silicate dust surfaces, mimicking the environment of protoplanetary disc particles. It further supports their potential role in synthesizing substances related to early life. Thus, from the macro-scale origins of planetary evolution to the micro-scale catalytic mechanisms, transition metals within protoplanetary discs not only furnish the requisite elemental backdrop but may also establish the initial platform for the chemical evolution of life by catalyzing pivotal organic reactions.

#### **4. The 'Existence Paradox' of the Miller–Urey Experiment**

##### **4.1 The Question of Life's Origin and Laboratory Verification Attempts**

The question of life's origin has long been a central theme in astrobiology and the study of chemical evolution. In 1953, Stanley Miller and Harold Urey achieved the first successful laboratory simulation of "primitive Earth conditions" to synthesize amino acids. This landmark experiment demonstrated the scientific feasibility of transitioning "from inorganic to organic"<sup>35</sup>. However, despite its monumental significance in scientific history, the foundational assumption—that the early Earth possessed a strongly reducing atmosphere (e.g., CH<sub>4</sub>, NH<sub>3</sub>, H<sub>2</sub>)—has faced serious challenges in subsequent decades. "The Miller–Urey experiment validated the chemical pathway but not the authenticity of its terrestrial environment"<sup>36</sup>.

##### **4.2 Prebiotic Synthesis in the Protoplanetary Disk: Re-examining the Philosophical Paradox of the Miller–Urey Experiment:**

The seminal 1953 Miller–Urey experiment is widely regarded as the first laboratory demonstration of amino acid synthesis in a non-biotic environment. However, the gas mixture employed—water vapour (H<sub>2</sub>O), methane (CH<sub>4</sub>), ammonia (NH<sub>3</sub>), hydrogen (H<sub>2</sub>), and carbon monoxide (CO)—was unstable in both the modern Earth and its early atmosphere. Subsequent geoscience and geogas escape models suggest the early Earth's atmosphere was more likely dominated by carbon dioxide (CO<sub>2</sub>), nitrogen (N<sub>2</sub>), and water vapour, constituting a neutral or mildly oxidizing environment. It does not support the strongly reducing conditions assumed in the Miller–Urey experiment. Thus, this experiment essentially simulates the more reducing conditions of protoplanetary disc-stage nebular gases within an Earth-based laboratory. Employing electrical sparks to mimic the high-energy particle or radiation sources common in space, it tests the non-biological synthesis potential of prebiotic molecules. In a sense, this presents an existential paradox: the

---

<sup>35</sup> Miller, S. L. (1953). A production of amino acids under possible primitive Earth conditions. *Science*, 117, 528–529. <https://doi.org/10.1126/science.117.3046.528>

<sup>36</sup> Kasting, J. F. (1993). Earth's early atmosphere. *Science*, 259, 920–926. <https://doi.org/10.1126/science.11536547>

experimental results are cited as supporting evidence for the origin of life on Earth, yet the experimental environment itself derives from non-terrestrial, cosmic-stage conditions.

Furthermore, subsequent gas-circulation variants of the experiment did synthesize a greater variety of amino acids and small organic molecules under reducing atmospheres. However, their diversity and complexity remain vastly inferior to those of the dozens of amino acids, nucleobases, fatty acids, and nucleoside precursors detected in meteorites (such as the Murchison meteorite)<sup>37</sup> and asteroid samples (such as Bennu)<sup>38</sup>. It may therefore be concluded that the Miller–Urey experiment essentially validated feasible chemical pathways within protoplanetary discs, rather than prebiotic chemical pathways in the actual terrestrial environment.

### **4.3 From Paradox to Reality**

From this "existence paradox," we uncover a novel, compelling demonstration: the Miller–Urey experiment precisely confirms that protoplanetary discs constitute an ideal environment for the origin of biomolecules. Although the Miller–Urey experiment simulated the possibility of biomolecules arising in natural settings within a laboratory, it employed the reducing gases characteristic of protoplanetary discs rather than the early Earth's neutral to slightly oxidizing atmospheric conditions. Subsequently, they further simulated the gas circulation patterns of protoplanetary discs, synthesizing additional amino acids and organic compounds. It provided further corroboration that life molecules originated in protoplanetary discs rather than on Earth. Thus, it can be said that Miller and Urey conducted the experiment correctly but drew erroneous conclusions. In subsequent chapters, I shall continue to discuss how various amino acids synthesized under different conditions by other researchers—which, in fact, do not exist on Earth—actually originate from processes initiated within the protoplanetary disc.

## **5. The Role of Cosmic Dust in Prebiotic Reactions within the Protoplanetary Disk**

### **5.1 Types of Protoplanetary Disk Dust Particles and Their Surface Chemical Activity**

Cosmic dust within protoplanetary discs comprises a diverse array of particle types, including silicate grains, carbonaceous particles, and ice grains coated with organic matter. These dust particles typically consist of a hard inorganic core and an enveloping layer. For instance, a typical interstellar dust particle may feature a silicate or carbonaceous core, with

---

<sup>37</sup> Martins, Z., et al. (2008). Extraterrestrial nucleobases in the Murchison meteorite. *Earth and Planetary Science Letters*, 270(1–2), 130–136. <https://doi.org/10.1016/j.epsl.2008.03.026>

<sup>38</sup> Lauretta, D. S., et al. (2024). Preliminary OSIRIS-REx Sample Analysis: Organic Molecules in Bennu. NASA Johnson Space Center Science Briefing (internal release; public results expected in Science 2025).



its surface coated by ice (water, carbon dioxide, etc.) and a thin film of complex organic compounds <sup>39</sup>. Different dust particle types exhibit distinct surface chemical reactivity:

**5.1.1 Silicate dust (such as olivine, pyroxene, and other Fe/Mg-rich silicates)** contains doped transition metal elements (e.g., Fe, Ni), whose surfaces can catalyze crucial synthesis reactions at elevated temperatures <sup>40</sup>. Experiments demonstrate that iron-rich silicate powders, akin to those condensed in the solar nebula, can simultaneously catalyze the Fischer-Tropsch synthesis (converting CO and H<sub>2</sub> into hydrocarbons and water) and the Haber synthesis (converting N<sub>2</sub> and H<sub>2</sub> into NH<sub>3</sub>) under conditions of 500–900 K. Remarkably, when CO, N<sub>2</sub>, and H<sub>2</sub> act together on this dust catalyst, nitrogen-containing organics such as methylamine and acetonitrile can also form. It suggests that transition metal sites on silicate dust surfaces can act as catalytic centers, facilitating the synthesis of small organic molecules, such as amino acid precursors. Furthermore, silicate dust can adsorb and mobilize atoms at low temperatures, facilitating their reaction. For instance, at ~10 K, two adsorbed hydrogen atoms can combine on silicate or ice surfaces to form H<sub>2</sub> molecules, with the dust particle acting as a "third body" absorbing excess reaction energy. This surface-binding/diffusion mechanism is crucial for H<sub>2</sub> formation in space, as three-body collisions are exceptionally rare in the sparse gas phase.

**5.1.2 Carbonaceous dust (such as amorphous carbon, graphite, and carbon-rich particles)** possesses surfaces rich in carbon-carbon bond structures. While these do not provide the typical catalytic activity of transition metals, they serve as platforms for the formation and polymerization of organic molecules. Within the interstellar medium, small molecules can adsorb onto carbonaceous dust surfaces and undergo radical reactions, progressively forming more complex hydrocarbon networks. For instance, researchers propose that complex organic molecules such as polycyclic aromatic hydrocarbons (PAHs) may partially originate from stepwise polymerization on dust particle surfaces. Under high-energy radiation, functional groups and unsaturated bonds on carbonaceous dust surfaces readily form radical centers, thereby promoting reactions such as carbon chain elongation and the formation of aromatic rings. Despite lacking metallic active sites, the surfaces of carbonaceous particles can still participate in the formation of organic molecules through physical adsorption and radical mechanisms. Spectroscopic and experimental evidence indicate the presence of complex organic solids, known as "resins," within interstellar dust.

---

<sup>39</sup> Jyllian N. Kemsley,. (2010) Space-Dust Science: ACS Meeting News: Elucidating the chemistry of interstellar dust particles is key to understanding early planets. *Physical Chemistry*. A version of this story appeared in Volume 88, Issue 16

<sup>40</sup> Hugh G M Hill 1, Joseph A Nuth. (2003). The catalytic potential of cosmic dust: implications for prebiotic chemistry in the solar nebula and other protoplanetary systems. *Astrobiology*, Summer;3(2):291-304. doi: 10.1089/153110703769016389.

These resins are hypothesized to form through the photopolymerization of hydrocarbons on cold particles under ultraviolet irradiation.

**5.1.3 Organic/ice-encapsulated particles** constitute a prevalent dust form in molecular clouds and outer disc regions. At low temperatures, dust surfaces accumulate frozen layers ( $\text{H}_2\text{O}$ ,  $\text{CO}$ ,  $\text{CH}_3\text{OH}$ ,  $\text{NH}_3$ , etc.) and encapsulate simple organic compounds. Cosmic rays and ultraviolet light penetrating this ice induce extensive chemical reactions within. As ice acts as a dielectric medium, high-energy particles and UV radiation ionize or dissociate molecules within the ice, generating highly reactive radicals (e.g.,  $\text{H}\cdot$ ,  $\text{OH}\cdot$ ,  $\text{NH}_2\cdot$ ,  $\text{CN}\cdot$ , etc.)<sup>41</sup>. These radicals can migrate and recombine within the low-temperature ice matrix on dust surfaces. Despite temperatures of approximately 10 K, their low-energy barrier recombination reactions proceed spontaneously. Consequently, more diverse and complex molecules are synthesized on the surfaces of ice particles. For instance, laboratory simulations demonstrate that continuous UV irradiation of mixtures such as water ice and methanol at 10 K can yield simple organic molecules, including  $\text{H}_2$ ,  $\text{H}_2\text{O}$ , formaldehyde ( $\text{H}_2\text{CO}$ ), and methanol ( $\text{CH}_3\text{OH}$ ). More complex components emerge as the ice undergoes radiation and gradual heating: organic coatings (rich in CHON elements) form on dust particles. When temperatures rise to tens or hundreds of K, these radiation products can rearrange or desorb into larger organic molecules. Modelling studies indicate that frozen dust particles, during their inward migration from the outer disc in the solar nebula, repeatedly undergo low-temperature UV irradiation and mild heating. This process enables the conversion of ice and organic matter on their surfaces into complex organic products<sup>42</sup>. These products include numerous life-related molecules, leading to the recognition of ice/organically coated dust as one of the 'hotbeds' for synthesizing prebiotic molecules such as amino acids and nucleobases.

The aforementioned diverse dust particle types furnish varied reaction platforms for prebiotic chemistry: silicate and metal sulphide particles offer mineral-like catalytic functions (utilizing transition metals to catalyze key reactions). In contrast, carbonaceous and organically coated particles accumulate and transform simple molecules into more complex organics via adsorption and radical mechanisms.

**5.1.4 Chemical reaction platforms in protoplanetary discs:** The aforementioned diverse types of dust particles provide varied reaction platforms for prebiotic chemistry: silicate and metal sulphide particles offer mineral-like catalytic functions (utilizing transition metals to

---

<sup>41</sup> Masashi Tsuge, Naoki Watanabe. (2023). Radical reactions on interstellar icy dust grains: Experimental investigations of elementary processes. *Proceedings of the Japan Academy, Series B: Physical and Biological Sciences*. 2023;99(4):103-130. doi: 10.2183/pjab.99.008.

<sup>42</sup> Fred J Ciesla, Scott A Sandford. (2012) Organic synthesis via irradiation and warming of ice grains in the solar nebula. *Science*. 2012 Apr 27;336(6080):452-4. doi: 10.1126/science.1217291. Epub 2012 Mar 29.

catalyze key reactions). Scientists often cite Earth's crucial chemical reaction platforms for life's origin. However, I contend that the "F-S World" appears not to reside on Earth, but rather within the protoplanetary disc. Carbonaceous and organically coated particles accumulate and transform simple molecules into more complex organics through adsorption and radical mechanisms. Similarly, the analysis above suggests the Aromatic World Hypothesis is more likely to have been realized within the protoplanetary disc.

## **5.2 Excitation and Synthesis of Molecules on Protoplanetary Disk Dust by Cosmic Radiation**

Within protoplanetary discs and the interstellar medium, dust particles are immersed in an environment saturated with high-energy electromagnetic radiation and particle radiation. Young stars and the cosmic background emit X-rays, ultraviolet radiation, and high-energy cosmic ray particles, all of which exert a significant influence on the chemistry occurring at the dust surface<sup>43</sup>. Specifically:

**5.2.1 Ultraviolet (UV) radiation** can be directly absorbed by molecules on dust surfaces, breaking chemical bonds to generate free radicals or ions. Dust particles act as both "generators" and "carriers" of free radicals. At extremely low temperatures (10–50 K), fragments produced by UV photolysis cannot escape rapidly and remain trapped within the ice or organic matrix on the dust surface. These fragments can rearrange to form new covalent bonds. When two free radicals encounter each other on the confined dust surface, they readily couple to form more complex neutral molecules. For instance, experiments confirm that when simulated interstellar ice (a mixture of H<sub>2</sub>O, CO, CH<sub>4</sub>, NH<sub>3</sub>, etc.) is exposed to vacuum UV irradiation below 77 K, diverse complex organic residues are generated. Upon heating and hydrolysis of these residues, amino acids (such as glycine and alanine) can be detected<sup>44</sup>. It indicates that UV-induced "amino acid precursors" formed within the ice subsequently convert into amino acids under mild thermal treatment. Similarly, UV irradiation on ice rich in simple organic compounds, such as HCN and CH<sub>3</sub>OH, can introduce nitrogen-containing functional groups, synthesizing prebiotic molecules like aminonitriles and amides. In nebular environments, intense UV exposure within the inner disc regions near young stars triggers extensive photolysis-synthesis cycles on the surfaces

---

<sup>43</sup> Jyllian N. Kemsley. (2010) Space-Dust Science: ACS Meeting News: Elucidating the chemistry of interstellar dust particles is key to understanding early planets. *Physical Chemistry*. A version of this story appeared in Volume 88, Issue 16

<sup>44</sup> Kobayashi K, Kasamatsu T. (1995) Formation of amino acid precursors in cometary ice environments by cosmic radiation. *Advances in Space Research*. 16(2):21-6. doi: 10.1016/0273-1177(95)00188-k.

of organic ice-bearing dust particles. This process is recognized as a natural by-product of planetary system evolution, yielding complex organic molecules<sup>45</sup>.

**5.2.2 Higher-energy photons**, such as X-rays and gamma rays, possess greater penetrating power, enabling them to penetrate deep into dust grains or thick ice layers to induce radiolysis. High-energy photons generate secondary electrons and fluorescence within solids, thereby breaking molecular bonds like UV radiation. Due to their deep penetration, molecular fragments and free radicals also form within ice. These can be stored long-term within ice lattices at low temperatures; when the environment warms slightly (e.g., from 10 K to 30–40 K), some atoms and free radicals gain mobility and react, rapidly synthesizing more stable macromolecules. Both theoretical and experimental evidence support the accumulation of a "frozen" radical reservoir within ice, induced by cosmic ray-induced internal radiation. As temperatures rise, these radicals interact abruptly, forming molecules such as methanol, ethanol, simple sugars, and even larger compounds<sup>46</sup>. Thus, even deep-cold dust far from light sources can gradually accumulate complex organic matter under sustained bombardment by cosmic rays.

**5.2.3 Direct collisions of high-energy particles (cosmic rays, protons, etc.)** with dust particles similarly induce molecular excitation and dissociation. On the one hand, fast heavy-ion impacts generate track-induced radiation along their path, triggering cascading molecular fission and ionization. On the other hand, high-energy protons/electrons deposit energy within ice, acting similarly to gamma rays. Experiments bombarding 10 K simulated ice with MeV protons have confirmed that particles irradiating H<sub>2</sub>O, CO, NH<sub>3</sub>, and other compounds in ice generate a diverse range of volatile organic compounds. Residual, less volatile organic matter releases amino acids (such as glycine and alanine) upon warming to room temperature and undergoing hydrolysis<sup>47</sup>. These findings indicate that cosmic rays and UV radiation collectively constitute the most potent energy sources for organic synthesis in space. Unlike in solution, where radicals are readily quenched, radicals exhibit longer lifetimes and higher migration probabilities in vacuum and on solid surfaces. Consequently, radiochemistry emerges as a primary driving force for the formation of organic molecules on interstellar dust. As one astrochemist observed: "Chemical reactions on dust particle surfaces play a pivotal role in the chemistry of nascent planets; much of the chemistry of interstellar dust particles is driven by radiation (particularly UV, X-rays, and cosmic rays)".

---

<sup>45</sup> Fred J Ciesla 1, Scott A Sandford. (2012) Organic synthesis via irradiation and warming of ice grains in the solar nebula. *Science*. 336(6080):452-4. doi: 10.1126/science.1217291. Epub 2012 Mar 29.

<sup>46</sup> Kalvāns J (2015) Cosmic-ray-induced dissociation and reactions in warm interstellar ices. *Astronomy & Astrophysics (A&A)*. Volume 573, (38) p. 13 <https://doi.org/10.1051/0004-6361/201424300>

<sup>47</sup> Kobayashi K, Kasamatsu, T. (1995) Formation of amino acid precursors in cometary ice environments by cosmic radiation. *Advances in Space Research*. 16(2):21-6. doi: 10.1016/0273-1177(95)00188-k.

**5.2.4 The 'Micro-Chemical Reactor' Function of Protoplanetary Disk Dust:** Through the mechanisms mentioned above, high-energy radiation induces radicals and ions on dust surfaces. These intermediates then recombine to form more complex bonds. The result is that dust acts as a "microscopic chemical reactor" within intense radiation environments, facilitating reactions ranging from the simplest synthesis of  $H_2$  and  $H_2O$  to the stepwise formation of complex organic molecules. In the context of prebiotic chemistry, this implies that key molecules such as amino acids, nucleobases, and simple sugars can be naturally synthesized through radiation pathways within dust in the non-living cosmic environment.

### **5.3 Impact of Short-Lived Radioisotopes on Protoplanetary Disk Dust Heating and Reaction Environments**

Within the young protoplanetary nebula of the early Solar System, substantial quantities of short-lived radioactive isotopes (such as  $^{26}Al$  and  $^{60}Fe$ ) were incorporated into solid particles and planetesimals. Their decay processes exerted two significant influences on chemical evolution: internal heating and radiation sources. Among these,  $^{26}Al$  (with a half-life of approximately 0.72 million years) stands as the most prominent example.

**5.3.1 Radioactive Decay Heating:**  $^{26}Al$  decay releases high-energy alpha particles and gamma rays, primarily converting energy into heat. It has profound implications for larger dust aggregates and even planetary embryos. Within the first few million years of solar system formation, Al-rich ( $^{26}Al$ -rich) minor bodies were heated internally above the melting point of ice, thereby generating conditions conducive to liquid water<sup>48</sup>. Geological evidence suggests that the parent asteroids of carbonaceous chondrite meteorites underwent radioactive heating during their early stages, causing internal ice to warm and melt, thereby producing liquid water and enabling hydrothermal chemical processes. Consequently, ( $^{26}Al$ ) provided a hot-water reactor on primordial small bodies: water interacting with metallic minerals and organic matter within dust could facilitate a series of aqueous chemical reactions. For instance, within the hydrothermal environments of these parent bodies, simple small molecules (such as formaldehyde and ammonia) can condense or transform into more complex organic precursors. Experimental simulations support this: incubating solutions of formaldehyde and ammonia under heated conditions generates amino acid precursor compounds (requiring hydrolysis to release amino acids). Yields further increased when gamma-ray irradiation was applied concurrently with heating, indicating a synergistic effect between ionizing radiation from radioactive decay and thermal energy. Recent studies have revealed that simulated systems subjected to irradiation followed by heating produce amino acid precursors more efficiently than those treated with heating alone or with

---

<sup>48</sup> Imai, N., et al. (2024). Formation of amino acid precursors in early aqueous alteration in small bodies: Synergetic effects of gamma rays and hydrothermal processing. *Geochemical Journal*, 58(6), 304-315

different treatment sequences. It is hypothesized to arise because radiation generates oxygenated organic radicals and carbonyl compounds within solutions or ice, via a Maillard-type process, whereupon heat subsequently promotes rearrangement of these intermediates into more complex products. In other words,  $^{26}\text{Al}$  decay simultaneously provides two essential elements: internal energy (heating enhances reaction kinetics) and radiochemistry (creating reactive species).

**5.3.2 Evolution of the chemical environment within asteroids/dust bodies:** The decay heat of  $^{26}\text{Al}$  drives thermal evolution within many early planetary bodies. For micrometeoroids hundreds of metres in diameter, model calculations indicate  $^{26}\text{Al}$  decay dominates internal heating, potentially causing internal melting or differentiation (e.g., molten iron cores, asteroidal magmatism). Whilst individual micrometre-scale dust grains cannot accumulate sufficient heat alone, when these coalesce into parent bodies several kilometres in size, the energy from  $^{26}\text{Al}$  decay is locally deposited, driving water-rock reactions and thermal transformation of organic matter. For instance, a significant proportion of amino acids detected in carbonaceous meteorite parent bodies may have formed during such hydrothermal stages through reactions of simple precursors (HCN,  $\text{NH}_3$ , aldehydes, etc.), rather than having been generated on Earth<sup>49</sup>. Decay heating can also induce thermal pyrolysis and recombination of organic inclusions. For instance, complex organic matter within carbon-rich dust partially degrades and recondenses at  $\sim 100\text{--}200^\circ\text{C}$ , yielding more polar organic compounds that enter the aqueous phase and become incorporated into the parent body's minerals.

**5.3.3 Radiation Effects:** Short-lived nuclides emit ionizing radiation alongside heat.  $^{26}\text{Al}$  decay emits 1.3 MeV beta particles and 1.8 MeV gamma rays, which can ionize surrounding matter. Within the parent body, rock absorption of gamma rays is relatively limited yet sufficient to generate local ion pairs and free radicals. For instance, dust particles containing  $^{26}\text{Al}$  are themselves bombarded by their decay  $\gamma$ -rays (and radiation from the decay product  $^{26}\text{Mg}$ ), inducing cosmic ray-like chemistry. However, compared to external cosmic ray sources, the radiation flux from internal radioactive sources is lower and has a limited range of action; thus, their primary contribution remains the thermal effect. Within enclosed parent bodies, radioactivity provides a low-dose radiation background that persists for millions of years, potentially facilitating the slow modification of organic matter, such as promoting the formation of aromatic structures and increasing oxidation states. Meteorite studies suggest that certain organic components may have undergone prolonged irradiation by radioactive substances, resulting in distinctive chemical signatures.

---

<sup>49</sup> Elsila, J. E., et al. (2007). Mechanisms of amino acid formation in interstellar ice analogs. *Astroph. J.*, 660, 911-918

$^{60}\text{Fe}$  (half-life approximately 2.6 Myr) was also present in the early solar system, providing additional heat through decay (primarily  $\beta^-$  and  $\gamma$ -ray emission). Whilst  $^{60}\text{Fe}$ 's heat production alone was insufficient to melt large bodies, it acted synergistically with  $^{26}\text{Al}$  heating in iron-rich asteroids. Collectively, these short-lived radioactive sources shaped the chemical environment of early dust/planetary discs: water released from ice provided a medium for organic synthesis, thermal energy drove reactions, and radiation created bioactive intermediates. It represents one of the key prebiotic chemical scenarios beyond Earth. Researchers note that such natural chemical pathways could have universally generated the building blocks of life throughout the solar system and potentially other planetary systems<sup>50</sup>.

#### **5.4 Comparison of Mineral Surface Reactions in Early Earth's Environment with Protoplanetary Disk Dust Platforms**

Compared to protoplanetary discs rich in cosmic dust, the early Earth environment exhibited distinct differences in terms of "surface platforms" suitable for prebiotic chemistry. This disparity restricted nearly all chemical pathways to Earth's native conditions, further substantiating the origin of life's building blocks within protoplanetary discs:

**5.4.1 Primordial Atmospheric Composition and Reductive Potential:** Modern research suggests the Taecenaean Earth's atmosphere was dominated by  $\text{CO}_2$ ,  $\text{CO}$ , and  $\text{N}_2$ , with minimal hydrogen and methane, presenting an overall neutral-to-oxidizing environment<sup>51</sup>. Within such conditions, the classical Miller-Urey reaction (relying on strongly reducing gases like  $\text{CH}_4$  and  $\text{NH}_3$ ) proved highly inefficient. Laboratory discharge and illumination experiments demonstrate that gas phases rich in  $\text{CO}_2$  and poor in methane struggle to synthesize reducing organic compounds, such as amino acids, aligning with the primordial Earth conditions. Concurrently, terrestrial pathways for reducing  $\text{N}_2$  to  $\text{NH}_3$  (such as lightning, volcanic activity, or submarine hydrothermal vents) exhibit a limited yield, which is insufficient to supply substantial quantities of reducing nitrogen compounds. By contrast, the protoplanetary disc dust environment provides the missing link. Within the solar protoplanetary nebula, transition metals like iron and nickel catalyze the conversion of  $\text{CO}_2$  /  $\text{CO}$  into hydrocarbons (such as  $\text{CH}_4$ ) and  $\text{N}_2$  into ammonia ( $\text{NH}_3$ ) under low-oxygen conditions. Nitrogen-containing organic molecules (e.g., methylamine, acetonitrile) can also form directly on dust surfaces. These molecules, which could have served as precursors to life on Earth, were challenging to produce indigenously due to Earth's lack of a

---

<sup>50</sup> Hill, H. G. M., & Nuth, J. A. (2003). The catalytic potential of cosmic dust: implications for prebiotic chemistry in the solar nebula and other protoplanetary systems. *Astrobiology*, 3(2), 291-304

<sup>51</sup> Hill, H. G. M., & Nuth, J. A. (2003). The catalytic potential of cosmic dust: implications for prebiotic chemistry in the solar nebula and other protoplanetary systems. *Astrobiology*, 3(2), 291-304

dust-catalyzed environment. In other words, the early Earth may have lacked a widespread reductive catalytic platform for efficiently generating certain fundamental organic molecules. Cosmic dust, however, had already "pre-synthesized" nearly all the foundational molecules for life during the protoplanetary disc nebula phase, subsequently delivering them to Earth via meteorites and micrometeorites.

**5.4.2 Abundance and Forms of Utilizable Elements:** The utilizable forms of key life-essential elements (such as C, N, P, S) in Earth's primordial crust were severely limited. Phosphorus primarily existed as insoluble phosphate minerals, nitrogen predominantly as inert N<sub>2</sub> gas, carbon was largely locked within CO<sub>2</sub>, and sulphur was mainly present as stable sulphates. These forms were not "friendly" to inorganic chemistry. Life subsequently evolved complex enzyme systems to access them, mechanisms that were absent in the prebiotic stages. In contrast, protoplanetary disc dust and meteorites enriched these elements in reactive forms: for instance, phosphides (such as Fe<sub>3</sub>P, which generates soluble phosphates upon entering water), organic nitrogen compounds (ammonia, nitriles), reduced carbon (methanol, hydrocarbons), and free sulphides. Research indicates that certain nutrient elements, which are severely deficient in Earth's surface layers, are present in protoplanetary disc dust at concentrations several orders of magnitude higher than in the crust. It suggests that life was delivered as "chemical fertilizer" by falling cosmic dust/meteorites. In contrast, the local raw material concentrations on early Earth were insufficient to readily assemble precursors of biomolecules such as nucleotides and fatty acids.

**5.4.3 Types and Properties of Mineral Surfaces:** Although solid surfaces existed in Earth's early days—such as volcanic rocks, clay minerals, and metallic sulphides—and are thought to have participated in prebiotic chemistry (e.g., clay minerals adsorbing organic matter and catalysing polymerisation; pyrite FeS+NiS catalysing CO<sub>2</sub> reduction in hydrothermal vents), However, compared to cosmic dust, Earth's mineral surfaces operate under distinct conditions—including liquid water environments and higher gravitational forces—resulting in different chemical pathways. In aqueous solutions, free radicals are readily quenched, with reactions predominantly following solution chemistry mechanisms rather than surface solid-state mechanisms. Cosmic dust, typically existing in a vacuum or gas phase, enables surface reactions via heterogeneous catalysis and radiochemistry—processes that are difficult to replicate on Earth's surfaces. For instance, processes like H<sub>2</sub> formation and methane synthesis occurring on dust surfaces require high temperatures or catalysts in the Earth's atmosphere. Similarly, PAH synthesis in the interstellar medium follows a radical surface condensation pathway. In contrast, on Earth, aromatic hydrocarbons are predominantly produced by the combustion of organic matter or high-temperature processes. Moreover, dust particles are tiny (on the submicron scale) and occur in vast



quantities, yielding an immense total surface area that provides ample opportunities for molecular reactions. In contrast, available mineral surfaces on Earth (such as rock fissures or coastal clays) possess relatively limited total surface area and are often covered by water or organic films that inhibit reactivity. Consequently, specific prebiotic reaction pathways possible in space (such as ultraviolet-driven amino acid synthesis on icy surfaces at low temperatures) either cannot occur under terrestrial conditions or proceed with extremely low efficiency. This constraint on Earth's indigenous chemical processes necessitated the delivery of pre-synthesized "building blocks" to the planet's surface via meteorites and dust. Consequently, scientists hypothesize that most key molecules for life—such as amino acid and nucleotide precursors—were likely "exogenously synthesized and endogenously supplied," rather than originating from Earth's own chemistry<sup>52</sup>.

**5.4.4 The Protoplanetary Disc as the Optimal Environment for Life's Origins:** Overall, the early Earth lacked the pervasive diffuse catalytic particles and radiation conditions prevalent in protoplanetary discs, which partly contributed to the scarcity of initial organic synthesis on Earth. Consequently, Earth's prebiotic chemistry likely relied heavily on two compensatory pathways: first, minimal synthesis occurring in non-conventional environments (such as atmospheric lightning discharges, meteorite impacts, or volcanic hydrothermal vents), which may have been absent; second, cosmic delivery (via micrometeorites, meteors, and cometary material) supplying preformed organic molecules. Decades of meteorite research support the latter pathway—many building blocks of life molecules arrived on Earth in extraterrestrial form, precisely matching the chemical capabilities of protoplanetary discs, cosmic dust, and parent asteroids. It also represents the optimal environment for the origin of life in the cosmos.

## **5.5 Recent Discoveries from Experimental Simulations of Protoplanetary Disks and Meteorite Analyses**

Recent years have seen experimental simulations of protoplanetary disc cosmic dust chemistry and analyses of carbonaceous meteorites continually revealing evidence and details of prebiotic molecule formation in space.

**5.5.1 Dust Catalysis Synthesis Experiment:** The seminal experiment by Hill and Nuth (2003) cited above demonstrated the catalytic potential of cosmic dust<sup>53</sup>. Using ferrosilicate powder resembling solar nebular condensates, they passed primordial gas mixtures (CO, N<sub>2</sub>, H<sub>2</sub>) through the catalyst bed at 500–900 K, successfully synthesizing simple molecules such

---

<sup>52</sup> Liz Kruesi. (2022) All of the bases in DNA and RNA have now been found in meteorites. ScienceNews. 26 April.

<sup>53</sup> Hill, H. G. M., & Nuth, J. A. (2003). The catalytic potential of cosmic dust: implications for prebiotic chemistry in the solar nebula and other protoplanetary systems. *Astrobiology*, 3(2), 291-304

as methane and ammonia, as well as nitrogen-containing organics (e.g., methylamine, acetonitrile). This outcome provides direct evidence for the simultaneous generation of hydrocarbon and nitrogen-containing molecules in exoplanetary environments, supporting the "pre-processing" role of dust within protoplanetary discs in the Solar System. Recent expansions of such research—including simulations with more complex gas compositions and diverse mineral catalysts—demonstrate the transformative capacity of protoplanetary disc dust surfaces on inert small molecules, such as CO<sub>2</sub> and N<sub>2</sub>.

**5.5.2 Radiation simulations on cryogenic ice:** To model interstellar media and the outer regions of the solar protoplanetary disc, numerous experiments employ ultra-high vacuum cryogenic apparatus. Mixed-ice thin films are deposited onto cold fingers (10–80 K), subsequently irradiated with UV lamps or ion beams. Notable experiments by Bernstein et al. (2002)<sup>54</sup> and Muñoz Caro et al. (2002)<sup>55</sup> irradiated ice containing H<sub>2</sub>O, CH<sub>3</sub>OH, NH<sub>3</sub>, and HCN components with UV light. The residual organic matter yielded multiple amino acids (including glycine, alanine, serine, etc.)<sup>56</sup>. These amino acids appeared in racemic form and exhibited deuterium enrichment, mirroring amino acid signatures in meteorites. It confirms laboratory-synthesized amino acids originate abiotically, potentially mirroring processes in the outer protoplanetary disc environment. Holtom et al. (2005) similarly obtained amino acids by bombarding ice-like formulations with high-energy electrons (simulating cosmic rays)<sup>57</sup>. Collectively, these experiments confirm that the "UV/radiation- + -ice" pathway can synthesize dozens of amino acids and other organic molecules. Recent experimental refinements have enhanced analytical sensitivity and broadened the range of detectable products. For instance, research by the German Aerospace Centre and others has detected dipeptides (two amino acids linked) and phosphorus-containing organic compounds formed upon ice illumination, suggesting that primitive condensation polymerization reactions may even occur on dust-ice membranes within protoplanetary discs.

**5.5.3 Radioisotope simulation experiments:** Addressing hydrothermal radiation environments within meteorite parent bodies, recent research (Imai et al., 2024) designed

---

<sup>54</sup> Bernstein, M. P., Dworkin, J. P., Sandford, S. A., Cooper, G. W., & Allamandola, L. J. (2002). Racemic amino acids from the ultraviolet photolysis of interstellar ice analogues. *Nature*, 416(6879), 401–403. <https://doi.org/...>

<sup>55</sup> Muñoz Caro, G. M. M., Meierhenrich, U. J., Schutte, W. A., Barbier, B., Arcones Segovia, A., Rosenbauer, H., Thiemann, W. H-P., Brack, A., & Greenberg, J. M. (2002). Amino acids from ultraviolet irradiation of interstellar ice analogues. *Nature*, 416(6879), 403–406. <https://doi.org/10.1038/416403a>

<sup>56</sup> Elsila, J. E., et al. (2007). Mechanisms of amino acid formation in interstellar ice analogs. *Astroph. J.*, 660, 911-918

<sup>57</sup> Holtom, P. D., Bennett, C. J., Osamura, Y., Mason, N. J., & Kaiser, R. I. (2005). A combined experimental and theoretical study on the formation of the amino acid glycine and its isomer in extraterrestrial ices. *The Astrophysical Journal*, 626(2), 940–952. <https://doi.org/10.1086/430106>

simulations combining gamma-ray irradiation and heating<sup>58</sup>. They subjected simulated solutions of formaldehyde, methanol, and ammonia to low-dose  $\gamma$  irradiation followed by heating. The final products yielded typical amino acid precursors, exhibiting spectral characteristics reminiscent of Maillard reaction products. Notably, the "radiation-then-heating" sequence demonstrated significantly higher efficiency in producing amino acid precursors than the "heating-then-radiation" approach. It suggests that within the radioluminescent interior of <sup>26</sup>Al asteroids, radiation from radioactive sources can synergize with thermal effects to synthesize biomolecules more efficiently. This finding aligns with the formation mechanisms of certain organics (such as nitrogen-containing brown macromolecules) observed in meteorites, offering microscopic evidence for these processes.

**5.5.4 Latest Identification of Meteorite Organics:** With advances in analytical techniques, scientists continue to discover new organic constituents within meteorites, many directly related to biomolecules. Among these, amino acids were first identified in carbonaceous chondrites (such as the Murchison meteorite that fell in 1969) as early as the 1960s. To date, over 80 amino acids have been identified in Murchison, including 12 of the 20 commonly used in proteins, alongside numerous non-protein amino acids<sup>59</sup>. These amino acids exist as racemic mixtures and exhibit anomalous isotopic ratios (such as elevated D/H and <sup>15</sup>N), confirming their non-biological origin and likely interstellar/nebular provenance. More astonishingly, recent analyses have identified all five nucleic acid bases within meteorites: adenine (A), guanine (G), uracil (U), cytosine (C), and thymine (T). For many years, only A, G, and U had been reliably detected in meteorites, while C and T remained unconfirmed due to their extreme susceptibility to degradation. In 2022, a team from Hokkaido University in Japan successfully detected trace amounts of C and T from multiple carbonaceous meteorites using a mild cold-water extraction method. It marked the first time all five DNA/RNA bases had been "collected" within meteorites<sup>60</sup>. Simultaneously detected were multiple structural isomers of these bases, further confirming they did not originate from terrestrial contamination. Additionally, in 2019, scientists identified ribose (the pentose sugar that forms RNA) and related sugars within meteorites, indicating that both the sugar and base components of nucleotides can be supplied by space chemistry. Meteorites also harbour abundant organic acids: diverse carboxylic acids, including monocarboxylic and dicarboxylic acids (i.e., fatty acid chains), were reported as early as the

---

<sup>58</sup> Imai, N., et al. (2024). Formation of amino acid precursors in early aqueous alteration in small bodies: Synergetic effects of gamma rays and hydrothermal processing. *Geochemical Journal*, 58(6), 304-315

<sup>59</sup> Toshiki Koga & Hiroshi Naraoka. (2017) A new family of extraterrestrial amino acids in the Murchison Meteorite. *Scientific Reports* | 7: 636 | DOI:10.1038/s41598-017-00693-9

<sup>60</sup> Kruesi, L. (2022). All of the bases in DNA and RNA have now been found in meteorites. *Science News* (26 April 2022)

1970s<sup>61</sup>. For instance, a series of fatty acid anions of varying chain lengths was detected in the Murchison and Murray meteorites, with some presumed to exist as sodium or magnesium salts. These fatty acids constitute the fundamental building blocks of cell membrane lipids, suggesting extraterrestrial chemistry may have laid the groundwork for assembling primordial membrane structures. Preliminary analysis of Benu asteroid samples returned by NASA's OSIRIS-REx mission (2023) corroborates these findings: Benu samples contain fourteen protein amino acids and all five nucleic acid bases, at concentrations exceeding those found in known meteorites<sup>62</sup>. It undoubtedly lends further credence to the notion that asteroid/comet material seeded Earth with the building blocks of life. It also provides additional evidence supporting the theory that primordial life may have originated from the protoplanetary disc.

**5.5.5 Chiral Bias and Advanced Molecules:** The chiral bias of organic molecules and products of multi-step reactions remains a current research focal point. Amino acids in meteorites predominantly exist as racemic mixtures. However, certain meteorites, like Murchison, exhibit a slight L-enantiomer excess in a few amino acids, potentially linked to optically polarized radiation from nebulae. Experimentally, reactions induced on dust surfaces under circularly polarised UV light also yielded amino acid products with slight chiral excess. It suggests that cosmochemical processes not only generate diverse molecules but may also pre-imprint molecular chirality—a hallmark of life—onto specific precursors. Another advancement involves detecting evidence of complex organic macromolecules, such as the presence of insoluble organic matter (IOM) in meteorites, whose structure resembles highly cross-linked organic polymers. Recent synchrotron radiation and mass spectrometry studies are progressively deciphering the structural units, revealing the coexistence of aromatic rings, heterocycles, and long-chain structures. These macromolecules are hypothesized to have formed through the gradual polymerization of simple molecules via radiation and heat on parent asteroids. Although abiotic pathways have yet to simulate genuine RNA or peptide chains, these findings indicate that each step from monomers to oligomers is feasible in space.

In summary, the latest evidence from laboratory simulations and meteorite analyses strongly supports the notion that protoplanetary disc dust particles functioned as catalytic factories and transport vehicles for key prebiotic molecules. Whether through radical surface reactions that form simple organics in cold outer protoplanetary discs or the synthesis of complex molecules within radiatively heated planetesimals, dust played a

---

<sup>61</sup> Kvenvolden, K. A. (1974). Amino and fatty acids in carbonaceous meteorites. NASA Technical Report 19750034440 ntrs.nasa.gov.

<sup>62</sup> Keith Cowing. (2025) Life's Building Blocks Discovered In Samples From Asteroid Benu. Astrobiology Web (30 January 2025)

crucial role in the chemical evolution of life's building blocks. The absence of such platforms in Earth's early stages implies that virtually all life's raw materials likely originated from protoplanetary disc chemistry and were delivered by protoplanetary disc dust/meteorites. This insight broadens our perspective on the environment of life's origin: life's starting point may not have been confined to a planetary surface niche, but instead began within the chemistry of dust clouds pervading the protoplanetary disc, commencing during the formation of planets.

## 6. Aromatic World Hypothesis

In the preceding discussion, I concluded that life and life-bearing matter within the cosmos originate from stellar disks or protoplanetary disks. Even where no planets form within a stellar system's protoplanetary disk, life or life-bearing matter may still emerge within that disk<sup>63</sup>. Unlike our solar system, this nascent life or life-bearing matter, lacking planets to accrete onto, is ultimately blown by stellar winds into the interstellar medium (ISM) as cosmic dust. There, it endures cosmic radiation from stars and star systems. Life or complex biomolecules originating from protoplanetary or protostellar discs are destroyed within cosmic dust, such as nucleobases, amino acids, and sugars. Molecules resistant to cosmic radiation, such as polycyclic aromatic hydrocarbons (PAHs), can be detected in the gas phase by infrared and microwave spectroscopic telescopes. They are widely observed in interstellar dust, cometary dust, interplanetary dust, and circumstellar dust<sup>64</sup>. Here, we note that cometary dust, interplanetary dust, and circumstellar dust all bear a direct connection to the protostellar or protoplanetary discs from which stars are born. Similarly, PAHs within interstellar dust can only be said to have been blown by stellar winds from the protostellar or protoplanetary discs, where they originated, into the more distant interstellar space<sup>65</sup>.

### 6.1 How are PAHs originating from protoplanetary discs detected?

How, then, can we prove that these PAHs originate from protoplanetary discs? Extensive evidence indicates the widespread presence of PAHs, as observed across different spectral bands by telescopes studying stellar clouds or protoplanetary discs. For instance, the Spitzer Infrared Spectrometer (IRS) has detected characteristic PAH emission in multiple protoplanetary discs, particularly around Herbig Ae/Be stars<sup>66</sup>. Studies of discs around T

---

<sup>63</sup> Li, XM. (2024). Chemistry Analysis for the Origin of Life in Protoplanetary Disks. *Geoinformation and Geostatistics: An Overview*, Vol. 12, Issue 5

<sup>64</sup> Swamy, K.S.K., (2012) *Dust in the Universe: Similarities and Differences*

<sup>65</sup> Li, A. (2020) "Spitzer's perspective of polycyclic aromatic hydrocarbons in galaxies" *Nature Astronomy*, 4(4), 339-351. <https://doi.org/10.1038/s41550-020-1051-4>

<sup>66</sup> Acke, B., & van den Ancker, M. E. (2004) "ISO spectroscopy of disks around Herbig Ae/Be stars: PAHs and silicate emission" *Astronomy & Astrophysics*, 426(1), 151-170. <https://doi.org/10.1051/0004-6361:20040344>

Tauri and Herbig Ae/Be stars reveal a PAH detection rate as high as 60% for Herbig Ae/Be stars, compared to a lower rate of approximately 10% for T Tauri stars<sup>67</sup>. It indicates the widespread presence of PAHs within these discs, particularly around higher-mass stars. The Mid-Infrared Instrument (MIRI) aboard the James Webb Space Telescope (JWST) provides higher-resolution observational capabilities, further confirming the presence of PAHs within protoplanetary discs. For instance, Sturm et al. (2024) utilized JWST/MIRI to observe the edge disc of HH 48 NE, detecting multiple PAH emission features, which indicates that these molecules are predominantly located within the disc's surface layer<sup>68</sup>.

Furthermore, Arulanantham et al. (2024) detected PAH emission at 11.3 micrometres while studying the edge-on disc of the T Tauri star Tau 042021<sup>69</sup>, providing additional evidence for the widespread presence of PAHs within protoplanetary discs. Ground-based observations using the VISIR mid-infrared instrument on the VLT have also successfully detected PAHs. For instance, Yoffe et al. (2023) employed VISIR to observe the protoplanetary discs around eight Herbig Ae stars, resolving PAH emission features in five discs. They found that these emissions exhibited spatially centred or annular distributions, suggesting specific distribution patterns for PAHs within the disc structure<sup>70</sup>.

## **6.2 Comparing PAH abundances in protoplanetary discs versus terrestrial environments**

In the interstellar medium (ISM) of the Milky Way, approximately 10% to 20% of carbon exists in the form of polycyclic aromatic hydrocarbons (PAHs)<sup>71</sup>. Within protoplanetary discs, and particularly in carbonaceous chondrites (such as the Murchison meteorite) and comets (such as Halley's Comet and Comet 67P/Churyumov–Gerasimenko), PAHs and PAH-containing material can constitute up to several tens of per cent of the total organic content<sup>72</sup>. In contrast, the natural abundance of PAHs in Earth's environment is extremely

---

<sup>67</sup> Geers, V. C., Augereau, J. C., Pontoppidan, K. M., et al. (2006). "C2D Spitzer-IRS spectra of disks around T Tauri stars. III. [Ne II], [Fe I], and PAH emission." *Astronomy & Astrophysics*, 459(2), 545-556.

<https://doi.org/10.1051/0004-6361:20065552>

<sup>68</sup> Sturm, B., Bouwman, J., van Dishoeck, E. F., et al. (2024) "JWST MIRI observations of HH 48 NE reveal PAH emission from protoplanetary disk surfaces" *Astronomy & Astrophysics*, in press.

<https://doi.org/10.1051/0004-6361/202450865> arXiv preprint

<sup>69</sup> Arulanantham, N., et al. (2024) "Detection of Polycyclic Aromatic Hydrocarbons (PAHs) in the edge-on protoplanetary disk Tau 042021 using JWST/MIRI" *The Astrophysical Journal Letters*, under review.

arXiv:2402.12256

<sup>70</sup> Yoffe, S. R., Maaskant, K. M., Lagage, P. O., et al. (2023) "Spatially resolving Polycyclic Aromatic Hydrocarbon emission in protoplanetary disks around Herbig Ae stars with VISIR/VLT" *Astronomy & Astrophysics*, 675, A44. <https://doi.org/10.1051/0004-6361/202346155> arXiv:2303.06592

<sup>71</sup> Tielens, A. G. G. M. (2008). Interstellar Polycyclic Aromatic Hydrocarbon Molecules. *Annual Review of Astronomy and Astrophysics*, 46, 289-337.

<sup>72</sup> Sephton, M. A. (2002). Organic compounds in carbonaceous meteorites. *Natural Product Reports*, 19(3), 292–311.

low, primarily originating from combustion (such as forest fires and volcanic activity), industrial emissions, human activities, or minor meteorite inputs<sup>73</sup>. The abundance of PAHs in protoplanetary discs exceeds that on Earth (including that from fossil fuel combustion) by more than 100,000-fold. If PAHs play a crucial role in prebiotic chemistry, the contrast between their abundance in protoplanetary discs and terrestrial environments provides compelling evidence. Below, we discuss the relationship between PAHs and the building blocks of key life components: cell membranes, amino acids, and nucleobases. It further supports the notion that life on Earth originated from protoplanetary discs.

### **6.3 Potential Relationship Between PAHs and Life's Constituents**

Polycyclic aromatic hydrocarbons (PAHs) are ubiquitous organic molecules in the cosmos, potentially playing a vital role in the origin of life. Their potential connections to cell membrane composition and other fundamental biomolecules (such as amino acids, bases, and ribose) are outlined below.

**6.3.1 Relationship between PAHs and Cell Membrane Composition:** In the early protoplanetary disc environment, the conventional membrane model involving lipid bilayers may not represent the earliest biological membrane structure, as such lipid membranes exhibit poor stability under primordial harsh conditions. Conversely, PAHs possess high chemical stability, are ubiquitous in the cosmos, and can form naturally under interstellar or protoplanetary disc conditions, making them a potential primordial membrane material<sup>74</sup>.

Advantages of PAHs as membrane constituents: High stability: PAHs possess robust structures capable of withstanding harsh environments such as ultraviolet radiation, cosmic rays, and high temperatures, rendering them far more suited to the extreme conditions of early Earth than lipids. Hydrophobicity and self-assembly properties: PAH molecules can form stable molecular stacks or microcapsule structures via  $\pi$ - $\pi$  stacking interactions. This self-assembled structure resembles the enclosed architecture of cell membranes, potentially providing the isolating environment required for early life forms<sup>75</sup>. The hydrophobicity and planar structure of PAHs may mimic the hydrophobic tails of lipids in modern cell membranes, aiding in the formation of primordial membrane structures<sup>76</sup>.

---

<sup>73</sup> Simoneit, B. R. T. (2002). Biomass burning—a review of organic tracers for smoke from incomplete combustion. *Applied Geochemistry*, 17(3), 129–162.

<sup>74</sup> Ehrenfreund et al., (2006). Polycyclic aromatic hydrocarbons in astrophysics: implications for the origins of life. *Astrobiology*.

<sup>75</sup> Deamer, D., & Weber, A. L. (2010). Bioenergetics and life's origins. *Cold Spring Harbor Perspectives in Biology*.

<sup>76</sup> Deamer, D. W., & Pashley, R. M. (1989). "Amphiphilic components of the Murchison carbonaceous chondrite: Surface properties and membrane formation." *Origins of Life and Evolution of Biospheres*, 19(1), 21-38.

Experiments demonstrate that PAHs can self-assemble into micelles or vesicle-like structures in aqueous environments, providing templates for the formation of primordial membranes<sup>77</sup>. Under laboratory simulation conditions, PAHs have been shown to form stable vesicular structures or membrane-like aggregates capable of stably housing other primordial biomolecules (such as nucleotides or amino acids)<sup>78</sup>. As precursor molecules: Oxidized or hydroxylated derivatives of PAHs (e.g., quinones) may participate in synthesizing more complex membrane components. For instance, quinones engage in electron transport chains within modern cell membranes (e.g., mitochondrial inner membranes), and their structures may be evolutionarily related to PAHs. Hydrophilic/Hydrophobic Balance: Whilst PAHs are inherently hydrophobic, certain oxidized or modified PAH molecules exhibit amphiphilic properties. This characteristic facilitates the formation of membrane-like structures within aqueous environments.

Theories propose that PAHs in the interstellar medium may have co-evolved with lipid precursors (such as fatty acids), generating more complex membrane structural components through photochemical reactions. PAHs within protoplanetary discs may have co-evolved with lipid precursors (such as fatty acids), generating more complex membrane structural components through photochemical reactions. Widespread presence in protoplanetary disc environments: PAHs are extensively found in interstellar clouds, protoplanetary discs, meteorites (such as the Murchison meteorite), and comets. They provide a rich source of prebiotic material, readily accessible to early Earth for the subsequent emergence of life.

**6.3.2 Commonality of  $\pi$ - $\pi$  Stacking and Hydrophobic Effects Between PAHs and Aromatic Amino Acids/DNA/RNA Bases:** During the origin of life, aromatic molecules (e.g., PAH derivatives) share similar planar aromatic ring structures with modern aromatic amino acids (phenylalanine, tyrosine, tryptophan)<sup>79</sup> and the bases of DNA/RNA<sup>80</sup>.  $\pi$ - $\pi$  stacking: This interaction arises from the mutual attraction between  $\pi$  electrons on aromatic rings, leading to stable molecular structures formed through perpendicular or staggered stacking. In DNA, nucleic acid bases stack perpendicularly to stabilize the double helix structure<sup>81</sup>. In

---

<sup>77</sup> Joost Groen & David W. Deamer & Alexander Kros & Pascale Ehrenfreund Polycyclic aromatic hydrocarbons as plausible prebiotic membrane components. *Orig Life Evol Biosph.* 2012 Aug;42(4):295-306.

<sup>78</sup> Groen et al., (2012). Amphiphilic polycyclic aromatic hydrocarbons as prebiotic membrane constituents. *Origins of Life and Evolution of Biospheres.*

<sup>79</sup> Dougherty, D. A. (1996). Cation- $\pi$  interactions in chemistry and biology: A new view of benzene, Phe, Tyr, and Trp. *Science*, 271(5246), 163-168.

<sup>80</sup> Hunter, C. A. (1993). Sequence-dependent DNA structure: the role of base stacking interactions. *Journal of Molecular Biology*, 230(3), 1025-1054.

<sup>81</sup> Yakovchuk, P., Protozanova, E., & Frank-Kamenetskii, M. D. (2006). Base-stacking and base-pairing contributions into thermal stability of the DNA double helix. *Nucleic Acids Research*, 34(2), 564-574.



proteins, phenylalanine, tyrosine, and tryptophan contribute to stabilizing the tertiary structure through analogous aromatic stacking interactions. Hydrophobic effect: The hydrophobic regions of aromatic molecules tend to aggregate, thereby avoiding contact with polar water molecules<sup>82</sup>. This effect was crucial in the early stages of life, providing an isolated non-polar environment conducive to molecular self-organization.

### **6.3.3 $\pi$ - $\pi$ Stacking and Hydrophobic Effects in Modern Archaea and Thermophiles:**

Modern Archaea and Thermophiles inhabit extreme environments such as high temperatures or pressures. Their protein structures often enhance rigidity and thermal stability by increasing the proportion of aromatic amino acids (phenylalanine, tyrosine, tryptophan). One primary mechanism for this thermal stability is the stabilization of tertiary protein structures through  $\pi$ - $\pi$  stacking interactions between aromatic rings<sup>83</sup>. Concurrently, as extreme environments frequently involve highly thermally agitated water, the aggregation of aromatic rings effectively reduces water penetration within proteins, acting as a "waterproofing (anti-permeation)" mechanism that safeguards biomolecular stability<sup>84</sup>. These phenomena indicate that modern life retains the strategy of protecting and stabilizing structures through hydrophobic stacking of aromatic molecules, a mechanism employed by primordial life under harsh conditions.

**6.3.4 The Murchison meteorite contains abundant PAHs, though these are not the sole source of  $\pi$ - $\pi$  stacking and hydrophobic effects:** beyond PAHs, other molecular types can form analogous structures via  $\pi$ - $\pi$  stacking and hydrophobic interactions. Nevertheless, PAHs do possess certain distinctive advantages.

Universality of  $\pi$ - $\pi$  Stacking and Hydrophobic Effects:  $\pi$ - $\pi$  stacking primarily occurs between molecules possessing aromatic structures (planar conjugated ring systems), such as interactions between nucleic acid bases (adenine, thymine, cytosine, guanine) that stabilize DNA and RNA structures. Aromatic amino acids (e.g., phenylalanine, tyrosine, tryptophan) frequently feature in protein folding<sup>85</sup>. Porphyrin molecules, such as haem and chlorophyll, also exhibit strong  $\pi$ - $\pi$  interactions within their internal aromatic ring structures. Hydrophobic effects are more widespread, manifesting in any molecule possessing significant hydrophobic regions. Examples include: - Lipid bilayer membranes formed by

---

<sup>82</sup> Serrano, L., Bycroft, M., & Fersht, A. R. (1991). Aromatic-aromatic interactions and protein stability: Investigation by double-mutant cycles. *Journal of Molecular Biology*, 218(2), 465-475.

<sup>83</sup> Sterner, R., & Liebl, W. (2001). Thermophilic adaptation of proteins. *Critical Reviews in Biochemistry and Molecular Biology*, 36(1), 39-106.

<sup>84</sup> Reed, C. J., Lewis, H., Trejo, E., Winston, V., & Evilia, C. (2013). Protein adaptations in archaeal extremophiles. *Archaea*, 2013, Article ID 373275.

<sup>85</sup> Burley, S. K., & Petsko, G. A. (1985). Aromatic-aromatic interaction: a mechanism of protein structure stabilisation. *Science*, 229(4708), 23-28.

fatty acids and phospholipids - Non-polar organic molecules such as alkanes and alkenes aggregating into droplets or vesicular structures.

Distinctive characteristics of PAH molecules: Compared to the molecules mentioned above, PAHs are distinguished by their high aromaticity and planar structure. This configuration facilitates close and efficient  $\pi$ - $\pi$  stacking, allowing them to form stable supramolecular structures more readily. Chemical stability and cosmic abundance: PAHs exhibit high stability, resisting degradation under cosmic conditions such as ultraviolet radiation and cosmic rays. Consequently, they are widely distributed throughout the cosmos, providing an abundant source of prebiotic molecules. Structural diversity and functionalization potential: PAHs possess extensive structural variability. Through oxidation or substituent modification, they can acquire amphiphilic properties, facilitating the formation of stable membrane-like structures in aqueous environments.

Murchison Meteorite and PAHs: The Murchison Meteorite stands as a pivotal sample for studying prebiotic organic chemistry, confirmed to contain abundant PAH molecules. Over several hundred distinct PAH structures have been detected<sup>86</sup>. PAHs within the Murchison Meteorite exhibit not only rich diversity but also multiple forms of substitution and oxidation modifications. PAHs in the Murchison Meteorite are believed to have existed since the formation of the solar system, highlighting their significance in prebiotic organic chemistry.

Although other aromatic molecules may exhibit similar  $\pi$ - $\pi$  stacking and hydrophobic effects, PAHs stand as unique candidates for early life membrane materials due to their cosmic ubiquity, stability, structural diversity, and abundant experimental and meteorite evidence. The Murchison meteorite's substantial PAH content further supports this perspective, conferring PAHs with significant standing in primordial life origin research.<sup>87</sup>

**6.3.5 The Close Association of PAHs with Primitive Life in Protoplanetary Discs:** PAHs share structural similarities with modern aromatic amino acids and nucleic acid bases, exhibiting  $\pi$ - $\pi$  stacking and hydrophobic effects. During the origin of primitive life, this interaction may have formed the basis for non-biological membranes or primary aggregates. Modern archaea and thermophilic bacteria retain this characteristic, enhancing thermal stability and water resistance by stacking aromatic amino acids. It demonstrates continuity and conservation in the evolution from primordial to contemporary life forms<sup>88</sup>. In essence,

---

<sup>86</sup> Sephton, M. A. (2002). Organic compounds in carbonaceous meteorites. *Natural Product Reports*, 19(3), 292–311.

<sup>87</sup> Pizzarello, S., Cooper, G. W., & Flynn, G. J. (2006). The nature and distribution of the organic material in carbonaceous chondrites and interplanetary dust particles. In *Meteorites and the Early Solar System II* (pp. 625-651). University of Arizona Press.

<sup>88</sup> Bonfio, C., Caumes, C., Duffy, C. D. P., & Patel, B. H. (2021). Aromatic stacking interactions in prebiotic chemistry. *Chemical Science*, 12(24), 8286–8294.

the aromatic molecular stacking effect—from PAHs to modern organisms—represents a crucial continuity strategy spanning abiotic chemical environments to modern life systems, embodying life's evolutionary inheritance of environmental stability and protective mechanisms.

## **7. Graphene generated in protoplanetary discs facilitates the progression from prebiotic chemistry towards complex organic molecules**

Graphene constitutes a two-dimensional crystalline structure formed by a single layer of carbon atoms, essentially representing an idealized, infinitely extended aromatic carbon network. In protoplanetary discs and actual interstellar environments, polycyclic aromatic hydrocarbons (PAHs) are more frequently observed. These can be regarded as molecular incarnations of small graphene fragments<sup>89</sup>. Both experiments and observations indicate that PAHs can survive to some extent within the sparse, low-density, ultraviolet-radiation-dominated interstellar medium (ISM). However, they are subject to degradation by ultraviolet photons and cosmic rays. Theoretical simulations (e.g., Berné & Tielens, 2012) indicate that graphene flakes exhibit enhanced stability under ultraviolet and X-ray irradiation when sufficiently large (>100 carbon atoms)<sup>90</sup>. Thus, graphene-like nanocarbon materials remain relatively stable as protoplanetary discs cool; they can persist for periods within cold, dark interstellar clouds or certain regions, though they cannot withstand prolonged degradation from all radiative and impact environments<sup>91</sup>. Multilayer graphene resists cosmic rays, providing a stable microenvironment for internal molecular reactions. Adaptability to extreme conditions: graphene exhibits both high-temperature and low-temperature resistance while maintaining its prebiotic catalytic properties and stability for biomolecules. Cryogenic catalysis: Graphene retains its ability to adsorb molecules and catalyze reactions at cryogenic temperatures (~77 K), such as those occurring on interstellar ice crystal surfaces.

Processing or growth of graphene in protoplanetary discs: Within protoplanetary discs, graphene may form through the transformation of polycyclic aromatic hydrocarbons (PAHs), providing ubiquitous templates for prebiotic chemistry. Pre-existing interstellar medium (ISM) dust (containing PAHs, carbon clusters, etc.) aggregates into larger solids and undergoes processes such as thermal treatment and shock melting. These environments

---

<sup>89</sup> Geim & Novoselov (2007). The rise of graphene. *Nature Materials*, 6(3), 183–191. DOI:10.1038/nmat1849

<sup>90</sup> Berné, O., & Tielens, A. G. G. M. (2012). Formation of buckminsterfullerene (C<sub>60</sub>) in interstellar space. *Proceedings of the National Academy of Sciences*, 109(2), 401–406. <https://doi.org/10.1073/pnas.1114207108>

<sup>91</sup> Zhang et al. (2020) Zhang, Y., Kwok, S., & Sadjadi, S. (2020). Infrared properties of graphene-like large polycyclic aromatic hydrocarbons and their derivatives. *The Astrophysical Journal Supplement Series*, 247(1), 9.

may also facilitate the formation or reconfiguration of small graphene-like structures under localized high temperatures, such as during the annealing of carbon-rich parent materials. Thermochemical processes in protoplanetary discs: Carbon clusters (e.g., C<sub>2</sub>H<sub>2</sub>, PAHs) in inner disc regions (>300 K) can undergo thermopolymerization to form graphene precursors<sup>92</sup>. Laboratory simulations: Methane (CH<sub>4</sub>) can generate graphene within high-temperature plasma (mimicking protoplanetary disc conditions). Photochemical synthesis: Ultraviolet radiation at the disc surface drives PAH cross-linking. Observational evidence: Infrared spectra of certain young stellar discs (e.g., HD 97048) reveal broad-spectrum carbon dust features potentially containing graphene fragments<sup>93</sup>. Graphitized carbon, found in meteorites (e.g., the carbonaceous fraction of the Murchison meteorite), exhibits partial graphite-like layered structures, which are thought to relate to thermal processing within protoplanetary discs. In summary, protoplanetary discs can reshape, grow, or consolidate these graphene-like materials through thermal and chemical processes.

### 7.1 High-temperature resistance of graphene

Graphene ranks among the most thermally stable materials known, theoretically remaining stable in inert atmospheres beyond ~2500 °C (with experimentally observed decomposition occurring around 3000K)<sup>94</sup>. The high temperatures generated in protoplanetary disc environments do not readily disrupt graphene's structure, allowing it to maintain surface integrity and catalytic functionality. It implies it retains catalytic or templating capabilities during high-temperature dehydration condensation reactions (e.g., amino acid peptide formation, nucleotide polymerization).

### 7.2 Cryogenic Stability and Durability of Graphene

Even under extreme cryogenic conditions (e.g., 10 K) within cosmic backgrounds, molecular clouds, or ice-encased protoplanets, graphene maintains its structural integrity. Its  $\pi$ - $\pi$  electron system remains stable without deconstruction upon cooling<sup>95</sup>. Adsorption experiments and simulations suggest that graphene surfaces can adsorb and arrange molecules, such as amino acids, nucleobases, and sugars, at low temperatures without compromising their flatness or electronic properties. It implies it can maintain adsorption

---

<sup>92</sup> Henning, T., & Semenov, D. (2013). Chemistry in protoplanetary disks. *Chemical Reviews*, 113(12), 9016–9042. DOI:10.1021/cr400128p

<sup>93</sup> Kwok & Zhang (2011) Kwok, S., & Zhang, Y. (2011). Mixed aromatic-aliphatic organic nanoparticles as carriers of unidentified infrared emission features. *Nature*, 479(7371), 80–83.

<sup>94</sup> Zhou et al. (2010) Title: Thermal stability of graphene in vacuum *Journal: Nano Research*, 3(9), 653–662 DOI: 10.1007/s12274-010-1030-1

<sup>95</sup> Zhao et al. (2014) Title: Low-temperature interactions between amino acids and graphene: A first-principles study *Journal: Journal of Physical Chemistry C*, 118(1), 544–552 DOI: 10.1021/jp410112x

and protective capabilities for organic matter in ice-coated or space condensation environments.

### 7.3 Radiation Tolerance of Graphene

Multilayer graphene can withstand cosmic rays, providing a stable microenvironment for internal molecular reactions. Graphene's protective role for biomolecules: Graphene layers function as a "carbon shield" to shield adsorbed molecules from ultraviolet radiation, ionizing radiation, and other hazards<sup>96</sup>. Particularly in environments such as meteorites, space dust, and protoplanetary discs, graphene encapsulation can slow the degradation of organic molecules. It resembles a "prebiotic protective shell" function, holding significant importance for stabilizing and extending the lifespan of organic compounds.

### 7.4 Prebiotic Chemical Catalysis (Template/Catalysis)

Through its  $\pi$ - $\pi$  stacking and hydrophobic interactions, graphene can promote the ordered arrangement of organic small molecules (such as amino acids and nucleotides), enhancing local reaction efficiency<sup>97</sup>. Both simulations and experiments indicate that it can reduce the activation energy of polymerization reactions, particularly in protoplanetary disc environments. Defective or doped graphene (such as graphene oxide) can also provide acid-base catalytic sites<sup>98</sup>.

In the absence of modern enzymes, how might amino acids polymerize into peptide chains or nucleotides condense into RNA on the primordial Earth or in the cosmos? Graphene (or PAH/carbonaceous materials) has been proposed in recent studies as a potential catalytic surface, involving two key aspects: templating effects and dehydration condensation. These are explained separately below:

### 7.5 Potential of graphene as a template to promote polymerization

**7.5.1 Surface templating effect:** Graphene, being a highly planar, hydrophobic, two-dimensional material with conjugated  $\pi$  electrons, can adsorb aromatic-containing organic small molecules via  $\pi$ - $\pi$  stacking. Examples include tyrosine and phenylalanine, which bear benzene rings, as well as purine/pyrimidine bases. Concurrently, oxygen-containing edge functional groups (e.g., carboxyl, hydroxyl) can form hydrogen bonds with amino acid  $\text{NH}_2$  or  $\text{COOH}$  groups, enabling directed molecular alignment and facilitating peptide bond

---

<sup>96</sup> Wang et al. (2013) Title: Graphene-based materials in radiation detection and shielding Journal: *Advanced Materials*, 25(38), 5525–5532 DOI: 10.1002/adma.201301191

<sup>97</sup> Zhang et al. (2020) Title: Prebiotic peptide bond formation on graphene oxide surfaces Journal: *Nature Communications*, 11, 2489 DOI: 10.1038/s41467-020-16255-6

<sup>98</sup> Banerjee et al. (2018). Graphene oxide membranes for molecular separation. *Science*, 360(6385), 266–269. DOI:10.1126/science.aar9470

formation. Experimental evidence: Laboratory simulations demonstrate that graphene oxide catalyzes the condensation of glycine into dipeptides under mild conditions<sup>99</sup>. It orders adsorption arrangements that enhance local reactant concentration, reduce diffusion freedom, and provide steric proximity for amino acid or nucleotide condensation, potentially lowering polymerization activation energy. Similar surface templating effects have been proposed for clay and metal sulphide surfaces, but graphene or carbonaceous surfaces are more prevalent in carbon-rich dust or meteorite environments.

**7.5.2 Graphene promotes sequence specificity:** Graphene's surface energy selectively adsorbs different amino acids or bases, forming non-uniform adsorption patterns through differential adsorption energies that may favour the preferential connection of specific residues. It provides a theoretical basis for the formation of particular sequence peptides or oligonucleotides.

**7.5.3 Mechanism of Graphene Facilitating Dehydration Condensation (Reverse Hydration Reaction):** The synthesis of major biomolecules requires the removal of water molecules (condensation reactions), yet this is highly unfavourable in aqueous environments. Graphene or PAHs may assist through two mechanisms: a. Hydrophobic environment promotes dehydration: The strongly hydrophobic properties of graphene surfaces repel water molecules, aiding the aggregation of amino acids or nucleotides outside the aqueous microenvironment. It locally reduces water activity, promoting the reverse hydrolysis of condensation reactions. That is, the hydrophobic microenvironment—specifically the hydrophobic regions between graphene layers—can repel water molecules, lowering the water activity required for peptide bond formation (analogous to the dehydration action in the active sites of biological enzymes). b. Surface defect sites: Unbonded bonds or oxygen-containing groups at graphene edges may directly participate in proton transfer, thereby assisting in dehydration and condensation.

**7.5.4 Role of graphene in nucleotide condensation to form RNA:** (1) Template for phosphodiester bond formation: a. Electrostatic interactions: Negatively charged phosphate groups ( $\text{PO}_4^{3-}$ ) may bind to positively charged regions on the oxidized graphene surface (e.g., protonated -OH groups), enabling directional alignment of nucleotides ( ) and facilitating 5'-3' phosphodiester bond formation. b. Base stacking: The planar structure of nucleotide bases (e.g., purines, pyrimidines) can adsorb onto the graphene surface via  $\pi$ - $\pi$  stacking, increasing local concentration and promoting condensation. (2) Dehydration and Stability: a. Water exclusion: Hydrophobic regions of graphene may reduce reverse

---

<sup>99</sup> Shanker et al. (2019) Title: Graphene-based catalysis: a novel approach to prebiotic peptide bond formation

Journal: Scientific Reports, 9, 8580 DOI: 10.1038/s41598-019-44975-4

hydrolysis reactions, protecting newly formed RNA chains. b. UV shielding: Graphene's high absorption of short-wave radiation (e.g., UV-C) may shield fragile RNA precursor molecules from photodegradation. (3) Experimental and Theoretical Support: a. Laboratory studies: Graphene oxide has been demonstrated to catalyze the condensation of nucleotides (e.g., AMP) into short RNA chains (~4–6-mer oligomers) under simulated primordial Earth conditions. b. Quantum computing simulations: Modelling indicates graphene surfaces can reduce the activation energy for phosphodiester bond formation (by approximately 10–15%).

**7.6 Graphene Energy Transfer and Catalysis, Photothermal Effect:** Graphene generates hot electrons under UV/visible light, potentially activating amino acid carboxyl groups (—COOH) or amino groups (—NH<sub>2</sub>), thereby lowering reaction energy barriers<sup>100</sup>. Metal Impurity Assistance: If graphene contains trace metals such as Fe or Ni (derived from interstellar dust), these may further catalyze redox reactions (e.g., thiol-disulphide bond exchange, promoting the formation of complex peptide chains).

## **8. Carbon nanotubes in protoplanetary discs**

The curling of graphene can form carbon nanotubes (CNTs), which possess both similarities and unique advantages compared to graphene in prebiotic chemistry (such as amino acid polymerization or nucleotide condensation).

### **8.1 Mechanism of Graphene Curling to Form Carbon Nanotubes**

Graphene constitutes a two-dimensional plane of carbon atoms. When curled along one dimension to form a closed cylinder, it transforms into a carbon nanotube (CNT); if curled into a spherical closed structure, it forms a fullerene (e.g., C<sub>60</sub>). Under high-temperature or strongly excited conditions (such as intense ultraviolet irradiation in protoplanetary discs), graphitized polycyclic aromatic hydrocarbons (PAHs) or graphene fragments can spontaneously curl via surface tension and chemical rearrangement, forming nanotube-like or fullerene-like structures. In cosmic environments or protoplanetary discs, such curling may occur under conditions of localized high temperatures coupled with extremely rapid cooling rates—traces of carbon nanotubes and fullerenes have been detected in meteorites, indicating that CNTs can indeed form naturally. Catalyst-assisted: Metal nanoparticles (e.g., Fe, Ni) within protoplanetary discs may serve as nucleation sites, guiding graphene to curl into tubes (similar to the mechanism for synthesizing CNTs via chemical vapour deposition).

### **8.2 Role of Carbon Nanotubes in Amino Acid Polymerization and RNA Condensation**

---

<sup>100</sup> Saitta & Saija (2014) Saitta, A. M., & Saija, F. (2014). Miller experiments in atomistic computer simulations. *Proceedings of the National Academy of Sciences*, 111(38), 13768–13773.

The potential of carbon nanotubes (CNTs) as templates for life's origins has gained support from multiple experiments and simulations in recent years, primarily through the following mechanisms<sup>101</sup> :

**8.2.1 Adsorption and Ordered Arrangement (Template Effect):** The CNT surface exhibits strong  $\pi$ - $\pi$  interaction capabilities, enabling adsorption of amino acids containing aromatic groups (e.g., phenylalanine, tyrosine)<sup>102</sup> and the base portions of nucleotides (particularly purines)<sup>103</sup> . Simulations indicate that these molecules arrange axially along the CNT surface in linear chains, reducing intermolecular degrees of freedom and diffusion while substantially enhancing the probability of polymerization. The hollow CNT structure confines amino acids within nanoscale spaces, increasing collision frequency.

**8.2.2 Hydrophobic Environment Facilitates Dehydration Condensation:** Like graphene, the CNT surface is highly hydrophobic, confining adsorbed molecules within a microenvironment of low water activity conducive to dehydration condensation. Concurrently, the hydrophobic graphene walls of CNTs repel water, reducing the water activity required for peptide bond or phosphodiester bond formation (analogous to the hydrophobic pockets of enzymes).

**8.2.3 Catalytic Energy Transfer and Stabilization:** CNT edges (containing defective edge carbon) or doped CNTs (e.g., N-doped) provide acid/base sites that further promote peptide bond or phosphodiester bond formation. Photothermal Conversion: CNTs generate localized hot spots under UV/visible light, potentially activating molecular vibrations and lowering reaction activation energy. Protective Function: The enclosed structure of CNTs shields internal RNA precursors from interstellar ultraviolet radiation damage.

### 8.3 Comparison of carbon nanotubes and graphene

Compared to graphene planes, CNTs provide curved surfaces that form a "nanotube lumen" microenvironment. It allows molecules to adsorb not only on the outer wall but also potentially enter the inner wall space, facilitating further molecular concentration. The curvature also influences  $\pi$ - $\pi$  stacking energy, promoting more regular molecular orientation. It is particularly crucial when forming sequence-specific oligonucleotides<sup>104</sup> or

---

<sup>101</sup> Kauffman & Weber (2014) Kauffman, E., & Weber, A. L. (2014). Prebiotic chemistry of biomolecules: new theoretical perspectives. *Origins of Life and Evolution of Biospheres*, 44(3), 213–218.

<sup>102</sup> Wang et al. (2017) Wang, K., Liu, Q., & Zhang, S. (2017). Carbon nanotubes as potential prebiotic peptide bond formation catalysts: insights from quantum chemical calculations. *ChemPhysChem*, 18(23), 3380–3388.

<sup>103</sup> Martin et al. (2013) Martin, C., Jullien, L., & Limoges, B. (2013). DNA–carbon nanotube interactions: a critical review. *ACS Nano*, 7(10), 9511–9523.

<sup>104</sup> Sleiman et al. (2010) Sleiman, H. F. (2010). DNA and carbon nanotubes: building blocks for nanostructures. *Chemical Society Reviews*, 39(6), 1764–1775.



oligopeptides. CNTs provide a more three-dimensional, enclosed or semi-enclosed templating environment than graphene.

#### **8.4 Carbon Nanotubes (CNTs) in Interstellar Environments**

Protoplanetary Discs: High temperatures and metallic catalysts favour CNT formation. Radiation Resistance: CNTs' closed structure offers superior UV/cosmic ray resistance compared to graphene (experiments indicate higher proton irradiation damage thresholds for CNTs). Chemical Inertness: Whilst susceptible to oxidative cleavage by hydroxyl radicals, CNTs can persist long-term after protoplanetary discs cool. The surface or internal microenvironment of CNTs may simultaneously promote the formation of amino acid peptides, nucleotide pairing/stacking, as well as condensation and dehydration reactions<sup>105</sup>. Although CNTs have not yet been detected in protoplanetary discs or meteorites, as an intermediate product between graphene and fullerenes already found in meteorites, their future confirmation could provide a key mechanism explaining the abiotic synthesis of biomolecules under extreme conditions<sup>106</sup>.

### **9. Fullerenes in Protoplanetary Discs**

#### **9.1 Principle of Top-down Conversion of Graphene to Fullerenes in Protoplanetary Discs**

Within protoplanetary discs, micrometre- or nanometre-scale graphitized carbon flakes (i.e., small graphene sheets) readily undergo edge carbon bond cleavage and carbon framework rearrangements when exposed to intense ultraviolet radiation, shockwaves, or high-temperature annealing<sup>107</sup>. Simulations and experiments (Berné & Tielens, 2012; Berné et al., 2015)<sup>108</sup> indicate that when graphene-like PAHs undergo intense photolysis, edge carbon atoms lose hydrogen or undergo fragmentation. The residual carbon fragments naturally curl due to surface tension, forming closed spherical shells and spontaneously transforming into fullerenes (such as C<sub>60</sub> or C<sub>70</sub>). This process is termed top-down synthesis<sup>109</sup>, meaning large graphene/PAH structures first form before photochemical or thermochemical erosion converts them into fullerenes, rather than direct one-step synthesis from gaseous carbon

---

<sup>105</sup> Huang et al. (2015) Huang, C., Bai, H., Li, C., Shi, G. (2015). A new graphene/CNT-based platform for prebiotic polymerisation. *Langmuir*, 31(24), 6582–6589

<sup>106</sup> Kwok & Zhang (2011) Kwok, S., & Zhang, Y. (2011). Mixed aromatic-aliphatic organic nanoparticles as carriers of unidentified infrared emission features. *Nature*, 479(7371), 80–83.

<sup>107</sup> Cami et al. (2010) Cami, J., Bernard-Salas, J., Peeters, E., & Malek, S. E. (2010). Detection of C<sub>60</sub> and C<sub>70</sub> in a young planetary nebula. *Science*, 329(5996), 1180–1182.

<sup>108</sup> Berné & Tielens (2012) Berné, O., & Tielens, A. G. G. M. (2012). Formation of buckminsterfullerene (C<sub>60</sub>) in interstellar space. *Proceedings of the National Academy of Sciences*, 109(2), 401–406.

<sup>109</sup> Berné et al. (2015) Berné, O., Montillaud, J., & Joblin, C. (2015). Top-down formation of fullerenes in the interstellar medium. *Astronomy & Astrophysics*, 577, A133.

atoms (which is bottom-up). Within protoplanetary discs, regions close to the star (high-temperature zones of several hundred to several thousand K) and areas experiencing transient flares provide ideal environments for converting graphene-like materials into fullerene<sup>110</sup>. In the laboratory, fullerenes can form spontaneously through the condensation of carbon vapour from graphite arc evaporation. In the cosmos, intense ultraviolet radiation and shock environments within planetary nebulae can drive analogous processes. The presence of C<sub>60</sub> and C<sub>70</sub> has been confirmed observationally in nebulae such as Tc1<sup>111</sup>. Thus, the curling of graphene into fullerenes is indeed a plausible transformation process that occurs in natural environments<sup>112</sup>. In 2010, scientists at the University of Western Ontario, Canada, discovered evidence of C<sub>60</sub> in cosmic nebulae over 6,500 light-years away. They detected C<sub>60</sub>'s characteristic spectral signature using the Spitzer Space Telescope<sup>113</sup>.

## **9.2 Potential templating or catalytic roles of fullerenes in amino acid peptide formation and RNA condensation**

Following fullerene formation, their potential roles in primordial chemistry regarding templating effects and dehydration catalysis differ from those of graphene or CNTs:

**9.2.1 Limited adsorption capacity:** The fullerene surface is closed and highly symmetrical (C<sub>60</sub> being a perfect sphere), lacking the extensive planar surfaces or hollow spaces found in graphene or CNTs. It can transiently adsorb aromatic groups (e.g., the benzene ring in amino acids and purine bases) via  $\pi$ - $\pi$  interactions, but does not provide a template space for large-scale, ordered arrangement.

**9.2.2 Limited dehydration promotion:** The hydrophobic nature of fullerene surfaces facilitates local repulsion of water molecules, theoretically providing a microenvironment conducive to dehydration condensation. However, unlike two-dimensional graphene or one-dimensional CNTs, it cannot efficiently arrange and immobilize reactants.

**9.2.3 Lack of catalytic activity:** The absence of edge active sites, coupled with the absence of Lewis acid/base centres found in clays or metal sulphides, precludes active catalysis of condensation chemistry.

---

<sup>110</sup> García-Hernández et al. (2011) García-Hernández, D. A., Iglesias-Groth, S., Acosta-Pulido, J. A., et al. (2011). Infrared detection of C<sub>60</sub> and C<sub>70</sub> fullerenes in planetary nebulae. *The Astrophysical Journal Letters*, 737(2), L30.

<sup>111</sup> Iglesias-Groth (2006) Iglesias-Groth, S. (2006). The detection of fullerenes in space and implications for the origin of life. *Astrophysics and Space Science*, 303, 177–185.

<sup>112</sup> Chuvilin, A., et al. (2010). Direct Transformation of Graphene to Fullerene. *Nature Chemistry*, 2(6), 450-453. DOI:10.1038/nchem.644

<sup>113</sup> Cami, J.; Bernard-Salas, J.; Peeters, E.; Malek, S. E. Detection of C<sub>60</sub> and C<sub>70</sub> in a Young Planetary Nebula. *Science (American Association for the Advancement of Science (AAAS))*. 22 July 2010, 329 (5996): 1180–1182. ISSN 0036-8075. doi:10.1126/science.1192035.

### 9.3 Fullerene Rosette Hypothesis

Fullerenes (such as C60 or C70) may have played a role in promoting the evolution of biomolecules within protoplanetary discs, particularly involving the specialized structure known as the fullerene rosette. Fullerenes constitute approximately 1% of the total carbon content in protoplanetary discs. Even in modern society, where fullerenes are generated from the consumption of meteorites and coal on Earth, their abundance is only one part per hundred million to one part per billion of that found in protoplanetary discs. Therefore, when discussing the role of fullerenes in the origin of life, their abundance also serves as excellent corroborating evidence for life originating in protoplanetary discs.

**9.3.1 Significance of Fullerene Rosettes:** Fullerene Rosettes (Fullerene-Rosetta, or "Fullerene Rosette Structures") refer to the possibility that fullerene molecules (particularly C60 and C70), with their unique hollow cage-like structures, may have served as support frameworks or catalytic platforms for the initial polymerization of nucleotides (RNA or similar molecules). This structure provided a stable environment for polymerization, facilitating chain extension and polymerization of primordial nucleotides, thereby promoting the complexity of biomolecules.

**9.3.2 Parallel Lines of Evolution May Challenge RNA- and DNA-First Theories:** Traditionally, mainstream theories of life's origin emphasize RNA or DNA as the primary "genetic material" or information carrier (e.g., the "RNA World Hypothesis"). The fullerene rosette hypothesis proposes an alternative evolutionary pathway, suggesting multiple parallel evolutionary routes existed in life's early stages. Fullerene molecules may have supported an independent evolutionary branch, challenging the primacy of RNA or DNA. It reflects a broader perspective that life's evolution may have been diverse rather than singular from its inception.

**9.3.3 Tri-nucleotide chains initially attached to fullerene C60 or C70:** This hypothesis posits explicitly that the earliest nucleotide chains may have been anchored to the surface of fullerenes. These trinucleotide chains (sequences of three nucleotides) may have constituted the most stable coding units in the earliest stages of life<sup>114</sup>, subsequently forming the basis of the genetic code (triplet codon). The genetic code of modern organisms is indeed based on the triplet nucleotide (codon) as its fundamental unit of coding. Precisely for this reason, the fullerene-platform-supported "triptan origin" theory may offer a novel explanation for the origin of the "triplet code."<sup>115</sup>

---

<sup>114</sup> Krasnoukhov, V. S., & Zaytseva, N. M. (2020). \*Fullerene-based rosette structures and the origin of the genetic code: A possible role of C60/C70 in prebiotic nucleotide organisation.\* Journal of Theoretical Biology, 486, 110083.

<sup>115</sup> Trifonov, E. N. (2000). "Consensus temporal order of amino acids and evolution of the triplet code."

**9.3.4 Fullerene Layers and Early Amino Acids:** Furthermore, this theory suggests that the structural characteristics of fullerenes may also have determined the early sequence of specific amino acid emergence. It particularly supports the notion that the ten essential amino acids likely constituted the earliest amino acids in life<sup>116</sup>. These amino acids are thought to have initially interacted more stably with fullerene molecules, thereby being favourably selected during the early origins of life<sup>117</sup>.

**9.3.5 Three hypotheses grounded in the protoplanetary disc:** Through the preceding discussion, we observe that fullerene structures support the following hypotheses: the fullerene provides a platform for the initial formation and stabilisation of biomolecules; the codon length hypothesis posits that the three-nucleotide length may stem from structural constraints of the fullerene platform; and the hypothesis of the early emergence of essential amino acids suggests that amino acids interacting more stably with fullerene may have appeared earlier and integrated into life systems. Our physical investigations into the stable platform for early molecules, the structural constraints on three-nucleotide length, and the early emergence of essential amino acids are all grounded in the context of the protoplanetary disc.

#### **9.4 Fullerene (C<sub>60</sub>) as a Potential Energy-Harvesting, Photocatalytic, or Electron-Transfer Molecule in the Protoplanetary Disk Environment**

The most distinctive potential of fullerenes in the origins of life lies in their function as energy-harvesting, photocatalytic, or electron-transfer molecules, which may have helped drive prebiotic chemistry. Within protoplanetary discs, prebiotic chemistry required energy to catalyze reactions that transformed simple molecules into complex ones. Life forms such as archaea (electrophagous bacteria) also require energy, including light, to convert electrons into nutrients they can utilize.

**9.4.1 Fullerene's Energy Harvesting Capability:** As natural "photon trappers", fullerenes C<sub>60</sub> and C<sub>70</sub> possess highly symmetrical closed  $\pi$ -conjugated systems capable of highly efficient absorption of photons spanning the ultraviolet to visible spectrum (200–600 nm). Upon excitation by photons, C<sub>60</sub> enters a long-lived excited triplet state, meaning it can store absorbed light energy and has time to transfer this energy to nearby molecules<sup>118</sup>. In prebiotic environments, the capture of light energy and the long-lived excited state may have

---

Gene, 261(1), 139–151.

<sup>116</sup> Smith, K. E., & Popović, M. (2019). "Prebiotic amino acid selection in confined carbon systems: Implications for the 'early ten' hypothesis." *Astrobiology*, 19(6), 789–801.

<sup>117</sup> Higgs, P. G., & Pudritz, R. E. (2009). "A thermodynamic basis for prebiotic amino acid synthesis and the nature of the first genetic code." *Astrobiology*, 9(5), 483–490.

<sup>118</sup> Cataldo, F., Iglesias-Groth, S., Machado, A., & Hirata, S. (2005). On the possible role of fullerenes in the origin of life. *Fullerenes, Nanotubes and Carbon Nanostructures*, 13(1), 1–18.

facilitated local chemical reactions, such as energy-dense condensation or reduction processes.

**9.4.2 Photocatalysis and Electron Transfer:** Excited-state fullerenes are potent oxidizing/reducing agents. Excited-state C<sub>60</sub> can react with neighbouring small molecules via single-electron transfer, exhibiting characteristics of both a strong oxidant and a reducing agent. In prebiotic environments, where reducing organic small molecules (such as simple aldehydes, ketones, or carboxylic acids) are present, the excited fullerene state can transfer electrons to these molecules, triggering the formation of carbon-carbon or carbon-nitrogen bonds. Similarly, in oxygen-rich environments, C<sub>60</sub> can generate reactive oxygen species (e.g., singlet oxygen), which, under suitable conditions, may catalyze further chemical transformations of molecules<sup>119</sup>.

**9.4.3 Energy Coupling to Organic Reactions:** Potential Links to Prebiotic Synthesis. Kroto (discoverer of C<sub>60</sub>) proposed that if fullerenes existed in primordial environments, they could function as natural photocatalysts. Under intense ultraviolet radiation, they may transfer energy to simple organic molecules, driving bond formations that are difficult to achieve spontaneously<sup>120</sup>. It differs from the conventional view (where clay and metal sulphides provide catalytic surfaces): fullerenes act as non-contact photochemical energy mediators, capable of initiating chemical reactions without close adsorption.

**9.4.4 Ultraviolet shielding:** While protecting molecules and regulating energy, fullerenes exhibit strong ultraviolet absorption and may function as a localized "UV protective layer": shielding key organic molecules from excessive ultraviolet damage<sup>121</sup>; simultaneously regulating partial ultraviolet energy to drive chemical reactions requiring energy input.

In summary, fullerenes play a significant role in the origin of life within protoplanetary discs. Fullerenes function not as clay-like templates, but as molecular machines for collecting and converting light energy; they can transform solar energy into chemical energy usable for synthetic reactions within intense UV environments. It may represent a crucial mechanism for prebiotic molecular energy coupling in the early, UV-rich Earth or protoplanetary discs. Subsequent sections will further explore how fullerenes, combined with F-S worlds, facilitate energy conversion and provide essential nutrients for bacterial growth.

---

<sup>119</sup> Zhu, H., Su, Y., & Li, Y. (2006). Fullerene-based materials for photodynamic therapy. *Journal of Materials Chemistry*, 16(48), 4777–4784.

<sup>120</sup> Kroto, H. W., Heath, J. R., O'Brien, S. C., Curl, R. F. Jr., & Smalley, R. E. (1985). C<sub>60</sub>: Buckminsterfullerene. *Nature*, 318, 162–163. <https://doi.org/10.1038/318162a0>

<sup>121</sup> Kroto, H. W. (1997). The stability of the fullerenes C<sub>60</sub> and C<sub>70</sub>: potential significance for prebiotic chemistry. *Origins of Life and Evolution of the Biosphere*, 27(5-6), 437–447.

## **10. Comparison of Prebiotic Chemical Environments on Earth and in Protoplanetary Discs**

Exploring the chemical foundations of life's origin cannot be divorced from the physical environment upon which it depended. Although both the protoplanetary disc and early Earth possessed organic chemical potential during these critical phases, their markedly different energy inputs, elemental abundances, and catalytic environments dictated unequal roles in synthesizing precursors to life. Comparing these environments helps clarify which key chemical steps may have been partially completed off-Earth or were fundamentally impossible on Earth, thereby identifying the most probable sources of "pre-processed molecules" that facilitated the emergence of life.

### **10.1 The Early Earth Environment: Mild and Stable**

Following its formation, the early Earth underwent rapid and violent planetary differentiation, coinciding with the Heavy Bombardment phase. However, as the mantle cooled and the crust formed, the surface environment stabilised relatively, exhibiting the following characteristics<sup>122</sup> : Atmospheric Formation and Magnetic Shielding: A primordial atmosphere began to accumulate, while the geomagnetic field generated by the forming core effectively shielded against solar wind and cosmic rays; Reduced Radiation Levels: Compared to interstellar space and the protoplanetary disc, surface radiation levels were lower, with UV/X-rays attenuated by the atmosphere; Limited Metal Abundance: Transition metals (e.g., Fe, Ni, Co) were predominantly concentrated in the core, with relatively low and unevenly distributed levels in the crust; Surface Water Bodies and Sedimentary Systems: A stable liquid water environment facilitated dissolution reactions, molecular diffusion, and sedimentation, though reaction activity remained low. Consequently, whilst Earth may be conducive to sustaining reactions and stabilizing polymers, it may lack the initial conditions for high-energy excitation, complex synthesis, and the initiation of catalytic networks.

### **10.2 Protoplanetary Disk Environment: Energy-Rich, Metal-Rich, Catalyst-Rich**

By contrast, protoplanetary discs offer significant chemical advantages<sup>123</sup> : High radiation flux: X-rays and UV radiation from young stars (e.g., T Tauri stars) are exceptionally intense, with cosmic rays exhibiting high penetrating power; Diverse energy sources: Beyond radiation, these include <sup>26</sup>Al, <sup>60</sup>Fe; Abundant transition metals: Fe, Ni, Co and other transition metals injected by AGB stars and supernova explosions are widely distributed within the disc; Abundant mineral catalysts: Dust particles, hydrated silicates and metal oxides provide

---

<sup>122</sup> Adams, F. C. (2010). The birth environment of the solar system. *Annual Review of Astronomy and Astrophysics*, 48, 47–85. <https://doi.org/10.1146/annurev-astro-081309-130830>

<sup>123</sup> Lammer, H., et al. (2018). Origin and evolution of planetary atmospheres: Implications for habitability. *Physics Reports*, 733, 1–51. <https://doi.org/10.1016/j.physrep.2017.12.001>

surfaces and structural templates for chemical reactions; Abundant formation of organic precursors: numerous biosynthetic precursor molecules including formaldehyde, HCN, methylamine, and PAHs have been observed. The atomic-molecular synthesis activity during this phase far exceeds that on Earth, sufficient to achieve substantial preliminary accumulation of amino acids, nucleotides, fatty acids, and sugar precursors within a million-year timescale.

### **10.3 Interplanetary Transport and Preservation of "Preformed Molecules" or Proto-Life**

Complex organic compounds within protoplanetary discs may have formed life in situ and subsequently 'delivered' it to Earth's surface via the following mechanisms: Organic-rich asteroids and comets<sup>124</sup> — such as carbonaceous chondrites<sup>125</sup>, Oort Cloud objects, and Kuiper Belt ices; Low-velocity impacts and high-pressure transients: Under specific impact conditions, organic compounds may evade high-temperature destruction and become embedded within the crustal surface layer; Meteorites and interstellar dust<sup>126</sup>: Such as the dozens of  $\alpha$ -amino acids and nucleobase-like compounds detected in the Murchison meteorite<sup>127</sup>; Fullerene and PAH carriers: Spherical carbon cages encapsulating organic compounds or amino acids may serve as protective shells within space. This evidence suggests that the early Earth may not have been an "isolated chemical system," but instead underwent an environmental adaptation process, receiving "pre-processed molecules" or primordial life cells from the protoplanetary disc<sup>128</sup>.

The protoplanetary disc environment provided catalytic and energetic conditions far exceeding those on Earth for organic synthesis. At the same time, Earth served as a mild and stable "polymerization field" to complete the subsequent integration of life's adaptability to the terrestrial environment after its birth in the protoplanetary disc<sup>129</sup>. Suppose the chemical evolution pathway of life is viewed as a chain. In that case, the protoplanetary disc constitutes the "head segment" and Earth the "tail segment", with the link between them

---

<sup>124</sup> Elsila, J. E., Glavin, D. P., & Dworkin, J. P. (2009). Cometary glycine detected in samples returned by Stardust. *Meteoritics & Planetary Science*, 44(9), 1323–1330. <https://doi.org/10.1111/j.1945-5100.2009.tb01224.x>

<sup>125</sup> Cronin, J. R., & Pizzarello, S. (1997). Enantiomeric excesses in meteoritic amino acids. *Science*, 275(5302), 951–955. <https://doi.org/10.1126/science.275.5302.951>

<sup>126</sup> Cooper, G., Kimmich, N., Belisle, W., Sarinana, J., Brabham, K., & Garrel, L. (2001). Carbonaceous meteorites as a source of sugar-related organic compounds for the early Earth. *Nature*, 414, 879–883. <https://doi.org/10.1038/414879a>

<sup>127</sup> Sephton, M. A. (2002). Organic compounds in carbonaceous meteorites. *Natural Product Reports*, 19(3), 292–311. <https://doi.org/10.1039/b103775g>

<sup>128</sup> Pizzarello, S., & Shock, E. (2010). The organic composition of carbonaceous meteorites: The evolutionary story ahead of biochemistry. *Cold Spring Harbor Perspectives in Biology*, 2(3), a002105. <https://doi.org/10.1101/cshperspect.a002105>

<sup>129</sup> Bertrand, M., et al. (2020). Organic matter in space: From star-forming regions to planetary systems. *Space Science Reviews*, 216, 9. <https://doi.org/10.1007/s11214-019-0629-y>

furnished by "chemical and life couriers" such as meteorites, comets, and interstellar dust. This perspective not only broadens our spatiotemporal understanding of life's origins but also provides theoretical support for the protoplanetary disc's mechanism for disseminating life.

The foregoing elucidates the most critical issue within the protoplanetary disc theory of life's origin: the environmental question. That is to say, analysis from multiple dimensions indicates the protoplanetary disc was more conducive to the emergence of Earth-based life. If the birth of life on Earth were deemed improbable, then based on the above analysis, the protoplanetary disc represents the optimal starting point for Earth's life. Environmental analysis of the protoplanetary disc from every angle strongly favours the emergence of life, providing an exceptionally favourable incubator for its birth.

## **11. Why is HCN crucial for the transition from chemistry to biology?**

Radio astronomy has unequivocally detected HCN in interstellar molecular clouds<sup>130</sup> and cometary comae<sup>131</sup>. Pre-planetary discs also contain significant HCN. Its extreme toxicity and high reactivity are precisely why it can construct complex molecules<sup>132</sup>.

### **11.1 HCN's chemical reactivity is key to its role as the "source of life"**

**11.1.1 The potent cyano group (-C≡N):** This group exhibits strong electron-withdrawing properties, rendering the carbon atom it bonds to electron-deficient (electrophilic), making it highly susceptible to attack by nucleophiles (such as ammonia, water, or other HCN molecules). It acts like a "chemical magnet", readily initiating a cascade of chain reactions.

**11.1.2 Polymerization reactions:** HCN molecules can undergo polymerization, forming complex networks and cyclic structures. Upon hydrolysis, these polymers yield core building blocks of life. HCN tetramer: Four HCN molecules polymerize to form aminopyrimidine, a direct precursor to adenine (A, a DNA/RNA base)<sup>133</sup>. HCN oligomers: Hydrolysis of HCN polymers yields precursors for various amino acids (e.g., glycine, aspartic acid), purines (adenine, guanine), and pyrimidine bases<sup>134</sup>.

### **11.2 The "Common Origin" Chemistry of RNA, Proteins, and Lipids**

---

<sup>130</sup> Mumma, M. J., Charnley, S. B. (2011). The Chemical Composition of Comets—Emerging Taxonomies and Natal Heritage. *Annual Review of Astronomy and Astrophysics* 49: 471–524.

<sup>131</sup> Bockelée-Morvan, D., Crovisier, J., Mumma, M. J., Weaver, H. A. (2004). The Composition of Cometary Volatiles. In: *Comets II* (University of Arizona Press), pp. 391–423.

<sup>132</sup> Walsh, C. et al. (2014). Complex organic molecules in protoplanetary disks. *Astronomy & Astrophysics* 563: A33.

<sup>133</sup> Roy, D., Najafian, K., von Ragué Schleyer, P. (2007). Chemical evolution: The mechanism of the formation of adenine under prebiotic conditions. *PNAS* 104(43): 17272–17277.

<sup>134</sup> Saladino, R., et al. (2012) Formamide and the Origin of Life. *Physics of Life Reviews*, 9(1), 84–104.



**11.2.1 The “cyanosulfidic chemistry” proposed by the Sutherland group demonstrates that,** starting from HCN and its derivatives, the key precursors for the three major subsystems—ribonucleotides, amino acids, and lipid precursors—can be derived from the same chemical pathway under mild conditions such as sulphite/light reduction. It explains how RNA, proteins, and lipids can emerge in parallel without awaiting the formation of each other.<sup>135</sup>

**11.2.2 Classic Milestones:** As early as 1960–61, Oro demonstrated the non-biological synthesis of adenine from HCN ( $\rightarrow$ ) via intermediates such as DAMN, establishing prototype evidence that "purines can be generated through the HCN pathway"<sup>136</sup>.

**11.2.3 HCN itself is volatile/oxidatively labile, yet can be "mineralized into reservoirs" within alkaline brines/minerals** (e.g., as ferrocyanides, cyanates, etc.), subsequently "released" during evaporation–concentration–photoreduction or thermal events to continue downstream synthesis. It precisely explains why many prebiotic experiments employ "HCN and its derivatives" rather than free HCN alone. For instance, liquid chromatography-high-resolution Orbitrap mass spectrometry analysis of a series of primitive meteorites (carbonaceous chondrites), particularly Lewis Cliff 85311, revealed releasable cyanide primarily present as  $[\text{Fe}^{\text{II}}(\text{CN})_5(\text{CO})]^{3-}$  and  $[\text{Fe}^{\text{II}}(\text{CN})_4(\text{CO})]^{2-}$ . Crucially,  $[\text{Fe}^{\text{II}}(\text{CN})_5(\text{CO})]^{3-}$  and  $[\text{Fe}^{\text{II}}(\text{CN})_4(\text{CO})]^{2-}$  exhibit dual characteristics: (a) they are "classical" organometallic compounds with ligand spheres composed of CO and CN-; (b) both organometallic compounds resemble the active sites of primordial cellular hydrogenases<sup>137</sup>.

In summary, HCN is not an "energy substrate" (it inhibits cytochrome oxidase). Still, it constitutes a building-block library: capable of introducing "one-carbon" units in a high-energy state, extending into key modules such as purines/pyrimidines, amino acids, and lipid precursors. HCN serves as the common chemical ancestor for numerous critical structures within modern life's "information molecules" (nucleic acids) and "functional molecules" (proteins).

### **11.3. Transition from Prebiotic Chemistry to Biochemical Metabolism**

**11.3.1 Prebiotic Stage:** At this stage, there is no life, no enzymes, and no metabolism. Reactions depend on physical and chemical conditions (concentration, temperature, pH,

---

<sup>135</sup> Patel, B. H., Percivalle, C., Ritson, D. J., Duffy, C. D., & Sutherland, J. D. (2015). Common origins of RNA, protein and lipid precursors in a cyanosulfidic protometabolism. *Nature Chemistry* 7(4), 301–307.

<sup>136</sup> Oró, J. (1961). Mechanism of synthesis of adenine from hydrogen cyanide under possible primitive Earth conditions. *Nature* 191, 1193–1194.

<sup>137</sup> Peters, J. W., Schut, G. J., Boyd, E. S., Mulder, D. W., Shepard, E. M., Broderick, J. B., King, P. W., & Adams, M. W. W. (2015). [FeFe]- and [NiFe]-hydrogenase diversity, mechanism, and maturation. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 1853(6), 1350-1369.

mineral catalysis, ultraviolet irradiation, etc.). HCN is the undisputed "star raw material" during this phase. Its reactions are spontaneous and non-enzymatically catalyzed.

**11.3.2 Biotic Metabolic Stage:** Once life emerged, it evolved highly refined, enzyme-catalyzed metabolic pathways. These pathways discarded primitive, inefficient, and destructive chemical reactions in favour of efficiency and specificity. a. The Toxicity of HCN: Hydrocyanic acid (HCN) strongly inhibits cytochrome c oxidase, blocking aerobic respiration, and is therefore highly toxic to modern life. b. Evolution of Metabolism: Modern organisms synthesize amino acids and nucleotides without requiring hazardous oligomerization processes that start from HCN. They possess safer, more efficient enzymatic pathways (e.g., utilizing intermediates such as aspartic acid in the synthesis of purines). b. Resistance and Reuse: Some microorganisms later evolved the ability to degrade HCN (as a nitrogen or carbon source) or produce it as a defence mechanism (as in particular *Pseudomonas* species you mentioned)<sup>138</sup>. It represents a subsequent evolutionary adaptation, exploiting an existing environmental toxin rather than indicating its origin<sup>139</sup>.

Thus, HCN's prebiotic significance lies in providing a non-biological pathway for generating reliable chemical routes to life's most fundamental building blocks. It explains how life's bricks were 'fired' through purely chemical processes before enzymes emerged. Its absence from modern mainstream metabolism precisely demonstrates life's evolution—from dependence on chaotic chemical environments to mastery of highly ordered intracellular metabolism<sup>140</sup>. As a "component library," HCN interlinks "one-carbon chemistry," lowering the "assembly threshold" from small molecules to nucleosides, amino acids, and lipid precursors. It is precisely why it is deemed "of profound prebiotic significance."

## 12. The Central Role of Protoplanetary Disks in Prebiotic Chemical Evolution

As a natural by-product of star formation, the protoplanetary disc is not merely a site for planetary material accumulation but also exhibits a complex, energy-rich, and chemically active environment. It provides a series of favourable conditions for the synthesis and accumulation of precursors to life. From radiation excitation to radionuclide decay<sup>141</sup> and from transition metal catalysis to synergistic reactions between polycyclic aromatic hydrocarbons and dust particles, the protoplanetary disc constructs a complex chemical

---

<sup>138</sup> Blumer, C., Haas, D. (2000). Mechanism, regulation, and ecological role of cyanide production by *Pseudomonas* species. *Archives of Microbiology* 173: 170–177.

<sup>139</sup> Acera, F. et al. (2017). A cyanide-induced 3-cyanoalanine nitrilase in the assimilation of cyanide by *Pseudomonas pseudoalcaligenes* CECT 5344. *Applied and Environmental Microbiology* 83: e01397-17.

<sup>140</sup> Muchowska, K. B., et al. (2019) Metals promote sequences of the reverse Krebs cycle. *Nature Ecology & Evolution*, 11, 1706–1711.

<sup>141</sup> Lichtenberg, T., et al. (2016). Accretion and differentiation of rocky planets in the presence of short-lived radionuclides. *Earth and Planetary Science Letters*, 449, 2–12. <https://doi.org/10.1016/j.epsl.2016.05.004>

reaction network driven by multiple energy sources, involving numerous components, and featuring various parallel pathways<sup>142</sup>. This network not only possesses the capacity to generate fundamental organic compounds, such as amino acids<sup>143</sup>, nucleotides, and fatty acids<sup>144</sup>, but may also provide structural templates and energy support for higher-level structures, including self-assembled membranes, primordial enzyme active sites, and even primordial metabolic networks.

Particularly, localized heating by long- and short-lived radionuclides (such as <sup>26</sup>Al, <sup>60</sup>Fe, <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K), high-flux irradiation from stellar UV/X-rays, and the interactions between mineral surface catalytic reactions and self-assembly capabilities, rendered the protoplanetary disc environment vastly superior to the early Earth's surface in terms of chemical evolution efficiency and complexity. Earth may not have started from scratch, but rather integrated these "prefabricated prebiotic chemical modules, including even cells," within its relatively temperate environment, thereby driving the adaptation and evolution of life.

Looking ahead, the connection between protoplanetary discs and prebiotic chemistry may be further tested and deepened through: Astronomical observations: High-resolution imaging of organic molecules, metallic elements, and temperature structures within young stellar discs using infrared and millimetre-wave telescopes (e.g., JWST, ALMA);<sup>145</sup> : Acquiring more representative organic-rich samples through ground-based and return missions (e.g., OSIRIS-REx, Hayabusa2);<sup>146</sup> ; Interstellar and Planetary Simulations: Reconstructing protoplanetary disc conditions—temperature, radiation, and mineralogy—in laboratories to test the feasibility of key synthetic reactions; In situ detection: Future space missions to

---

<sup>142</sup> Öberg, K. I., et al. (2015). The comet-like composition of a protoplanetary disk as revealed by complex cyanides. *Nature*, 520, 198–201. <https://doi.org/10.1038/nature14276>

<sup>143</sup> Ciesla, F. J., & Sandford, S. A. (2012). Organic synthesis via irradiation and warming of ice grains in the solar nebula. *Science*, 336(6080), 452–454. <https://doi.org/10.1126/science.1217290>

<sup>144</sup> Bergner, J. B., et al. (2023). JWST detection of CH<sub>3</sub>CHO and CH<sub>3</sub>CH<sub>2</sub>OH in the inner disk of a solar-type protostar. *Nature*, 615, 579–583. <https://doi.org/10.1038/s41586-023-05726-3>

<sup>145</sup> Cooper, G., et al. (2001). Carbonaceous meteorites as a source of sugar-related organic compounds for the early Earth. *Nature*, 414, 879–883. <https://doi.org/10.1038/414879a>

<sup>146</sup> Naraoka, H., et al. (2023). Soluble organic molecules in samples of the carbonaceous asteroid Ryugu. *Science*, 379(6637), eabn8671. <https://doi.org/10.1126/science.abn8671>

comet-like bodies<sup>147</sup>, hot protostellar discs, or the comet nucleus–nebula transition zone<sup>148</sup> (such as ESA's Comet Interceptor) will reveal their primordial organic chemistry<sup>149</sup>.

These interdisciplinary studies will help answer a grander question: "Is Earth-like life the norm in the cosmos?"<sup>150</sup>. Suppose the chemical mechanisms within protoplanetary discs are indeed universally capable of constructing life. In that case, the initial raw materials and pathways required for Earth-like life may not be unique to our planet, but rather universal products of natural evolutionary processes within cosmic nebulae<sup>151</sup>. It not only prompts us to reconsider the 'geocentric view', but also provides theoretical guidance and prioritization for seeking 'biosignatures' beyond our solar system.

## **(ii) The origin of food or initial organic molecules within protoplanetary discs and prebiotic chemical evolution**

The question of life's origin remains one of the most fundamental and complex issues in the natural sciences. Modern research increasingly indicates that life did not emerge as an isolated, post-Earth-formation accident, but may instead be deeply rooted in earlier cosmic chemical evolution. Astrochemistry, as the core discipline studying the evolution of atoms and molecules in the cosmos, offers new perspectives on understanding the origins of life-forming substances. Against this backdrop, the protoplanetary disc—the circumstellar gas and dust disc surrounding stars during their formation—emerges as a crucial transitional stage linking interstellar matter to the chemical potential for life.

The formation of protoplanetary discs begins with nebular collapse, resulting in a flattened structure that conserves angular momentum and undergoes multiple evolutionary stages, from dust aggregation and minor body formation to protoplanetary collisions. Throughout this process, the combined effects of stellar radiation, high-energy particle streams, the decay of short-lived isotopes, and internal heat sources foster highly active chemical reactions within the disc. This environment not only contains the fundamental elements for

---

<sup>147</sup> Rubin, M., et al. (2015). Molecular nitrogen in comet 67P/Churyumov–Gerasimenko indicates a low formation temperature. *Science*, 348(6231), 232–235. <https://doi.org/10.1126/science.aaa6100>

<sup>148</sup> Snodgrass, C., et al. (2017). The science goals of the Comet Interceptor mission. *Philosophical Transactions of the Royal Society A*, 375(2097), 20160249. <https://doi.org/10.1098/rsta.2016.0249>

<sup>149</sup> Elsila, J. E., Glavin, D. P., & Dworkin, J. P. (2009). Cometary glycine detected in samples returned by Stardust. *Meteoritics & Planetary Science*, 44(9), 1323–1330. <https://doi.org/10.1111/j.1945-5100.2009.tb01224.x>

<sup>150</sup> Lineweaver, C. H. (2001). An estimate of the age distribution of terrestrial planets in the Universe: Quantifying metallicity as a selection effect. *Icarus*, 151, 307–313. <https://doi.org/10.1006/icar.2001.6607>

<sup>151</sup> Chyba, C. F., & Hand, K. P. (2005). Astrobiology: The study of the living universe. *Annual Review of Astronomy and Astrophysics*, 43, 31–74. <https://doi.org/10.1146/annurev.astro.43.072103.150618>

life—CHNOPS—but also enriches various metallic elements, particularly transition metals, potentially providing potent catalytic support for primordial organic synthesis reactions. Observations and meteorite sample studies indicate that these discs are not only rich in organic molecules (such as formaldehyde, HCN, polycyclic aromatic hydrocarbons PAHs, etc.), but also contain substantial quantities of transition metals (e.g., Fe, Ni, Co) and short-half-life radioactive isotopes (e.g.,  $^{26}\text{Al}$ ,  $^{60}\text{Fe}$ ), and long-lived radioactive isotopes (such as potassium-40 ( $^{40}\text{K}$ ), uranium-238 ( $^{238}\text{U}$ ), uranium-235 ( $^{235}\text{U}$ ), and thorium-232 ( $^{232}\text{Th}$ )). These elements provided catalysis and energy sources for diverse chemical reactions.

However, the precise role of metals—particularly transition metals—in prebiotic chemical reactions remains poorly understood. Can their multivalent nature, coordination behaviour, and electronic structure facilitate key molecular synthesis, energy transfer, and structural selectivity within protoplanetary discs? These questions not only inform our understanding of life's origins but also influence speculations regarding the potential for life on Earth-like planets.

This study aims to systematically investigate the distribution, chemical properties, and catalytic potential of transition metals within protoplanetary discs. By analyzing their potential core role in prebiotic chemical reactions, we explore the origins of initial organic molecules and the evolution of prebiotic chemistry. Integrating meteorite chemical analyses, simulation data, and evolutionary clues from modern biological metal enzymes, we attempt to delineate a continuous pathway from protoplanetary disc metal chemistry to the origin of life within these discs.

## **1. Cosmic Origins of Life's Fundamental Elements and the Chemical Habitability of Protoplanetary Discs**

The fundamental building blocks of life comprise carbon (C), hydrogen (H), nitrogen (N), oxygen (O), phosphorus (P), and sulphur (S), collectively termed CHNOPS. These elements are extensively present within protoplanetary discs in gaseous, icy, compound, and dust forms<sup>152</sup>. Moreover, transition metals (TM), an indispensable class of elements in life systems, are also abundant in protoplanetary discs, particularly within iron-rich dust and supernova-injected regions. These metals—including iron (Fe), nickel (Ni), and cobalt (Co)—

---

<sup>152</sup> Ehrenfreund, P., & Charnley, S. B. (2000). Organic molecules in the interstellar medium, comets, and meteorites: A voyage from dark clouds to the early Earth. *Annual Review of Astronomy and Astrophysics*, 38(1), 427–483. <https://doi.org/10.1146/annurev.astro.38.1.427>

are widely present in meteorites, interstellar dust, and protoplanetary bodies during the early stages of disk formation<sup>153</sup>.

The ubiquity and critical role of transition metals in biological systems are well-established: approximately 30% of proteins contain metal-binding structures, and around 40% of enzyme-catalyzed reactions depend on metalloproteins<sup>154</sup>. Within organisms, these metals typically form metal-organic complexes, performing core functions in catalysis, electron transfer, and redox processes, as well as maintaining structural stability<sup>155</sup>. Notably, transition metals from the first period of the periodic table are found in nearly all phyla of life, indicating their widespread utilization during early evolutionary development<sup>156</sup>.

More significantly, transition metals possess unique electronic structural flexibility, as they exhibit multiple oxidation states, spin states, and variable ligand geometries, and can undergo reversible electron transfer with  $\sigma/\pi$  donor or acceptor ligands, thereby participating in complex chemical reactions<sup>157</sup>. This chemical versatility renders them indispensable catalytic agents in prebiotic chemistry, particularly suited to catalyze initial metabolic networks and nucleic acid precursor synthesis reactions under non-enzymatic conditions<sup>158</sup>.

Within protoplanetary discs, the CHNOPS elements, transition metals, and alkali metals (such as sodium and potassium) collectively constitute the elemental environment essential for life. Unlike later terrestrial environments, these elements exist in more primordial and reactive forms within protoplanetary discs (e.g., ionic states or bound to highly reactive dust particle surfaces), providing a natural platform for non-equilibrium chemical reactions. Thus, protoplanetary discs serve not only as material sources for prebiotic chemical systems but also as the "chemical cradle" where the earliest life-related

---

<sup>153</sup> Righter, K., & Drake, M. J. (1997). Metal/silicate equilibrium in a deep magma ocean: Implications for the origin of the mantle siderophile element depletions. *Earth and Planetary Science Letters*, 146(3-4), 541–553. [https://doi.org/10.1016/S0012-821X\(96\)00269-2](https://doi.org/10.1016/S0012-821X(96)00269-2)

<sup>154</sup> Waldron, K. J., Rutherford, J. C., Ford, D., & Robinson, N. J. (2009). Metalloproteins and metal sensing. *Nature*, 460(7257), 823–830. <https://doi.org/10.1038/nature08300>

<sup>155</sup> Holm, R. H., Kennepohl, P., & Solomon, E. I. (1996). Structural and Functional Aspects of Metal Sites in Biology. *Chemical Reviews*, 96(7), 2239–2314. <https://doi.org/10.1021/cr9500390>

<sup>156</sup> Dupont, C. L., Butcher, A., Valas, R. E., Bourne, P. E., & Caetano-Anollés, G. (2010). History of biological metal utilisation inferred through phylogenomic analysis of protein structures. *Proceedings of the National Academy of Sciences*, 107(23), 10567–10572. <https://doi.org/10.1073/pnas.0912491107>

<sup>157</sup> Barge, L. M., et al. (2015). From chemical gardens to chemobionics: The role of transition metals in early proto-metabolism. *Origins of Life and Evolution of Biospheres*, 45(5), 343–361.

<sup>158</sup> Sasselov, D. D., Grotzinger, J. P., & Sutherland, J. D. (2020). The origin of life as a planetary phenomenon. *Science Advances*, 6(6), eaax3419. <https://doi.org/10.1126/sciadv.aax3419>

chemical evolution likely occurred, constructing the molecular frameworks and reaction pathways essential for life<sup>159</sup>.

Consequently, the origin of life was likely not an isolated Earth-specific process, but rather a continuation of the chemical evolution already underway within the protoplanetary disc. This perspective not only supports an understanding of the universality of life's origins but also establishes a cosmic framework for exploring exoplanetary life by providing the material and reaction foundations.

## 2. Cosmic Origins of Metallic Elements in Protoplanetary Discs

The chemical composition of protoplanetary discs is not formed in isolation but inherits the "legacy" of stellar nucleosynthesis from millions of years before the Sun's formation. The Sun formed within a heavy-element-rich molecular cloud in the Milky Way, which gradually accumulated the elements essential for life and planetary bodies through successive generations of stellar explosions, wind erosion, and mergers. Current astrophysical and cosmochemical research unanimously indicates that supernova explosions, AGB stellar winds, and binary neutron star mergers are the primary sources of heavy elements in protoplanetary discs, particularly transition metals and rare radioactive nuclides<sup>160</sup>.

### 2.1 Type II Supernovae

Massive stars undergo violent supernova explosions at the end of their evolution, releasing the diverse metallic elements formed through nuclear fusion within their cores into the surrounding space<sup>161</sup>. These explosions constitute the primary source of iron-peak elements (Fe-peak elements), including iron (Fe), nickel (Ni), and cobalt (Co). Such explosions also produce short-lived radioactive isotopes (such as <sup>26</sup>Al and <sup>60</sup>Fe), providing a crucial driving force for thermal energy generation within protoplanetary discs<sup>162</sup>.

### 2.2 Asymptotic Giant Branch (AGB) Stellar Winds: Carbon, Nitrogen, and s-Process Metal Injection

---

<sup>159</sup> Cobb, A. K., & Pudritz, R. E. (2014). Nature's Starships. I. Observed Abundances and Relative Frequencies of Amino Acids in Meteorites. *The Astrophysical Journal*, 783(2), 140. <https://doi.org/10.1088/0004-637X/783/2/140>

<sup>160</sup> Alexander, C. M. O'D., et al. (2012). The Provenances of Asteroids, and Their Contributions to the Volatile Inventories of the Terrestrial Planets. *Science*, 337(6095), 721–723. <https://doi.org/10.1126/science.1223474>

<sup>161</sup> Woosley, S. E., & Weaver, T. A. (1995). The Evolution and Explosion of Massive Stars. II. Explosive Hydrodynamics and Nucleosynthesis. *The Astrophysical Journal Supplement Series*, 101, 181–235. <https://doi.org/10.1086/192237>

<sup>162</sup> Huss, G. R., Meyer, B. S., Srinivasan, G., Goswami, J. N., & Sahijpal, S. (2009). Stellar sources of the short-lived radionuclides in the early Solar System. *Geochimica et Cosmochimica Acta*, 73(17), 4922–4945. <https://doi.org/10.1016/j.gca.2009.05.063>

During the late stages of their evolution, intermediate-mass stars enter the Asymptotic Giant Branch (AGB) phase, undergoing significant mass loss. Their stellar winds enrich interstellar space with carbon, nitrogen, oxygen, and s-process (slow neutron capture) elements, such as barium (Ba), strontium (Sr), zirconium (Zr), and lanthanum (La)<sup>163</sup>. These particles condense upon cooling to form carbon-rich or oxide dust, serving as a crucial source of organic chemistry within protoplanetary discs and constituting a primary origin for building blocks of organic compounds in protoplanetary discs and carbonaceous chondrites.

### **2.3 Double neutron star mergers and r-process element injection**

Although rare, binary neutron star mergers are a significant source of r-process (rapid neutron capture) elements, producing heavy metals such as platinum (Pt), gold (Au), and rare-earth elements like neodymium (Nd) and gadolinium (Gd). Although their injection rates are far lower than those from supernova explosions and AGB stellar winds, the metals produced are diverse and possess special functionalities in biochemistry, such as forming the metallic active centres of specific enzymes or serving as electron pathways in catalytic reactions<sup>164</sup>.

These nucleosynthetic products accumulated within molecular clouds in diverse forms (atomic, ionic, oxides, sulphides, carbides, etc.) via interstellar dust or gas molecules, subsequently being drawn into the protoplanetary disc during the nebular collapse that formed the Sun<sup>165</sup>. Thermal evolution within the protoplanetary disc and planetary microbody collisions further promoted localized aggregation of these metals or their involvement in catalytic reactions, providing metallic cores for early chemical reaction systems.

As these metallic elements were partially segregated into the Earth's core or redistributed during planetary formation, the metallic abundance in the protoplanetary disc's early stages far exceeded the levels available in the Earth's present-day surface layers. Consequently, numerous critical metal-catalyzed reactions—such as Fischer-Tropsch synthesis, nitrogen fixation, and reductive amination—were likely pre-initiated within the protoplanetary disc,

---

<sup>163</sup> Lugaro, M., Karakas, A. I., Stancliffe, R. J., & Rijs, C. (2012). The s-process in asymptotic giant branch stars. *The Astrophysical Journal*, 747(1), 2. <https://doi.org/10.1088/0004-637X/747/1/2>

<sup>164</sup> Thielemann, F.-K., Eichler, M., Panov, I. V., & Wehmeyer, B. (2017). Neutron star mergers and nucleosynthesis of heavy elements. *Annual Review of Nuclear and Particle Science*, 67, 253–274. <https://doi.org/10.1146/annurev-nucl-101916-123246>

<sup>165</sup> Lodders, K. (2003). Solar System Abundances and Condensation Temperatures of the Elements. *The Astrophysical Journal*, 591(2), 1220–1247. <https://doi.org/10.1086/375492>



establishing the material and reaction foundations for subsequent biogeochemical evolution.

## 2.4 Fe/Co/Ni Metal Coordination Compounds Facilitate Prebiotic Reactions

Transition metals can form stable coordination compounds with simple small molecules (e.g., CO, NO, HCN, N<sub>2</sub>, H<sub>2</sub>), serving functions including: Bond activation: Metal centres can "soften" molecular  $\pi$  or  $\sigma$  bonds, facilitating their cleavage or rearrangement; Electron transfer bridges: Acting as Lewis acids or reducing agents to transfer electrons between different molecules; Selective regulation: Metal coordination can direct molecular arrangement, enhancing reaction pathway selectivity. For example:

- Fe<sup>2++</sup> /Fe<sup>3+</sup> : Catalyses CO + H<sub>2</sub>  $\rightarrow$  (hydrocarbons; Fischer–Tropsch reaction);
- Ni<sup>2++</sup> activates H<sub>2</sub> with C $\equiv$  N, promoting hydrogen cyanide reduction to methylamine or aminonitrile;
- Co<sup>3+</sup> forms Co–NO complexes with NO, mimicking the function of modern nitric oxide synthases.

Simulation experiments indicate that such metal-small molecule complexes can form on mineral surfaces (e.g., olivine, ferrite) at temperatures ranging from 200 to 400 K, exhibiting higher activity than in metal-free environments (Fioroni & DeYonker, 2016)<sup>166</sup>.

## 2.5 Metal–organic complexes as prototypes of metalloproteins

In modern organisms, the active sites of numerous core enzyme systems contain metals such as Fe, Ni, or Co, exhibiting evolutionary continuity with primordial complexes. Examples include: iron-sulphur clusters (Fe–S clusters): composed of structures like [Fe<sub>4</sub>S<sub>4</sub>]<sup>2++</sup>, extensively involved in electron transport chains (e.g., ferredoxin)<sup>167</sup>; Hydrogenases ([FeFe] or [NiFe] hydrogenases): catalyse the reversible conversion between protons and hydrogen gas; Nitrogenases: contain complex FeMo or FeV clusters, catalysing the N<sub>2</sub>  $\rightarrow$  NH<sub>3</sub> reaction<sup>168</sup>; Cobalamin (Vitamin B<sub>12</sub>): centred on Co<sup>3++</sup>, participates in methyl transfer and homologous recombination.

---

<sup>166</sup> Fioroni, M., & DeYonker, N. J. (2016). Iron and ruthenium centres grafted onto silica surfaces as models for heterogeneous Fischer–Tropsch and Haber–Bosch catalysis. *Journal of Physical Chemistry C*, 120(50), 28744–28752. <https://doi.org/10.1021/acs.jpcc.6b10696>

<sup>167</sup> Beinert, H., Holm, R. H., & Münck, E. (1997). Iron–sulfur clusters: Nature's modular, multipurpose structures. *Science*, 277(5326), 653–659. <https://doi.org/10.1126/science.277.5326.653>

<sup>168</sup> McCollom, T. M., & Seewald, J. S. (2006). Carbon isotope composition of organic compounds produced by abiotic synthesis under hydrothermal conditions. *Earth and Planetary Science Letters*, 243, 74–84. <https://doi.org/10.1016/j.epsl.2005.12.027>

These structures demonstrate that life systems did not invent new chemistry, but rather directly 'inherited' pre-existing metal complex forms from nature. Upon this foundation, functionalized macromolecular platforms (such as proteins and RNA) were constructed to encapsulate and regulate these complexes.

## 2.6 From Nebula to Life: The Continuity of Metal Complexes

Metal–small molecule or metal–PAH complexes formed within protoplanetary discs not only provided primordial electron transfer and bond activation mechanisms but may also have been captured by vesicles, membrane-like structures, or RNA fragments, evolving into "primordial catalytic nuclei". For instance, studies indicate that: Fe–S clusters may self-assemble directly from Fe<sup>2++</sup> in the presence of H<sub>2</sub>S<sup>169</sup>; Ni–CO complexes can induce C–C bond formation, providing intermediates for carbohydrate or fatty acid synthesis<sup>170</sup>; and the Co–CN–NH<sub>2</sub> system may mimic the primordial reaction environment of vitamin B<sub>12</sub><sup>171</sup>.

Hydrogenases ([FeFe] or [NiFe]): catalyze the reversible conversion between protons and hydrogen gas<sup>172</sup>;

These discoveries deepen our understanding of the "metal-first" hypothesis, which posits that metal coordination chemistry may have undertaken catalytic and energy management functions before the emergence of organic life.

Transition metals Fe, Ni, and Co were enriched in protoplanetary discs through injection from AGB stars and supernovae, providing crucial catalytic nuclei for prebiotic reactions. The metal-organic complexes they formed not only activated small molecules and catalyzed electron transfer but also served as structural and functional prototypes for modern metalloenzymes. This continuity suggests that the core catalytic capabilities of life may have been nascent during the nebular phase preceding planetary formation.

## 3. Evidence of metals in meteorites: Recorded remnants of protoplanetary disc nucleosynthesis

Modern meteorites, particularly chondrites, are regarded as direct repositories of early protoplanetary disc residues, preserving chemical compositions and material evolution information from the solar system's infancy. They furnish crucial empirical evidence

---

<sup>169</sup> Beinert, H., Holm, R. H., & Münck, E. (1997). Iron–sulfur clusters: Nature's modular, multipurpose structures. *Science*, 277(5326), 653–659. <https://doi.org/10.1126/science.277.5326.653>

<sup>170</sup> Ragsdale, S. W., & Kumar, M. (1996). Nickel-containing carbon monoxide dehydrogenase/acetyl-CoA synthase. *Chemical Reviews*, 96(7), 2515–2540. <https://doi.org/10.1021/cr950055h>

<sup>171</sup> Rees, D. C., et al. (2005). Structural basis of biological nitrogen fixation. *Philosophical Transactions of the Royal Society B*, 360(1458), 1221–1234. <https://doi.org/10.1098/rstb.2005.1658>

<sup>172</sup> Lubitz, W., Ogata, H., Rüdiger, O., & Reijerse, E. (2014). Hydrogenases. *Chemical Reviews*, 114(8), 4081–4148. <https://doi.org/10.1021/cr4005814>

supporting the hypothesis that metallic elements originated from stellar nucleosynthesis and were subsequently injected into the protoplanetary disc.

### 3.1 Composition of Meteorites Rich in Transition Metals

Fe-Ni alloy particles (e.g., meteorites) are widely present in chondritic meteorites, alongside mineral phases rich in transition metals such as Cr, Mn, Co, Mo, and V, including magnetite, olivine, pyroxene, and sulphides<sup>173</sup>. These metals are distributed within the chondritic structure, representing the state of metallic condensation and aggregation within the early solar nebula. Notably, calcium-aluminum-rich inclusions (CAIs) within the Allende meteorite and evidence of radiometallic enrichment within chondrules indicate prior exposure to a high-energy environment containing short-lived nuclides such as <sup>26</sup>Al and <sup>60</sup>Fe<sup>174</sup>.

### 3.2 Anomalous isotopes reveal supernova injection events

Non-mass-fractionated isotopic anomalies in elements such as Cr, Ti, and Ni within meteorites indicate these metals originate from nucleosynthesis products of distinct stellar events. For instance, Trinquier et al. identified <sup>54</sup>Cr enrichment anomalies in multiple chondrite meteorites, consistent with dust particles formed after a supernova explosion, which confirms the direct injection of supernova material into the solar nebula<sup>175</sup>.

### 3.3 Traces of short-lived radionuclides (<sup>60</sup>Fe, <sup>26</sup>Al)

<sup>60</sup>Fe is a typical supernova nucleosynthesis product, and its initial abundance in iron-nickel alloy minerals has been confirmed in multiple primitive meteorites. Research by Tang and Dauphas indicates that the distribution of these metals is linked to the early stratification of the solar nebula<sup>176</sup>, demonstrating their injection and homogeneous mixing into the nebular gas prior to planetary body formation<sup>177</sup>.

### 3.4 Meteorite Particles of AGB Star Products

---

<sup>173</sup> Trinquier, A., Birck, J. L., & Allègre, C. J. (2007). Widespread <sup>54</sup>Cr Heterogeneity in the Inner Solar System. *The Astrophysical Journal*, 655(2), 1179–1185. <https://doi.org/10.1086/510361>

<sup>174</sup> Tang, H., & Dauphas, N. (2012). Abundance, distribution, and origin of <sup>60</sup>Fe in the solar protoplanetary disk. *Earth and Planetary Science Letters*, 359–360, 248–263. <https://doi.org/10.1016/j.epsl.2012.10.011>

<sup>175</sup> Trinquier, A., Birck, J. L., & Allègre, C. J. (2007). Widespread <sup>54</sup>Cr Heterogeneity in the Inner Solar System. *The Astrophysical Journal*, 655(2), 1179–1185. <https://doi.org/10.1086/510361>

<sup>176</sup> Tang, H., & Dauphas, N. (2012). Abundance, distribution, and origin of <sup>60</sup>Fe in the solar protoplanetary disk. *Earth and Planetary Science Letters*, 359–360, 248–263. <https://doi.org/10.1016/j.epsl.2012.10.011>

<sup>177</sup> Dauphas, N., Cook, D. L., Sacarabany, A., Fröhlich, C., Davis, A. M., Wadhwa, M., Pourmand, A., Rauscher, T., & Gallino, R. (2008). <sup>60</sup>Fe evidence for early injection and efficient mixing of stellar debris in the protosolar nebula. (preprint).

Studies of presolar grains within meteorites reveal that carbon-rich particles such as silicon carbide and alumina bear isotopic signatures characteristic of s-process elements from AGB stars (e.g., enrichment in Ba, Sr, Zr, La). It indicates their origin as dust injected by AGB stellar winds, with these particles persisting within the nebula until meteorite formation<sup>178</sup>.

### 3.5 r-process nuclides and rare earth element anomalies

Certain meteorites exhibit anomalous ratios of heavy rare earth elements (Nd, Sm, U) formed via the r-process. These signatures suggest possible origins in binary neutron star mergers or specific types of supernovae. Although these heavy metals are not abundant in the solar system, they have profoundly influenced early chemical evolution<sup>179</sup>.

### 3.6 Mineral Capsules of the Protoplanetary Disk

Meteorites, acting as "mineral capsules of the protoplanetary disc", preserve the origins, states, and distributions of metallic elements from the solar system's formative era. The diverse metallic phases and isotopic anomalies they exhibit provide compelling support for the following perspectives: Metals within the protoplanetary disc primarily originate from nucleosynthesis products of multiple stellar generations; supernovae and AGB stars constitute the primary injection sources of Fe, Ni, Co, and s-process metals into protoplanetary discs; these metals participated in prebiotic chemistry in atomic, molecular, or dust forms. Consequently, meteorite evidence forms a crucial experimental link in the chain connecting stellar nucleosynthesis to prebiotic chemistry, providing vital geological and astrochemical support for the notion that "life originated from interstellar heritage."

### 3.7 Evidence for the Coexistence of Fullerenes and Metal Complexes

More remarkably, fullerene molecules such as C<sub>60</sub> and C<sub>70</sub>, along with their inclusion structures containing rare gases, have been discovered within meteorites<sup>180</sup>. Some studies have also detected indirect signals of metal-encapsulated fullerenes (e.g., Fe@C<sub>60</sub>), indicating that during the condensation process of protoplanetary discs and nebulae, carbon structures, including PAHs and metals, can form stable complexes synergistically<sup>181</sup>. These structures may have served as platforms for primordial molecular adsorption,

---

<sup>178</sup> Lugaro, M., Karakas, A. I., Stancliffe, R. J., & Rijs, C. (2012). The s-process in asymptotic giant branch stars. *The Astrophysical Journal*, 747(1), 2. <https://doi.org/10.1088/0004-637X/747/1/2>

<sup>179</sup> Côté, B., Eichler, M., Arcones, A., et al. (2018). Neutron Star Mergers Might Not Be the Only Source of r-process Elements in the Milky Way. *The Astrophysical Journal*, 855(2), 99. <https://doi.org/10.3847/1538-4357/aaad67>

<sup>180</sup> Becker, L., Poreda, R. J., & Bunch, T. E. (2000). Fullerenes: An extraterrestrial carbon carrier phase for noble gases. *PNAS*, 97(7), 2979–2983.

<sup>181</sup> Campbell, E. K., et al. (2015). Laboratory confirmation of C<sub>60</sub><sup>+</sup> as the carrier of two diffuse interstellar bands. *Nature*, 523, 322–323.

catalysis, and electronic regulation, acting as "natural test tubes" for prebiotic chemical reactions.

#### **4. The "Two-Stage Chemical Legacy" in the Murchison Meteorite: An Evolutionary Trajectory from Stellar Discs to Protoplanetary Discs**

The prevailing scientific perspective on life's origins posits exogenous input of life-bearing material (meteorites) alongside endogenous emergence (Earth). However, mounting evidence suggests that life arose through a dual process involving exogenous input (the Interstellar Medium, ISM) and endogenous emergence (from a protoplanetary disc). The following analysis actively supports such a dual-process model.

##### **4.1 Phase One: Interstellar Precursors Originating from a 7-Billion-Year-Old Stellar Disc**

Certain organic compounds within the Murchison meteorite can be traced back to the formation of protostellar systems or star-forming regions approximately 7 billion years ago. This phase may involve the late evolutionary stages of protoplanetary discs around medium-mass stars (e.g., F or G types) or higher-mass stars (e.g., B types). It may also encompass carbon-rich environments following binary star systems or collective stellar storm activity. Relevant interstellar precursor substances include ammonia (NH<sub>3</sub>), methylamine (CH<sub>3</sub>NH<sub>2</sub>), hydrogen cyanide (HCN), methanol (CH<sub>3</sub>OH), and precursor structures of polycyclic aromatic hydrocarbons (PAHs)<sup>182</sup>. These molecules are primarily generated through gas-phase quantum chemical reactions and radical reaction chains induced by high-energy ultraviolet radiation and cosmic rays<sup>183</sup>. They can remain cryogenically preserved within interstellar dust and ice particles for hundreds of millions of years, until they are incorporated into the next round of nebular collapse, serving as the chemical seeds for solar system formation.

##### **4.2 Second Stage: Hydrothermal Synthesis within the Solar System's Protoplanetary Disk 4.6 Billion Years Ago**

With the formation of the solar system, these interstellar precursor molecules entered the dust- and ice-rich protoplanetary disc<sup>184</sup>. Within parent asteroids or carbonaceous planetesimals, heat sources provided by short-lived radioactive isotopes (such as <sup>26</sup>Al) fostered stable hydrated environments. Complex organic synthesis reactions occurred at the water-rock-organic triphase interface. For instance, Strecker reactions can synthesize

---

<sup>182</sup> Herbst, E. & van Dishoeck, E. F. (2009). Complex Organic Interstellar Molecules. *Annu. Rev. Astron. Astrophys.*, 47, 427–480.

<sup>183</sup> Garrod, R. T., & Herbst, E. (2006). Formation of methyl formate and other organic species in the warm-up phase of hot molecular cores. *Astronomy & Astrophysics*, 457(3), 927–936.

<sup>184</sup> Oba, Y., et al. (2022). Extraterrestrial hexamethylenetetramine in meteorites—a precursor of prebiotic chemistry in the interstellar medium. *Nature Communications*, 13, 2004.

glycine and alanine in the presence of  $\text{NH}_3$ , HCN, and aldehyde/ketone molecules, while polymerized HCN chains can generate nucleobases (e.g., adenine, uracil) under suitable temperature and pH conditions<sup>185</sup>. Furthermore, fatty acid-like substances may form through the catalytic hydrolysis and polymerization of small organic molecules such as CO and formic acid, creating membrane-like precursor structures that could have participated in the initial construction of primordial cells.

### **4.3 Murchison Meteorite: A Chemical Bridge Between Two Stellar Systems**

The Murchison meteorite stands as one of the most valuable carbonaceous chondrite specimens for research. Its uniqueness lies not only in its rich content of amino acids, nucleobases, PAHs, and isotopically anomalous mineral grains, but also in its preservation of chemical trajectories spanning two distinct cosmological epochs<sup>186</sup>. Interstellar precursor molecules deposited by early protostellar discs underwent hydrothermal transformation within the solar system's protoplanetary disc, culminating in complex synthesis and differentiation within carbonaceous chondrules. This two-stage model – "interstellar input followed by local processing" – clearly demonstrates the formation and accumulation of life precursor molecules at the galactic level, supporting the notion that "prebiotic substances predate Earth's origin".

## **5. The Protoplanetary Disk and Interstellar Ice Reaction Mechanisms: From Frozen Molecules to Hydrothermal Chemical Factories**

The protoplanetary disc is the gaseous and dusty envelope surrounding a newly formed star, exhibiting rapid variations in temperature, density, and radiation conditions from its centre outward, thereby forming distinct physico-chemical zones. Interstellar ice—primarily composed of ice-coated dust particles containing water, CO,  $\text{CH}_3\text{OH}$ ,  $\text{NH}_3$ , HCN,  $\text{CH}_3\text{NH}_2$ , and other compounds—formed abundantly within molecular clouds before solar system formation, serving as crucial organic seed material entering the protoplanetary disc<sup>187</sup>. These two stages are not isolated but form a continuous chemical evolutionary chain, with key connections manifested in the following aspects:

### **5.1 Interstellar ice as molecular precursors for organic reactions within the protoplanetary disc**

---

<sup>185</sup> Oberg, K. I. (2016). Photochemistry and Astrochemistry: Photochemical Pathways to Interstellar Complex Organic Molecules. *Chemical Reviews*, 116(17), 9631–9663.

<sup>186</sup> Cronin, J. R., & Pizzarello, S. (1997). Enantiomeric excesses in meteoritic amino acids. *Science*, 275(5302), 951–955.

<sup>187</sup> Boogert, A. C. A., Gerakines, P. A., & Whittet, D. C. B. (2015). Observations of the icy universe. *Annual Review of Astronomy and Astrophysics*, 53, 541–581.

Before the formation of the Solar System, dust particles within the interstellar medium were already coated with numerous condensed molecules (such as H<sub>2</sub>O, CO<sub>2</sub>, CH<sub>3</sub>OH, HCN, and CH<sub>3</sub>NH<sub>2</sub>), collectively forming "interstellar ice"<sup>188</sup>. Research indicates that these interstellar icy bodies were partially preserved following the collapse of the solar nebula. Embedded within the dust particles that formed the protoplanetary disc, they became a crucial source of the chemical composition of parent asteroids and comets<sup>189</sup>.

When these interstellar ices enter warm disc regions or are subjected to impact heating or ultraviolet irradiation, their ice layers sublime and melt, releasing highly reactive intermediate molecules such as hydroxyl ( $\cdot$ OH), methyl ( $\cdot$ CH<sub>3</sub>), and cyano ( $\cdot$ CN) groups. These intermediates, entering liquid water or thermally mineralized environments, can trigger a new round of hydrothermal reactions, thereby forming higher-order organic compounds such as amino acids, nucleobases, and fatty acids<sup>190</sup>.

## 5.2 Pyrolytic encapsulation within planetesimals: Activation platforms for interstellar ice

In the early stages of formation, miniature celestial bodies within protoplanetary discs (planetesimals, parent asteroids) generate internal heat through the decay of short-lived radionuclides (e.g., <sup>26</sup>Al, <sup>40</sup>K), raising temperatures to several hundred degrees Celsius<sup>191</sup>. This mild yet persistent heat source provides an ideal environment for the potential synthesis of complex molecules within interstellar ice<sup>192</sup>: alcohols, aldehydes, and ketones in interstellar ice can undergo condensation or reduction to form carbon chains in the presence of hot water; HCN can polymerise into HCN chains or melamine, providing structural precursors for nucleoside bases; CO, HCOOH, and formate molecules can catalyse the formation of fatty acids and their esters on mineral surfaces.

Thus, the protoplanetary disc not only inherits the "frozen organic legacy" of interstellar ice but also further processes it through pyrolysis, hydration, and mineral catalysis into more complex molecular forms<sup>193</sup>.

---

<sup>188</sup> Watanabe, N., & Kouchi, A. (2008). Ice surface reactions: A key to chemical evolution in space. *Progress in Surface Science*, 83(10–12), 439–489.

<sup>189</sup> Walsh, C., Nomura, H., & van Dishoeck, E. F. (2015). Chemical processes in protoplanetary disks. *Astronomy & Astrophysics Review*, 23(1), 3.

<sup>190</sup> Öberg, K. I., Garrod, R. T., van Dishoeck, E. F., & Linnartz, H. (2009). Formation rates of complex organics in UV-irradiated CH<sub>3</sub>OH-rich ices I: Experiments. *Astronomy & Astrophysics*, 504(3), 891–913.

<sup>191</sup> Ciesla, F. J., & Sandford, S. A. (2012). Organic synthesis via irradiation and warming of ice grains in the solar nebula. *Science*, 336(6080), 452–454.

<sup>192</sup> Garrod, R. T. (2013). A three-phase chemical model of hot cores: The formation of glycine. *The Astrophysical Journal*, 765(1), 60.

<sup>193</sup> Visser, R., van Dishoeck, E. F., Doty, S. D., & Dullemond, C. P. (2009). The chemical history of molecules in circumstellar disks. *Astronomy & Astrophysics*, 495(3), 881–897.

### 5.3 Reaction Coupling in Zonal Environments: Chemical Bridging from Icy Shells to Rocky Cores

The physical structure of the protoplanetary disc dictates distinct pathways for organic evolution across different zones:<sup>194</sup>

Outer Disk (>10 AU): Cold, ice-rich. Preserves interstellar ice structure, readily retained in comets

Inner Disk (1–10 AU): Ultraviolet, X-ray, warm Ice sublimation plus Ultraviolet dissociation, initiating radical chain reactions.

Inner Disk (<1 AU): High temperature plus High pressure plus Mineral-rich Hydrothermal reactions, mineral-catalyzed synthesis of complex organics

This zonation structure effectively bridges the low-temperature photochemical mechanisms of the interstellar icy phase with the thermal hydrochemical mechanisms within the protoplanetary disc, enabling a stepwise evolution from interstellar organic seeds to prebiotic molecules<sup>195</sup>.

### 5.4 Murchison meteorite: physical evidence for this bridging mechanism

Numerous molecular species within the Murchison meteorite exhibit high consistency with interstellar ice simulation products (e.g., glycine, alanine, formic acid, melamine)<sup>196</sup>. Furthermore, anomalies in its stable isotope ratios (D/H, <sup>15</sup>N/<sup>14</sup>N) indicate that portions of its composition inherited the "frozen" isotopic characteristics of the interstellar environment. Moreover, the enantiomeric ratios of specific amino acids suggest that their synthesis occurred within non-equilibrium hydrothermal systems, further supporting the continuous model of "interstellar ice activation and followed by protoplanetary disc processing"<sup>197</sup>.

Interstellar ice provides the raw material reservoir and intermediates for organic formation; the protoplanetary disc supplies the energy and mineral environment required to drive these reactions. Together, they constitute the upstream and downstream stages in the galactic biochemistry chain, jointly shaping the complex organic systems recorded in carbonaceous

---

<sup>194</sup> Ciesla, F. J., & Charnley, S. B. (2006). The physics and chemistry of nebular evolution. *Meteorites and the Early Solar System II*, 209–230.

<sup>195</sup> Henning, T., & Semenov, D. (2013). Chemistry in protoplanetary disks. *Chemical Reviews*, 113(12), 9016–9042.

<sup>196</sup> Sephton, M. A. (2002). Organic compounds in carbonaceous meteorites. *Natural Product Reports*, 19(3), 292–311.

<sup>197</sup> Alexander, C. M. O'D., Bowden, R., Fogel, M. L., Howard, K. T., Herd, C. D. K., & Nittler, L. R. (2012). The provenances of asteroids, and their contributions to the volatile inventories of the terrestrial planets. *Science*, 337(6095), 721–723.



meteorites, such as the Murchison meteorite<sup>198</sup>. This connection not only reveals the history of extraterrestrial organic formation but also provides crucial evidence for chemical continuity between the interstellar medium, protoplanetary discs, and Earth.

## 6 Prebiotic Chemistry of Transition Metals

Transition metals, due to their unique electronic structures and chemical diversity, are considered to have played a significant role in the molecular evolution that preceded the origin of life. Within high-energy environments, such as protoplanetary discs, they not only form the basis of structural minerals but may also act as catalysts in diverse organic reactions, driving the generation and transformation of prebiotic molecules<sup>199</sup>.

### 6.1 Multivalency and Redox Reactions

The d-orbital electronic structure of transition metals confers multiple stable oxidation states. For instance, iron (Fe) can switch between +2 and +3 states, copper (Cu) between +1 and +2, while manganese (Mn) possesses stable states ranging from +2 to +7<sup>200</sup>. This multivalency enables them to facilitate electron transfer reactions without the need for enzymes, making them pivotal catalysts for primordial redox processes<sup>201</sup>.

In experiments simulating early Earth and astrochemical environments, Fe<sup>2+</sup> and Fe<sup>3+</sup> are frequently employed to catalyze the transformation of organic acids, aldehydes, nitro compounds, and phosphoric acid derivatives. For instance, Fe<sup>3+</sup> catalyze H<sub>2</sub>O<sub>2</sub>-mediated oxidation reactions, whilst Fe<sup>2+</sup> drive molecular reduction to form amine, alcohol or hydrocarbon intermediates. This redox capability is scarce in enzyme-free systems, highlighting the potential of transition metals as primordial "electron carriers"<sup>202</sup>.

### 6.2 Coordination Chemistry Properties and Catalytic Microenvironment Construction

---

<sup>198</sup> Pizzarello, S., Cooper, G. W., & Flynn, G. J. (2006). The nature and distribution of the organic material in carbonaceous chondrites and interplanetary dust particles. In *Meteorites and the Early Solar System II* (pp. 625–651).

<sup>199</sup> Cody, G. D., Alexander, C. M. O'D., & Tera, F. (2000). Organo-metallic compounds in the Murchison meteorite: Implications for the origin of life. *Geochimica et Cosmochimica Acta*, 64(2), 333–347. [https://doi.org/10.1016/S0016-7037\(99\)00295-1](https://doi.org/10.1016/S0016-7037(99)00295-1)

<sup>200</sup> Kitadai, N., Nakamura, R., Yamamoto, M., Takai, K., & Yoshida, N. (2021). Metals likely promoted protometabolism in early ocean alkaline hydrothermal systems. *Scientific Reports*, 11, 10971. <https://doi.org/10.1038/s41598-021-90261-7>

<sup>201</sup> Russell, M. J., Hall, A. J., & Martin, W. (2010). Serpentinisation as a source of energy at the origin of life. *Geobiology*, 8(5), 355–371. <https://doi.org/10.1111/j.1472-4669.2010.00249.x>

<sup>202</sup> Morowitz, H. J., Kostelnik, J. D., Yang, J., & Cody, G. D. (2000). The origin of intermediary metabolism. *Proceedings of the National Academy of Sciences*, 97(14), 7704–7708. <https://doi.org/10.1073/pnas.110153997>

Transition metal ions possess extensive coordination capabilities, forming coordination complexes with diverse geometric structures—such as tetra-coordinate, hexa-coordinate, and octahedral configurations with water, carboxylic acids, amines, phosphates, hydroxyl groups, and sulphur-containing functionalities. These coordination interactions not only spatially position reaction substrates but also significantly reduce reaction activation energy by modifying microenvironmental factors, such as charge distribution, acidity/alkalinity, and nucleophilicity<sup>203</sup>. Hydrated silicate, metal sulphide, or oxide particles within protoplanetary discs provided natural binding substrates for metal ions, creating reaction "hotspots" nanoreactors between transition metals and adsorbed organic small molecules. This environment may have mimicked later enzyme active sites, providing a template and catalytic foundation for primordial polymerization and condensation reactions<sup>204</sup>.

### 6.3 Stabilizing Role of Catalytic Intermediates

In modern enzyme catalysis, reactions often proceed via the formation of highly energetic intermediates (e.g., carboxylation intermediates, peroxides, radicals, enol intermediates) that are extremely unstable in their free states. Transition metals can temporarily 'capture' these intermediates through coordination, reducing their decomposition rates and guiding the reaction along favourable pathways.

For instance,  $\text{Ni}^{2+}$  and  $\text{Fe}^{2+}$  stabilize adsorbed CO and H intermediates in Fischer–Tropsch-type reactions, thereby promoting carbon chain elongation.  $\text{Zn}^{2+}$  stabilizes pentacoordinate phosphate intermediates, playing a crucial role in modern nucleic acid polymerases. Thus, the "intermediate trapping" function of such metals holds profound significance in prebiotic chemistry as well<sup>205</sup>.

### 6.4 Traces of Metal Enzyme Evolution in Contemporary Life Systems

Modern life systems contain numerous metalloproteins with transition metals as active centers, exhibiting a high dependence on metal ions. It may reflect "genetic traces" of early metal involvement in reactions. Common examples include iron–sulphur clusters (Fe–S clusters), which are present in electron transport chains, nitrate reductases, carboxylases, and other key enzymes, enabling redox reactions and electron transfer. Zinc enzymes (Zn

---

<sup>203</sup> Holm, N. G., & Andersson, E. M. (2005). Hydrothermal simulation experiments as a tool for studies of the origin of life on Earth and other terrestrial planets: A review. *Astrobiology*, 5(4), 444–460.

<https://doi.org/10.1089/ast.2005.5.444>

<sup>204</sup> Dupont, C. L., Butcher, A., Valas, R. E., Bourne, P. E., & Caetano-Anollés, G. (2010). History of biological metal utilisation inferred through phylogenomic analysis of protein structures. *Proceedings of the National Academy of Sciences*, 107(23), 10567–10572. <https://doi.org/10.1073/pnas.0912491107>

<sup>205</sup> Maret, W. (2013). Zinc biochemistry: From a single zinc enzyme to a key element of life. *Advances in Nutrition*, 4(1), 82–91. <https://doi.org/10.3945/an.112.003038>

enzymes): Such as carbonic anhydrase, DNA polymerase, and insulin dehydrogenase, which utilize  $Zn^{2+}$  to provide Lewis acid centres, stabilizing intermediates and promoting deprotonation. Cu enzymes participate in cellular processes such as pigment oxidation and superoxide dismutase activity, serving as primary metal centers for oxygen activation<sup>206</sup>. Ni enzymes: Examples include methanesequioxigenase and hydrogenases, demonstrating potent catalytic capabilities for reduction reactions and small-molecule transformations.

These enzymes commonly possess complex metal coordination environments that are highly similar to those formed in protoplanetary discs or Earth-like environments, suggesting they may share a common evolutionary origin<sup>207</sup>.

## 6.5 From Environmental Catalysis to Enzymatic Catalysis

Transition metals, owing to their multivalency, coordination capabilities, and ability to stabilize intermediates, may not only have driven prebiotic organic reactions within protoplanetary discs but were also 'borrowed' and optimized by biological systems during evolution, becoming cornerstones of modern life's core reaction systems. This continuity 'from environmental catalysis to enzyme catalysis'<sup>208</sup> may well represent a profound manifestation of the deep connection between life and the cosmic environment.

## 7. Analysis of Prebiotic Chemical Conditions in Protoplanetary Discs

Although interstellar dust (ISM) can undergo chemical reactions under cosmic ray irradiation, producing precursor molecules for amino acids and nucleic acids, Earth itself lacks the conditions for possessing dust and conducting pre-chemical reactions. Nevertheless, the prebiotic conditions necessary for generating key pre-life substances, such as amino acids, nucleic acids, and the precursors for forming cell membranes, can only be realized within protoplanetary discs. Consequently, the physicochemical reactions occurring within protoplanetary disc dust must be repeatedly emphasized as constituting the core of prebiotic chemical reactions. It can be stated that without the physicochemical reactions of protoplanetary disc dust, life would not exist.

### 7.1 Dust Particles and Surface Catalysis

---

<sup>206</sup> Gray, H. B., & Winkler, J. R. (2005). Long-range electron transfer. *Proceedings of the National Academy of Sciences*, 102(10), 3534–3539. <https://doi.org/10.1073/pnas.0408029102>

<sup>207</sup> Raanan, H., et al. (2018). Metal-centred prebiotic chemistry and the evolution of metalloproteins. *Nature Ecology & Evolution*, 2, 1703–1710. <https://doi.org/10.1038/s41559-018-0670-y>

<sup>208</sup> Morowitz, H. J., Kostelnik, J. D., Yang, J., & Cody, G. D. (2000). The origin of intermediary metabolism. *Proceedings of the National Academy of Sciences*, 97(14), 7704–7708. <https://doi.org/10.1073/pnas.110153997>



Within protoplanetary discs, dust particles not only serve as the fundamental building blocks for planetary embryos, but their surfaces may also function as crucial platforms for prebiotic chemical reactions. These particles are predominantly composed of iron-rich silicate minerals, such as olivine ( $(\text{Mg, Fe})_2\text{SiO}_4$ ) and pyroxene ( $(\text{Mg, Fe})\text{SiO}_3$ ). Their crystal structures, surface activity, and responsiveness to energy inputs (such as light and heat) provide diverse physicochemical microenvironments conducive to early organic synthesis<sup>209</sup>.

## 7.2 Catalytic Functionality of Iron-Rich Silicate Powders

Peridotite and pyroxene minerals belong to the earliest-condensing refractory phases in the solar nebula's condensation sequence. They are thus widely distributed in calcium-aluminum-rich inclusions (CAIs), chondrites, and protoplanetary disc dust. These mineral structures typically contain  $\text{Fe}^{2+}$  or  $\text{Fe}^{3+}$ . At surface defect sites, they provide Lewis acid centres capable of adsorbing gas molecules such as  $\text{CO}$ ,  $\text{H}_2$ , and  $\text{N}_2$ . Through surface electron transfer, they induce dissociative adsorption and subsequent reactions. Studies indicate that under simulated solar protoplanetary nebula conditions, these iron-rich silicate powders can simultaneously catalyze two classical reaction pathways<sup>210</sup>.

## 7.3 Fischer–Tropsch-type synthesis reactions ( $\text{CO} + \text{H}_2 \rightarrow \text{hydrocarbons} + \text{H}_2\text{O}$ )

This reaction pathway occurs on metallic iron, iron oxide, or ferrosilicate surfaces, serving as a crucial mechanism for synthesizing lower alkanes, alkenes, and aromatics:



Within the mid-planetary disc region (1–10 AU, with temperatures approximately 200–600 K)<sup>211</sup>, this reaction may proceed on the surfaces of Fe-rich dust particles<sup>212</sup>, particularly when the energy barrier is lowered and reaction rates increase upon excitation by X-rays or UV radiation.

---

<sup>209</sup> Hill, H. G. M., & Nuth, J. A. (2003). The catalytic potential of cosmic silicates: Implications for prebiotic chemistry in the solar nebula. *Astrobiology*, 3(2), 291–304. <https://doi.org/10.1089/153110703769016488>


<sup>210</sup> Loeffler, M. J., Raut, U., Baragiola, R. A., & Carlson, R. W. (2008). Synthesis of hydrogen peroxide in water ice by ion irradiation. *Icarus*, 197(1), 291–295. <https://doi.org/10.1016/j.icarus.2008.03.013>

<sup>211</sup> Kress, M. E., & Tielens, A. G. G. M. (2001). Surface chemistry on interstellar dust grains: The formation of organic molecules. *Astrophysical Journal*, 551(1), 775–785. <https://doi.org/10.1086/320239>

<sup>212</sup> McCollom, T. M., & Seewald, J. S. (2006). Carbon isotope composition of organic compounds produced by abiotic synthesis under hydrothermal conditions. *Earth and Planetary Science Letters*, 243, 74–84. <https://doi.org/10.1016/j.epsl.2005.12.027>

#### 7.4 Hubble-type synthesis reaction ( $\text{N}_2 + \text{H}_2 \rightarrow \text{NH}_3$ )

Despite the high triple bond energy of  $\text{N}_2$  molecules (941 kJ/mol), Fe-rich mineral surfaces provide electron cloud displacement and  $\pi$ -antienergetic orbital coordination capabilities, inducing  $\text{N} \equiv \text{N}$  bond activation. Simulation experiments and DFT studies indicate that at low pressure and temperature,  $\text{Fe}_2\text{SiO}_4$  or FeO-coated mineral surfaces can achieve the following transformation:



It implies that iron-rich dust particles within the solar nebula could not only catalyze hydrocarbon synthesis<sup>213</sup> but also facilitate nitrogen reduction processes, providing a natural platform for the generation of ammonia precursors.

#### 7.5 Multicomponent Reaction: $\text{CO} + \text{N}_2 + \text{H}_2 \rightleftharpoons$ Nitrogen-containing Organics

More remarkably, when  $\text{CO}$ ,  $\text{H}_2$ , and  $\text{N}_2$  coexist within protoplanetary discs, these catalytic surfaces can further catalyze the formation of nitrogen-containing organics (such as methylamine, acetonitrile, and carbamates). Studies indicate that C–N bonds can form via the following steps in the presence of Fe-Si-O active surfaces or FeO clusters:

- 1.)  $\text{CO}$  adsorption +  $\text{H}_2 \rightarrow$  HCO or  $\text{CH}_x$  surface intermediates
- 2.) Partial dissociation of  $\text{N}_2 \rightarrow$  N or  $\text{NH}_x$  active species
- 3.)  $\text{CH}_x + \text{NH}_x \rightarrow$   $\text{CH}_3\text{NH}_2$  (methylamine) or  $\text{CH}_3\text{CN}$  (acetonitrile), etc.

Such reactions require high temperatures and pressures, or artificial catalysts, under terrestrial conditions. However, in protoplanetary discs, they may proceed spontaneously on iron-rich silicate dust surfaces, assisted by UV or radiation<sup>214</sup>. It offers a novel interpretative pathway for the formation of nitrogen-containing organic molecules—precursors to fundamental building blocks of life such as amino acids and nucleobases.

#### 7.6 Dust Particles: Natural Reactors and Selective Reaction Containers

From a prebiotic chemical perspective, these dust particles serve not only as catalytic platforms but also as "microscopic reaction vessels" with the following advantages:

---

<sup>213</sup> Charaya, S., Delgass, W. N., Ribeiro, F. H., & Agrawal, R. (2016). Haber–Bosch synthesis of ammonia over iron-based catalysts: Mechanistic studies. *ACS Catalysis*, 6(9), 5947–5953. <https://doi.org/10.1021/acscatal.6b01260>

<sup>214</sup> Fioroni, M., & DeYonker, N. J. (2016). Iron and ruthenium centres grafted onto silica surfaces as models for heterogeneous Fischer–Tropsch and Haber–Bosch catalysis. *The Journal of Physical Chemistry C*, 120(50), 28744–28752. <https://doi.org/10.1021/acs.jpcc.6b10696>

Localised concentration effect: Reagents can be concentrated at the nanoscale, increasing contact opportunities; Reaction site diversity: Surface lattice defects, metal ions, and charge inhomogeneities induce multi-path chemical selectivity; Thermal stability and structural protection: Dust particles buffer radiation and thermal fluctuations, safeguarding intermediates from destruction; Ice-enveloped dust systems: Can coexist with water ice or organic ice, facilitating interfacial reactions and phase-change-induced reaction chains.

### **7.7 Early Formation of "C-H-N" Skeletons Provides Crucial Chemical Pathways**

Iron-rich silicate dust particles within protoplanetary discs possess not only Fischer–Tropsch-type hydrocarbon synthesis capabilities but also nitrogen reduction functions akin to the Haber process, even driving the synthesis of nitrogen-containing organic compounds within complex gas systems. This concurrent catalysis of CO, H<sub>2</sub>, and N<sub>2</sub> systems positions them as one of the most promising natural reaction platforms in prebiotic chemistry, providing a crucial chemical pathway for constructing the "C-H-N" skeleton during the early stages of life<sup>215</sup>.

## **8. Experimental simulations and theoretical models: validating metal-catalyzed prebiotic mechanisms**

Understanding the specific role of metals in prebiotic organic reactions within protoplanetary discs necessitates reconstructing plausible reaction pathways through experimental simulations and theoretical modelling. Over recent decades, multiple simulation experiments utilizing metals such as Fe, Co, and Ni as catalytic centres have demonstrated the potential for forming complex organic molecules under enzyme-free, non-biological conditions. These studies not only elucidate reaction mechanisms at the molecular level but also enable their extrapolation for comparative analysis within protoplanetary disc chemical environments.

### **8.1 Fe/Co/Ni Catalysis in Fischer–Tropsch-Type Reactions**

Fischer–Tropsch-type synthesis (FTT) was initially employed industrially to synthesize alkanes and alcohols from carbon monoxide (CO) and hydrogen (H<sub>2</sub>) in the presence of metal catalysts. In prebiotic chemistry research, FTT reactions have been extensively utilized to

---

<sup>215</sup> Sabbah, H., et al. (2010). Gas-phase synthesis of nitriles in the interstellar medium: Reaction of CN with unsaturated hydrocarbons. *Astrophysical Journal*, 718, 1001–1007. <https://doi.org/10.1088/0004-637X/718/2/1001>

simulate hydrocarbon chain growth processes under primordial planetary or interstellar conditions<sup>216</sup>.

Experiments demonstrate that Fe, Co, and Ni catalysts can convert CO + H<sub>2</sub> mixtures into straight-chain hydrocarbons, alcohols, and fatty acids under low pressure and elevated temperatures (200–400°C)<sup>217</sup>. In experiments simulating meteorite impact or protoplanetary disc thermal environments, these metals are frequently supported on silicate particles or oxide surfaces, such as Fe<sub>3</sub>O<sub>4</sub>/SiO<sub>2</sub> or Ni/Al<sub>2</sub>O<sub>3</sub>, to mimic the physical properties of protoplanetary dust.

FTT products encompass organic compounds with chain lengths ranging from C1 to C18, some of which are structurally analogous to modern fatty acids, esters, and membrane structural precursors, providing potential chemical precursors for the formation of membranes<sup>218</sup>.

## 8.2 Haber Synthesis and Nitrogen Precursor Formation

Beyond hydrocarbon synthesis, nitrogen fixation and transformation in primordial environments constituted a critical challenge. The Haber–Bosch process, utilized in modern industry for synthesizing ammonia (NH<sub>3</sub>), follows the reaction pathway  $N_2 + 3H_2 \rightleftharpoons 2NH_3$ , relying on metal catalysts such as iron or molybdenum.

In simulated protoplanetary disc experiments, metal sulphides such as Fe, Fe<sub>3</sub>O<sub>4</sub>, and FeS can generate ammonia or other nitrogen-containing intermediates under high temperatures and H<sub>2</sub> atmospheres, without requiring high-pressure conditions<sup>219</sup>. Some studies indicate that metal oxides present in protoplanetary discs, including Ti, Fe, and Mo, may also facilitate precursor transformations by adsorbing and activating N<sub>2</sub> molecules on their surfaces. The resulting NH<sub>3</sub> or amine compounds can subsequently condense with organic acids to form amino acid precursors<sup>220</sup>.

---

<sup>216</sup> McCollom, T. M., & Seewald, J. S. (2006). Carbon isotope composition of organic compounds produced by abiotic synthesis under hydrothermal conditions. *Earth and Planetary Science Letters*, 243(1-2), 74–84. <https://doi.org/10.1016/j.epsl.2005.12.027>

<sup>217</sup> Yamanaka, M., et al. (2020). Hydrothermal Fischer-Tropsch-type synthesis of long-chain hydrocarbons up to C35 with NiFe catalysts. *ACS Earth and Space Chemistry*, 4(5), 732–743. <https://doi.org/10.1021/acsearthspacechem.9b00302>

<sup>218</sup> Yamanaka, M., et al. (2020). Hydrothermal Fischer-Tropsch-type synthesis of long-chain hydrocarbons up to C35 with NiFe catalysts. *ACS Earth and Space Chemistry*, 4(5), 732–743. <https://doi.org/10.1021/acsearthspacechem.9b00302>

<sup>219</sup> Brandes, J. A., et al. (1998). Abiotic nitrogen reduction on the early Earth. *Nature*, 395, 365–367. <https://doi.org/10.1038/26441>

<sup>220</sup> Lerner, N. R., & Peterson, E. (1994). The photochemical production of amino acids from mixtures of ammonia, formaldehyde and hydrogen cyanide. *Origins of Life and Evolution of the Biosphere*, 24, 113–116. <https://doi.org/10.1007/BF01582004>

This reaction system provides an abiotic starting point for amino acid synthesis and may explain the co-occurrence of amino acids and amines in certain meteorites. The Haber synthesis is detectable only within protoplanetary discs, as the conditions favourable for this process were absent in Earth's early environment. Consequently, the Haber synthesis does not apply to the environmental conditions that prevailed on Earth during its primordial era.

### **8.3 Experiments on metal coordination catalysis on protoplanetary disc dust or mineral surfaces**

Beyond gas-phase reactions, simulation experiments extensively investigate the catalytic capabilities of metal ions on mineral or dust surfaces. By grafting metals (such as  $\text{Zn}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{Cu}^{2+}$ ) onto natural or synthetic silicate, phosphate, or alumina materials, these studies mimic the porous structure and surface coordination sites of protoplanetary disc dust.

For instance, Fioroni and DeYonker (2016) modelled the adsorption of  $\text{Fe}^{2+}$  onto  $\text{SiO}_2$  surfaces and its catalysis of the  $\text{H}_2$  decomposition and reduction reactions<sup>221</sup>. Similar structures have been observed in iron-rich dust, serpentine, or pyrite. Such models reveal multi-coordination structures between metal centres and hydroxyl groups or organic anions, forming microenvironments analogous to enzyme active sites. This surface catalytic system effectively promotes the following reaction types: esterification of carboxylic acids with alcohols, condensation of amines with carboxylic acids, and dehydration polymerization of phosphates<sup>222</sup>.

**8.4 Although experimental conditions often require moderate "optimization" to obtain reaction products** (e.g., elevated temperatures, increased catalyst concentrations), these results retain practical astrochemical comparability. Inner regions of protoplanetary discs (<10 AU) can reach temperatures of several hundred K, harbouring abundant gases such as  $\text{CO}$ ,  $\text{H}_2$ , and  $\text{N}_2$ , alongside high-surface-area, metal-rich dust particles.

Furthermore, energy inputs for reactions are provided by the decay of radioactive isotopes (e.g.,  $^{26}\text{Al}$ ), stellar wind heating, and ultraviolet and X-ray irradiation. ALMA observations and comet missions (e.g., Rosetta) have also detected numerous reaction products—including alkanes, ketones, amines, and amino acids—widely present in interstellar ice and protoplanetary material.

---

<sup>221</sup> Fioroni, M., & DeYonker, N. J. (2016). Iron and ruthenium centres grafted onto silica surfaces as models for heterogeneous Fischer–Tropsch and Haber–Bosch catalysis. *The Journal of Physical Chemistry C*, 120(50), 28744–28752. <https://doi.org/10.1021/acs.jpcc.6b10696>

<sup>222</sup> Fioroni, M., & DeYonker, N. J. (2016). Iron and ruthenium centres grafted onto silica surfaces as models for heterogeneous Fischer–Tropsch and Haber–Bosch catalysis. *The Journal of Physical Chemistry C*, 120(50), 28744–28752. <https://doi.org/10.1021/acs.jpcc.6b10696>



Consequently, experiments such as the FTT and Haber reactions, alongside mineral surface catalysis<sup>223</sup>, not only replicate theoretical models but also further validate the actual capabilities of protoplanetary discs in prebiotic synthesis<sup>224</sup>.

## 9. The transition from metallic elements to biological functions

Metallic elements, particularly transition metals, play a pivotal role in enzymeless prebiotic chemistry due to their unique redox activity, coordination chemistry, and ability to stabilize intermediates. The widespread reliance of modern life systems on metalloproteins suggests these metallic functions represent not recent evolutionary innovations, but rather a "hereditary" inheritance of early non-living reaction mechanisms. Understanding the transformative pathways from protoplanetary metallic environments to biochemical functional modules is crucial for elucidating the "continuity" mechanisms underlying life's origins.

### 9.1 The 'Inheritance' of Metals: From Non-living Catalysis to Retention in Living Systems

Modern life systems exhibit a profound dependence on transition metals as active centres within key enzymes. It includes the role of Fe–S clusters in electron transfer, the activity regulation of Zn<sup>2</sup>S in nucleic acid polymerization and deprotonation, and the catalytic functions of elements such as Mo and Ni in nitrogen fixation and methanogenesis<sup>225</sup>. These metalloproteins are conserved across all known life forms, indicating an extremely ancient evolutionary origin predating the establishment of cellular mechanisms.

Experimental simulations demonstrate that during the abiotic prebiotic stage, these metals could already perform functions such as energy conversion, substrate activation, and polymerization reactions on mineral or dust surfaces<sup>226</sup>. Such metallic reaction centers stabilize high-energy intermediates through coordination structures, exhibiting functional characteristics analogous to those of modern enzymes. It suggests that life systems likely "inherited" these non-living catalytic strategies, which were progressively encapsulated and optimized by organic frameworks (such as peptide chains) during molecular evolution,

---

<sup>223</sup> Goesmann, F., et al. (2015). Organic compounds on comet 67P/Churyumov–Gerasimenko revealed by COSAC mass spectrometry. *Science*, 349(6247), aab0689. <https://doi.org/10.1126/science.aab0689>

<sup>224</sup> Öberg, K. I., et al. (2015). The comet-like composition of a protoplanetary disk as revealed by complex cyanides. *Nature*, 520, 198–201. <https://doi.org/10.1038/nature14276>

<sup>225</sup> Dupont, C. L., et al. (2010). History of biological metal utilisation inferred through phylogenomic analysis. *PNAS*, 107, 10567–10572. <https://doi.org/10.1073/pnas.0912491107>

<sup>226</sup> Cody, G. D., et al. (2000). Organometallic compounds in the Murchison meteorite. *Geochim. Cosmochim. Acta*, 64, 333–347. [https://doi.org/10.1016/S0016-7037\(99\)00295-1](https://doi.org/10.1016/S0016-7037(99)00295-1)

ultimately evolving into today's highly specialized metalloproteins<sup>227</sup>. This "continuous inheritance" from non-enzymatic metal centres to biocatalytic systems constitutes a distinct "chemical inheritance" phenomenon in the origin of life. This transformation of metallic functionality "from non-living to living" may be regarded as a "molecular memory" mechanism in life's evolution: effective chemical strategies were preserved and transmitted through genetic systems, thereby establishing the fundamental framework for metabolism and structure.

## **9.2 Comparison of Metal Availability in the Protoplanetary Disk and Primitive Earth**

The primordial Earth, though rich in metallic resources, possessed an early environment (>4 billion years ago) characterized by strongly reducing oceans and an unstable crust. This environment severely limited the bioavailability of numerous transition metals (such as Mo, Zn, Cu)<sup>228</sup>. Particularly under anaerobic conditions, these elements proved difficult to dissolve and incorporate into reaction systems, resulting in a significant deficiency of certain metals within the early geochemical environment.

By contrast, the protoplanetary disc exhibited more uniform elemental distribution and a more complete array of metals. Elements such as Fe, Ni, Mg, and Zn, commonly present in meteorites, planetesimals, and comets in metallic, oxide, and sulphide forms, demonstrate high reactivity under the influence of ultraviolet radiation, radioactive decay, and high-temperature heating<sup>229</sup>. Moreover, these metals often deposit as nanoparticles onto high-surface-area dust or silicate particles, providing natural conditions for the formation of active catalytic surfaces.

Consequently, protoplanetary discs may have provided superior metal availability and catalytic environments for prebiotic synthesis compared to early Earth. It also suggests that some key metallic components essential for life may not have formed endogenously on Earth, but instead completed their birth and evolution within protoplanetary discs before being transported to Earth for further biological evolution.

## **9.3 The Concept of the Protoplanetary Disk as a "Prebiotic Chemical Reactor"**

Synthesising analyses from preceding sections, the protoplanetary disc may be conceptualised as a natural "prebiotic chemical reactor" exhibiting several core characteristics: Highly enriched CHNOPS and metallic elements, injected by supernovae,

---

<sup>227</sup> Raanan, H., et al. (2018). Metal-centred prebiotic chemistry and the evolution of metalloproteins. *Nat. Ecol. Evol.*, 2, 1703–1710. <https://doi.org/10.1038/s41559-018-0670-y>

<sup>228</sup> Pasek, M. A., & Lauretta, D. S. (2005). Phosphide corrosion and reactive phosphorus on the early Earth. *Astrobiology*, 5, 515–535. <https://doi.org/10.1089/ast.2005.5.515>

<sup>229</sup> Öberg, K. I., et al. (2015). The comet-like composition of a protoplanetary disk. *Nature*, 520, 198–201. <https://doi.org/10.1038/nature14276>

AGB stars, and comets, forming a catalytically potent mixed system; Diverse energy-driving mechanisms, including stellar radiation, high-energy particle irradiation, radioactive isotope decay, and interstellar impacts; A multi-scale reaction environment, ranging from nanoscale dust coordination sites to thermal reaction spaces within planetesimals, capable of satisfying the conditions for complex reaction chains to form<sup>230</sup>; Continuous material cycling and combinatorial experiments, with physical perturbations of the protoplanetary disc (such as disc winds, migration, and collisions) constantly rearranging matter, providing a diverse backdrop for chemical evolution<sup>231</sup>.

Such environments are not only conducive to individual reactions but particularly suited to combinatorial chemical evolution. Molecules such as fatty acids, peptide chains, and nucleotides can progressively accumulate and undergo selection within the same system, marking the first step toward the origin of life. These conditions converge to form a dynamic, multi-level, modular natural reaction system, akin to a "prototype reactor" preceding biological cells. Within this system, metal catalysis simultaneously influences key reaction pathways across different spatial scales, including carbon chain elongation, nitrogen reduction, fatty acid polymerization, and phosphoester bond formation, thereby enhancing the probability of co-occurrence and selective aggregation of prebiotic precursors.

This hypothesis differs from both Directed Panspermia and any other panspermia theory. The protoplanetary disk origin of life posits that life emerged exclusively within protoplanetary disks, not in other star systems or planetary bodies. Earth's life originated from such a disk, with its core components potentially synthesized and pre-packaged within celestial matter before being delivered to our planet. I term this panspermia theory "planetary disks Panspermia".

## **10. Polycyclic Aromatic Hydrocarbons (PAHs) and Complex Organic Compounds**

Polycyclic aromatic hydrocarbons (PAHs) constitute a class of hydrocarbons widely distributed throughout the cosmos, typically comprising two or more fused benzene rings. Owing to their structural stability, strong optical activity, and high reactivity diversity, PAHs are regarded as among the most abundant complex organic molecules in the universe. They may play a significant role in the origin of life.

### **10.1 Infrared Characteristics of PAHs and Protoplanetary Disk Observations**

---

<sup>230</sup> Fioroni, M., & DeYonker, N. J. (2016). H<sub>2</sub> formation at Fe-grafted SiO<sub>2</sub> surfaces. *J. Phys. Chem. C*, 120, 8182–8189. <https://doi.org/10.1021/acs.jpcc.6b00192>

<sup>231</sup> Morowitz, H. J., et al. (2000). The origin of intermediary metabolism. *PNAS*, 97, 7704–7708. <https://doi.org/10.1073/pnas.110153997>

PAHs undergo electronic excitation upon ultraviolet irradiation, subsequently releasing energy via infrared radiation. They exhibit characteristic emission features in the mid-infrared region<sup>232</sup>, including: 3.3  $\mu\text{m}$  (C–H stretching vibration); 6.2  $\mu\text{m}$  (C=C skeletal vibration); 7.7  $\mu\text{m}$  (aromatic C–C bond mixed vibration); 11.3  $\mu\text{m}$  (C–H bending vibration)

These features have been observed in protoplanetary discs surrounding numerous young stars, such as Herbig Ae/Be stars<sup>233</sup> and T Tauri discs (Geers et al., 2006; Acke et al., 2010). The intensity and spatial distribution of PAHs provide insights into the disc's radiation field, gas density, and dust particle evolution<sup>234</sup>.

## 10.2 Capacity to form $\pi$ -complexes with elements such as Fe, Si, and Ti

The aromatic  $\pi$  electron cloud of PAHs can form  $\pi$ -coordinated complexes with transition metals (e.g., Fe, Ni, Ti) or metalloid elements (e.g., Si). These complexes exhibit the following characteristics: Enhanced stability: Metal ions embed within the PAH ring plane, either donating or accepting electrons to form synergistic bonding; Catalytic centre function: Fe–PAH complexes catalyze electron transfer, radical reactions, or small-molecule reduction; Photoactivity regulation: Metal coordination induces red shifts in PAH absorption bands, broadening the excitation reaction window.

Relevant experimental and theoretical studies indicate: Fe–PAH complexes can mediate redox reactions in the excited state (Szczepanski & Vala, 2006)<sup>235</sup>; Ti–Fullerene/PAH complexes have been proposed as the source of the 21  $\mu\text{m}$  infrared feature (Kimura et al., 2005)<sup>236</sup>; Si–PAH  $\pi$  complexes exhibit high spectral matching with interstellar dust infrared features (Joalland et al., 2009)<sup>237</sup>. Such complexes may also be regarded as the chemical prototypes of modern metalloproteinase active sites.

## 10.3 PAH oxidation products (quinones) and electron transfer potential

---

<sup>232</sup> Hudgins, D. M., & Allamandola, L. J. (1999). Infrared spectroscopy of matrix-isolated polycyclic aromatic hydrocarbon cations. 1. PAH<sup>+</sup> bands and diagnostics. *Astrophysical Journal*, 516, L41–L44. <https://doi.org/10.1086/311976>

<sup>233</sup> Geers, V. C., Augereau, J.-C., Pontoppidan, K. M., et al. (2006). PAH emission characteristics in Herbig Ae/Be systems: Evidence for evolution. *Astronomy & Astrophysics*, 459(2), 545–556. <https://doi.org/10.1051/0004-6361:20065411>

<sup>234</sup> Acke, B., Bouwman, J., Juhász, A., et al. (2010). Probing the PAH emission in Herbig Ae/Be stars with Spitzer. *Astrophysical Journal*, 718(1), 558–573. <https://doi.org/10.1088/0004-637X/718/1/558>

<sup>235</sup> Szczepanski, J., & Vala, M. (2006). Infrared spectra and structures of iron–PAH complexes: A matrix isolation study. *Astrophysical Journal*, 646, 666–675. <https://doi.org/10.1086/504872>

<sup>236</sup> Kimura, H., Ishimoto, H., Yamamoto, T., & Mukai, T. (2005). The 21  $\mu\text{m}$  feature in post-AGB stars and titanium-containing fullerene derivatives. *Astronomy & Astrophysics*, 432, 737–743. <https://doi.org/10.1051/0004-6361:20041961>

<sup>237</sup> Joalland, B., Simon, A., & Marsden, C. J. (2009). Structures and IR spectra of silicon–PAH  $\pi$ -complexes: A theoretical study. *Astronomy & Astrophysics*, 494, 969–975. <https://doi.org/10.1051/0004-6361:200810579>

Under illumination, ionization, or oxidative conditions, PAHs may form quinone structures, such as naphthoquinone and benzoquinone. These molecules possess the following biochemical significance<sup>238</sup> : Efficient electron acceptors: Quinones can participate in single- or double-electron transfer; Formation of electrochemical gradients: When coordinated with Fe<sup>2++</sup> / Fe<sup>3++</sup> or other metals, they can generate primordial electron flows; Precursor function: In modern organisms, quinones (e.g., ubiquinone, coenzyme Q) serve as pivotal electron carriers in respiratory and photosynthetic chains. Thus, oxidized derivatives of PAHs formed within protoplanetary discs may have underpinned early primordial metabolic networks.

#### **10.4 PAHs as Structural Templates for Prebiotic Membranes or Primitive Enzyme Systems**

Owing to their high planarity, hydrophobicity, and  $\pi$ - $\pi$  stacking capacity, PAHs and their complexes are proposed to participate in the following structures and functions: Membrane-like self-assembly: PAHs form monolayers or multilayered films at the water-organic interface<sup>239</sup> , resembling precursor lipid membranes<sup>240</sup> ; Ordered reaction environments:  $\pi$ - $\pi$  stacking arranges small molecules, enhancing reaction selectivity; Model for metal-coordinated enzymes: Fe-PAH or Ni-PAH structures may mimic the active centres of primordial reductases (e.g., pre-iron-sulphur clusters). Some studies even propose PAHs as prototypical templates for RNA bases, promoting directional formation or polymerization of purine and pyrimidine structures under UV irradiation.

PAHs rank among the earliest stable complex organic molecules formed within protoplanetary discs. Their high abundance, multi-spectral observational signatures, and capacity for  $\pi$ -coordination with metallic elements, alongside their redox activity, confer a multifaceted potential for catalysis, energy conversion, and structural templating. From complex organic synthesis to protometabolic prototypes and the formation of membrane-like structures, PAHs may represent pivotal intermediaries bridging cosmic chemistry and prebiotic biochemistry.

### **11. Polycyclic Aromatic Hydrocarbons (PAHs) and Transition Metals (TMs) in Protoplanetary Disks**

---

<sup>238</sup> Ehrenfreund, P., Rasmussen, S., Cleaves, H. J., & Chen, L. (2006). Astrophysical and astrochemical insights into the origin of life. *Reports on Progress in Physics*, 69(3), 871–932. <https://doi.org/10.1088/0034-4885/69/3/R03>

<sup>239</sup> Dworkin, J. P., Deamer, D. W., Sandford, S. A., & Allamandola, L. J. (2001). Self-assembling amphiphilic molecules: Synthesis in simulated interstellar/precometary ices. *Proceedings of the National Academy of Sciences*, 98(3), 815–819. <https://doi.org/10.1073/pnas.98.3.815>

<sup>240</sup> Ricardo, A., Carrigan, M. A., Olcott, A. N., & Benner, S. A. (2004). Borate minerals stabilise ribose. *Science*, 303(5655), 196–196. <https://doi.org/10.1126/science.1092321>

In protoplanetary discs, polycyclic aromatic hydrocarbons (PAHs) and transition metals (TMs) jointly participate in multiple pathways of prebiotic synthesis. The formation of metal-organic complexes—such as Fe-PAH, Si-PAH, and TM-carbides/nitrides—has profound implications for cosmochemistry, circumstellar chemistry, and the precursors of life. Within protoplanetary discs and star-forming regions, PAHs interact with abundant transition metals (e.g., Fe, Ni, Co, Cr, Ti) to form complex compound systems. These systems may represent: - Prebiotic organometallic precursors; - Key constituents of protoplanetary disc circumstellar gas/solid matter; - Precursors to early Earth organometallic enzymes and metabolic centres.

### 11.1 Specific Metal-PAH and Prebiotic Chemical Complex Types

Fe-PAH complexes: Fe-C<sub>16</sub>H<sub>10</sub><sup>++</sup> (cationic) Szczepanski et al., 2006 Potentially serving as templates for early Fe-organic coordination systems involved in electron transfer<sup>241</sup> Si-PAH  $\pi$ -complexes: Si<sup>+</sup>-PAH composite structures Joalland et al., 2009 Studies on the stability of silicon-aromatic systems in protoplanetary discs, potentially precursors to silicene structures<sup>242</sup> Ti-fullerene adduct: Ti-C<sub>60</sub><sup>++</sup> Kimura et al., 2005, 21  $\mu$ m spectral line

Suggests formation of interstellar Ti-organic complexes, potentially linked to early catalysis or electronic structural stability<sup>243</sup>; Oxides such as FeO, Fe<sub>2</sub>O<sub>3</sub> Iron oxide clusters formed in stellar winds/disks (Li et al., 2013) provide precursors for life's metal enzyme centres, potentially serving as oxygen reduction reaction platforms (<sup>244</sup>). Fe-SiO<sub>2</sub> surface H-grafted structures: Fe<sup>+</sup>-SiO<sub>2</sub>-H (Fioroni & DeYonker, 2016) indicate chemical catalytic capability on dust surfaces, favouring formation of small molecules like H<sub>2</sub> and HCO (<sup>245</sup>); TM-carbides/nitrides: CrCN, FeC, NiN, etc., Brewster & Ziurys (2001)<sup>246</sup>; Sheridan

---

<sup>241</sup> Szczepanski, J., Oomens, J., van Dongen, M., Polfer, N. C., & Eyler, J. R. (2006). Infrared Spectroscopy of Gas-Phase Iron-Polycyclic Aromatic Hydrocarbon Complexes: Fe<sup>+</sup>-C<sub>16</sub>H<sub>10</sub> and Fe<sup>+</sup>-C<sub>24</sub>H<sub>12</sub>. *The Journal of Physical Chemistry A*, 110(44), 13741–13750. <https://doi.org/10.1021/jp063171m>

<sup>242</sup> Joalland, B., Simon, A., Canosa, A., et al. (2009). Gas phase reactions of Si<sup>+</sup>-ions with polycyclic aromatic hydrocarbons: Formation of  $\pi$ -complexes and potential astrophysical implications. *Chemical Physics Letters*, 481(4–6), 193–197. <https://doi.org/10.1016/j.cplett.2009.09.025>

<sup>243</sup> Kimura, H., Mann, I., & Jessberger, E. K. (2005). Composition, structure, and size distribution of dust in the Local Interstellar Cloud. *Astronomy & Astrophysics*, 442(2), 667–674. <https://doi.org/10.1051/0004-6361:20053027>

<sup>244</sup> Li, A., Zhang, K., & Jiang, B. W. (2013). FeO in Interstellar Space: Optical Properties and Astrophysical Implications. *The Astrophysical Journal Letters*, 765(2), L43. <https://doi.org/10.1088/2041-8205/765/2/L43>

<sup>245</sup> Fioroni, M., & DeYonker, N. J. (2016). Astrochemistry of H-Atom Formation on Fe<sup>+</sup> Sites Grafted onto Silicate Dust: A DFT Study. *The Journal of Physical Chemistry C*, 120(33), 18712–18719. <https://doi.org/10.1021/acs.jpcc.6b04200>

<sup>246</sup> Brewster, M. A., & Ziurys, L. M. (2001). Millimetre-wave spectroscopy of FeC and CoC: New transition metal carbide molecules in the gas phase. *The Astrophysical Journal*, 548(1), L139–L142. <https://doi.org/10.1086/318933>

et al. observed antenna fingerprint molecules<sup>247</sup>, which may reveal interstellar gas-phase synthesis pathways<sup>248</sup>.

## 11.2 Prebiotic significance of these structures

### 11.2.1 Precursors of Primitive Enzyme-Like Structures Comprising Metal Centres:

Modern enzymes, such as iron-sulphur clusters (Fe-S clusters), cobalt-vitamin B<sub>12</sub>, and nickel-carboxylases, all contain transition metals. The aforementioned PAH-TM complexes or oxide clusters have demonstrated pro-enzymatic potential in both molecular structure and function, such as catalyzing CO<sub>2</sub> reduction and facilitating electron transfer.

### 11.2.2 Formation of electron donor/acceptor structures facilitates primitive metabolic network development:

$\pi$ -conjugated PAHs may serve as electron cloud platforms, interacting with ion centres such as Fe or Ni to form natural redox platforms. In the absence of cell membranes, such structures may facilitate the formation of primordial potential gradients (analogous to modern "energy-coupling membranes").

**11.2.3 Catalysis on dust particle surfaces:** The binding of TM with PAHs, silica, and ice layers provides physical platforms for interstellar catalytic reaction centres<sup>249</sup>. It facilitates the synthesis of key prebiotic small molecules such as H<sub>2</sub>, HCN, and HCOOH.

## 11.3 Why this constitutes the astrochemical frontier of 'astro-organometallic chemistry'

The formation of these complexes falls within the broader domain of astro-organometallic chemistry: whereas past astrochemistry emphasized the inorganic synthesis of small molecules, these results demonstrate genuine chemical interactions—such as coordination, insertion, and adsorption—between organic structures (e.g., PAHs) and metals within protostellar regions and protoplanetary discs.<sup>250</sup> They are no longer "inert molecules", but may form precursor structures akin to biomolecules under electron, photon, and dust catalysis. Such systems exhibit potential structural and functional continuity with coenzymes and enzyme active sites in living organisms.

---

<sup>247</sup> Sheridan, P. M., Pulliam, R. L., & Ziurys, L. M. (2002). Laboratory Spectroscopy of CrCN and CrNC: First Detection of Chromium Nitride Radicals. *The Astrophysical Journal*, 568(1), 597–605.

<https://doi.org/10.1086/338983>

<sup>248</sup> Sheridan, P. M., & Ziurys, L. M. (2003). Laboratory measurement and analysis of NiN and NiC rotational spectra. *The Astrophysical Journal*, 589(1), 486–492. <https://doi.org/10.1086/374332>

<sup>249</sup> Schwander, D., Huss, G. R., & Bizzarro, M. (2014). The early solar system's transition disk: Constraints from the fractionation of refractory elements in meteorites. *Geochimica et Cosmochimica Acta*, 147, 115–132. <https://doi.org/10.1016/j.gca.2014.03.010>

<sup>250</sup> Thiabaud, A., Marboeuf, U., Alibert, Y., et al. (2014). From stellar nebula to planets: The refractory components. *Astronomy & Astrophysics*, 562, A27. <https://doi.org/10.1051/0004-6361/201322011>

## 11.4 From "Celestial Inorganic Chemistry" to "Celestial Organometallic Chemistry"

Polycyclic aromatic hydrocarbons (PAHs) within protoplanetary discs and star-forming regions interact with transition metals such as iron, cobalt, and nickel, forming a series of organometallic complexes and oxide clusters. Examples include Fe-PAH and Si-PAH  $\pi$  complexes, FeO clusters, and fullerene metal adducts. Evidence for these compounds is not only clearly demonstrated in infrared and millimetre-wave spectral observations, but their structural and functional characteristics also reveal potential key mechanisms involved in primordial prebiotic chemical reactions. This series of discoveries extends traditional "astronomical inorganic chemistry" into "astronomical organometallic chemistry", enabling us to reinterpret the metal active centres of enzymes, energy metabolism systems, and even selective synthetic pathways for primordial molecules. This perspective is becoming an increasingly vital component within the cosmochemical framework for the origin of life.

## 12. Strecker Reaction Pathways and Amino Acid Synthesis Mechanisms in Protoplanetary Discs

### 12.1 The Significance of Strecker Synthesis in Origins of Life Research

The Strecker reaction represents a classic amino acid synthesis pathway involving a three-step reaction between an aldehyde (or ketone), ammonia ( $\text{NH}_3$ ), and hydrogen cyanide (HCN), ultimately yielding  $\alpha$ -amino acids. This reaction is widely employed in prebiotic chemistry research due to its structural versatility and high efficiency under relatively mild conditions. Research indicates that the Strecker reaction can synthesize common amino acids such as glycine, alanine, and valine under experimental conditions. Moreover, its intermediates (e.g.,  $\alpha$ -aminonitrile) and end products have been detected in multiple carbonaceous chondrite meteorites, providing direct evidence for its cosmic feasibility.

### 12.2 Strecker Synthesis Reaction Pathway and Reagent Conditions

The fundamental reaction steps of the Strecker synthesis are as follows<sup>251</sup> :

A. The aldehyde ( $\text{R-CHO}$ ) reacts with ammonia to form an imine:  $\text{R-CHO} + \text{NH}_3 \rightarrow \text{R-CH=NH}$

B. Addition of imine with HCN to form imidocarbons:  $\text{R-CH=NH} + \text{HCN} \rightarrow \text{R-CH(NH}_2\text{)-CN}$

C. Aminonitrile hydrolyses under hydrothermal conditions to yield amino acids:  $\text{R-CH(NH}_2\text{)-CN} + 2\text{H}_2\text{O} \rightarrow \text{R-CH(NH}_2\text{)-COOH} + \text{NH}_3$

---

<sup>251</sup> Oro, J. (1961). Mechanism of synthesis of adenine from hydrogen cyanide under possible primitive Earth conditions. *Nature*, 191(4794), 1193–1194.



The raw materials required for the above reaction—HCN, NH<sub>3</sub>, aldehydes, and water—have all been observed or inferred to exist within protoplanetary discs, being particularly prevalent in ice–dust particles, interstellar molecular clouds, cometary nuclei, and cometary comas<sup>252</sup>.

### 12.3 Reaction Platforms and Energy Sources in Protoplanetary Discs

Protoplanetary discs provide highly dynamic and structurally complex reaction platforms: Dust surfaces and interstellar ice shells: In cold environments, HCN and ammonia can form aminonitrile on the surfaces of ice-encapsulated dust particles<sup>253</sup>, exhibiting adsorption capacity and spatial confinement effects<sup>254</sup>; Ultraviolet irradiation and radiation activation: Ultraviolet and X-rays emitted by protostars (such as the proto-Sun) excite molecules within ice to form free radicals, promoting initial reactions<sup>255</sup>; Particle bombardment and radioactive decay:  $\alpha$  particles and short-lived nuclides (e.g., <sup>26</sup>Al, <sup>60</sup>Fe) generate heat sources within planetesimals, sustaining local hydrothermal environments to complete the final hydrolysis steps in the reaction chain; Solar infrared thermal radiation: provides fundamental driving forces for temperature gradients and material transport within the protoplanetary disc, facilitating material exchange between low-temperature and high-temperature regions.

### 12.4 Co-occurrence of Strecker intermediates and products in carbonaceous chondrites

Samples such as the Mochisen meteorite and Tagish Lake meteorite reveal not only diverse  $\alpha$ -amino acids but also their precursor compounds, including  $\alpha$ -aminonitriles, carboxynitriles, and imine derivatives. This chemical fingerprint of "co-occurrence of intermediates and products" indicates that all stages of the Strecker reaction can not only be initiated in the cold cosmic environment but also undergo final conversion within hydrothermal systems inside early protoplanetary bodies (e.g., planetesimals)<sup>256</sup>.

---

<sup>252</sup> Cody, G. D., et al. (2011). Organic thermochemistry of the Tagish Lake meteorite. *PNAS*, 108(48), 19171–19176.

<sup>253</sup> Elsila, J. E., et al. (2007). Mechanisms of amino acid formation in interstellar ices: Insight from isotopic analysis. *Astrobiology*, 7(1), 163–180.

<sup>254</sup> Danger, G., et al. (2013). Characterisation of amino acid precursors in interstellar ice analogs: Stability under thermal processing. *Geochimica et Cosmochimica Acta*, 118, 184–201.

<sup>255</sup> Bernstein, M. P., et al. (2002). UV irradiation of polycyclic aromatic hydrocarbons in ices: Production of alcohols, quinones, and ethers. *Astrophysical Journal*, 576(1), 1115.

<sup>256</sup> Pizzarello, S., et al. (2006). The organic content of the Tagish Lake meteorite. *Science*, 293(5527), 2236–2239.

## **12.5 Material Cycling and Environmental Regeneration Mechanisms in Protoplanetary Discs**

Protoplanetary discs are not static structures but multi-scale matter-energy circulation systems. Dust particles undergo radial and vertical mixing within the disc due to disc winds, gravitational perturbations, and magnetic fields. Cold, icy dust from the outer disc can enter the inner disc, where it receives thermal activation. In contrast, reaction products from the inner disc may migrate hydrodynamically to the outer disc's cold condensation zones, where they can refreeze. The planetesimal formation process encapsulates dust and reaction intermediates within the planetesimal, providing thermal shielding and a water-sealed environment conducive to molecular accumulation and transformation.

Thus, within this dynamic chain of "condensation–activation–migration–transformation–freezing", the Strecker reaction not only has the opportunity to proceed fully, but its distinct stages can occur sequentially at different spatial locations, ultimately forming complete product aggregates.

## **12.6 The Integrative Nature of Strecker Synthesis in the Protoplanetary Disk**

The Strecker reaction possesses complete reaction conditions, reactant sources, and energy excitation mechanisms within protoplanetary discs, rendering its feasibility far superior to earlier models positing that "interstellar ice forms only intermediates, requiring Earth's aqueous environment for completion." The disc's illumination, particles, radioactive decay, and temperature gradients constitute a highly integrated chemical reactor capable of in situ synthesis spanning the entire chain from small molecules to amino acids. The product assemblages carried by carbonaceous chondrites represent a "frozen record" of this reaction system. It is therefore reasonable to infer that substantial in situ amino acid synthesis occurred within the protoplanetary disc. These products were subsequently encapsulated within planetesimals, asteroids, and even comets, ultimately being delivered to the early Earth, laying the molecular foundation for the emergence of life.

## **13. Environmental integration for complete amino acid synthesis in the protoplanetary disc and realization of the Strecker reaction**

Analyses of carbonaceous chondrites, such as Murchison, have repeatedly revealed the coexistence of intermediates (e.g.,  $\alpha$ -aminonitriles) and final products (e.g.,  $\alpha$ -amino acids) within the Strecker synthesis pathway. This chemical co-occurrence strongly suggests that the Strecker reaction could not only be partially realized in interstellar ice but also likely

underwent its complete three-step reaction chain within the protoplanetary disc<sup>257</sup>. Notably, the liquid water environment required for later hydrolysis can be established within the disc by numerous thermal sources, no longer relying on conditions provided by the early Earth.

The protoplanetary disc harbours an exceptionally diverse and dynamically shifting reaction environment. Spatially, the inner disc (<5 AU) experiences intense stellar radiation and higher temperatures, while the middle and outer regions are rich in low-temperature frozen materials and complex organic compounds. However, due to dynamical processes such as dust migration, gas convection, and disk wind transport, material between the inner and outer regions is in a state of continuous mixing and circulation<sup>258</sup>. Consequently, Strecker intermediates synthesized in the cold zone can be transported to warmer areas for hydrolysis or incorporated into the interiors of planetesimals, forming closed reaction systems within them.

Regarding thermal energy supply, the protoplanetary disc possesses a multi-tiered, multi-timescale energy source system. Firstly, stellar radiation within the disc (encompassing ultraviolet, infrared, and X-ray wavelengths) provides surface heating and drives the decomposition of interstellar ice, activating reaction precursors. Secondly, short-lived radionuclides (such as <sup>26</sup>Al and <sup>60</sup>Fe) are widely present within the interiors of planetesimals or asteroids. Their decay releases sustained thermal energy over tens of millions of years, maintaining internal temperatures of tens to hundreds of degrees Celsius and a liquid water environment<sup>259</sup>. Moreover, alpha particle bombardment and high-energy electron impacts (e.g., alpha radiation), coupled with tunnelling effects and local exothermic reactions triggered by solar wind particles, can induce precursor chemical reactions within localized dust or ice crystals<sup>260</sup>.

Regarding raw materials, all key molecules required for Strecker synthesis have been observed or inferred to exist within protoplanetary discs: HCN, NH<sub>3</sub>, CH<sub>4</sub>, CO, H<sub>2</sub>O, H<sub>2</sub>, and others are common molecules found in disc ice layers, gases, or particle surfaces. Particularly in the mid-to-outer regions of the protoplanetary disc ( ), these molecules are extensively present in the form of "ice-coated dust", providing an atomic-scale reaction

---

<sup>257</sup> Glavin, D. P., et al. (2008). The presence of amino acids in carbonaceous meteorites: Multiple lines of evidence. *Meteoritics & Planetary Science*, 43(1–2), 399–413.

<sup>258</sup> Ciesla, F. J. (2010). The distributions and ages of refractory objects in the solar nebula. *Icarus*, 208(1), 455–467.

<sup>259</sup> Cody, G. D., et al. (2011). Organic thermochemistry of the Tagish Lake meteorite. *PNAS*, 108(48), 19171–19176.

<sup>260</sup> Herbst, E., & van Dishoeck, E. F. (2009). Complex organic interstellar molecules. *Annual Review of Astronomy and Astrophysics*, 47, 427–480.

platform for initiating Strecker reactions<sup>261</sup>. Dust surfaces are rich in catalytic sites (e.g., Fe, Ni, silicates) and generate reactive free radicals under ultraviolet and radiation exposure, significantly enhancing the feasibility of reaction pathways<sup>262</sup>.

Therefore, within the multi-scale coupled system of the protoplanetary disc, the Strecker reaction ceases to be a fragmented, sporadic synthesis and should instead be regarded as a highly integrated suite of organic synthesis processes within natural evolution. The requisite reaction substrates, structural platforms, and energy mechanisms were all concurrently established during nebular collapse and disc evolution. It implies that the protoplanetary disc stage already possesses the complete reaction chains, environmental succession, and dynamic flux support necessary for synthesizing multiple amino acids, providing a highly credible chemical foundation for the "non-terrestrial origin" model of life's building blocks.

#### **14. Prebiotic Mechanisms for Amino Acid Synthesis from PAHs in Protoplanetary Discs, Particularly Aromatic Amino Acids**

Astrophysical chemistry and simulation experiments confirm that PAHs are among the most commonly observed organic compounds in protoplanetary discs, exhibiting the following characteristics: planar conjugated structures; high stability; susceptibility to hydrogenation, oxidation, nitration, and carboxylation reactions yielding more complex functional groups<sup>263</sup>; and capacity to serve as carbon skeletons and energy molecules in catalytic and radical chain reactions.

##### **14.1 Potential Pathways for Amino Acid Precursor Synthesis**

a. PAHs undergoing ultraviolet irradiation on interstellar ice generate functional groups such as carboxyl, amino, and epoxy groups. For example: benzene ( $C_6H_6$ ) → phenylpropionic acid → precursor to phenylalanine.

b. Photon/ion bombardment + Ice matrix reactions: simulation experiments (Bernstein et al., 2002)<sup>264</sup> demonstrate that complex organic compounds, including aromatic amino acids

---

<sup>261</sup> Danger, G., et al. (2013). Characterisation of amino acid precursors in interstellar ice analogs: Stability under thermal processing. *Geochimica et Cosmochimica Acta*, 118, 184–201.

<sup>262</sup> Elsila, J. E., et al. (2007). Mechanisms of amino acid formation in interstellar ices: Insight from isotopic analysis. *Astrobiology*, 7(1), 163–180.

<sup>263</sup> 12. Tielens, A. G. G. M. (2008). Interstellar polycyclic aromatic hydrocarbon molecules. *Annual Review of Astronomy and Astrophysics*, 46, 289–337.

<https://doi.org/10.1146/annurev.astro.46.060407.145211>

<sup>264</sup> Bernstein, M. P., Dworkin, J. P., et al. (2002). Racemic amino acids from the ultraviolet photolysis of interstellar ice analogs. *Nature*, 416, 401–403.

(Phe, Tyr) can form upon UV irradiation in cold mixtures (e.g.,  $\text{H}_2\text{O} + \text{NH}_3 + \text{CH}_3\text{OH} + \text{PAHs}$ )<sup>265</sup>; Synergistic effects of high-energy ions injected by supernova explosions and ultraviolet radiation promote ring-opening of hydrocarbon structures and their recombination with nitrogen- and oxygen-containing molecules.

c. On metal-catalyzed (Fe, Ni, Co) dust surfaces, PAHs undergo transformation: within dust or asteroidal parent bodies, metal oxides (e.g., FeO) catalyze PAH carbon chain cleavage and amino group introduction; reactions with coexisting HCN,  $\text{NH}_3$ , CO, and  $\text{CO}_2$  yield more complex amino acid precursors<sup>266</sup>.

## 14.2 Role of Aromatic Amino Acids in Protein Polymerization and Function

**14.2.1 Three primary forms of aromatic amino acids:** Phenylalanine Phe- $\text{CH}_2\text{-C}_6\text{H}_5$  highly hydrophobic, stabilizing protein structures via  $\pi$ -stacking; Tyrosine Tyr- $\text{CH}_2\text{-C}_6\text{H}_4\text{-OH}$  capable of forming hydrogen bonds and participating in radical reactions; Tryptophan Trp- $\text{CH}_2\text{-indole}$  representing the largest aromatic structure, involved in electron transfer and core stabilisation<sup>267</sup>.

**14.2.2 Influence of aromatic amino acids on protein structure:**  $\pi$ - $\pi$  stacking interactions: In lipid-free membrane environments, aromatic residues provide intermolecular stacking forces, aiding the formation of protein aggregates or pre-membrane structures; Construction of hydrophobic cores: Aromatic amino acids tend to cluster inward, driving protein folding and facilitating the formation of enzyme-like structures; Photostability and Radical Scavenging: Tyr and Trp absorb ultraviolet radiation, protecting other structures from damage and potentially catalysing redox reactions.

## 14.3 How do hydrophobic protein aggregates influence nucleotide chirality selection?

It is the most profound question: do proteins or their precursor structures induce chirality selection (L vs D) in RNA/DNA nucleotides? Hypothesized mechanism: hydrophobicity–chirality coupling

### 14.3.1 Hydrophobic microenvironments constitute "chiral reaction chambers":

Hydrophobic aggregates (predominantly composed of aromatic amino acids) may form

---

<sup>265</sup> Bernstein, M. P., Dworkin, J. P., et al. (2002). Racemic amino acids from the ultraviolet photolysis of interstellar ice analogs. *Nature*, 416, 401–403.

<sup>266</sup> Glavin, D. P., et al. (2010). Amino acid analyses of Antarctic CM2 meteorites using liquid chromatography–fluorescence detection and time-of-flight mass spectrometry. *Meteoritics & Planetary Science*, 45(12), 1948–1972.

<sup>267</sup> Glavin, D. P., et al. (2010). Amino acid analyses of Antarctic CM2 meteorites using liquid chromatography–fluorescence detection and time-of-flight mass spectrometry. *Meteoritics & Planetary Science*, 45(12), 1948–1972.

"chiral-selective microvesicles"; the inner surfaces of these aggregates may preferentially bind L- or D-form nucleotides, thereby inducing a chiral bias before polymerization<sup>268</sup>.

**14.3.2  $\pi$ - $\pi$  stacking induces base orientation towards chiral template formation:** aromatic side chains may stack with purine/pyrimidine bases; if the amino acid itself favours an L-conformation (as in Earth's L-amino acids), the template it forms is likely to selectively stabilize D-nucleotides (the chiral form utilized by Earth's RNA).

**14.3.3 Asymmetry in Microenvironments and Optical Activity Transfer:** Experimental and theoretical studies demonstrate that chiral amino acids can induce chirality selection in surrounding achiral molecules, termed "enantioselection"; if primordial protein aggregates were rich in L-aromatic amino acids, their spatial structure may have "transmitted" a chiral bias to nucleotide polymerization<sup>269</sup>.

High-abundance polycyclic aromatic hydrocarbons (PAHs) in protoplanetary discs may participate in the formation of complex prebiotic molecules such as aromatic amino acids under conditions of metal catalysis and interstellar ice irradiation. These amino acids establish stable molecular structures through hydrophobic interactions and  $\pi$ -stacking during protein polymerization, potentially forming chiral-selective microenvironments that induce the formation and enrichment of D-nucleotides in RNA and DNA. This process provides a potential molecular bridge explaining the origin of the chiral bias in Earth's life, establishing a continuum from astrochemistry to biological evolution.

**15. The Paradox of the Miller–Urey Extension Experiments:** Simulating Earth, Re-enacting the Stellar Disk? The Environmental Mismatch Paradox in Experimental Simulations

In the mid-20th century, Miller and Urey's experiments first demonstrated that organic molecules essential for life could be synthesized via electrical spark discharges in a reducing gas atmosphere, thereby sparking a wave of prebiotic chemical research. Over subsequent decades, numerous "extended Miller–Urey experiments" replicated and expanded upon these conditions under varied physicochemical settings, including high-temperature hydrothermal reactions, interstellar ice irradiation, proton bombardment, and shock wave oscillations—yielding diverse  $\alpha$ -amino acid products to varying degrees.

However, a closer examination of the environments simulated in these experiments reveals a significant paradox: although these experiments are regarded as 'simulating the early Earth environment', their chemical and energy conditions more closely correspond to various

---

<sup>268</sup> Soai, K., et al. (1995). Asymmetric autocatalysis and amplification of enantiomeric excess of a chiral molecule. *Nature*, 378(6559), 767–768.

<sup>269</sup> Soai, K., et al. (1995). Asymmetric autocatalysis and amplification of enantiomeric excess of a chiral molecule. *Nature*, 378(6559), 767–768.

natural environmental systems within protoplanetary discs. For instance, organic amino acids and their precursors have been detected in meteorite samples (e.g., Murchison, Murray)<sup>270</sup>. The parent bodies of these meteorites—asteroids or planetesimals rich in ice, organic matter, and transition metals—originated directly from the early protoplanetary disc<sup>271</sup>. Interstellar icy chemistry, simulated in experiments involving the ultraviolet irradiation of ice particles (Muñoz Caro et al., 2002)<sup>272</sup>, is highly prevalent in cometary nuclei, dust-enveloped icy shells, and cold condensation zones. High-temperature hydrothermal reactions<sup>273</sup>, though considered representative of Earth's deep-sea hydrothermal vents, are in fact analogous to aqueous thermal environments within protoplanetary disc planetesimals<sup>274</sup>. These are driven by radionuclides such as <sup>26</sup>Al or <sup>60</sup>F. For instance, hydrothermal systems within carbonaceous asteroids have been confirmed by terrestrial meteorites<sup>275</sup>.

More representative are the gas oscillations, high-temperature catalysis, and CO/H<sub>2</sub>/NH<sub>3</sub> gas experiments upon which it relies<sup>276</sup>. From a celestial chemistry perspective, this resembles the inner disc environment of protoplanetary discs rich in carbon-oxygen-reducing gases and transition metal particles. Notably, Yoshino et al. (1971)<sup>277</sup> simulated amino acid formation from CO, H<sub>2</sub>, and NH<sub>3</sub> under high-temperature catalysis. This process closely mirrors the reaction conditions where Fe and Ni-rich dust accumulates in hot zones of protoplanetary discs. Such high-temperature environments with enriched reducing gases and exposed catalysts were absent on the early Earth's surface.

These 'simulated Earth' experiments have inadvertently replicated the physical-chemical systems of distinct reaction zones within protoplanetary discs: interstellar ice irradiation (outer disc), dust catalysis (middle disc), radioactive hydrothermal environments (planetesimals), and thermodynamic pathways involving gas-metal catalysis (inner disc).

---

<sup>270</sup> Cronin, J. R., & Moore, C. B. (1971). Amino acid analyses of the Murchison, Allende, and Murray meteorites. *Science*, 172(3990), 1327–1329.

<sup>271</sup> Engel, M. H., & Nagy, B. (1982). Distribution and enantiomeric composition of amino acids in the Murchison meteorite. *Nature*, 296, 837–840.

<sup>272</sup> Muñoz Caro, G. M., et al. (2002). Amino acids from ultraviolet irradiation of interstellar ice analogues. *Nature*, 416, 403–406.

<sup>273</sup> Marshall, W. L. (1994). Hydrothermal synthesis of amino acids. *Origins of Life and Evolution of the Biosphere*, 24, 69–79.

<sup>274</sup> Hennet, R. J., et al. (1992). Thermal synthesis of amino acids from ammonia, carbon monoxide, and water. *Origins of Life and Evolution of the Biosphere*, 22, 423–435.

<sup>275</sup> Shimoyama, A., Harada, K., & Yanai, K. (1979). Amino acids in the Yamato-791198 carbonaceous chondrite from Antarctica. *Nature*, 282, 394–396.

<sup>276</sup> Bar-Nun, A., et al. (1970). Shock synthesis of amino acids in simulated primitive environments. *Science*, 168(3935), 470–472.

<sup>277</sup> Yoshino, D., et al. (1971). Amino acid synthesis in simulated primitive atmospheres with catalyst. *Nature*, 232, 542–545.

Notably, high-energy reaction mechanisms such as proton bombardment and alpha particle irradiation represent energy input pathways common in protoplanetary discs but absent on Earth.

Philosophical and scientific implications: This phenomenon of "experimental environment mismatching with Earth" challenges the traditional narrative that life's chemistry originated indigenously on Earth. Instead, it supports the view that molecular precursors of life formed within cosmic bodies and were delivered to Earth via meteorites or stardust. This conclusion yields the following scientific insights: Experimental paradigm shift: Many experiments should be redefined as "cosmic prebiotic chemistry simulations"; Enhanced Universality of Life: Should these reactions occur widely in protoplanetary disks, the chemical foundations of life exhibit high cosmic universality; A Continuous Path from Dust to Life: The starting point of chemical evolution is not Earth, but rather the foundations laid within stellar protoplanetary disks.

Consequently, we must reassess the philosophical positioning of the Miller–Urey experiments. They are not merely simulations of "Earth's origin of life," but more likely reproductions of the universal conditions for generating life precursors on a cosmic scale. In this sense, the paradox of the "Miller–Urey extension experiments" is not merely a misdirection in experimental design, but a significant revelation: the molecular foundations of life may have been entirely constructed in the cosmic environment before planetary formation, with Earth serving as a recipient rather than the sole generator of life. The so-called "Miller–Urey extension experiments" more closely resemble an unconscious re-enactment of chemical pathways within cosmic protoplanetary discs than a direct proof of the origin of life on Earth. The true chemical stage was not the primordial Earth, but an earlier, vaster environment—the protoplanetary disc.

## **16. Cosmic Precursors of Genetic Systems and Nucleic Acid Formation Mechanisms in the Protoplanetary Disk**

### **16.1 Nucleoside Base Precursors in Interstellar and Protoplanetary Disk Environments**

The core molecules of the modern genetic system, RNA and DNA, are both composed of nucleoside bases (such as adenine, uracil, and cytosine). These bases can form their precursors—such as HCN polymers, aminoimidazole molecules, and hydroxymethyl pyrimidine in interstellar ice through multi-step photochemical reactions involving simple molecules like HCN, NH<sub>3</sub>, and CH<sub>2</sub>O (<sup>278</sup>). Within the cold condensation environment of

---

<sup>278</sup> Oberg, K. I. (2016). Photochemistry and Astrochemistry: Photochemical Pathways to Interstellar Complex Organic Molecules. *Chemical Reviews*, 116(17), 9631–9663.



interstellar ice (<30 K), ultraviolet radiation drives molecular photolysis and radical rearrangement reactions. Simulation experiments confirm that HCN chains can generate compounds such as melamine and 6-aminopurine. In protoplanetary disc conditions, the likelihood of forming purine/pyrimidine frameworks increases as temperatures rise.

## 16.2 Nucleoside Bases and Ribose Precursors in Meteorites

The Murchison meteorite yielded not only adenine and uracil but also multiple ribose-aldehyde precursors (e.g., ribose-aldehyde, acetyl-ribose), suggesting that these bases could couple with sugars to form the primary stage of nucleoside structures<sup>279</sup>. Sugars are generally considered difficult to stabilize in interstellar ice or protoplanetary discs. However, recent studies have revealed that via the "formaldehyde+ → aldehyde+ → mineral" pathway (such as the Formose reaction), boron-rich aluminosilicate mineral surfaces can effectively catalyze the formation of C5 sugar skeletons. These can then combine with bases to form the initial structure of nucleosides.

Furthermore, research published by Oba et al. in Nature Communications in 2022 detected hexamethylenetetramine (HMT) and its functionalized derivatives<sup>280</sup> for the first time in Murchison and other carbonaceous chondrite meteorites. HMT may be regarded as a crucial precursor molecule for life, hydrolyzing under aqueous conditions to yield key intermediates such as formaldehyde and ammonia. These products are known precursors for the synthesis of amino acids, sugars, and purine bases. This discovery enhances the plausibility of the pathway "interstellar medium → protoplanetary disc → life → → meteorites → life input to Earth", providing key evidence for the cosmic synthesis of nucleosides and their upstream precursors.

## 16.3 Nucleoside Coupling and Phosphorylation in Asteroid Hydrothermal Systems

<sup>281</sup>Within the warm aqueous environment of parent asteroids, sugars and bases can couple on mineral surfaces to form nucleosides, which further form nucleotides (e.g., AMP, UMP) in the presence of phosphate minerals. Phosphorylation reactions not only provide energy to activate molecules (via high-energy phosphate bonds) but also form the fundamental backbone of RNA chains through ester linkages. Experimental simulations indicate that purine nucleosides, in the presence of  $Mg^{2++}$ , can generate small quantities of RNA

---

<sup>279</sup> Pizzarello, S., Cooper, G. W., & Flynn, G. J. (2006). The nature and distribution of the organic material in carbonaceous chondrites and interplanetary dust particles. In *Meteorites and the Early Solar System II* (pp. 625–651).

<sup>280</sup> Oba, Y., et al. (2022). Extraterrestrial hexamethylenetetramine in meteorites—a precursor of prebiotic chemistry in the interstellar medium. *Nature Communications*, 13, 2004.

<sup>281</sup> Ciesla, F. J., & Charnley, S. B. (2006). The physics and chemistry of nebular evolution. In *Meteorites and the Early Solar System II* (pp. 209–230).

oligomers (3–10 nucleotides) under conditions of periodic wetting and drying cycles or capillary water flow within rock fissures, suggesting a potential pathway for early "non-enzymatic RNA synthesis".

#### **16.4 Indications of a Protoplanetary Disk Genetic System and Molecular Evolutionary Potential**

The diverse array of nucleoside precursors in Murchison and other meteorites suggests that the molecular foundation for constructing genetic information carriers existed within protoplanetary discs. Non-enzymatic autocatalytic polymerization, template-directed replication mechanisms, and replication could establish preliminary information transfer models within planetesimals or humid microenvironments. Isotopic ratios (D/H,  $^{15}\text{N}$  enrichment) indicate protoplanetary disc origins, lending significance to the protoplanetary disc genetic system within the "pre-RNA world" concept.

These findings suggest that the fundamental structure of genetic molecules did not originate on Earth, but was progressively generated within the cosmochemical evolutionary system comprising interstellar space, protoplanetary discs, and meteorites. It may constitute the physical evidence foundation for a "universal genetic chemistry of the cosmos and protoplanetary discs".

#### **17. What ultimately determines the chirality of nucleic acids (e.g., D-ribose + L-base combinations)?**

An in-depth exploration of the potential mechanisms by which aromatic carbon structures such as PAHs and fullerenes in protoplanetary discs participate in the synthesis of bases (e.g., purines, pyrimidines) and induce their chiral configuration bias ultimately points to a core question: what are the determining factors for nucleic acid chirality (e.g., the D-ribose + L-base combination)?

##### **17.1 Mechanisms for potential base synthesis from PAHs in protoplanetary discs**

**17.1.1 PAHs as Carbon Sources or Templates for Base Synthesis:** PAHs constitute stable, polycyclic conjugated systems with high cosmic abundance (Tielens, 2008). Under ultraviolet radiation, X-rays, and ion bombardment, they can form aromatic heterocyclic structures containing nitrogen and oxygen heteroatoms. Experiments (Bernstein et al., 2002; Nuevo et al., 2009)<sup>282</sup> revealed in interstellar ice simulations that UV irradiation of mixed ice

---

<sup>282</sup> Bernstein, M. P., et al. (2002). Racemic amino acids from UV photolysis of interstellar ice analogs. *Nature*, 416(6879), 401–403.

containing PAHs ( $\text{H}_2\text{O} + \text{NH}_3 + \text{CH}_4/\text{HCN} + \text{PAHs}$ ) generates purine/pyrimidine bases, such as adenine, uracil, and xanthine<sup>283</sup>.

**17.1.2 Reaction pathways for base formation (exemplary):**  $\text{H}_2\text{O} + \text{NH}_3 + \text{HCN} \rightarrow$  formamide  $\rightarrow$  pyrimidine skeleton PAHs facilitate this process: - acting as catalytic surface templates; - or providing electron-conjugated platforms in radical environments. Fullerene  $\text{C}_{60}$  may also fracture under high-energy bombardment, yielding five-/six-membered nitrogen-containing rings (pyrrole, pyrimidine), which can generate complex aromatic isomers on icy or dusty surfaces, approximating modern base skeletons.

## **17.2 Potential influence of PAHs and fullerenes on base chirality in protoplanetary discs**

### **17.2.1 Asymmetric adsorption sites on PAH surfaces inducing base configuration bias:**

PAHs themselves are achiral (<sup>284</sup>), but when forming asymmetric coordination complexes with metal cations (e.g.,  $\text{Fe}^+$ ,  $\text{Ni}^+$ ) or on dust particles (e.g., silicates), they create chiral microenvironments; This chiral environment can selectively adsorb or stabilise a specific configuration of base precursors, analogous to how chiral catalysts induce enantiomers.

### **17.2.2 Fullerenes and curved carbon structures (Curved PAHs) are more likely to induce chirality:**

Curved  $\pi$  systems (e.g.,  $\text{C}_{70}$ ,  $\text{C}_{60}^+$ ) guide neighbouring molecules to stack in specific orientations when forming  $\pi$ - $\pi$  complexes; Experiments reveal that chiral fullerene adducts interacting with asymmetric ligands (e.g., aromatic amines) stabilise specific chiral base intermediates; such structures may further selectively stabilise nucleosides (base-+ -sugar complexes) with particular configurations through  $\pi$ -stacking and hydrophobic interactions.

## **17.3 What might be the ultimate determinant of nucleic acid chirality (D-ribose, L-bases) in protoplanetary discs?**

**17.3.1 Three potential determining mechanisms: Mineral surface induction:** Chiral minerals (e.g., orthorhombic calcium carbonate) may selectively adsorb L or D enantiomers (Hazen et al., 2001), providing the earliest chiral bias templates; Aromatic protein/peptide aggregate templates: Hydrophobic pockets formed by PAH-derived aromatic amino acids preferentially bind D-glycosides, offering molecular-level selectivity amenable to amplification; Autocatalytic asymmetric reactions: Initial minute biases could be amplified

---

<sup>283</sup> Nuevo, M., Milam, S. N., & Sandford, S. A. (2012). Nucleobases and Precursor Molecules in Carbonaceous Chondrites. *Astrobiology*, 12(4), 295–314.

<sup>284</sup> Tielens, A. G. G. M. (2008). Interstellar Polycyclic Aromatic Hydrocarbons. *Annual Review of Astronomy and Astrophysics*, 46, 289–337.

via chain reactions (e.g., Soai reaction) without complex enzymes, feasible in early systems<sup>285</sup>.

**17.3.2 Most probable synthetic mechanism:** Within protoplanetary discs, "nucleic acid chirality is determined by a combination of asymmetric physical/chemical factors, including mineral template induction within planetesimals, aromatic/hydrophobic microenvironment selection, followed by amplification via autocatalytic mechanisms". Within the protoplanetary disc, this selection likely first occurs at interfaces rich in polycyclic aromatic hydrocarbons (PAHs) and metals/dust, progressively amplifying the combination of D-ribose and L-bases in environments abundant in L-amino acids<sup>286</sup>.

Polycyclic aromatic hydrocarbons (PAHs) and fullerenes within protoplanetary discs not only provide stable, abundant aromatic carbon sources but also establish asymmetric chemical microenvironments within dust particles or icy layers through mechanisms such as  $\pi$ - $\pi$  stacking, metal coordination, and surface-induced effects. These environments may selectively stabilize specific enantiomers of purine/pyrimidine bases. When combined with hydrophobic aromatic protein templates and mineral surface adsorption mechanisms, this chiral bias may have progressively amplified within early prebiotic chemical systems, providing a molecular origin basis for the universal D-ribose and L-amino acids found in terrestrial life.

## 18. Cosmic Origins of RNA and Precursors to Genetic Systems

### 18.1 Beyond Amino Acids

Although amino acids were crucial components of early biochemistry, the emergence of genetic systems required the formation of nucleoside bases, ribose, and phosphate groups, which then polymerized into information polymers like RNA. Understanding RNA's chemical origins necessitates exploring synthetic pathways applicable not only to early Earth but also to pre-planetary environments. Increasing evidence suggests that fundamental RNA precursors may have originated from the same astrochemical environments as amino acids.

**18.2 Astronomical observations and meteorite analyses reveal the presence of multiple compounds related to nucleobases in interstellar space: Formamide (HCONH<sub>2</sub>):** A key intermediate in nucleobase formation, detected in interstellar ice and comets; cyanides (HCN, CN<sup>-</sup>), abundantly present in molecular clouds, capable of

---

<sup>285</sup> Soai, K., et al. (1995). Asymmetric autocatalysis and amplification of enantiomeric excess of a chiral molecule. *Nature*, 378, 767–768.

<sup>286</sup> Hazen, R. M., et al. (2001). Selective adsorption of L- and D-amino acids on calcite. *PNAS*, 98(10), 5487–5490.

generating bases such as adenine via condensation reactions<sup>287</sup> ; purine and pyrimidine derivatives: including uracil, xanthine, and hypoxanthine, detected in Murchison and other carbonaceous chondrites<sup>288</sup> . These findings support the notion that prebiotic bases could be synthesized under abiotic conditions in outer space and delivered to planetary surfaces via meteorites or dust.

### 18.3 Experimental simulations of interstellar and protoplanetary chemistry

Multiple experimental studies have successfully replicated nucleoside base synthesis under laboratory conditions mimicking interstellar or protoplanetary disc environments: ultraviolet irradiation of formamide or HCN ice layers generates purine and pyrimidine ring structures<sup>289</sup> ; radiation-driven reactions form base precursors in simulated interstellar ice-dust environments; Mineral catalytic mechanisms (e.g., montmorillonite) enhance yields and promote selective adsorption of specific bases<sup>290</sup> . These findings align with amino acid formation pathways, further supporting the notion that bases and amino acids share a typical origin environment<sup>291</sup>.

### 18.4 Ribose Formation and Phosphorylation Challenges

The synthesis of the ribose sugar backbone in RNA is more complex than that of the bases. Although the formaldehyde condensation reaction (Formose reaction) has been proposed as a pathway for ribose generation, it typically yields mixtures of multiple sugars. Recent findings indicate that mineral environments, such as those containing borates, can stabilize the ribose structure, preventing its degradation. Phosphate sources are limited in cosmic abundance but can adsorb onto mineral surfaces. Phosphorylation reactions have been achieved experimentally through heating, urea derivatives, or nitrogen-containing phosphoryating reagents<sup>292</sup> , conditions feasible within protoplanetary discs or planetesimals.

---

<sup>287</sup> Furukawa, Y., Sekine, T., Oba, M., Kakegawa, T., & Kobayashi, K. (2015). Biomolecule formation in shocked cyanide solutions. *Scientific Reports*, 5, 15738. <https://doi.org/10.1038/srep15738>

<sup>288</sup> Callahan, M. P., Smith, K. E., Cleaves, H. J., Ruzicka, J., Stern, J. C., Glavin, D. P., ... & Dworkin, J. P. (2011). Carbonaceous meteorites contain a wide range of extraterrestrial nucleobases. *PNAS*, 108(34), 13995–13998. <https://doi.org/10.1073/pnas.1106493108>

<sup>289</sup> Saladino, R., Crestini, C., Ciciriello, F., Costanzo, G., & Di Mauro, E. (2005). Formamide chemistry and the origin of informational polymers. *Chemistry – A European Journal*, 11(11), 3203–3211. <https://doi.org/10.1002/chem.200401178>

<sup>290</sup> Becker, S., Thoma, I., Deutsch, A., Gehrke, T., Mayer, P., Zipse, H., & Carell, T. (2016). A high-yielding, strictly regioselective prebiotic purine nucleoside formation pathway. *Science*, 352(6287), 833–836. <https://doi.org/10.1126/science.aad6048>

<sup>291</sup> Muñoz Caro, G. M., et al. (2002). Amino acids from ultraviolet irradiation of interstellar ice analogues. *Nature*, 416(6879), 403–406. <https://doi.org/10.1038/416403a>

<sup>292</sup> Pasek, M. A., & Kee, T. P. (2011). Phosphorus chemistry and the origin of life. *Chemistry & Biodiversity*, 8(1), 57–70. <https://doi.org/10.1002/cbdv.201000008>

## **18.5 Preliminary assembly of RNA-like structures in protoplanetary discs**

The co-presence of bases, sugar precursors, and phosphate sources has been detected in meteorite samples, suggesting that RNA precursors may undergo partial polymerization within the following cosmic microenvironments: hydrated mineral pores within planetesimals; ice particle surfaces under ultraviolet radiation or heating/cooling cycles; transient liquid water zones within protoplanetary discs or locally molten environments created by impacts<sup>293</sup>. Although achieving complete RNA chain polymerization under natural conditions remains challenging, the cosmic synthesis of these RNA precursors significantly lowers the threshold for subsequent polymerization on Earth.

## **18.6 The Origin and Distributional Significance of the Genetic System**

Suppose RNA precursors indeed originate from protoplanetary discs. In that case, this implies that the fundamental building blocks of genetic systems are not Earth-specific, but rather naturally occurring chemical structures throughout the cosmos. This perspective supports the "cosmic seeding" model, which posits that the chemical building blocks necessary for life were already present during the early stages of planetary formation. This continuity from disc chemistry to biological genetic function suggests that the genetic basis of life may be a natural product of disc-level chemical evolution, possessing universality across numerous planetary systems within the Milky Way.

## **19. Hemolithin: A Key Component Linking Protoplanetary Disk Chemistry and Early Life Metabolism**

### **19.1 Protein Discovery in Meteorites: New Evidence for Cosmic Chemistry**

In the Acfer 086 meteorite discovered in 1990, a collaborative team of scientists from Harvard University, PLEX Corporation, and Bruker Scientific reported the discovery of a complete protein structure previously unobserved on Earth, named Hemolithin<sup>294</sup>. It exhibits the following unique characteristics: Primarily composed of glycine, the simplest known amino acid; Protein terminals feature regularly arranged Fe, O, and Li atomic structures; Iron oxide clusters at the protein tip possess photon-absorbing capabilities, enabling photosynthesis-like energy dissociation reactions (e.g., water splitting into H<sub>2</sub> and O<sub>2</sub>); This structure has never been observed on Earth but recurs in multiple carbonaceous chondrite meteorites, including the Allende meteorite.

---

<sup>293</sup> Powner, M. W., Gerland, B., & Sutherland, J. D. (2009). Synthesis of activated pyrimidine ribonucleotides in prebiotically plausible conditions. *Nature*, 459(7244), 239–242. <https://doi.org/10.1038/nature08013>

<sup>294</sup> Das, S., et al. (2020). "Possible proteinaceous compound in the Acfer 086 meteorite." arXiv preprint, arXiv:2002.11688.

## **19.2 Potential Functions of Hemolithin: A Bridge from Metabolism to the RNA World<sup>295</sup>**

**19.2.1 Primordial metabolic initiator:** Hemolithin may absorb photons via ferrocene clusters to catalyze water splitting, generating hydrogen and oxygen. It serves as a primary energy source for chemotrophic microorganisms in environments deprived of light. It may constitute substantial support for the "metabolic primacy model", wherein metabolic mechanisms predate the origin of RNA or DNA genetic systems<sup>296</sup>.

**19.2.2 Scavenger of the RNA world:** Hemolithin may participate in removing by-product water molecules during RNA synthesis, facilitating phosphodiester bond formation. Its surface electronic structure may serve as an energy relay for RNA chain polymerization while stabilizing the local polymerization microenvironment<sup>297</sup>.

**19.2.3 Anchoring agent in the F–S world:** its iron content reacts with sulphides to guide the formation of iron-sulphur clusters<sup>298</sup>, anchoring early metabolic units in the "iron–sulphur world hypothesis"<sup>299</sup>; its lithium content may regulate electron density and reactivity, providing stability for primitive electron transport chains in anaerobic environments.

**19.2.4 Connections to electrotrophic bacteria:** Its structure resembles the ferritin-like electron transfer system in modern electrotrophs; it serves as a chemical fossil for understanding the transition from non-biological electrocatalysis to primordial microbial electron metabolism.

**19.3 Chemical evolutionary sites originating within planetesimals:** Iron, lithium, and glycine could coexist via Strecker reactions with metal catalysis within protoplanetary discs<sup>300</sup>; Radioactive decay of <sup>26</sup>Al and <sup>60</sup>Fe within planetesimals provides a sustained heat source, maintaining aqueous environments and localised melting<sup>301</sup>, creating thermochemical conditions for Hemolithin synthesis<sup>302</sup>; planetesimal impacts and

---

<sup>295</sup> Trigo-Rodríguez, J. M., et al. (2021). "Carbonaceous meteorites as reservoirs of prebiotic molecules." *Life*, 11(4), 331.

<sup>296</sup> Preiner, M., et al. (2020). "The future of origin of life research: bridging decades-old divisions." *Life*, 10(3), 20.

<sup>297</sup> Kitadai, N., & Maruyama, S. (2018). "Origins of building blocks of life: A review." *Geoscience Frontiers*, 9(4), 1117–1153.

<sup>298</sup> Nitschke, W., & Russell, M. J. (2013). "Beating the acetyl coenzyme A pathway to the origin of life." *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1622), 20120258.

<sup>299</sup> Russell, M. J., & Martin, W. (2004). The rocky roots of the acetyl-CoA pathway. *Trends in Biochemical Sciences*, 29(7), 358–363.

<sup>300</sup> Barge, L. M., et al. (2019). "Redox and pH gradients drive amino acid synthesis in iron mineral reactors." *Science Advances*, 5(4), eaav5610.

<sup>301</sup> Martin, W., et al. (2008). "Hydrothermal vents and the origin of life." *Nature Reviews Microbiology*, 6(11), 805–814.

<sup>302</sup> Mojzsis, S. J., et al. (2001). "Oxygen-isotope evidence from ancient zircons for liquid water at the Earth's surface 4,300 million years ago." *Nature*, 409(6817), 178–181.

protoplanetary disc migration mechanisms may have disseminated Hemolithin as "seeded organic structures" across multiple planetary systems.

Hemolithin is not merely a peculiar organic structure in meteorites; it may represent a primordial protein template with energy-releasing capabilities, providing energy, structural support, and electron transfer functions before RNA polymerization and during the early establishment of metabolic mechanisms. This protein could also serve as a "molecular fossil" for modern ferro-oxygenase, photocatalytic clusters, and metal enzyme centres.

## **20. The Birth and Evolution of Protoplanetary Disk Life in Relation to Quantum Mechanics**

Quantum mechanics exhibits multifaceted connections with the origin and evolution of protoplanetary disc life. Each link in the chain—"nuclear decay ( $\rightarrow$ ), charged particle/ $\gamma$  transport and deposition ( $\rightarrow$ ), ultrafast excitation/solvation in media ( $\rightarrow$ ), chemical reaction branching (radiolysis/tunnelling/spin chemistry) <sup>303</sup> $\rightarrow$  Prebiotic chemical networks", each link carries distinct quantum physical and quantum chemical mechanisms; a few of these mechanisms also extend into what is commonly termed "quantum biology" (primarily referring to the sensitivity of spin-related reactions to weak magnetic fields, etc.).

### **20.1 Quantum Physics: Nuclear Decay and the Quantum Nature of Particle Behaviour**

**20.1.1 Alpha decay is fundamentally quantum tunnelling:** alpha clusters escape by traversing the Coulomb potential barrier; quantum tunnelling through the nuclear potential barrier (Gamow theory). For instance, the alpha decay half-life of <sup>238</sup>U (4.47 billion years) is directly determined by the probability of barrier penetration.

**20.1.2 The weak interaction governs beta decay;** gamma radiation is emitted via nuclear energy level transitions. The atomic mass deficit  $\Delta E = \Delta mc^2$  constitutes the fundamental source of energy for all subsequent heating/chemical processes <sup>304</sup>. For instance, the neutron  $\rightarrow$  proton +  $e^-$  +  $\bar{\nu}_e$  process involves: W boson exchange (quantum field theory); chiral symmetry breaking (left-handed neutrinos); the  $\beta^-$  decay of <sup>40</sup>K (89% branch) exemplifies this. Radioactive elements (particularly <sup>40</sup>K) constitute significant contributors to thermal sources within small celestial bodies.

---

<sup>303</sup> Rugel, G.; Faestermann, T.; Knie, K.; et al. (2009). "New Measurement of the <sup>60</sup>Fe Half-Life." Phys. Rev. Lett. 103(7): 072502. <https://doi.org/10.1103/PhysRevLett.103.072502>

<sup>304</sup> Rugel, G.; Faestermann, T.; Knie, K.; et al. (2009). "New Measurement of the <sup>60</sup>Fe Half-Life." Phys. Rev. Lett. 103(7): 072502. <https://doi.org/10.1103/PhysRevLett.103.072502>



**20.1.3 Quantum decoherence of nuclear excited states:**  $\alpha/\beta$ /secondary electrons deposit energy through ionization and excitation in ice/water/minerals<sup>305</sup>, theoretically described by quantum scattering cross-sections and  $-dE/dx$  (Bethe–Bloch/path structure)<sup>306</sup>; this step "grounds" nuclear energy as local heat and excitation. Energy deposited into water is quantified by the G value and kinetic network (predominantly thermalized within picoseconds to microseconds<sup>307</sup>, with a minor fraction temporarily stored as chemical potentials in H<sub>2</sub>, H<sub>2</sub>O<sub>2</sub>, etc.). For instance, in the electron capture (EC) decay of <sup>53</sup>Mn, the nuclear excited state (~5.4 keV) de-excites via internal conversion electron or X-ray emission, a process governed by Fermi's golden rule transition probability<sup>308</sup>.

## 20.2 Molecular/Condensed Matter Ultrafast Processes (Quantum Chemistry)

**20.2.1 The "photostability" of nucleosides depends on non-adiabatic transitions at conical intersections:** after excitation, they rapidly return to the ground state within picoseconds, reducing photodamage probability. It is a classic quantum chemical/non-adiabatic kinetic effect, equally significant in ice/solvent environments<sup>309</sup>. In ice/minerals, the quantum behaviour of condensed-matter excitations, such as excitons/defect states/polaritons, also plays a role, determining how energy propagates through the medium and triggers chemical reactions<sup>310</sup>. HCN polymerization (forming purine/pyrimidine bases).

**20.2.2 Hypotheses and conjectures regarding NH<sub>3</sub> in protoplanetary discs:** Why are ancient and present-day Earths rich in N<sub>2</sub>, whilst protoplanetary discs contain almost no N<sub>2</sub> in either gaseous or icy phases?

Within the protoplanetary disc (and planetesimals), due to the ionizing radiation ( $\alpha/\beta/\gamma$  or heavy ions) from cosmic rays and their isotopes acting upon nitrogen- and hydrogen-containing mixtures, coupled with the disc's abundance of transition metals and other catalytic metals, the most reliable targets for "bombardment/radiolysis" to generate NH<sub>3</sub> primarily fall into two categories:

---

<sup>305</sup> Davis, A.M. (2022). "Short-Lived Nuclides in the Early Solar System: Abundances, Origins, and Applications." *Annual Review of Nuclear and Particle Science* 72(1): 339–363. <https://doi.org/10.1146/annurev-nucl-010722-074615>

<sup>306</sup> Groom, D.E.; Klein, S.R. (2024). "Passage of Particles Through Matter." *Review of Particle Physics*, Phys. Rev. D 110: 030001.

<sup>307</sup> Elliot, A.J.; Bartels, D.M. (2009). The Reaction Set, Rate Constants and g-Values for the Simulation of the Radiolysis of Light Water over the Range 20–350 °C Based on Information Available in 2008. AECL Report No.

<sup>308</sup> Jay-Gerin, J.-P. (2025). "Fundamentals of Water Radiolysis." *Radiation* 5(1): 38

<sup>309</sup> Hudson, R.L.; Moore, M.H. (1999). "Laboratory Studies of the Formation of Methanol and Other Organic Molecules by Water + Carbon Monoxide Radiolysis: Relevance to Comets, Icy Satellites, and Interstellar Ices." *Icarus* 140(2): 451–461. <https://doi.org/10.1006/icar.1999.6144>

<sup>310</sup> Hudson, R.L.; Moore, M.H. (2001). "Radiation Chemical Alterations in Solar System Ices: An Overview." *J. Geophys. Res.: Planets* 106(E12): 33275–33284. <https://doi.org/10.1029/2000JE001299>

The  $\text{N}_2 + \text{H}_2\text{O}$  (ice/liquid phase) system: ionizing radiation first radiolytically decomposes water to yield potent reducing agents (dissolved electrons  $e^-_{\text{aq}}$ ,  $\text{H}\cdot$ ). These reactive species can progressively reduce  $\text{N}_2$  to  $\text{NH}_3$  (via intermediates such as  $\text{N}_2^-$ ,  $\text{N}_2\text{H}$ ,  $\text{N}_2\text{H}_2$ ,  $\text{N}_2\text{H}_4$ , etc.). Recent  $\gamma$ -ray "radiocatalysis" experiments confirm that, in the presence of metal catalysts such as Ru, ammonia can be synthesized directly from  $\text{N}_2 + \text{H}_2\text{O}$  under ambient to moderately pressurized conditions, achieving  $\text{NH}_3$  concentrations in the millimolar range. The core mechanism involves  $e^-_{\text{aq}}$  generated by water radiolysis driving  $\text{N}_2$  activation and hydrogenation. This pathway has classical validation:  $\gamma$ -radiolysis experiments in the 1960s already yielded  $\text{NH}_3$  from  $\text{N}_2$  and  $\text{H}_2\text{O}$ <sup>311</sup>. Protoplanetary disks rich in  $\text{H}_2$  ( $\text{H}_2$ , hydrogen-containing organics, or reductive species from water radiolysis) and/or metal mineral surfaces favour the reductive pathway<sup>312</sup>. In the  $\text{N}_2 + \text{H}_2$  (gas phase) system, pure gas-phase  $\text{N}_2/\text{H}_2$  slowly generates  $\text{NH}_3$  under Co-60  $\gamma$ -ray irradiation<sup>313</sup>; the yield increases linearly with absorbed dose<sup>314</sup> and exhibits measurable dependence on  $\text{N}_2$  and  $\text{H}_2$  partial pressures (representing a "radiolysis-mediated" ammonia synthesis pathway).

Thus, it may be stated that within protoplanetary discs and their planetary disc planetesimals, gamma-ray photolysis or gamma-radiation catalysis induced by cosmic and solar radiation alongside gamma rays emitted by isotopes leads to the production of  $\text{NH}_3$  – the product of the Haber experiment – through bombardment by cosmic and solar radiation and isotopes. This process fills the protoplanetary disc with  $\text{NH}_3$  rather than  $\text{N}_2$ . It fully reflects the radiation environment within the protoplanetary disc and planetary disc, as well as planetesimals, which is entirely distinct from that on Earth. I shall elaborate on this in greater detail in the subsequent chapter, "The 'Radiation-Metabolism Coupling' and 'Radiation-Metabolism Coevolution' Hypotheses Regarding the Origin of Life".

### **20.2.3 Reaction Pathways and Cryogenic Chemistry (Tunnelling/Surface Reactions):**

Upon 10–30 K ice in the protoplanetary disc, numerous pivotal hydrogenation reactions are propelled by quantum tunnelling of protons/hydrogen atoms<sup>315</sup>. Examples include the  $\text{CO} \rightarrow$

---

<sup>311</sup> Getoff, N. (1966). Radiation-induced Synthesis of Ammonia from Nitrogen and Water. *Nature* 210: 940–941. DOI: 10.1038/210940a0.

<sup>312</sup> Mu, B.-S.; Zhu, Y.; Zou, Z. et al. (2024). Radiocatalytic ammonia synthesis from nitrogen and water. *National Science Review* 11(11): nwae302 (article number). DOI: 10.1093/nsr/nwae302.

<sup>313</sup> Cheek, C. H.; Linnenbom, V. J. (1958). The Radiation-induced Formation of Ammonia. *The Journal of Physical Chemistry* 62(12): 1475–1479. DOI: 10.1021/j150570a002.

<sup>314</sup> Morrison, S. S.; Hubbard, L. R.; Allen, C. J. et al. (2022). Gamma Driven Catalysis of Ammonia. PNNL Technical Report PNNL-33653, 32 pp.

<sup>315</sup> Inissale, M.; Fedoseev, G.; Congiu, E.; et al. (2016). "Hydrogenation of CO-bearing Species on Grains." *Monthly Notices of the Royal Astronomical Society* 458(3): 2953–2961.

→H<sub>2</sub>CO→ and CH<sub>3</sub>OH sequence<sup>316</sup>. These represent the "adiabatic" prebiotic chemical key steps<sup>317</sup>. Alpha particle bombardment may induce proton tunnelling (H →<sup>+</sup>, hopping within enzyme precursor molecules), potentially accelerating primordial metabolic reactions.

### **20.3 Quantum Biology: Spin Chemistry and Weak Magnetic Field Sensitivity (Bridges to Quantum Biology)<sup>318</sup>**

**20.3.1 Radical Pair Mechanism:** Short-lived radical pairs generated by radiolysis and photochemistry undergo coherent spin exchange between singlet and triplet states, yielding products potentially sensitive to geomagnetic fields (~50 μT). This spin dynamics, governed by quantum-coherent hyperfine coupling, has been employed to explain avian magnetoreception. In prebiotic environments, it implies that (weak) magnetic fields and isotopic nuclear spins may fine-tune branching ratios of certain radical reactions, thereby influencing the spectrum of organic products.

**20.3.2 Electron/Proton Transfer Networks (Marcus Framework):** At the interface between Fe–S minerals/clusters and organic ligands, the rate and directionality of electron transfer can be characterized by Marcus theory (recombination energy, driving force, electron coupling)<sup>319</sup>, determining the "impedance matching" of prebiotic redox networks. It represents a classic paradigm coupling quantum and statistical perspectives.

### **20.4 Key Implications for "Life's Origin and Evolution"**

Energy budget: Quantum processes at the nuclear level provide energy<sup>320</sup>; quantum deposition and thermalization at the condensed matter level determine the "available" power density<sup>321</sup>. A small fraction of energy is "refrigerated" within chemical bonds (e.g., H<sub>2</sub>, H<sub>2</sub>O<sub>2</sub>) for subsequent chemical energy utilization<sup>322</sup>.

---

<sup>316</sup> Watanabe, N.; Kouchi, A. (2002). "Efficient Formation of Formaldehyde and Methanol by the Addition of Hydrogen Atoms to CO in H<sub>2</sub>O–CO Ice at 10 K." *Astrophys. J.* 571(2): L173–L176.

<https://doi.org/10.1086/341412>

<sup>317</sup> Fuchs, G.W.; Cuppen, H.M.; Ioppolo, S.; et al. (2009). "Hydrogenation Reactions in Interstellar CO Ice Analogues." *A&A* 505: 629–639.

<sup>318</sup> Hore, P.J.; Mouritsen, H. (2016). "The Radical-Pair Mechanism of Magnetoreception." *Annu. Rev. Biophys.* 45: 299–344.

<sup>319</sup> Ritz, T.; Thalau, P.; et al. (2004). "Resonance Effects Indicate a Radical-Pair Mechanism in Avian Magnetoreception." *Nature* 429: 177–180.

<sup>320</sup> McDonough, W.F.; Sun, S.-s. (2020). "Radiogenic Power and Geoneutrino Luminosity of the Earth." *Geochem. Geophys. Geosyst.* 21(7): e2019GC008865. <https://doi.org/10.1029/2019GC008865>

<sup>321</sup> Carlson, R.W.; Johnson, R.E.; Anderson, M.S. (1999). "Hydrogen Peroxide on the Surface of Europa." *Science* 283(5410): 2062–2064.

<sup>322</sup> Teolis, B.D.; et al. (2017). "Water Ice Radiolysis: Synthesis of O<sub>2</sub>, H<sub>2</sub> and H<sub>2</sub>O<sub>2</sub> and Implications for Icy Worlds." *J. Geophys. Res.: Planets* 122:

Reaction Accessibility: Tunnelling opens pathways at low temperatures (e.g., methanol precursor synthesis); Non-adiabatic ultrafast relaxation enhances photostability of key biomolecules (facilitating accumulation); Spin chemistry may fine-tune product selectivity in weak fields. Network Directionality: Electron transfer between mineral/metal centres and organisms follows Marcus rules, determining which pathways are more "favourable" and influencing the sustainability of premetabolic prototypes.

In summary, the isotope decay–radiolysis–prebiochemistry chain<sup>323</sup> is founded end-to-end upon quantum principles; yet "quantum biology" typically denotes quantum effects within living systems (e.g., radical-induced magnetic sensing). For the prebiotic stage, it is more apt to state that quantum physical/chemical mechanisms determine energy deposition, reaction pathways, and selectivity, thereby establishing boundary conditions and preferences for life's origin<sup>324</sup>.

### **(iii) Source of life's energy: $F=ma$ and $\Delta E = \Delta m \times c^2$**

Dust within protoplanetary discs undergoes continuous dynamic motion. Dust particles interact with one another while being influenced by the disc's overall physical and chemical conditions, which vary due to differences in particle mass, density, and inherent physicochemical properties. Protoplanetary disc dust is subjected to diverse cosmic and protosolar radiation, including but not limited to infrared, visible light, ultraviolet, X-rays, gamma rays, and alpha and beta rays from isotopes. Consequently, the birth of life within the protoplanetary disc—its intricate physicochemical transformations—can be quantified, and this quantification is reproducible. This quantification differs from the crude descriptions of volcanic eruptions, celestial lightning, deep-sea hydrothermal vents, and so-called 'black smokers' on ancient Earth. The energy source for life can be broadly described by two equations:  $F=ma$  and  $\Delta E = \Delta m \times c^2$ . These equations are not merely simplistic descriptions of physical phenomena; they also provide profoundly insightful interpretations of the origin of life.

#### **1. $F=ma$ on the protoplanetary disc**

Mechanical-Chemical Coupling in Dust Collisions within Protoplanetary Discs:

From  $F=ma$  to Prebiotic Chemistry: This paper systematically summarizes the kinetic factors governing dust particle collisions in protoplanetary discs alongside their chemical

---

<sup>323</sup> Dauphas, N.; Chaussidon, M. (2011). "A Perspective from Extinct Radionuclides on a Young Stellar Object: The Sun and Its Accretion Disk." *Annu. Rev. Earth Planet. Sci.* 39(1): 351–386. <https://doi.org/10.1146/annurev-earth-040610-133428>

<sup>324</sup> Lichtenberg, T.; Golabek, G.J.; Burn, R.; et al. (2019). "A Water Budget Dichotomy of Rocky Protoplanets from  $^{26}\text{Al}$ -powered Heating." *Nature Astronomy* 3: 307–313.

consequences (mechanochemistry, radicals, and plasma). It provides applicable equations and illustrative examples<sup>325</sup>.

### 1.1 Physical Framework and Variables

- Mass:  $m = (4/3)\pi r^3 \rho$ ;  $\rho$  (density) for silicates typically taken as  $3000 \text{ kg}\cdot\text{m}^3/\text{m}^3$ ; <sup>326</sup>.

Peak deceleration:  $a_{(\text{max})} \approx v_{(\text{rel})}^2 / (2\delta)$ ; (where  $a$  is contact radius, material-dependent). Note:  $\delta$  denotes contact compression displacement ( $\sim 10^{-9}$ – $10^{-6}$  m),  $v_{(\text{rel})}$  is relative velocity, and  $r$  is particle radius.

Peak force:  $F_{(\text{max})} \approx mv_{(\text{rel})}^2 / (2\delta)$ ; Relative velocity:  $v_{(\text{rel})}$ : Induced by turbulence, differential drift and gas drag, typically  $10^{-9}$  m/s to  $10^{-6}$  m/s (spanning mild adhesion to violent fragmentation).

- Compression displacement (contact flattening scale):  $\delta$ ; nanometre–micrometre scale ( $10^{-9}$ – $10^{-6}$  m)<sup>327</sup>.
- Effective curvature:  $R^*$ ; collision of two spheres of equal radius  $r$ ,  $R^* = r/2$ .
- Effective elastic modulus:  $E^*$ ; given by two materials  $E$ ,  $\nu$  as  $E^* = [(1-\nu_1^2)/E_1 + (1-\nu_2^2)/E_2]^{-1}$ , with silicate approximated as 50 GPa<sup>328</sup>.

### 1.2 $F = ma$ and impact dimension ( $F=ma$ and three core quantities)

Peak acceleration:  $a_{\text{max}} \approx v_{\text{rel}}^2 / (2\delta)$

Peak force:  $F_{\text{max}} \approx m v_{\text{rel}}^2 / (2\delta) = m \cdot a_{(\text{max})}$ <sup>329</sup>

Kinetic energy (reduced mass  $\mu$ ):  $E_k = \frac{1}{2} \mu v_{\text{rel}}^2$  (for identical particles,  $\mu = m/2$ )<sup>330</sup>

Momentum and impulse:  $J = \int F dt \approx \mu \cdot \Delta v$ ; Contact time  $t(c)$ . See Hertz approximation<sup>331</sup>.

<sup>325</sup> Becker, T. et al. (2024). Tribocharged solids in protoplanetary discs: internal and external discharge time-scales, MNRAS 533, 413–422.

<sup>326</sup> Balduin, T. et al. (2023). Size-dependent charging of dust particles in protoplanetary discs, A&A 679, A101.

<sup>327</sup> Blum, J., & Wurm, G. (2008). The Growth Mechanisms of Macroscopic Bodies in Protoplanetary Disks. A&A 46, 21–56.

<sup>328</sup> Güttler, C., et al. (2010). The outcome of protoplanetary dust growth... I. Mapping the zoo of laboratory collision experiments. A&A 513, A56.

<sup>329</sup> Stronge, W.J. (2000/2018). Impact Mechanics, Cambridge Univ. Press

<sup>330</sup> Ormel, C.W.; Cuzzi, J.N. (2007). Closed-form expressions for particle relative velocities induced by turbulence, A&A 466, 413–420

<sup>331</sup> Jungmann, F., et al. (2021). Observation of bottom-up formation for charged grain aggregates. A&A 649, A72.

Upper limit of body temperature rise (if all kinetic energy is converted to heat):  $\Delta T_{\max} \approx v_{\text{rel}}^2 / (2 c_p)$ , where for silicate  $c_p \approx 800 \text{ J}\cdot\text{kg}^{-1}\cdot\text{K}^{-1}$ .

Typical parameters:  $\rho(\text{silicate}) \approx 3000 \text{ kg}\cdot\text{m}^{-3}$ ;  $r$ :  $\mu\text{m}$ – $\text{cm}$ ;  $v(\text{rel})$ :  $10^{-2}$ – $10 \text{ m}\cdot\text{s}^{-1}$  (from turbulence, drift and differential speed);  $\delta$ :  $10^{-9}$ – $10^{-6} \text{ m}$ ; contact time  $\sim \mu\text{s}$ ; volume heating is typically negligible, but flash heating and high shear may occur in the contact zone.

### 1.3 Hertz contact approximation and impact timescale

Contact radius:  $a = (3 F R^* / 4 E^*)^{1/3}$  <sup>332</sup>

Peak contact pressure:  $p_0 = 3F / (2\pi a^2)$

Displacement-contact radius relationship:  $\delta \approx a^2 / R^*$  <sup>333</sup>

Impact duration (empirical approximation):  $t_c \approx C \cdot (m_{\text{eff}})^2 / (R^* E^{*2} v_{\text{rel}})$  <sup>(1/5)</sup>,  $C \approx 2.9$ ,  $m_{\text{eff}} \approx \mu$

Note: The above represents the dimensional relationships for elastic contact; when plastic/brittle fracture occurs, the values of  $a$ ,  $p_0$ , and  $t_c$  will deviate, but the order-of-magnitude assessment remains valid <sup>334</sup>.

### 1.4 The Causal Chain from Mechanics to Chemistry

From Mechanics to Chemistry: Three-Stage Velocity–Particle Size Effect Spectrum

**1.4.1 Low-velocity/microparticles:** Bonding/sintering, slight compaction, minimal fragmentation; surface fluctuations  $\rightarrow$  expose fresh active sites. Chemistry: Rapid surface area growth; low-coordination sites exposed in Fe–S/ metal oxides  $\uparrow$ ; microporous networks facilitate water/ice film microreactors; supports prebiotic processes like FTT/CO hydrogenation.

**1.4.2 Medium-speed/millimetre scale:** Bouncing/compaction/partial fragmentation ("bounce barrier"); contact point pressures (ppp) reaching hundreds of MPa–GPa. Chemistry: Coexistence of mechanochemistry (bond breaking/defects, E'/F centres) and triboelectricity; ice/organic ice: generation of  $\text{H}\cdot$ ,  $\text{OH}\cdot$ ,  $\text{H}_2$ ,  $\text{H}_2\text{O}_2$  and other radiolysis-induced homologous radicals (synergizing with radiochemistry). PAHs/Organics: Dehydrogenation/cationization/oligomerization; Fe–S: Cracking exposes  $\text{Fe}^{2+}/\text{Fe}^{3+}$  and sulphur radical sites, promoting surface reactions of  $\text{CO}/\text{CO}_2 \rightarrow$  organics. Charged dust enhances discharge/microplasma probability, further driving precursor formation (CN/HCN/ $\text{NO}_x$ , etc.).

<sup>332</sup> Lacks, D.J.; Sankaran, R.M. (2011). Contact electrification of insulating materials, J. Phys. D 44, 453001

<sup>333</sup> Johnson, K. L. (1985). Contact Mechanics. Cambridge Univ. Press.

<sup>334</sup> Popov, V. L. Handbook of Contact Mechanics.

**1.4.3 High-velocity/millimetre-centimetre scale:** Severe fragmentation/jeting, contact pressures reaching GPa; bulk heating remains low, but flash temperatures in contact zones+ exhibit pronounced high shear. Chemistry: Surface area of newly formed debris increases dramatically →, causing a leap in catalytic site density; Mechanistic chemical chain reactions: Free radicals generated by fragmentation undergo secondary reactions within micro-pores/films, promoting methylation, carbonylation, alcoholation, etc.; Volatile re-evaporation: Trapped CO, CO<sub>2</sub>, H<sub>2</sub>, CH<sub>4</sub> desorb, replenishing gas-solid phase reactions.

Note: Particle-particle collisions alone are insufficient to achieve "globule melting" (which requires larger-scale energy sources, such as intra-disk shock waves or lightning); however, collisions prepare reactive species and structures, coupling with radiative/photochemical drivers.

## 1.5 Net effects and impacts on prebiotic chemistry

**1.5.1 Structural Effects:** Crushing + Compaction → Hierarchical pore formation and "microchamber" creation, enhancing reactant enrichment and selectivity, and improving diffusion-retention equilibrium<sup>335</sup>. GPa-level contact pressures + Microsecond-scale shear → Lattice defects, bond breakage, and active site exposure (e.g., E'/F centres)<sup>336</sup>.

**1.5.2 Surface Chemistry:** Freshly Exposed Surfaces and Defects → Low-coordination metal/sulphur sites, enhancing reaction rates for FTT, CO/CO<sub>2</sub> reduction, nitration/sulphidation pathways.

**1.5.3 Radical/Plasma Coupling:** Mechanochemistry + Triboelectricity → Parallel radical and electrochemistry<sup>337</sup>, generating key precursors like HCN, H<sub>2</sub>O<sub>2</sub>, peroxy radicals<sup>338</sup>. Mechanochemistry: Generation of radicals including H·, ·OH, H<sub>2</sub>, H<sub>2</sub>O<sub>2</sub> on ice/organic ice surfaces<sup>339</sup>; dehydrogenation/ionization/polymerization of PAHs<sup>340</sup>. Friction Electrification

---

<sup>335</sup> Dominik, C.; Tielens, A.G.G.M. (1997). The Physics of Dust Coagulation and the Structure of Dust Aggregates in Space, *ApJ* 480, 647–673.

<sup>336</sup> Parkes, R. J., et al. (2018). Rock-crushing derived hydrogen directly supports a methanogenic community.

<sup>337</sup> Michalchuk, A.A.L. et al. (2021). Tribochemistry, Mechanical Alloying, Mechanochemistry, *Frontiers in Chemistry* 9:685789

<sup>338</sup> Cuccu, F., et al. (2022). Mechanochemistry: New Tools to Navigate the Uncharted Routes... *ChemSusChem*.

<sup>339</sup> Stone, J. R., et al. (2023). Flash heating boosts the potential for mechanochemical reactions. *Frontiers in Geochemistry*.

<sup>340</sup> He, H., et al. (2023). A mineral-based origin of Earth's initial hydrogen peroxide. *PNAS* 120, e2221984120.

and Microdischarges: Charged dust-dust/dust-gas discharges<sup>341</sup>, yielding electrons/ions<sup>342</sup>, NO<sub>x</sub>/HCN precursors<sup>343</sup>, and peroxy radicals<sup>344</sup>.

**1.5.4 Synergy with Radiochemistry:** Collision-released active sites and radicals provide "landing sites" for radiolysis products from <sup>26</sup>Al/<sup>60</sup>Fe isotopes, jointly driving a dual-track prebiotic environment of H<sub>2</sub>/CO supply + ROS pressure. Radiolysis-generated H<sub>2</sub>, ·OH, H<sub>2</sub>O<sub>2</sub> from <sup>26</sup>Al, <sup>60</sup>Fe interact synergistically with collision-activated surfaces/radicals<sup>345</sup>.

## 1.6 Velocity–Particle Size–Chemical Outcome Matrix

Interval	Mechanical Outcomes	Surface/Structure Effects	Chemical and Prebiotic Effects
Low-speed/microparticles: $v \lesssim 0.1 \text{ m}\cdot\text{s}^{-1}$ , $r \sim \mu\text{m}$	Bonding/compaction; minimal fragmentation	Exposure of fresh active sites; formation of microporous networks/thin-film water layers	Enhanced surface catalysis; favourable for FTT, CO/CO <sub>2</sub> hydrogenation; reactant enrichment
Medium-speed/millimetre scale: $v \sim 0.1\text{--}1 \text{ m}\cdot\text{s}^{-1}$ , $r \sim 10^{-4}\text{--}10^{-3} \text{ m}$	Bouncing/compaction/partial fragmentation; contact pressure $p$ reaching $10^2\text{--}10^3 \text{ MPa}$	Defects and bond breakage (E'/F centres); triboelectric charging	Mechanochemical radicals (H·, ·OH, etc.); PAH dehydrogenation/ionisation; Fe–S site exposure → CO reduction; micro-discharges → HCN/NO <sub>x</sub> precursors
High-velocity/millimetre–centimetre: $v \sim 1\text{--}10 \text{ m}\cdot\text{s}^{-1}$	Intense fragmentation/jet; contact zone "flash"	Massive new surface area; hierarchical porosity	Chain radical reactions (methylation/carbonylation/alcoholation); volatiles desorption replenishing CO/H <sub>2</sub> /CH <sub>4</sub> ;

<sup>341</sup> Desch, S. J., & Cuzzi, J. N. (2000). The Generation of Lightning in the Solar Nebula. *Icarus* 143, 87–105.

<sup>342</sup> Mizzi, C.A. et al. (2019). Does Flexoelectricity Drive Triboelectricity?, *Phys. Rev. Lett.* 123, 116103

<sup>343</sup> James, S.L. et al. (2012). Mechanochemistry: opportunities for new and cleaner synthesis, *Chem. Soc. Rev.* 41, 413–447

<sup>344</sup> Maia, G. P., et al. (2025). Mechanochemical Reactivity of Ribonucleosides... *Applied Sciences*.

<sup>345</sup> Johansen, A., et al. (2018). Harvesting the decay energy of β-Al to drive lightning. *A&A* 611, L1.



	temperature" and high shear		synergistic effects with radiation/photochemistry
--	-----------------------------	--	---

## 2. Radiation Environment and Energy Injection Mechanisms

In the quest to understand the origins of life, the source and transfer of energy remain pivotal to constructing prebiotic chemical reaction networks. The protoplanetary disc's cosmic environment was highly dynamic, saturated with diverse high-energy radiation and radioactive nuclides, delivering energy far exceeding most natural sources in the modern terrestrial environment. This energy, conveyed through radiation, ionization, and thermal transfer, provided essential conditions for the synthesis, excitation, and transformation of organic molecules.

### 2.1 Thermal Energy Released by the Decay of Short-lived Isotopes such as $^{26}\text{Al}$ and $^{60}\text{Fe}$

Protoplanetary discs harbour several short-lived radionuclides (SLRs), notably:  $^{26}\text{Al}$  (half-life  $\approx 0.7$  Myr)<sup>346</sup>,  $^{60}\text{Fe}$  (half-life  $\approx 2.6$  Myr)<sup>347</sup>. These nuclides are believed to originate from supernova explosions or the injection of Wolf-Rayet stellar winds. They are widely distributed throughout the disc, being incorporated into planetesimals, dust, or ice grains. Their decay pathways involve  $\beta^+$  or  $\beta^-$ , releasing high-energy electrons,  $\gamma$ -rays, and heat, adhering to the mass-energy relationship:  $\Delta E = \Delta m \times c^2$ .

Estimates indicate that  $^{26}\text{Al}$  decay can deliver approximately  $6 \times 10^5$  J/kg of localized thermal energy, equivalent to sustaining heating on a planetoid scale for several million years. It induces the following reaction environments: localized hydration reactions ( $\text{H}_2\text{O}$  + minerals), thermal restructuring of organic matter, melting of the ice layer, and the formation of complex organic compounds (such as amines, alcohols, and aldehydes) within enclosed environments<sup>348</sup>. Thus, SLRs provide a crucial, temporally controllable, spatially localized, and energetically concentrated driving force for "chemically active zones" within protoplanetary discs.

<sup>346</sup> Cameron, A. G. W., & Truran, J. W. (1977). The supernova trigger for formation of the solar system. *Icarus*, 30, 447–461. [https://doi.org/10.1016/0019-1035\(77\)90101-6](https://doi.org/10.1016/0019-1035(77)90101-6)

<sup>347</sup> Rugel, G.; Faestermann, T.; Knie, K.; et al. (2009). "New Measurement of the  $^{60}\text{Fe}$  Half-Life." *Phys. Rev. Lett.* 103(7): 072502. <https://doi.org/10.1103/PhysRevLett.103.072502>

<sup>348</sup> Materese, C. K., Nuevo, M., Sandford, S. A., et al. (2015). Amino acid precursors from the UV irradiation of astrophysical ice analogs. *Astrobiology*, 15(6), 333–350. <https://doi.org/10.1089/ast.2014.1253>

## 2.2 Exposure of the Protoplanetary Disk to Cosmic Rays, X-rays, Ultraviolet and Gamma Radiation<sup>349</sup>

Protoplanetary discs are exposed to intense interstellar radiation environments, primarily comprising:<sup>350</sup> Cosmic rays (CRs): High-energy protons and heavy ions with formidable penetrating power, capable of penetrating hundreds of metres of solid material; X-rays and ultraviolet radiation (X-UV): Intense radiation originating from the central young star (T Tauri or Herbig Ae/Be)<sup>351</sup>, with intensities hundreds of times greater than the present-day Sun; Gamma rays ( $\gamma$ -rays): introduced by supernova remnants or neutron star winds, possessing extremely high ionising capability.

These rays can induce diverse primordial reactions within dust particles, ice layers, and organic films, such as molecular excitation and dissociation, radical generation, carbon chain cleavage and restructuring<sup>352</sup>, and alterations in surface electronic states (e.g., PAH oxidation). Particularly in environments where dust particles coexist with water ice, radiation-induced reactions can trigger complex pathways, such as H-abstraction<sup>353</sup>, radical-radical coupling, and photoreduction, thereby forming biologically relevant groups, including amino acid precursors, aldoses, and amines (Ciesla & Sandford, 2012)<sup>354</sup>.

## 2.3 Radiation-Induced Reactions and Radical Mechanisms

Experimental and theoretical studies indicate that the following reaction pathways may occur concurrently under radiative and photochemical conditions:

1.  $\text{CO} + \text{H}_2\text{O} + \text{UV} \rightarrow \rightarrow \text{HCO}\cdot, \cdot\text{OH} \rightarrow \rightarrow \text{H}_2\text{CO} \rightarrow \rightarrow \text{CH}_3\text{OH}$
2.  $\text{HCN} + \gamma \rightarrow \rightarrow \text{CN}\cdot + \text{H}\cdot \rightarrow \text{NH}_2\text{CN}$  (aminonitrile)
3.  $\text{H}_2\text{O ice} + \text{high-energy particle} \rightarrow \cdot\text{OH} + \text{H}\cdot \rightarrow \text{H}_2\text{O}_2, \text{HO}\cdot, \text{H}_2$

These radical reactions require neither high temperatures nor catalysts, being driven solely by high-energy electrons or photons. Consequently, they can occur in the cold outer regions

---

<sup>349</sup> Bergin, E. A., et al. (2007). The chemical evolution of protoplanetary disks: Observations and modelling. *Protostars and Planets V*, 751–766.

<sup>350</sup> Adams, F. C. (2010). The birth environment of the solar system. *Annual Review of Astronomy and Astrophysics*, 48, 47–85. <https://doi.org/10.1146/annurev-astro-081309-130830>

<sup>351</sup> Glassgold, A. E., Najita, J. R., & Igea, J. (1997). X-ray ionisation of protoplanetary disks. *Astrophysical Journal*, 480, 344–350. <https://doi.org/10.1086/303958>

<sup>352</sup> Hudson, R. L., & Moore, M. H. (2001). Radiation chemical alterations in solar system ices: An overview. *Journal of Geophysical Research: Planets*, 106(E12), 33275–33284. <https://doi.org/10.1029/2000JE001359>

<sup>353</sup> Cleeves, L. I., Bergin, E. A., Alexander, C. M. O., et al. (2014). The ancient heritage of water ice in the solar system. *Science*, 345(6204), 1590–1593. <https://doi.org/10.1126/science.1258055>

<sup>354</sup> Ciesla, F. J., & Sandford, S. A. (2012). Organic synthesis via irradiation and warming of ice grains in the solar nebula. *Science*, 336(6080), 452–454. <https://doi.org/10.1126/science.1217290>

of protoplanetary discs. Such conditions are virtually absent on Earth's surface but are the norm in protoplanetary discs, particularly beyond the 'snow line'.<sup>355</sup>

## 2.4 The high-energy environment of protoplanetary discs is more chemically active than Earth

Compared to modern Earth, protoplanetary discs possess several significant high-energy advantages:

Conditions	Protoplanetary Disk	Early Earth
Ultraviolet intensity	High (young stellar activity)	Attenuated by atmosphere
X-ray flux	High	Virtually absent
Radiation-induced radical formation	Abundant	Scarce
Dust catalysis/ice-encapsulated particle structure	Widespread	Rare
Internal heat sources within SLRs	Present ( $^{26}\text{Al}/^{60}\text{Fe}$ ) <sup>356</sup>	Primarily terrestrial core radioactivity

These discrepancies suggest that the radiation environment within protoplanetary discs is not merely destructive but also synthetic—providing a unique reaction space and excitation mechanism for the formation of primordial organic systems.

The decay of radionuclides within the protoplanetary disc, combined with stellar radiation and cosmic rays, collectively form a high-energy, sustainable, multi-modal energy input system. Whether through heat-induced reactions triggered by  $^{26}\text{Al}$  decay<sup>357</sup> or radical reaction chains induced by ultraviolet radiation/cosmic rays, these processes could facilitate the transition of primordial organic chemistry in environments devoid of enzymes,

<sup>355</sup> Cruikshank, D. P., et al. (2005). Organic matter in the solar system: From molecules to planetary systems. *Advances in Space Research*, 36(2), 178–183. <https://doi.org/10.1016/j.asr.2005.02.019>

<sup>356</sup> Lugaro, M., et al. (2018). Radioactive nuclei from cosmochemistry to habitability. *Progress in Particle and Nuclear Physics*, 102, 1–47. <https://doi.org/10.1016/j.ppnp.2018.05.002>

<sup>357</sup> Lichtenberg, T., et al. (2016). Accretion and differentiation of rocky planets in the presence of short-lived radionuclides. *Earth and Planetary Science Letters*, 449, 2–12. <https://doi.org/10.1016/j.epsl.2016.05.004>

high pressure, or biological involvement. Such energetic conditions may have held greater potential for creating "pre-life systems" than those present on early Earth.

### 3. Radioactive Isotopes in the Protoplanetary Disc: A Core Factor in Life's Origin and Evolution

Radioactive isotopes within protoplanetary discs play a central role in research on the origin of life within these discs. Due to their predominantly heavy metallic nature, both short- and long-half-life radioactive isotopes had migrated mainly into the Earth's core or mantle during the early formation of the primordial planet. Those present in the crust exhibited extremely low abundances. When the primordial Earth's crust was still molten, these short-lived radioactive isotopes had already been depleted, rendering them insignificant for the origin of life. However, within the protoplanetary disc, both short- and long-lived radioactive isotopes played a pivotal role in the emergence and evolution of life within the disc. The critical role of isotopes in life's evolution will be discussed in detail in Part VI.

The "short-lived (SLR) and long-lived radioactive isotopes frequently discussed in early solar system studies" are organized into two tables, detailing: half-lives, primary decay types/radiation, and common radiochemical products when this radiation bombards different targets (H<sub>2</sub>O, C–O, (CO/CO<sub>2</sub>)/carbonyl-containing), Fe–S mineral clusters, and PAHs.

The SLR list is based on the isotopic review in Protostars & Planets VII, supplemented with standard geochronological systems; half-lives are based on recent measurements. Radiochemical "products" refer to the typical major products, with specific distributions dependent on experimental conditions such as temperature, phase state (ice/aqueous solution/solid), dose, and mixing with other molecules<sup>358</sup>. Summary of Radioisotopes and Radiochemical Products in Protoplanetary Discs:

#### 3.1.1 Short-Lived Radioactive Isotopes (SLRs)

Isotope	Half-life	Decay / Radiation	Typical Irradiation Products (by Target)
<sup>7</sup> Be	53 days	EC, γ	H <sub>2</sub> O: e <sup>-</sup> aq, H·, ·OH, H <sub>2</sub> , H <sub>2</sub> O <sub>2</sub> ; C–O ice: HCO, H <sub>2</sub> CO, CH <sub>3</sub> OH; Fe–S: Fe <sup>3+</sup> /S·; CO <sub>2</sub> ; PAHs:

<sup>358</sup> Castillo-Rogez, J. C., & Young, E. D. (2017). Origin of short-lived radionuclides in the early Solar System. *Annual Review of Earth and Planetary Sciences*, 45, 61–87.

			ionisation, CO, CH <sub>4</sub> dehydrogenation <sup>359360361</sup>
<sup>10</sup> Be	1.387 Myr	$\beta^-$ , $\gamma$ (weak)	As above
<sup>26</sup> Al	0.717 Myr	$\beta^+$ /EC, $\gamma$ (1.809 MeV)	As above, enhanced high-energy $\gamma$ radiation decay
<sup>36</sup> Cl	0.301 Myr	$\beta^- \rightarrow ^{36}\text{Ar}$ ; EC $\rightarrow$ <sup>36</sup> S	Ibid.
<sup>41</sup> Ca	0.10 Myr	Electron capture (X-ray/Auger)	Ibid.
<sup>53</sup> Mn	3.7 Myr	$\beta^- \rightarrow ^{53}\text{Cr}$	Ibid.
<sup>60</sup> Fe	2.6 Myr	$\beta^- (\rightarrow ^{60}\text{Co} \rightarrow ^{60}\text{Ni})$ , $\gamma$	Ibid.
<sup>107</sup> Pd	6.5 Myr	$\beta^- \rightarrow ^{107}\text{Ag}$	Ibid.
<sup>129</sup> I	15.7 Myr	$\beta^- \rightarrow ^{129}\text{Xe}$ , $\gamma$	Ibid.
<sup>146</sup> Sm	68 Myr	$\alpha \rightarrow ^{142}\text{Nd}$	Ibid.
<sup>182</sup> Hf	8.9 Myr	$\beta^- \rightarrow ^{182}\text{W}$	Ibid.
<sup>205</sup> Pb	Variable	EC	Ibid.
<sup>244</sup> Pu	80 Myr	$\alpha$	Ibid.
<sup>247</sup> Cm	15.6 million years	$\alpha$	Ibid.

### 3.1.2 Long-lived Radioisotopes

Isotope	Half-life	Decay / Radiation	Typical Irradiation Products (by Target)
<sup>238</sup> U	4.47 Gyr	Alpha chain, beta, gamma	H <sub>2</sub> O: e <sup>-</sup> aq, H $\cdot$ , $\cdot$ OH, H <sub>2</sub> , H <sub>2</sub> O <sub>2</sub> ;

<sup>359</sup> Sophie Le Caër. (2011) Water Radiolysis: Influence of Oxide Surfaces on H<sub>2</sub> Production under Ionising Radiation. *Water*, 3, 235–253; doi:10.3390/w3010235

<sup>360</sup> Dartnell, L. R. (2011). Ionising radiation and life. *Astrobiology*, 11(6), 551–582.

<sup>361</sup> Draganic, I. G., Draganic, Z. D., & Nikolic, M. (1991). *Radiation Chemistry of Water*. CRC Press.

			C–O: HCO, H <sub>2</sub> CO, CH <sub>3</sub> OH; CO, CH <sub>4</sub> CO <sub>2</sub> , Fe–S: Fe <sup>3+</sup> , S <sup>·</sup> ; PAHs: ionisation <sup>362,363</sup>
<sup>235</sup> U	0.704 Gyr	α chain	Ibid.
<sup>232</sup> Th	14.05 Gyr	Alpha chain	As above
<sup>40</sup> K	1.25 Gyr	β <sup>-</sup> , EC	As above
<sup>87</sup> Rb	48.8 Gyr	β <sup>-</sup>	Ibid.
<sup>147</sup> Sm	106 Gyr	α	Ibid.
<sup>176</sup> Lu	37.8 Gyr	β <sup>-</sup> , EC	Ibid.

### 3.2 Brief introduction to isotope "products from bombardment of different targets"

- H<sub>2</sub>O (liquid/ice): e<sup>-</sup> <sub>aq</sub>, H<sup>·</sup>, ·OH, H<sub>3</sub>O<sup>+</sup>, H<sub>2</sub>, H<sub>2</sub>O<sub>2</sub>; at high doses/in mixtures, O<sub>2</sub>, HO<sub>2</sub>·/O<sub>2</sub><sup>-</sup>. These constitute the elementary product spectrum under all ionizing radiation (α, β, γ, and secondary electrons)<sup>364</sup>.
- C–O systems (CO/CO<sub>2</sub> with H<sub>2</sub>O/CH<sub>4</sub>/CH<sub>3</sub>OH ice mixtures): HCO, H<sub>2</sub>CO, HCOOH, CH<sub>3</sub>OH, CH<sub>4</sub>, CO<sub>2</sub> (mutually interconverting), and more complex organics; products strongly influenced by composition and temperature<sup>365</sup>.
- Fe–S (pyrite/FeS/biogenic sulphide clusters): ·OH, H<sub>2</sub>O<sub>2</sub>-induced Fe<sup>2++</sup> → Fe<sup>3++</sup>, S(0)/polysulphide anions, surface radicals and electron transfer; under anoxic conditions, radiolysis products may drive "anoxic pyrite oxidation" and ROS generation<sup>366</sup>.

<sup>362</sup> Sophie Le Caër. (2011) Water Radiolysis: Influence of Oxide Surfaces on H<sub>2</sub> Production under Ionising Radiation. *Water*, 3, 235–253; doi:10.3390/w3010235

<sup>363</sup> Draganic, I. G., Draganic, Z. D., & Nikolic, M. (1991). *Radiation Chemistry of Water*. CRC Press.

<sup>364</sup> Sophie Le Caër. (2011) Water Radiolysis: Influence of Oxide Surfaces on H<sub>2</sub> Production under Ionising Radiation. *Water*, 3, 235–253; doi:10.3390/w3010235

<sup>365</sup> Maria Magdalena Ramirez Corredores. (2020) Radiation-Induced Chemistry of Carbon Dioxide: A Pathway to Close the Carbon Loop for a Circular Economy. *Front. Energy Res. Front. Energy Res.*, 8, <https://doi.org/10.3389/fenrg.2020.00108>

<sup>366</sup> Corey A Cohn. (2006) Role of pyrite in formation of hydroxyl radicals in coal: possible implications for human health. *Part Fibre Toxicol.* 2006 Dec 19;3:16. doi: 10.1186/1743-8977-3-16

- PAHs: Ionization ( $\text{PAH}^+$  /  $\text{PAH}^-$ ), dehydrogenation/rehydrogenation, fragmentation (commonly  $\text{C}_n^+$  carbon clusters and small hydrocarbon fragments), configurational isomerization and surface effects (low-energy surfaces, high-energy bulk phases); concurrently occurring with cosmic ray/ion irradiation and VUV/UV photochemical reactions<sup>367</sup>.
- Mixed ice ( $\text{H}_2\text{O}:\text{CO}/\text{CO}_2:\text{CH}_4:\text{NH}_3$  etc.): Cosmic ray/heavy ion irradiation leads to the destruction and resynthesis of precursor molecules, with observed formation of  $\text{H}_2\text{CO}$ ,  $\text{H}_2\text{O}_2$ ,  $\text{CH}_4$ ,  $\text{HCOOH}$ , etc., accompanied by sputtering/desorption<sup>368</sup>.

**3.3 Within protoplanetary and planetary disks, isotopes of varying half-lives—both long-lived and short-lived**—not only induce diverse chemical reactions in surrounding matter by emitting different types of radiation and particles, but also generate internal heating effects within the disk particles themselves. It enables reactions and subsequent life forms to adapt and evolve within the disk particles. Isotopes heat disk particles through two distinct mechanisms:<sup>369</sup>.

**3.3.1 Radioactive decay:** The spontaneous transformation of atomic nuclei releases energy ( $\alpha/\beta$  particles,  $\gamma$  photons, nuclear recoil, and trace neutrinos).

**3.3.2 Radiolysis:** These radiation species ionize/excite surrounding matter (commonly water, ice, or organic compounds), breaking chemical bonds and generating new chemical species ( $\cdot\text{OH}$ ,  $\text{H}\cdot$ ,  $\text{H}_2$ ,  $\text{H}_2\text{O}_2$ , etc.)<sup>370</sup>.

The two represent a cause-and-effect relationship: decay provides energy and particles, while radiolysis constitutes the chemical and thermal effects resulting from these particles depositing energy into the medium<sup>371</sup>.

### **3.4 At which stage does the heat generation mechanism occur?**

---

<sup>367</sup> J W L Lee, . (2021) Time-resolved relaxation and fragmentation of polycyclic aromatic hydrocarbons investigated in the ultrafast XUV-IR regime. *Nat Commun.* 2021 Oct 20;12:6107. doi: 10.1038/s41467-021-26193-z

<sup>368</sup> C. Mejía. (2024) Modifications of astrophysical ices induced by cosmic rays. *A&A*;A.687, <https://doi.org/10.1051/0004-6361/202349120>

<sup>369</sup> Spinks, J. W. T.; Woods, R. J. (1990). *An Introduction to Radiation Chemistry* (3rd ed.). John Wiley & Sons, New York. ISBN: 0-471-61403-3.

<sup>370</sup> Le Caër, S. (2011). Water Radiolysis: Influence of Oxide Surfaces on  $\text{H}_2$  Production under Ionising Radiation. *Water*, 3(1), 235–253. <https://doi.org/10.3390/w3010235>.

<sup>371</sup> Knoll, G. F. (2010). *Radiation Detection and Measurement* (4th ed.). John Wiley & Sons, Hoboken, NJ. ISBN: 978-0470131480.

**3.4.1 At the instant of decay,** energy is released in the form of kinetic energy of charged particles ( $\alpha/\beta$ ),  $\gamma$  photon energy, subatomic nucleus recoil energy, and neutrino energy. These are not yet heat (they are the energy of ordered particles/photons).

**3.4.2 Particle "braking" (energy deposition) within the medium:** Charged particles transfer energy to surrounding molecules via ionization/excitation (described by "linear energy transfer" or LET,  $dE/dx$ ), rapidly (on the femtosecond–picosecond timescale) converting to random thermal motion and generating heat. If  $\gamma$  photons are absorbed via photoelectric effect/Compton scattering/pair production, their energy is first transferred to electrons, which then deposit energy as described above to produce heat. Neutrinos interact scarcely with matter, their energy largely escaping without generating heat. Recoiling subatomic nuclei in solids also rapidly transfer kinetic energy to the lattice, producing heat.

Conclusion: Primary heat is generated during the process of "particle/photon interaction with the medium and subsequent halting", not "at the moment of emission". Decay merely "loads energy into the bullet/photon"; the actual heat generation occurs when the bullet strikes the target and dissipates its energy as heat.

**3.4.3 Radiolysis and Heat Distribution:** Within the same deposited energy, a portion is used for bond breaking/generating free radicals and molecules (chemical work), yet the vast majority ultimately dissipates as heat. <sup>372</sup>Chemical energy "stored" (e.g.,  $H_2$  produced by water radiolysis, or the separation of  $H_2O_2/O_2$  in ) will release heat upon subsequent reaction/recombination; this merely retrieves chemical potential energy previously "deposited" by radiation, without creating additional energy ex nihilo.

### **3.5 How should "decay heat" versus "radiolysis heat" be understood?**

**3.5.1 Decay-induced heating:** Calculating the energy budget from the source perspective = (number of decays per unit time)  $\times$  (energy deposited per decay in the medium)<sup>373</sup>

**3.5.2 Radiolysis heating:** Describing the consequences of the same energy deposition from the target material's perspective: Ionization/excitation  $\rightarrow$  Free radicals and chemical products + Heat. Chemically, the G value<sup>374</sup> (number of molecules produced per 100 eV)<sup>375</sup>

---

<sup>372</sup> Jay-Gerin, J.-P. (2025). Fundamentals of Water Radiolysis. Encyclopedia, 5(1), 38. <https://doi.org/10.3390/encyclopedia5010038>.

<sup>373</sup> U.S. Department of Energy. (1993). DOE Fundamentals Handbook: Nuclear Physics and Reactor Theory (Vol. 1 & Vol. 2). DOE-HDBK-1019/1-93 and DOE-HDBK-1019/2-93, Washington, DC.

<sup>374</sup> Elliot, A. J.; Bartels, D. M. (2009). The Reaction Set, Rate Constants and G-Values for the Simulation of the Radiolysis of Light Water over the Range 20 to 350 °C (Based on Information Available in 2008). Atomic Energy of Canada Limited (AECL), Report No. 153-127160-450-001, Mississauga, ON.

<sup>375</sup> Macdonald, D.D. "A Critical Review of Radiolysis Issues in Water-Cooled Reactors"



is commonly used to describe yield. So-called "radiation-induced heat generation" either involves the immediate conversion of the entire energy deposit into heat or the temporary storage of part of the energy as chemical potential, later released as reaction heat. The conservation of energy constrains the entire process<sup>376</sup>.

### 3.6 Relationship between particle type and heat generation:

Alpha particles: Short range, high linear energy transfer (LET), with energy almost entirely deposited within micrometre scales→. Localized intense radiation+Significant heat generation<sup>377</sup>. Beta particles (electrons/positrons): Longer range, moderate linear energy transfer (LET), with energy deposited over larger volumes→. Heat generation is dispersed<sup>378</sup>. Gamma rays: Uncharged, requiring prior absorption/scattering before transferring energy to electrons; Significant heat production in high-density/thick media; substantial escape possible in sparse media<sup>379</sup>. Neutrinos: Virtually no heat production, negligible<sup>380</sup>.

## 4. Protoplanetary Disk vs. Early Earth: Comparative Energy Advantages for Life Origins

"Which environment—protoplanetary discs or the early Earth—favoured life's emergence?" requires quantitative comparison across four dimensions: energy source sustainability, diversity, intensity, and chemical driving capacity. The following presents a comprehensive analysis:

### 4.1 Persistence of Energy Sources

**4.1.1 Protoplanetary Disk and Disk Planetesimals, Radioactive Isotope Decay (e.g., <sup>26</sup>Al, <sup>60</sup>Fe, <sup>40</sup>K):** Time Scale: Short-lived nuclides (e.g., <sup>26</sup>Al's 710,000-year half-life)<sup>381</sup> provide burst energy early on<sup>382</sup>; long-lived nuclides (e.g., <sup>40</sup>K's 1.25-billion-year half-life) sustain

---

<sup>376</sup> Janik, D.; Janik, I.; Bartels, D. M. (2007). Neutron and  $\beta/\gamma$  Radiolysis of Water up to Supercritical Conditions. 1.  $\beta/\gamma$  Yields for H<sub>2</sub>, H• Atom, and Hydrated Electron. *The Journal of Physical Chemistry A*, 111(32), 7777–7786. <https://doi.org/10.1021/jp071751r>.

<sup>377</sup> Ziegler, J. F.; Ziegler, M. D.; Biersack, J. P. (2010). SRIM—The Stopping and Range of Ions in Matter (2010). *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 268(11–12), 1818–1823. <https://doi.org/10.1016/j.nimb.2010.02.091>.

<sup>378</sup> International Commission on Radiation Units and Measurements (ICRU). (2011). Report 85: Fundamental Quantities and Units for Ionising Radiation. *Journal of ICRU*, 11(1), 1–31. <https://doi.org/10.1093/jicru/ndr011>.

<sup>379</sup> Berger, M. J.; Coursey, J. S.; Zucker, M. A.; Chang, J. (1999; updated versions). ESTAR, PSTAR, and ASTAR: Computer Programs for Calculating Stopping-Power and Range Tables for Electrons, Protons, and Helium Ions (version lineage; online database). National Institute of Standards and Technology (NIST), Gaithersburg, MD.

<sup>380</sup> Ruedas, T. (2017). Radioactive Heat Production of Six Geologically Important Nuclides. *Geochemistry, Geophysics, Geosystems (G<sup>3</sup>)*, 18(9), 3530–3541. <https://doi.org/10.1002/2017GC006997>.

<sup>381</sup> Gounelle & Meynet (2012). Solar System genealogy revealed by extinct short-lived radionuclides in meteorites. *Astronomy & Astrophysics* 545, A4. DOI: 10.1051/0004-6361/201219031

<sup>382</sup> Lichtenberg et al. (2021) A water budget dichotomy of rocky protoplanets from <sup>26</sup>Al-heating. *Nature Astronomy*. 5, 111–121. DOI: 10.1038/s41550-020-01218-7

prolonged heating. Spatial distribution: Uniformly distributed within the planetesimal, unaffected by surface conditions. Stellar radiation (protoplanetary Sun): Continuous ultraviolet and X-ray bombardment of dust particles drives photochemical reactions (e.g., PAH dissociation).

**4.1.2 Early Earth, transient energy events:** Volcanic eruptions (hours-months), lightning (microseconds), meteorite impacts (instantaneous). Submarine hydrothermal vents (lifetime ~ tens of thousands of years, highly localized)<sup>383</sup>. Solar energy: Surface ultraviolet radiation (but the early atmosphere lacked the ozone layer, which most scattered/absorbed)<sup>384</sup>.

Conclusion: Energy release from the protoplanetary disc exhibits global and sustained characteristics (on a million-year timescale), far surpassing the transient events of the early Earth.

## 4.2 Diversity of Energy Forms

### 4.2.1 Radiation Types in the Protoplanetary Disc:

Form of Energy	Mechanism	Life-related products
$\alpha/\beta$ rays (isotopes)	Bombardment of $\text{H}_2\text{O} \rightarrow \text{H}_2 + \text{H}_2\text{O}_2$	Energy source ( $\text{H}_2$ ) and oxidising agent ( $\text{H}_2\text{O}_2$ ) <sup>385</sup>
Ultraviolet/X-rays (stellar)	Dissociation of $\text{CO}/\text{CH}_4$ to generate free radicals	Activation of organic molecules (e.g., carboxylic acids)
Infrared radiation (dust)	Heating ice particles to 100–300 K	Promotion of ice-phase organic reactions (e.g., formaldehyde polymerization)

<sup>383</sup> Damer & Deamer (2020). The hot spring hypothesis for an origin of life. *Astrobiology* 20(4), 429–452. DOI: 10.1089/ast.2019.2045

<sup>384</sup> Stüeken et al. (2016) Isotopic evidence for biological nitrogen fixation in Earth's earliest sediments. *Geobiology* 14(5), 483-501. DOI: 10.1111/gbi.12192

<sup>385</sup> Draganic et al. (1991). Radiolysis of water: Primary and secondary processes. *Radiation Physics and Chemistry* 38(3), 317–326. DOI: 10.1016/0969-806X(91)90096-6

Form of Energy	Mechanism	Life-related products
Cosmic rays	Induction of deep-matter radiolysis	Continuous H <sub>2</sub> generation within the star

- Synergistic effect: Coupling of radioactive decay heating ( $\Delta E = \Delta m \times c^2$ ) with dust dynamics ( $F = ma$ ) drives material cycling.

**4.2.2 Early Earth: Primary energy sources: geothermal (localized), lightning (random), solar (attenuated by atmosphere). Limitations:** Single energy form with uneven spatial distribution (e.g., hydrothermal vents confined to mid-ocean ridges).

Conclusion: The protoplanetary disc offers richer energy forms whilst simultaneously supplying both reducing agents (H<sub>2</sub>) and oxidizing agents (H<sub>2</sub>O<sub>2</sub>), fulfilling the redox coupling requirements of life chemistry.

### 4.3 Energy Intensity and Chemical Driving Capacity

#### 4.3.1 Quantitative Comparison

Parameters	Protoplanetary Disk	Early Earth
H <sub>2</sub> Production Rate	10 <sup>8</sup> -10 <sup>10</sup> molecules/g (rock)/yr (α decay) <sup>386</sup>	10 <sup>6</sup> -10 <sup>7</sup> (serpentinization)
Temperature gradient	50-200°C within planetesimals (persistent)	Hydrothermal vent 350°C (localized)
Free radical flux	OH·-concentration~10 <sup>-5</sup> M (photolysis of water)	<10 <sup>-7</sup> M (lightning/ultraviolet)

**4.3.2 Prebiotic chemical efficiency, protoplanetary disc:** Synergistic action of radioactive+ and ultraviolet radiation can enhance PAHs→amino acid conversion rate by 100-

<sup>386</sup> McCollom (2013) Serpentinites, hydrogen, and life. Reviews in Mineralogy and Geochemistry 75, 365-410. DOI: 10.2138/rmg.2013.75.12

fold (experimental simulation see *Huang et al., 2021*)<sup>387</sup>. Fe-S cluster minerals (e.g., greigite) catalyze  $\text{CO} \rightarrow \text{CH}_4$  under  $\alpha$  irradiation (*Russell et al., 2014*)<sup>388</sup>. Early Earth: Hydrothermal vent Fischer-Tropsch reactions required high temperatures and pressures, yielding limited products (primarily alkanes). Conclusion: Protoplanetary discs possessed greater energy intensity and could drive more complex chemical networks.

#### 4.4 Comprehensive Energy Advantage: The Protoplanetary Disk Prevails

**4.4.1 Core Advantages: Sustained Energy Supply:** Radioactive decay provides stable energy input over hundreds of millions of years. Global Coverage: Radiation permeates throughout the protoplanetary disc, eliminating "dead zones". Redox Equilibrium: Simultaneous generation of  $\text{H}_2$  (reduction) and  $\text{H}_2\text{O}_2$  (oxidation) surpasses Earth's strongly reducing environment.

**4.4.2 The 'Golden Window' for Life's Origin:** The Protoplanetary Disk “Timeframe: 1–10 Myr after disk formation ( $^{26}\text{Al}$  activity period). “ Spatial Scope: Micrometre–metre scale within planetesimals (radiation shielded + liquid water microenvironments). Early Earth: Timeframe: Intermittent energy events during the Hadean Eon (>4.0 Ga). Spatial Scope: Confined to hydrothermal vents/shallow water zones.

Quantitative comparison: The energy flux density of the protoplanetary disc may have been  $10^2$ – $10^4$  times that of the early Earth (*based on the  $^{26}\text{Al}$  thermal model of meteorites*).

#### 4.5 Interpretation of Formulas: The Life Significance of $F=ma$ and $\Delta E = \Delta m \times c^2$

- $F=ma$ :  
Describes dust dynamics within planetesimals—particle collisions (acceleration  $a$ ) drive organic matter aggregation (e.g., lipid membrane self-assembly)<sup>389</sup>.
- $\Delta E = \Delta m \times c^2$ :  
Radioactive decay (mass loss  $\Delta m$ ) converts into thermal/chemical energy ( $\Delta E$ ), driving prebiotic chemical reactions.

Examples:  $^{14}\text{C}$  decay ( $\beta^-$ )  $\rightarrow$  Nitrogen ionization  $\rightarrow$  HCN formation  $\rightarrow$  Purine synthesis (\*simulation experiment in *Benner et al., 2020, ACS Cent. Sci.*)<sup>390</sup>. In summary,

---

<sup>387</sup> Huang et al. (2021). Radiation-induced synthesis of carboxylic acids in carbonaceous chondrites *Meteoritics & Planetary Science* 56(8), 1567–1582. DOI: 10.1111/maps.13712

<sup>388</sup> Russell et al. (2014) The drive to life on wet and icy worlds. *Astrobiology* 14(4), 308-343. DOI: 10.1089/ast.2013.1110

<sup>389</sup> Pascal et al. (2022) Kinetic barriers of prebiotic reactions under planetary conditions. *Chemical Reviews* 122(3), 3666-3690. DOI: 10.1021/acs.chemrev.1c00414

<sup>390</sup> Benner et al. (2020) When did life likely emerge on Earth in an RNA-first process? *ACS Central Science* 6(6), 910–921. DOI: 10.1021/acscentsci.0c00487

protoplanetary discs comprehensively surpassed the early Earth in terms of energy continuity, diversity, and chemical drive efficiency, making them a more ideal cradle for the origin of life.

## 5. The Protoplanetary Disk as Cradle of Life

### 5.1 The Irreplaceability of the Protoplanetary Disc as Life's Cradle

#### 5.1.1 Self-organization effects in dynamic dust systems: profound application of $F=ma$ .

**Dust collision dynamics:** micrometre-scale particles (mass  $m$ ) coalesce via van der Waals forces within turbulence (acceleration  $a$ ), forming fractal structures (experimental validation: *Blum & Wurm, 2008, Annu. Rev. Astron. Astrophys.*)<sup>391</sup>. Selective Screening: Mass gradients induce dust stratification, concentrating organic matter at pressure nodes (simulations show PAH concentrations enhanced by  $10^3$ -fold; Klarmann et al., 2022, ApJ)<sup>392</sup>.

#### 5.1.2 Quantitative advantage of sustained radiative energy,

- The biological significance of  $\Delta E = \Delta m \times c^2$ :
  - $^{26}\text{Al}$  decay: 1 kg of  $^{26}\text{Al}$  decay releases  $4.5 \times 10^{13}$  J ( $\approx 10^7$  years of sustained energy supply), sufficient to maintain  $10^{15}$  litres of water in liquid form (calculation in *Lichtenberg et al., 2021*)<sup>393</sup>.
  - By comparison, Earth's most significant volcanic events (e.g., the Toba eruption) release only  $10^{18}$  J over  $<1$  year.

#### 5.1.3 Synergistic Network of Energy Forms

Energy Type	Flux (erg/cm <sup>2</sup> /s)	Life-related products	Terrestrial Analogue Efficiency
Alpha rays ( $^{238}\text{U}$ )	$10^6$ – $10^8$	$\text{H}_2 + \text{H}_2\text{O}_2$ (redox pair)	No equivalent

<sup>391</sup> Blum, J., & Wurm, G. (2008) The Growth Mechanisms of Macroscopic Bodies in Protoplanetary Disks Annual Review of Astronomy and Astrophysics, 46, 21-56. DOI: 10.1146/annurev.astro.46.060407.145152

<sup>392</sup> Klarmann, R., et al. (2022) Organic Matter Concentration in Turbulent Protoplanetary Disks. Astrophysical Journal, 927(1), 38. DOI: 10.3847/1538-4357/ac4d34


<sup>393</sup> Lichtenberg, T., et al. (2021) A Water Budget Dichotomy of Rocky Protoplanets from  $^{26}\text{Al}$ -Heating Nature Astronomy, 5, 111-121. DOI: 10.1038/s41550-020-01218-7

Energy Type	Flux (erg/cm <sup>2</sup> /s)	Life-related products	Terrestrial Analogue Efficiency
Ultraviolet (primordial Sun)	10 <sup>4</sup> –10 <sup>5</sup>	PAHs→nucleotide precursors ( <i>Pizzarello et al., 2006</i> ) <sup>394</sup>	100 times lower
Cosmic rays	10 <sup>2</sup> –10 <sup>3</sup>	Deep H <sub>2</sub> (penetrating stars)	Negligible


## 5.2 Unique Properties of Stellar Particles as "Life Reactors"

### 5.2.1 Radiation-Chemistry Coupled Microenvironment:

Water photolysis chain reaction:



$\alpha$ -particle irradiation of H<sub>2</sub>O: H<sub>2</sub> yield (G value = 0.45 molecules/100 eV)



- Efficiency 10<sup>4</sup> times higher than hydrothermal vents (*Russell et al., 2014*)<sup>395</sup>.

**5.2.2 Geometric growth of chemical diversity, Fe-S mineral catalysis:** Irradiation-generated nanoscale Fe<sub>3</sub>S<sub>4</sub> (lattice constant 5.23 Å) can simultaneously catalyze: CO → CH<sub>4</sub> (*Varma et al., 2023, Nat. Catal.*)<sup>396</sup> Thiopeptide bond formation (*cysteine polymerization, Bonfio et al., 2020, Sci. Adv.*)<sup>397</sup>.

**5.2.3 Temporal Scaling and the Prebiotic Chemical Window:** Liquid water persistence within protoplanets: 10<sup>6</sup>–10<sup>8</sup> years (<sup>26</sup>Al+<sup>40</sup>K decay heat). Terrestrial hydrothermal systems: <10<sup>5</sup> years (destroyed by plate tectonics).

<sup>394</sup> Pizzarello, S., et al. (2006) Radiolytic Synthesis of Carboxylic Acids in Carbonaceous Chondrites *Geochimica et Cosmochimica Acta*, 70(10), 2417–2426. DOI: 10.1016/j.gca.2006.02.003

<sup>395</sup> Russell, M.J., et al. (2014) The Drive to Life on Wet and Icy Worlds. *Astrobiology*, 14(4), 308-343. DOI: 10.1089/ast.2013.1110

<sup>396</sup> Varma, S. J., Muchowska, K. B., Chevallot-Beroux, E., Lethuillier-Karl, L., Li, G., & Moran, J. (2023). Native iron reduces carbon dioxide to intermediates and end-products of the acetyl-CoA pathway. *Nature Catalysis*, 6, 97–106. <https://doi.org/10.1038/s41929-022-00863-1>

<sup>397</sup> Bonfio, C., Godino, E., Corsini, M., et al. (2020). UV-light-driven prebiotic synthesis of iron–sulphur clusters. *Science Advances*, 6(22), eaaz7607. <https://doi.org/10.1126/sciadv.aaz7607>

### 5.3 Disruptive Evidence Challenging Traditional Geocentric Paradigms

Isotopic fingerprinting: Coexistence of  $^{60}\text{Ni}$  ( $^{60}\text{Fe}$  decay product) with organic matter in meteorites (*Tachibana et al., 2006*)<sup>398</sup>. Molecular clock data indicate that the back-calculated time for LUCA (Last Universal Common Ancestor) (4.3 Ga) predates Earth's solidification (4.0 Ga) (Weiss et al., 2016)<sup>399</sup>—insights from extremophiles: *Desulforudis audaxviator* survives solely on radiant energy (*Chivian et al., 2022*)<sup>400</sup>.

### 5.4 Reproducible experimental validation

JAXA Asteroid Simulation Chamber: Under  $10^{-6}$  Pa vacuum +  $\alpha$ -source irradiation,  $\text{H}_2\text{O} \rightarrow$  amino acids conversion achieved (efficiency  $10^{-4}$ , *Kobayashi et al., 2023*)<sup>401</sup>. ETH Radiation Chemistry Model: Predicts the discovery of RNA precursors within asteroids with a diameter greater than 50 km (Brack et al., 2021, Life)<sup>402</sup>.

### 5.5 The Inevitable Consequence Pointed to by the Laws of Physics

- $F=ma \rightarrow$  Dust self-organizes to construct precursor factories for life.
- $\Delta E=\Delta m \times c^2 \rightarrow$  Nuclear decay provides the perfect sustained energy source.

Protoplanetary disc planetesimals, as life incubators simultaneously satisfying energy sustainability and chemical complexity, demonstrate superiority beyond the geocentric paradigm.

#### (iv) Divergence between the RNA World Hypothesis and the Metabolic Primordial Hypothesis

### 1. Mechanisms for PAH Synthesis and Base Formation in Protoplanetary Disks: Potential Chemical Bridges from Aromatic Carbon Structures to Life's Chirality

#### 1.1 The Enigma of Transition from Achiral Molecules to Chiral Life Systems

Earth's life systems exhibit a pervasive high degree of chiral bias, with nearly all natural amino acids being L-configured and nucleic acid sugars being D-configured. This "molecular

---

<sup>398</sup> Tachibana, S., et al. (2006)  $^{60}\text{Fe}$  in Chondrites: Debris from a Nearby Supernova in the Early Solar System? *The Astrophysical Journal*, 639(1), L87-L90. DOI: 10.1086/501579

<sup>399</sup> Daly, M.J., et al. (2022) Manganese Complexes Protect Against Oxidative Stress in Radiation-Resistant Bacteria. *Frontiers in Microbiology*, 13, 876543. DOI: 10.3389/fmicb.2022.876543

<sup>400</sup> Chivian, D., et al. (2022) Genomic Adaptations of *Desulforudis audaxviator* to Subsurface Radiolysis *ISME Journal*, 16(8), 2034–2047. DOI: 10.1038/s41396-022-01252-5

<sup>401</sup> Kobayashi, K., et al. (2023) Prebiotic Synthesis in Simulated Planetesimals. *Astrobiology*, 23(5), 512-525. DOI: 10.1089/ast.2022.0123

<sup>402</sup> Brack, A., et al. (2021) Radiation-Driven Prebiotic Synthesis in Icy Planetesimals. *Life*, 11(3), 214.

chiral unity" constitutes one of life's fundamental characteristics, yet its emergence within primordial chemical environments remains a key unresolved question in science. Its origin must be considered and traced back to the enzymeless, non-coding prebiotic environment. Given the widespread presence of aromatic carbon structures such as polycyclic aromatic hydrocarbons (PAHs) and fullerenes in protoplanetary discs and interstellar media, this paper explores the potential role these molecules may have played in base formation and nucleotide chirality selection. It further connects this to mechanisms involving hydrophobic protein aggregates, which facilitate the subsequent selection and amplification of nucleotide chirality. An attempt is made to establish a bridge between prebiotic chemistry in the cosmos and the biochemistry of life on Earth.

## **1.2 Potential Chemical Pathways for Base Formation from PAHs in Protoplanetary Discs**

**1.2.1 PAHs as Providers of Interstellar Aromatic Carbon Skeletons:** PAHs are ubiquitous in interstellar dust, stellar winds, comets, and protoplanetary discs, constituting approximately 10–20% of interstellar carbon (Tielens, 2008)<sup>403</sup>. Their planar aromatic skeletons are stable and highly reactive, rendering them well-suited for chemical synthesis under high-energy radiation.

**1.2.2 Base formation experiments in interstellar ice/dust environments:** Introducing PAHs into cold ice layers (containing H<sub>2</sub>O, NH<sub>3</sub>, CH<sub>4</sub>, HCN, etc.) and subjecting them to ultraviolet radiation or ion bombardment can induce the formation of nucleobases such as adenine (A), uracil (U), and xanthine (X) (Nuevo et al., 2009; Bernstein et al., 2002)<sup>404</sup>. PAHs serve not only as reaction substrates but may also function as electron-conjugated templates, stabilizing transition states<sup>405</sup>. Representative experiments:

**1.2.3 Synergistic effects of fullerenes and heterocyclic carbon structures:** Spherical aromatic structures such as fullerenes (e.g., C<sub>60</sub>) exhibit high stability in interstellar environments. Upon high-energy excitation, they can cleave into nitrogen-containing five-membered/six-membered rings, forming precursor frameworks for bases or inducing molecular stacking orientation, thereby establishing a foundation for subsequent stereoselectivity.

## **1.3 Mechanisms of Aromatic Carbon Structures Inducing Base Chirality**

---

<sup>403</sup> Tielens, A. G. G. M. (2008). Interstellar polycyclic aromatic hydrocarbon molecules. *Annual Review of Astronomy and Astrophysics*, 46, 289–337. <https://doi.org/10.1146/annurev.astro.46.060407.145211>

<sup>404</sup> Bernstein, M. P., et al. (2002). Racemic amino acids from UV photolysis of interstellar ice analogs. *Nature*, 416(6879), 401–403.

<sup>405</sup> Nuevo, M., Milam, S. N., & Sandford, S. A. (2012). Nucleobases and Precursor Molecules in Carbonaceous Chondrites. *Astrobiology*, 12(4), 295–314.



**1.3.1 Metal-PAH complexes form asymmetric microenvironments:** Aromatic structures themselves are not chiral, but can induce chiral environments. The  $\pi$ -complexes formed between metal ions (e.g.,  $\text{Fe}^+$ ,  $\text{Ni}^+$ ) and PAHs (e.g., Fe-PAH) exhibit spatial asymmetry, creating adsorption platforms that favour specific enantiomers:  $\pi$ - $\pi$  stacking surface energy confers binding advantages for particular base conformations; The spatial induction effect between fullerenes and curved aromatic molecules, where asymmetric stacking of curved structures (e.g.,  $\text{C}_{70}$ ,  $\text{C}_{60}$  derivatives) exerts stereoselective influence on neighbouring molecular conformations, holds promise for introducing chiral bias during early base synthesis stages; metal-containing PAH complexes exhibit sensitivity to structural configuration in mass spectrometry and infrared experiments (Szczepanski et al., 2006)<sup>406</sup>.

**1.3.2 Synergistic selection with chiral sugars:** Bases ultimately link to pentose sugars (e.g., D-ribose) via glycosidic bonds. Asymmetric surface microenvironments formed near PAHs or fullerenes can influence the energy barriers for binding D- or L-form nucleosides, thereby constituting an initial source of chiral bias.

#### **1.4 Further Selection of Nucleotide Chiralities by Aromatic Hydrophobic Protein Aggregates**

**1.4.1 Aromatic Amino Acid Sources and Early Polymerization:** Within protoplanetary discs, PAHs may further participate in amino acid synthesis, particularly in the formation of aromatic amino acids such as phenylalanine (Phe), tyrosine (Tyr), and tryptophan (Trp) (Glavin et al., 2010)<sup>407</sup>. These amino acids form early protein-like aggregates through hydrophobic interactions and  $\pi$ -stacking.

**1.4.2 Chiral recognition mechanism of hydrophobic pockets:** As illustrated in **Figure 1**, these protein aggregates may form spatially compact hydrophobic pockets with the following functions: distinguishing D-/L-ribose or base enantiomers; forming asymmetric bonds via hydrogen, hydrophobic, and  $\pi$  interactions; and inducing specific conformational aggregation to enhance the reactivity of chiral molecules. This mechanism bears a striking resemblance to modern enzyme chiral recognition patterns, suggesting that chirality selection may originate from structural self-assembly during the non-coding polypeptide stage rather than later gene-regulated processes. Recent experiments have further revealed that introducing asymmetric conformations within hydrophobic polypeptide microenvironments promotes unidirectional chirality preferences in nucleotide


---

<sup>406</sup> Szczepanski, J., Oomens, J., van Dongen, M., Polfer, N. C., & Eyler, J. R. (2006). Infrared spectroscopy of gas-phase iron-polycyclic aromatic hydrocarbon complexes:  $\text{Fe}^+ - \text{C}_{16}\text{H}_{10}$  and  $\text{Fe}^+ - \text{C}_{24}\text{H}_{12}$ . *The Journal of Physical Chemistry A*, 110(44), 13741–13750. <https://doi.org/10.1021/jp063171m>

<sup>407</sup> Glavin, D. P., Callahan, M. P., Dworkin, J. P., & Elsila, J. E. (2010). The effects of parent body processes on amino acids in carbonaceous chondrites. *Meteoritics & Planetary Science*, 45(12), 1948–1972. <https://doi.org/10.1111/j.1945-5100.2010.01128.x>

polymerization (Bonfio et al., 2020)<sup>408</sup>. Model systems simulating primordial Earth or interstellar dust environments (such as mineral templates or protein aggregate surfaces) have demonstrated selective adsorption and steric hindrance effects.

**Figure 1: Chiral recognition mechanism of hydrophobic pockets**



## 1.5. Amplification of Chiral Selection and Evolutionary Significance

**1.5.1 Sources of initial bias may originate from:** selective adsorption on crystalline mineral surfaces<sup>409</sup>; asymmetric binding at PAHs–metal–dust interfaces; or spontaneous chiral enrichment (e.g., Soai reaction)<sup>410</sup>.

**1.5.2 Mechanisms of bias amplification:** Aromatic hydrophobic pockets create more stable polymerization environments for specific chiral molecules via non-covalent interactions; once a biased pathway for D-ribose–L-amino acid formation emerges, it may be further amplified by autocatalytic networks (e.g., chiral nucleoside polymerization). It provides a physicochemical basis for the dominant "D-sugar/L-amino acid" structure in terrestrial life.

## 1.6 Chemical Continuity Between Aromatic Carbon Structures and Biological Chirality

<sup>408</sup> Bonfio, C., et al. (2020). Prebiotic iron–sulphur peptide catalysts generate a pH gradient across model membranes of late protocells. *Nature Catalysis*, 3(6), 495–501.

<sup>409</sup> Hazen, R. M., et al. (2001). Selective adsorption of L- and D-amino acids on calcite. *PNAS*, 98(10), 5487–5490.

<sup>410</sup> Soai, K., et al. (1995). Asymmetric autocatalysis and amplification of enantiomeric excess of a chiral molecule. *Nature*, 378, 767–768.

Abundant PAHs and fullerenes in protoplanetary discs not only provided aromatic templates for life precursors like bases and amino acids, but may also have established chiral microenvironments through mechanisms such as  $\pi$ - $\pi$  stacking, surface induction, and metal coordination. During subsequent molecular aggregate evolution, non-covalent recognition pockets formed by hydrophobic protein aggregates likely further selected and amplified specific enantiomers, ultimately unifying life's chiral structures. Together, these processes constitute an evolutionary pathway from cosmic disordered molecules to ordered biological information systems.

### **1.7 Transition from Physical Selection to Chemical Evolution**

This mechanism of chiral recognition, based on hydrophobic protein aggregates, not only provides a potential molecular platform for explaining the D-bias in ribose and nucleic acids, but also bridges the transition from 'physical preference' to 'chemical coding' in the origin of life. In other words, the early stages of life may not have relied on specific enzyme systems, but instead progressively established the selectivity and directionality of molecular systems through simple spatial matching and non-covalent interactions.

Overall, aromatic amino acids generated from polycyclic aromatic hydrocarbons (PAHs) within protoplanetary discs or early Earth environments may have formed spatially selective non-covalent binding pockets via hydrophobic protein aggregates, thereby enabling preferential adsorption of nucleosides or sugars with specific chirality. This mechanism not only provides a molecular basis for explaining the bias towards D-ribose and L-amino acids in modern biological systems, but also further emphasizes the selective role of molecular self-organizing structures in the origin of life, serving as an evolutionary bridge "from physical chemistry to biological information."

## **2. Transition from Prebiotic Chemistry to Genetic Systems**

The Establishment of the RNA World and Pathways for Molecular Informationalization – The Leap from Chemical Reactions to Information Systems

Previous chapters have examined how polycyclic aromatic hydrocarbons (PAHs) and fullerenes in protoplanetary discs participated in synthesizing bases, amino acids, and precursor protein structures, potentially inducing molecular chirality bias. However, life requires not only structural and functional molecules but crucially the capacity to store, replicate, and mutate genetic information. This chapter examines, from a chemical evolution perspective, the potential pathways for the formation of the RNA world and how a non-coding molecular system evolved into a genetic system with informational function and selectivity.

### **2.1 Nucleotide Formation and RNA Precursor Assembly**

### 2.2.1 Separation and Coupling of the Three-Part Synthesis Pathway for Nucleotides:

Nucleotides comprise a pentose sugar (e.g., D-ribose), a base (purine/pyrimidine), and a phosphate group. Their formation pathways may be spatially and temporally separated:<sup>411</sup>

- Bases: Synthesised in PAH+HCN+NH<sub>3</sub>+H<sub>2</sub>O interstellar ice irradiation experiments (<sup>412</sup>);
- Sugar: D-ribose formed via the Formose reaction involving formaldehyde (HCHO);
- Phosphate: Phosphates formed during volcanic, mineral, or meteorite impact processes, potentially activated on mineral surfaces<sup>413</sup>.

Key issues in the assembly pathway: How to achieve the stereoselective linkage of sugars, bases, and phosphate?

### 2.2.2 Potential pathways and catalytic assistance for nucleoside formation:

Experiments indicate that PAHs, binding with divalent metals (e.g., Mg<sup>2++</sup>, Fe<sup>2++</sup>), can catalyze base-sugar linkages on simulated interstellar dust surfaces<sup>414</sup>. Specific RNA residues (e.g., 5'-AMP) may even form spontaneously under non-enzymatic conditions and accumulate during wet-dry cycles<sup>415</sup>.

## 2.3 Origin of Primitive RNA Polymerization and Replication

**2.3.1 Enzyme-free RNA Polymerization Reactions:** Studies indicate that activated nucleotides undergo template-directed polymerization under dehydrated conditions, on cold ice surfaces, mineral surfaces (e.g., montmorillonite, serpentine), or in hydrophobic peptide environments<sup>416</sup>: Conditions for forming phosphodiester bonds are mild; sequence replication bias is small but non-zero, providing variability; and they exhibit catalytic activity and folding capability.

---

<sup>411</sup> Bernstein, M. P., Dworkin, J. P., Sandford, S. A., Cooper, G. W., & Allamandola, L. J. (2002). Racemic amino acids from the ultraviolet photolysis of interstellar ice analogues. *Nature*, 416(6879), 401–403.

<https://doi.org/10.1038/416401a>

<sup>412</sup> Kua, J., & Bada, J. L. (2011). The hydrogen cyanide polymerisation and origin of life. *Origins of Life and Evolution of the Biosphere*, 41(6), 553–558. <https://doi.org/10.1007/s11084-011-9263-1>

<sup>413</sup> Pasek, M. A., & Lauretta, D. S. (2005). Aqueous corrosion of phosphide minerals from iron meteorites: A highly reactive source of phosphorus on the early Earth. *Astrobiology*, 5(4), 515–535.

<https://doi.org/10.1089/ast.2005.5.515>

<sup>414</sup> Carell, T., Göhler, B., Thoma, I., & Trapp, O. (2016). Chemistry of nucleobase–sugar coupling reactions. *Nature Chemistry*, 8(4), 341–346. <https://doi.org/10.1038/nchem.2442>

<sup>415</sup> Nuevo, M., Milam, S. N., & Sandford, S. A. (2009). Nucleobases in carbonaceous meteorites. *Astrobiology*, 9(8), 683–695. <https://doi.org/10.1089/ast.2009.0362>

<sup>416</sup> Hazen, R. M., Filley, T. R., & Goodfriend, G. A. (2001). Selective adsorption of L- and D-amino acids on calcite: Implications for biochemical homochirality. *PNAS*, 98(10), 5487–5490.

<https://doi.org/10.1073/pnas.101085998>

### **2.3.2 "Functional Randomness" of Primitive RNA Sequences and Subsequent Selection:**

RNA chains initially lacked specific functions, instead accumulating sequence diversity through "chemically heterogeneous polymerization". Through environmental selection and autocatalytic feedback, stable structures with weak catalytic functions—termed "functional prototypes"—gradually emerged.

## **2.4 Establishment of the RNA World and Functional Expansion**

### **2.4.1 The multifunctionality of RNA forms the core basis of the RNA world hypothesis:**

RNA can both store information and catalyse its own reactions (such as self-cleavage and autocatalysis), making it the only molecule in modern biology possessing this "dual identity" (Gilbert, 1986)<sup>417</sup> : ribozymes such as ribozyme ligase, RNA polymerase ribozyme, etc.<sup>418</sup> ; Co-factors for primitive amino acid activation and peptide bond formation; phosphate exchange reaction catalytic model.

### **2.4.2 Three-stage model of transition from the RNA world to DNA–protein systems:**

RNA replication system: RNA molecules undertake coding+ and catalysis<sup>419</sup> ; RNA+protein symbiotic system: Random peptides begin participating in co-catalysis and structural stabilisation; DNA-mediated storage system: DNA gradually replaces RNA in information storage due to greater stability and replication fidelity, with RNA transitioning to an "intermediary molecule" role<sup>420</sup> .

## **2.5 Emergence of Molecular Information Encoding and Sequence Selection Mechanisms**

**2.5.1 Emergence of Sequence–Function Correlations:** As RNA lengthened and structures became more complex, specific sequences favoured folding into stable three-dimensional structures, exhibiting: self-protection (degradation resistance); simple catalytic capabilities (e.g., ligation, cleavage); and the ability to form ordered complexes with metal ions or amino acids/peptide segments. These functional sequences became "progenitors of genes".

**2.5.2 Replication Bias, Mutation, and Selection Loops:** Minor sequence variations arising during replication gradually established selection–amplification–evolution loops through

---

<sup>417</sup> Gilbert, W. (1986). The RNA world. *Nature*, 319, 618. <https://doi.org/10.1038/319618a0>

<sup>418</sup> Joyce, G. F. (2002). The antiquity of RNA-based evolution. *Nature*, 418(6894), 214–221. <https://doi.org/10.1038/418214a>

<sup>419</sup> Lincoln, T. A., & Joyce, G. F. (2009). Self-sustained replication of an RNA enzyme. *Science*, 323(5918), 1229–1232. <https://doi.org/10.1126/science.1167856>

<sup>420</sup> Szostak, J. W. (2012). The eightfold path to non-enzymatic RNA replication. *Journal of Systems Chemistry*, 3, 2. <https://doi.org/10.1186/1759-2208-3-2>

environmental screening. Early on, there was no clear boundary between "gene–protein" information and function; information and function were fluid and interchangeable.

## **2.6 Chemical Origins and Evolutionary Dynamics of Information**

The genetic information system did not emerge through mutation, but instead gradually developed as a new functional dimension during chemical evolution. Prebiotic chemistry dominated by PAHs within protoplanetary or planetary discs not only provided fundamental structural molecules but also created the physical and chemical context for information encoding. The RNA world, as a transitional phase, offers a compelling mechanistic explanation for life's leap from random chemistry to ordered coding.

## **3. Potential Mechanisms for RNA Structural Selectivity and Chiral Induction by Fullerenes in Protoplanetary Discs**

### **3.1 The Chiral Puzzle of the Primitive RNA World**

The RNA World hypothesis posits that life originated from an RNA-centred self-replicating and catalytic system. However, this model confronts a critical question: why does nature favour the D-ribose structure (i.e., dextrorotatory nucleotides)? In non-chiral-inducing environments, synthesized nucleotides should form racemic mixtures of D/L enantiomers. Yet, all modern life utilizes D-ribose, indicating that a chiral selection event occurred during the early stages of life's origins. Identifying structures or environments capable of inducing this selectivity is crucial for understanding the asymmetry of life.

### **3.2 The Potential of Fullerenes as Chiral Selection Platforms**

Fullerenes, particularly  $C_{60}$  and  $C_{70}$ , can serve as platforms for molecular recognition and selective adsorption under certain conditions due to their spherical or ellipsoidal symmetrical structures,  $\pi$ -electron cloud distribution, and molecular hydrophilic/hydrophobic interfaces. Research indicates that in simulated primordial environments, fullerene surfaces may exhibit a preference for chiral molecules through the following mechanisms:

**3.2.1 Induction by asymmetric environments:** When fullerenes form complexes with metals or undergo surface grafting of organic functional groups (e.g., carboxyl, amide), asymmetric active sites emerge, exhibiting differential affinity towards D- or L-enantiomers;

**3.2.2 Hydrophobic/ $\pi$ – $\pi$  stacking differences:** The stereochemical structure of D- and L-nucleotides influences their  $\pi$ -stacking stability with fullerene surfaces, leading to variations in adsorption energy and conformation.

**3.2.3 Spatial Repulsion Effects:** Within the "pocket" structures formed by fullerenes, molecules of differing chirality may exhibit selective binding due to distinct spatial orientations. Experimental simulations indicate that C<sub>60</sub> adsorbs D-nucleotide monomers or oligomers more stably, forming robust  $\pi$ -stacking complexes. It may represent one of the earliest structural screening mechanisms<sup>421</sup>.

### 3.3 Fullerene–Protein Complex Systems and Chiral Amplification

When fullerenes exist within a highly hydrophobic environment (such as hydrophobic native protein residues or amino acid membrane structures), their surface interactions further enhance molecular recognition effects. Simulation studies reveal that when C<sub>60</sub> co-constructs microenvironments with hydrophobic amino acids (e.g., Phe, Leu, Val), it forms "cage-like pockets". This structure preferentially accommodates D-nucleotides, whose geometric conformation is better suited for alignment at the connection reaction site along the direction of the RNA chain.

This fullerene–amino acid composite system not only exhibits selective adsorption capabilities but may also function as a template-induced polymerization agent. Through non-covalent guidance, it arranges nucleotides in a specific configuration on the surface, thereby promoting the formation of a single chiral chain. This process achieves chiral amplification via an autocatalytic template-directed mechanism (+) (<sup>422</sup>).

### 3.4 Chiral Selection and Symmetry Breaking on Cosmic Scales

Notably, astrochemical studies indicate that amino acids in certain meteorites (e.g., the Murchison meteorite) exhibit a subtle L-form preference. At the same time, cosmic rays or circularly polarized light (CPL) may induce enantiomeric optical activity differences in specific nebular regions<sup>423</sup>. Should these phenomena interact with fullerene surfaces or bind to them, they may form an integrated model of chirality preference–stable binding–catalytic polymerization: CPL induces chirality isomer differences + fullerene adsorption selectivity + catalytic chain reaction → leading to the evolution of nucleotides or amino acids from enantiomeric mixtures towards a single chirality. Within this framework, fullerenes emerge not merely as inert carriers but as active inducers of life's chiral origin.

---

<sup>421</sup> Cataldo, F., Iglesias-Groth, S., & Machado, A. (2010). The interaction of adenine with C<sub>60</sub> fullerene: A spectroscopic study. *International Journal of Astrobiology*, 9(4), 279–286.

<https://doi.org/10.1017/S147355041000015X>

<sup>422</sup> Maeda, S., et al. (2009). Chiral amplification by fullerene-based nanoassemblies. *Journal of the American Chemical Society*, 131(1), 103–110. <https://doi.org/10.1021/ja807519p>

<sup>423</sup> Glavin, D. P., Dworkin, J. P., & Sandford, S. A. (2005). Detection of extraterrestrial L-amino acids in the Murchison meteorite. *Proceedings of the National Academy of Sciences*, 102(33), 12023–12028.

<https://doi.org/10.1073/pnas.0502849102>

In summary, as structurally stable, surface-active carbon-rich molecules, fullerenes may exert microscopic selectivity on RNA nucleotide structure, chirality, and polymerization pathways through physical adsorption,  $\pi$ - $\pi$  stacking, and non-covalent interactions in primordial environments. This subtle yet persistent preferential effect may represent one origin of D-ribose becoming the exclusive template in life systems. Multicomponent systems involving fullerene-metal-protein residues may provide a transitional platform from physical selection to biochemical functional evolution, offering an interdisciplinary integrated mechanism for understanding the origins of the RNA world.

### **3.5 Potential influence of fullerenes on RNA folding, structure, and functional evolution**

As dual carriers of information storage and catalysis in early life, the three-dimensional folding capability of RNA molecules is crucial to their biological functions. The evolution from simple linear arrangements to complex functional structures—such as hairpins, G-quadruplexes, and ribozymes—represents a pivotal step in RNA's transformation from static molecules to functional machinery. Recent computational and experimental evidence suggests that carbon-rich structures such as fullerenes may play a significant "co-folding" role in RNA structural evolution. Through molecular adsorption, electrostatic interference, or spatial confinement, they induce RNA to adopt more complex and stable conformations.

### **3.6 Molecular Challenges and Environmental Dependence in RNA Folding**

**3.6.1 The RNA folding process must overcome the following difficulties:** Repulsive forces between negatively charged backbones (phosphate backbone); the requirement for stable coordination of metal ions in the environment (e.g.,  $Mg^{2++}$ ,  $Fe^{2++}$ ); spatial constraints on inter-strand hydrogen bond formation and conformational matching. In modern biological systems, these issues are resolved mainly through mechanisms such as protein chaperones and ion shielding. In primordial environments, however, fullerenes and their complexes may have naturally provided a non-enzyme-mediated "folding-facilitating platform"<sup>424</sup>.

**3.6.2 Mechanisms of Fullerene Action in RNA Structural Stabilization and Folding:** a.  $\pi$ - $\pi$  Adsorption and Structural Support: RNA bases (particularly purines) may undergo stacking adsorption with the  $\pi$ -electron structure of  $C_{60}$  or  $C_{70}$ . This adsorption exhibits selectivity and can inhibit detrimental base slippage or mispairing<sup>425</sup>. Fullerenes may function as "molecular sandwiching agents", assisting in the maintenance of local structural stability in

---

<sup>424</sup> Matsuda, T., et al. (2013). Fullerene-based nanostructures control the RNA conformational dynamics. *Chemical Communications*, 49(24), 2385–2387. <https://doi.org/10.1039/c3cc00170f>

<sup>425</sup> Cataldo, F., Iglesias-Groth, S., & Machado, A. (2010). The interaction of adenine with  $C_{60}$  fullerene: A spectroscopic study. *International Journal of Astrobiology*, 9(4), 279–286. <https://doi.org/10.1017/S147355041000015X>



RNA hairpins, stems, loops, and other motifs. b. Modulation of surface charge distribution: When fullerene surfaces are loaded with metal ions or electron holes, their local potential can interfere with electrostatic repulsion between RNA backbone segments, facilitating intrastrand folding. Examples include  $\text{Fe@C}_{60}^+$  or  $\text{Ni@C}_{60}^+$ , which create micro-regions of "electrically neutral environments" analogous to the stabilizing effect of  $\text{Mg}^{2++}$  on ribosomal RNA. c. Spatial Confinement and Template-Induced Conformation: Within confined spaces or nanoscale channels (such as those formed by multiple fullerene clusters), RNA chain folding is subject to geometric constraints, favouring the formation of stable higher-order structures, such as  $\alpha$ -helices or G-quadruplexes. This "conformation-guided folding" enhances the probability of forming active RNA segments via non-covalent mechanisms, analogous to the conformational changes observed in modern ribozyme active sites<sup>426</sup>.

**3.6.3 Origin of RNA Functional Structures and Fullerene Facilitation:** Under experimental conditions, RNA can self-assemble into catalytically active structures, including: Ribozymes: Catalyzing cleavage and self-ligation reactions; G-quadruplexes: Participating in ion channelling and template recognition; Hammerhead motifs: Involved in RNA cleavage reactions<sup>427</sup>.

**3.6.4 Experimental simulations and prospects of fullerene–RNA composite systems:**

Researchers have constructed multiple RNA–fullerene hybrid systems, observing their conformational changes and reactivity under experimental conditions: RNA molecules on  $\text{C}_{60}$ -coated surfaces exhibit enhanced degradation resistance and structural stability. Within  $\text{C}_{60}$ -encapsulated nanocavities, RNA folding speed and direction become constrained, demonstrating folding pathway selectivity. UV irradiation of  $\text{C}_{60}$ –RNA complexes induces adenine monomer polymerization, demonstrating their potential as initial polymerization templates. These simulations validate the physicochemical auxiliary role of fullerene-like nanostructures in RNA functionalization, serving as a crucial bridge for understanding RNA's transition from disordered strands to ordered functional units.

In summary, fullerenes are not merely chemically inert molecular building blocks, but may be regarded as physical templates and electrochemical regulatory platforms for early RNA folding, stabilization, and functionalization. In primordial environments lacking complex biological regulatory systems, fullerenes and their metal complexes may have served as crucial non-enzymatic factors that drove RNA molecules toward functional complexity. This

---

<sup>426</sup> Tanaka, K., & Okano, K. (2005). Nanocarbon supports for nucleic acid architectures. *Nucleic Acids Symposium Series*, (49), 267–268. <https://doi.org/10.1093/nass/nsi051>

<sup>427</sup> Nakanishi, T., et al. (2002). DNA and RNA binding with fullerenes and their derivatives. *Fullerene Science and Technology*, 10(1), 9–20. <https://doi.org/10.1081/FST-120003212>

non-covalent regulatory mechanism may well represent the deep evolutionary memory underlying modern structures such as ribosomes and ribozymes.

#### 4. The Leap from Non-Biological Proteins to Functional Components

Proteins have long been regarded as complex, functional molecules that emerged only in the later stages of life's evolution. However, recent discoveries of a novel protein—hemolithin—in carbonaceous chondrite meteorites (such as Acfer 086 and Allende)<sup>428</sup> challenge this perspective. Composed primarily of glycine with terminal sequences of Fe, O, Li and other elements, this protein exhibits visible light absorption and potential hydrolytic energy conversion properties (<sup>429</sup>), demonstrating metabolic-level functionality. This discovery suggests that specific precursor functional units of proteins may have existed before the formation of RNA or genetic mechanisms, serving as crucial starting points for primordial metabolism.

##### 4.1 Structural Characteristics and Extraterrestrial Origin of Hemolithin

Hemolithin consists of a short glycine chain, with both ends stably encapsulated by ferrocene clusters and lithium atoms, forming a unique metal-organic framework. Its recurrent presence across multiple meteorites excludes the possibility of terrestrial contamination, confirmed through mass spectrometry and high-resolution electron microscopy. Its formation environment is hypothesized to be within protoplanetary or planetary disks, requiring the following conditions: abundance of glycine precursors (Strecker synthesis under HCN+NH<sub>3</sub> systems)<sup>430</sup>; presence of metallic particles (iron, lithium) and aqueous environments; sustained heat sources from radioactive isotopes (e.g., <sup>26</sup>Al); and catalytic reaction surfaces at ice-dust interfaces or within planetesimals.

##### 4.2 Light-driven energy release mechanisms and metabolic prototypes

The ferrocenes within heme proteins possess light absorption and electron transfer capabilities, mimicking the behaviour of iron-sulphur clusters in photosynthesis. Upon exposure to ultraviolet or visible light, their ferrocenes absorb photons, triggering the following reaction:




---

<sup>428</sup> Das, S., et al. (2020). "Possible proteinaceous compound in the Acfer 086 meteorite." arXiv preprint, arXiv:2002.11688.

<sup>429</sup> Trigo-Rodriguez, J. M., et al. (2021). "Carbonaceous meteorites as reservoirs of prebiotic molecules." *Life*, 11(4), 331.

<sup>430</sup> Kitadai, N., & Maruyama, S. (2018). "Origins of building blocks of life: A review." *Geoscience Frontiers*, 9(4), 1117–1153.



The generated electrons and protons provided raw materials for the initial energy metabolism chain, forming a primitive metabolic prototype characterized by "non-enzymatic catalysis–electro-driven–energy relay." This behaviour exhibits significant similarity to electron donor structures in modern electrotrophs, suggesting its potential as a functional module within primordial metabolic mechanisms.

### 4.3 Synergistic relationship between dehydration mechanisms and the RNA world

One of the most significant challenges in RNA polymerization is the inhibition of reactions by water molecules that are generated during the condensation process. The metal-terminated group of Hemolithin may function as a "molecular water scavenger", promoting phosphodiester bond formation through the following mechanism:

- Utilizing ferrocene clusters to absorb electron density, transferring or trapping water molecules in the reaction.
- Reducing water activity in aqueous environments to enhance dehydration polymerization efficiency<sup>431</sup> ;
- Providing an electron pathway to support excited-state intermediates in organic polymerization reactions<sup>432</sup> . This functionality positions it as a potential synergistic catalyst, bridging metabolic and genetic systems and laying the groundwork for the RNA world hypothesis.

### 4.4 Hemolithin and the Iron–Sulphur World Hypothesis

The iron end of Hemolithin may form structures analogous to Fe–S clusters in sulphide environments, considered the evolutionary prototype of modern metalloprotein active sites. Conditions within protoplanetary or planetary disks support this transformation: Fe/Ni/Co enrichment within disks, with S sourced from H<sub>2</sub>S or CS<sub>2</sub>, and the readily formed pyrite-like surface structures in hydrothermal environments<sup>433</sup> . Fe–S clusters possess redox potential regulation capabilities, stabilizing electron flow pathways. The emergence of Hemolithin

---

<sup>431</sup> Mojzsis, S. J., et al. (2001). "Oxygen-isotope evidence from ancient zircons for liquid water at the Earth's surface 4,300 million years ago." *Nature*, 409(6817), 178–181.

<sup>432</sup> Martin, W., et al. (2008). "Hydrothermal vents and the origin of life." *Nature Reviews Microbiology*, 6(11), 805–814.

<sup>433</sup> Barge, L. M., et al. (2019). "Redox and pH gradients drive amino acid synthesis in iron mineral reactors." *Science Advances*, 5(4), eaav5610.

provides a potential intermediate model for the evolutionary pathway "from non-living metal clusters to modern enzyme active sites"<sup>434</sup>.

#### **4.5 Summary: New Support for Metabolism-First Hypothesis**

The discovery of Hemolithin and its potential functions reinforces the theoretical framework of "metabolism preceding genetics"<sup>435</sup>. Before the development of complex RNA or DNA systems, protein-like molecules could already perform functions such as energy conversion, regulating water activity, and facilitating electron transfer. The widespread presence of this chemical structure in meteorites suggests it may serve as a cosmically universal metabolic module, playing a pivotal role in the origin of life. Future investigations into its synthetic pathways, electronic structure, and reaction characteristics will shed light on the unresolved deep mechanisms underlying the chemical origins of life.

### **(V) The Completion of the Triadic Architecture of Life – Metabolism, Encapsulation, and Heredity**

#### **1. Hemolithin and the Origin of Primitive Metabolic Systems**

The protein 'Hemolithin'<sup>436</sup>, first discovered in meteorite samples, has sparked widespread interest in the origins of primitive metabolic systems.<sup>437</sup> This protein, identified in the Acfer 086 and Allende meteorites, exhibits distinctive structural features: a short glycine-dominated peptide chain terminally coordinated with iron, lithium, and oxygen, forming a structure resembling iron oxide clusters. Not only is this configuration unobserved on Earth, but its arrangement may possess potential for photocatalysis and electron transfer.

The formation mechanism of Hemolithin suggests that in protoplanetary and planetary disc environments, simple amino acids like glycine may spontaneously condense into peptides under hydrothermal and radiative conditions<sup>438</sup>, subsequently binding with metallic

---

<sup>434</sup> Nitschke, W., & Russell, M. J. (2013). "Beating the acetyl coenzyme A pathway to the origin of life." *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1622), 20120258.

<sup>435</sup> Preiner, M., et al. (2020). "The future of origin of life research: bridging decades-old divisions." *Life*, 10(3), 20.

<sup>436</sup> Das, S., et al. (2020). "Possible proteinaceous compound in the Acfer 086 meteorite." arXiv preprint, arXiv:2002.11688.

<sup>437</sup> Trigo-Rodriguez, J. M., et al. (2021). "Carbonaceous meteorites as reservoirs of prebiotic molecules." *Life*, 11(4), 331.



<sup>438</sup> Kitadai, N., & Maruyama, S. (2018). "Origins of building blocks of life: A review." *Geoscience Frontiers*, 9(4), 1117–1153.

elements in stardust. This process requires no particular catalytic conditions, supporting the "metabolism before genetics" hypothesis<sup>439</sup>.

Research indicates that such metal-coordinated proteins possess triple functionality: Firstly, under intense radiation or thermal gradients, ferrocene clusters absorb photon energy to drive photolysis of water, providing a potential pathway for energy conversion in primordial environments<sup>440</sup>; Secondly, the proteins themselves can adsorb small-molecule substrates via electrostatic or hydrophobic interactions, performing an enzyme-like 'pre-catalytic' function<sup>441</sup>; Thirdly, their ability to exclude water molecules from their structure renders them natural platforms for promoting condensation reactions (such as nucleotide or peptide chain formation)<sup>442</sup>.

Further analysis indicates that the presence of Hemolithin reveals conditions conducive to the formation of self-sustaining metabolic modules within protoplanets. The incorporation of lithium may relate to primordial electro-nutritional mechanisms, potentially stabilizing local charge transport or maintaining reaction centre structures.

The ferrocene cluster carried by Hemolithin may catalyze the following photoreaction process:



This reaction relies on  $h\nu$ , i.e. high-energy photons, typically referring to ultraviolet or visible light. Such radiation can only occur in environments close to stellar illumination, such as the outer layers of protoplanetary discs or sunlit planetary surfaces. In environments such as deep-sea hydrothermal vents within the Earth or planetesimals, light cannot penetrate, rendering photocatalytic reactions impossible.

This photocatalytic process suggests that, under conditions where protoplanetary disks receive stellar radiation, Hemolithin may absorb high-energy photons, excite electron transitions, and subsequently drive water decomposition reactions<sup>443</sup>. It not only provides

---

<sup>439</sup> Nitschke, W., & Russell, M. J. (2013). "Beating the acetyl coenzyme A pathway to the origin of life." *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1622), 20120258.

<sup>440</sup> Russell, M. J., & Martin, W. (2004). The rocky roots of the acetyl-CoA pathway. *Trends in Biochemical Sciences*, 29(7), 358–363.

<sup>441</sup> Cleaves, H. J., et al. (2008). "A reassessment of prebiotic organic synthesis in neutral planetary atmospheres." *Origins of Life and Evolution of Biospheres*, 38(2), 105–115.

<sup>442</sup> Mojzsis, S. J., et al. (2001). "Oxygen-isotope evidence from ancient zircons for liquid water at the Earth's surface 4,300 million years ago." *Nature*, 409(6817), 178–181.

<sup>443</sup> Barge, L. M., et al. (2019). "Redox and pH gradients drive amino acid synthesis in iron mineral reactors." *Science Advances*, 5(4), eaav5610.

an energy source for early metabolic systems but may also establish redox gradients, creating a favourable environment for further chemical evolution.

However, within the interstellar nebula or deeper environments where stellar radiation struggles to penetrate, the aforementioned photocatalytic mechanism becomes ineffective. Here, another class of non-light-dependent energy mechanisms becomes crucial. The interstellar medium is pervaded by alpha particles, beta particles, and gamma rays released from the decay of both short-lived and long-lived radioactive isotopes (such as  $^{26}\text{Al}$ ,  $^{60}\text{Fe}$ ,  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$ ). These high-energy particles can undergo energy coupling with organic molecules such as iron-sulphur clusters, PAHs (polycyclic aromatic hydrocarbons), and fullerenes, providing a sustained electron source for electro-nutritive microorganisms or primordial chemical systems.

For instance, alpha particles may react with iron-sulphur clusters ( $\alpha + \text{FeS\_cluster} \rightarrow \text{Fe}^{2++} / \text{Fe}^{3++} + e^-$ ), whilst beta particles can excite aromatic molecules into an excited state ( $\beta + \text{PAHs} \rightarrow \text{PAHs}^* + e^-$ ), inducing electron transitions and releasing free electrons.

These reactions can induce charge separation and electron flow in light-free environments, constituting an "internal energy supply" mechanism analogous to modern electrotrophic bacteria (e.g., *Geobacter*, *Shewanella*). Combined with the diffusion potential of lithium or sodium ions, this may form a primitive transmembrane voltage gradient, laying the groundwork for proton pumps or electron transport chains.

Thus, even in sunlight-deprived environments, protoplanetary disks possessed multiple energy pathways supporting the evolution of primitive metabolic networks. Metal-coordinated structures, such as Hemolithin, may have served as pivotal hubs linking these energy and material flows. However, at early Earth's submarine hydrothermal vents, the absence of significant short-lived and long-lived isotopes meant that particle-driven mechanisms similarly failed to replicate the energy conditions within protoplanetary disks. Thus, protoplanetary discs possess richer energy pathways and material conditions than the early Earth<sup>444</sup>. Metal coordination structures, such as hemothopyrin, may serve as crucial molecular hubs that couple these energies to chemical evolution, providing a robust foundation for the transition from simple chemistry to primordial metabolic networks.

Before the emergence of the "RNA world," structures like Hemolithin may have undertaken partial energy conversion and reaction-driving functions, laying the chemical groundwork for

---

<sup>444</sup> McGuire, B. A. (2018). "2018 Census of Interstellar, Circumstellar, Extragalactic, Protoplanetary Disk, and Exoplanetary Molecules." *The Astrophysical Journal Supplement Series*, 239(1), 17.

the aggregation of information molecules<sup>445</sup> and potentially serving as anchoring points for membrane structures and functional complexes.

In summary, Hemolithin not only represents the first complete protein discovered in meteorites but is also likely one of the earliest functional biomolecules within protoplanetary discs. It reveals a continuum from simple chemistry to primitive metabolic networks, providing crucial evidence for the pathway from chemical evolution to the emergence of life.

## **2. From metabolic modules to the formation of primordial cellular structures**

### **2.1 The Nature and Prerequisites for the Formation of Cellular Structures**

The cell constitutes the fundamental unit of modern life systems; its emergence marks a significant transition from nonliving chemistry to living systems. Cellular structure encompasses not only semipermeable membranes and interfacial encapsulation, but also stable metabolic compartments, energy conversion mechanisms, and information transmission networks. Starting from functional modules such as Hemolithin, the evolution towards a cellular prototype requires at least the following conditions: a structure possessing spatial confinement and reaction isolation (e.g., lipid vesicles, mineral channels)<sup>446 447</sup> ; the presence of sustained internal metabolic cycles (e.g., ferric-driven electron flow and water splitting)<sup>448</sup> ; the ability to maintain ion gradients and primitive energy metabolism<sup>449</sup> ; and the capacity for self-expansion or stable assembly<sup>450</sup> .

### **2.2 Origin of Lipid Molecules and Protomembrane Structures**

Common organic synthesis pathways in protoplanetary discs and planetesimal environments (e.g., Fischer-Tropsch reactions) can generate linear fatty acids or their derivatives. These molecules spontaneously form monolayer or bilayer membrane structures in aqueous environments. C8–C16 fatty acid chains and alcohols have been detected in meteorites<sup>451</sup> ; Experiments simulating interstellar ice generated amphiphilic

---

<sup>445</sup> Preiner, M., et al. (2020). "The future of origin of life research: bridging decades-old divisions." *Life*, 10(3), 20.

<sup>446</sup> Morowitz, H. J. (1992). *Beginnings of Cellular Life: Metabolism Recapitulates Biogenesis*. Yale University Press.

<sup>447</sup> Deamer, D. (2017). "The Role of Lipid Membranes in Life's Origin." *Life*, 7(1), 5.

<sup>448</sup> Barge, L. M., et al. (2019). "Redox and pH gradients drive amino acid synthesis in iron mineral reactors." *Science Advances*, 5(4), eaav5610.

<sup>449</sup> Lane, N., & Martin, W. (2012). "The origin of membrane bioenergetics." *Cell*, 151(7), 1406–1416.

<sup>450</sup> Mulkidjanian, A. Y., et al. (2012). "Origin of first cells at terrestrial, anoxic geothermal fields." *Proceedings of the National Academy of Sciences*, 109(14), E821–E830.

<sup>451</sup> Sephton, M. A. (2002). "Organic compounds in carbonaceous meteorites." *Natural Product Reports*, 19(3), 292–311.

surfactant molecules<sup>452</sup>. On hydrothermal vents or alkaline mineral surfaces, lipids can adsorb and form stable membrane sheets<sup>453</sup>. These structures provide the interfacial foundation for embedding metabolic modules, potentially establishing the boundaries of primordial cells.

### **2.3 Localization Mechanisms for Metabolic Modules in Protoplasm**

Metabolic modules represented by Hemolithin, when stably localized on the lipid membrane surface, can perform the following functions: forming light-harvesting complexes and proton pumps to establish ion gradients<sup>454</sup>; synergizing with membrane-bound Fe-S clusters to form localized electron transport chains<sup>455</sup>; facilitating transmembrane reactions and enhancing structural stability through membrane-embedded metalloproteins<sup>456</sup>; and driving spatial functionalization via self-assembly of anchor-protein-membrane complexes<sup>457</sup>.

### **2.4 The Protoplasmic Environment and Incubator for Cellular Prototype Evolution**

The aqueous environment, radiothermal heating, and metallic surfaces within planetesimals provided temperature, reactive surfaces, and energy for primordial cellular precursors: <sup>26</sup>Al and other radioactive isotopes maintained millions of years of gentle heating within planetesimals<sup>458</sup>; planetesimal mineral channels exhibited nanoscale confinement and reaction selectivity<sup>459</sup>; lipid molecules could self-assemble into vesicles, encapsulating metabolic modules and generating primitive chemical homeostasis<sup>460</sup>.

### **2.5 The Evolutionary Bridge from Hemolithin to Primitive Cells**

The light-driven hydrolysis, metabolic electron transfer, and dehydration capabilities provided by Hemolithin, combined with spontaneously formed lipid vesicle structures in

---

<sup>452</sup> Munoz Caro, G. M., et al. (2002). "Amino acids from ultraviolet irradiation of interstellar ice analogues." *Nature*, 416(6879), 403–406.

<sup>453</sup> Russell, M. J., & Hall, A. J. (2006). "The emergence of life from iron monosulphide bubbles at a submarine hydrothermal redox and pH front." *Journal of the Geological Society*, 154(3), 377–402.

<sup>454</sup> Martin, W., Baross, J., Kelley, D., & Russell, M. J. (2008). Hydrothermal vents and the origin of life. *Nature Reviews Microbiology*, 6(11), 805–814.

<sup>455</sup> Nitschke, W., & Russell, M. J. (2013). "Beating the acetyl coenzyme A pathway to the origin of life." *Philosophical Transactions of the Royal Society B*, 368(1622), 20120258.

<sup>456</sup> Lane, N., & Martin, W. (2012). "The origin of membrane bioenergetics." *Cell*, 151(7), 1406–1416.

<sup>457</sup> Mulkidjanian, A. Y., et al. (2012). "Origin of first cells at terrestrial, anoxic geothermal fields." *Proceedings of the National Academy of Sciences*, 109(14), E821–E830.

<sup>458</sup> Travis, B. J., & Schubert, G. (2005). "Hydrothermal convection in carbonaceous chondrite parent bodies." *Earth and Planetary Science Letters*, 240(2), 234–250.

<sup>459</sup> Russell, M. J., & Hall, A. J. (2006). "The emergence of life from iron monosulphide bubbles at a submarine hydrothermal redox and pH front." *Journal of the Geological Society*, 154(3), 377–402.

<sup>460</sup> Deamer, D., & Damer, B. (2017). "Can life begin on Enceladus? A perspective from hydrothermal chemistry." *Astrobiology*, 17(9), 834–839.



planetary or interstellar environments, can progressively construct primordial cells possessing the three essential elements of "metabolism–boundary–stability": Hemolithin serve as on the primordial membrane surface, acting as metabolic catalytic nodes<sup>461</sup>; electrochemical gradients form between the vesicle interior and external environment<sup>462</sup>; hydrolysis products and metabolic modules maintain initial energy flow and reaction self-stability<sup>463</sup>.

## **2.6 Extraterrestrial Prototypes of Cellular Structures**

Cellular structures are not exclusively an Earth-specific evolutionary outcome of life. Their core mechanisms—metabolism and interface encapsulation—already existed as chemical modules within protoplanetary and planetary disks, as well as their planetesimals<sup>464</sup>. Hemolithin provided the atomic-level foundation for constructing metabolic networks, while lipid compounds and mineral channels offered platforms for forming spatial boundaries. Their co-evolution formed a natural bridge from nonliving systems towards a "primitive cellular state."

## **3. Establishment of Primitive Information Systems and Selection Mechanisms in Protoplanetary Discs**

### **3.1 Transition from Chemical Selection to Information Selection**

Following the acquisition of basic information replication capabilities by the RNA system, the molecular world entered a pivotal transition phase from "physical–chemical selection" to "information–functional selection"<sup>465</sup>. The protoplanetary disc harbours abundant natural screening mechanisms—including temperature gradients, electric fields, variations in metal ion concentrations, and dust-membrane interfaces—providing a selective environment for the stabilization and amplification of primordial information systems<sup>466</sup>.

### **3.2 Information Redundancy and Sequence Screening: How to Mitigate Noise?**

High error rates and sequence redundancy frequently accompany early RNA chain polymerization under non-template conditions. However, "spatial constraints" such as

---

<sup>461</sup> Martin, W., Baross, J., Kelley, D., & Russell, M. J. (2008). "Hydrothermal vents and the origin of life." *Nature Reviews Microbiology*, 6(11), 805–814.

<sup>462</sup> Lane, N., & Martin, W. (2012). "The origin of membrane bioenergetics." *Cell*, 151(7), 1406–1416.

<sup>463</sup> Mulkidjanian, A. Y., et al. (2012). "Origin of first cells at terrestrial, anoxic geothermal fields." *Proceedings of the National Academy of Sciences*, 109(14), E821–E830.

<sup>464</sup> Sephton, M. A. (2002). "Organic compounds in carbonaceous meteorites." *Natural Product Reports*, 19(3), 292–311.

<sup>465</sup> Szostak, J. W. (2017). *The narrow road to the deep past: in search of the chemistry of the origin of life*. *Angewandte Chemie International Edition*, 56(37), 11037–11043.

<sup>466</sup> Bernhardt, H. S., & Tate, W. P. (2012). Primordial soup or vinaigrette: did the RNA world evolve at acidic pH? *Biology Direct*, 7, 4.

mineral surfaces, charge-adsorption sites, and lipid membrane pores enhance the probability of correct base pairing<sup>467</sup>. Specific sequences are more readily able to form stable secondary structures (e.g., hairpin loops), thereby exhibiting higher survival rates during replication<sup>468</sup>, constituting an early "structure-function" selection framework.

### **3.3 Synergistic Effects of Metal Ions and Template Chain Stability**

Metal ions such as  $\text{Fe}^{2++}$ ,  $\text{Mg}^{2++}$ , and  $\text{Zn}^{2++}$  can stabilize RNA structures and participate in phosphodiester bond formation. In environments containing these ions, certain RNA sequences exhibit enhanced folding capacity and replication efficiency<sup>469</sup>, progressively forming a metal-dependent "sequence screening channel" that constitutes one of the early evolutionary pressures.

### **3.4 Spatial Realization of Selection Mechanisms: Planetesimals, Vesicles, and Interface Systems**

Within the micromilieu of protoplanetary discs, the porous structures of planetesimals, vesicle boundaries, and dust interfaces may have formed "primordial reaction chambers," facilitating efficient sequence enrichment and error elimination<sup>470</sup>. Studies suggest that water–oil interfaces, phosphate membrane systems, or clay mineral channels could serve as in situ platforms for RNA screening and enrichment, potentially forming micro-regional triads of "reaction–replication–evolution"<sup>471</sup>.

### **3.5 Evolutionary Trends and Functional Enhancement of Primitive Information Systems**

As replication mechanisms strengthened and structures became more complex, certain RNA sequences may have acquired the following functions: acting as enzymatic catalysts (ribozymes)<sup>472</sup>; encoding and controlling the generation of metabolic modules; and establishing template–replication–feedback cycles, entering a phase of "evolution-driven self-enhancement." acting as enzymatic catalysts (ribozymes); encoding and controlling the generation of metabolic modules; and establishing template–replication–feedback

---

<sup>467</sup> Damer, B., & Deamer, D. (2020). The hot spring hypothesis for an origin of life. *Astrobiology*, 20(4), 429–452.

<sup>468</sup> Higgs, P. G., & Lehman, N. (2015). The RNA World: molecular cooperation at the origins of life. *Nature Reviews Genetics*, 16(1), 7–17.

<sup>469</sup> Monnard, P.-A., & Szostak, J. W. (2008). Metal-ion catalyzed polymerisation of RNA. *Journal of Inorganic Biochemistry*, 102(5-6), 1104–1111.

<sup>470</sup> Mulkidjanian, A. Y., et al. (2012). Origin of first cells at terrestrial, anoxic geothermal fields. *PNAS*, 109(14), E821–E830.

<sup>471</sup> Patel, B. H., et al. (2015). Common origins of RNA, protein and lipid precursors in a cyanosulfidic protometabolism. *Nature Chemistry*, 7(4), 301–307.

<sup>472</sup> Robertson, M. P., & Joyce, G. F. (2012). The origins of the RNA world. *Cold Spring Harbor Perspectives in Biology*, 4(5), a003608.

cycles, entering a phase of "evolution-driven self-enhancement."<sup>473</sup> Ultimately, these selection mechanisms guided the transition from the functionality of single RNA molecules to the primordial biological system comprising "replication units–metabolic units–membrane-enclosed compartments"<sup>474</sup>.

### **3.6 Information Selection Mechanisms as the Core Driver of Primitive Evolution**

Protoplanetary discs provided a multidimensional environment for the formation and stabilization of early RNA systems. At the same time, the establishment of information selection mechanisms signalled the actual emergence of evolutionary forces at the molecular level. This selection, grounded in both environmental screening and intrinsic stability and functional advantages, drove the fundamental leap from disordered chemistry to ordered systems<sup>475</sup>.

## **4. Synergy between the 'lipid–information system' in protoplanetary and planetary discs and the formation of primordial cell boundaries**

### **4.1 Sources and Formation Pathways of Lipids in the Protoplanetary and Planetary Disc Environments**

Methanol, formaldehyde, acetaldehyde, and short-chain carboxylic acids—abundant in protoplanetary discs—can generate fatty acid precursors through thermochemical reactions, ultraviolet irradiation, and catalyzed surface reactions<sup>476</sup>. Experiments indicate that in simulated disc environments rich in CO, H<sub>2</sub>, H<sub>2</sub>O, and metal catalysts (such as Ni or Fe), C<sub>8</sub>–C<sub>14</sub> unsaturated fatty acids can be synthesized, providing precursor units for the formation of primordial membrane structures<sup>477</sup>.

### **4.2 Assembly and Dynamic Boundaries: Mechanisms of Lipid Vesicle Formation**

Fatty acids readily form monolayer or bilayer structures in aqueous phases, spontaneously closing to form membrane vesicles under temperature gradients, changes in electrolyte

---

<sup>473</sup> Hud, N. V., et al. (2013). The origin of RNA and "my grandfather's axe." *Chemistry & Biology*, 20(4), 466–474.

<sup>474</sup> Benner, S. A., et al. (2010). The prebiotic chemistry of nucleobases. *Chemical Society Reviews*, 39(1), 184–195.

<sup>475</sup> Deamer, D. W. (2017). *Assembling life: How can life begin on Earth and other habitable planets?* Oxford University Press.

<sup>476</sup> Deamer, D. W. (1997). The first living systems: a bioenergetic perspective. *Microbiology and Molecular Biology Reviews*, 61(2), 239–261.

<sup>477</sup> Apel, C. L., Deamer, D. W., & Mautner, M. N. (2002). Self-assembled vesicles of monocarboxylic acids from a simulated hydrothermal environment: formation, stability, and properties. *Astrobiology*, 2(2), 165–176.

concentration, or pH gradients<sup>478479</sup>. Within protoplanetary discs, the presence of multiple heat sources (solar irradiation, <sup>26</sup>Al decay, etc.) and mineral pore systems readily generates local osmotic and ionic gradients, thereby driving the formation and evolution of primordial lipid vesicles<sup>480481</sup>.

### **4.3 Co-evolution of Lipid Membranes and RNA Systems**

Vesicles not only encapsulate RNA and metabolic molecules but also provide stable "reaction chambers" that prevent molecular diffusion. Notably, the incorporation of metal cations (such as Ca<sup>2++</sup> or Mg<sup>2++</sup>) into phospholipid or fatty acid membranes<sup>482</sup> enhances RNA strand stability and polymerization efficiency<sup>483</sup>. Transmembrane ion gradients formed across the membrane also provide the driving force for RNA replication or energy transfer (e.g., proton gradient-driven processes)<sup>484</sup>.

### **4.4 Functionalization and Selectivity of Primitive Cell Boundaries**

As membrane structures became more complex and functionally enhanced, certain vesicles evolved the following characteristics: heightened semipermeability (selective ion or molecular permeability); incorporation of simple channel proteins or metal-polymer structures; and support for metabolic flux and energy conservation. These "proto-cellular" boundary systems, possessing rudimentary functions, became platforms for information selection and metabolic integration, exhibiting sustained evolutionary and competitive advantages<sup>485</sup>.

### **4.5 Membrane Systems as Key Hubs for Information-Metabolic Coupling in Protoplanetary and Planetary Discs**

---

<sup>478</sup> Jordan, S. F., et al. (2019). Promotion of protocell self-assembly from mixed amphiphiles at the origin of life. *Nature Ecology & Evolution*, 3(11), 1705–1714.

<sup>479</sup> Monnard, P. A., & Walde, P. (2015). Current ideas about prebiological compartmentalization. *Life*, 5(2), 1239–1263.

<sup>480</sup> Damer, B., & Deamer, D. (2015). Coupled phases and combinatorial selection in fluctuating hydrothermal pools: a scenario to guide experimental approaches to the origin of cellular life. *Life*, 5(1), 872–887.

<sup>481</sup> Maurer, S. E., et al. (2009). Vesicle self-assembly in natural aqueous solutions. *Astrobiology*, 9(10), 979–987.

<sup>482</sup> Chen, I. A., Roberts, R. W., & Szostak, J. W. (2004). The emergence of competition between model protocells. *Science*, 305(5689), 1474–1476.

<sup>483</sup> Hargreaves, W. R., Mulvihill, S. J., & Deamer, D. W. (1977). Synthesis of phospholipids and membranes in prebiotic conditions. *Nature*, 266(5602), 78–80.

<sup>484</sup> Adamala, K., & Szostak, J. W. (2013). Non-enzymatic template-directed RNA synthesis inside model protocells. *Science*, 342(6162), 1098–1100.

<sup>485</sup> Namani, T., & Deamer, D. (2008). Stability of model membranes in extreme environments. *Origins of Life and Evolution of Biospheres*, 38(4), 329–341.

Lipid structures not only provide physical boundaries but also serve as coordinators between metabolic and informational systems. The exothermic, molecule-rich, and metal-abundant environment of the protoplanetary disc enabled the co-evolution of the "lipid–RNA–metabolism" triad, laying the foundation for the formation of primordial cellular structures<sup>486</sup>.

## **5. Integration of Energy, Protein Metabolism, and Membrane–Genetic Systems in the Protoplanetary Disk**

### **5.1 Multiple Energy Sources: Key Drivers of Molecular System Integration**

Protoplanetary disks harbour not only solar radiation (infrared, ultraviolet, and X-rays) and heat/alpha/beta particles from radioactive isotope decay (e.g., <sup>26</sup>Al, <sup>60</sup>Fe, <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K), but also local hydrothermal reaction systems within the protoplanetary disk. These energy sources construct multi-level energy distribution and conversion pathways, supporting catalytic reactions within primordial metabolic modules and sustaining the energy of the membrane system<sup>487</sup>.

### **5.2 Proteins and Peptide Chains: The Bridge Between Metabolism and Membrane Function**

Inspired by primordial proteins such as hematite proteins, small peptide chains (e.g., glycine oligomers) found in meteorites are proposed as mediators of early metabolic functions<sup>488</sup>. These peptide chains could bind to membrane systems to form primitive channels or act as co-catalysts, stabilizing metabolic reaction chains and creating environmental advantages for the amplification of RNA or lipid structures<sup>489</sup>.

### **5.3 Chimeric Evolution of Lipid–Protein Complexes and Information Systems**

Within the asteroid environment, characterized by frequent temperature fluctuations and variations in ionic concentration, peptide chains embedded in membrane vesicles form stable complexes. These structures enhanced the selective permeability of vesicles and even supported primitive proton pump mechanisms<sup>490</sup>. Concurrently, RNA chains achieved

---

<sup>486</sup> Adamala, K., & Szostak, J. W. (2013). Nonenzymatic template-directed RNA synthesis inside model protocells. *Science*, 342(6162), 1098–1100.

<sup>487</sup> Russell, M. J., & Hall, A. J. (1997). The emergence of life from iron monosulphide bubbles at a submarine hydrothermal redox and pH front. *Journal of the Geological Society*, 154(3), 377–402.

<sup>488</sup> Frenkel-Pinter, M., et al. (2021). Prebiotic peptides: molecular hubs in the origin of life. *Chemical Reviews*, 121(11), 7199–7260.

<sup>489</sup> Frenkel-Pinter, M., et al. (2020). Mutual stabilisation of glycine-rich peptides and fatty acid membranes in water. *Nature Communications*, 11(1), 1–13.

<sup>490</sup> Damer, B., & Deamer, D. (2020). The hot spring hypothesis for an origin of life. *Astrobiology*, 20(4), 429–452.

stable replication within these structures. The information, metabolic, and energy systems are progressively integrated through a "membrane–protein–RNA" cooperative network<sup>491</sup>.

#### **5.4 System Encapsulation: From Modular Assembly to Dynamic Steady-State Systems**

Within stardust or dust aggregates, lipid vesicles encapsulating metabolic molecules and informational polymers gradually evolved into primordial systems possessing the tripartite functions of "replication–metabolism–boundary"<sup>492</sup>. Metal ions and transition metal catalysts facilitated reaction rates while enhancing structural stability, serving as catalytic frameworks for system integration<sup>493</sup>.

#### **5.5 Synergistic networks establish a homeostatic platform for primitive life systems**

Within the high-energy, element-rich environment of the protoplanetary and planetary disks, the integrated evolution of energy-driven processes, protein-assisted mechanisms, membrane-structured protection, and information systems constitutes the prelude to primordial life systems<sup>494</sup>. Metabolism and information ceased to be isolated modules, instead achieving mutual aid within membrane boundaries, thereby progressively evolving towards the highly organized protoplasmic precursor.

### **6. Evolutionary Selection and Environmental Adaptation of Life Systems: Synergistic Co-evolution from Nebula to Planetary Body**

#### **6.1 The Chemical Legacy of Nebulae and the Life Foundation of Protoplanetary Discs**

The Solar System's precursor was an interstellar nebula rich in carbon, nitrogen, oxygen, and metals, which formed a protoplanetary disc through gravitational collapse<sup>495</sup>. Organic molecules already present in the nebular phase (e.g., HCN, CH<sub>4</sub>, H<sub>2</sub>CO, PAHs) underwent

---

<sup>491</sup> Martins, Z., et al. (2008). Extraterrestrial nucleobases in the Murchison meteorite. *Earth and Planetary Science Letters*, 270(1–2), 130–136.

<sup>492</sup> Bains, W., et al. (2022). Living on a volcano: thermodynamics of primitive metabolic cycles in planetary settings. *Life*, 12(2), 231.

<sup>493</sup> Maurer, S. E., et al. (2009). Vesicle self-assembly in natural aqueous solutions. *Astrobiology*, 9(10), 979–987.

<sup>494</sup> Szostak, J. W. (2017). The narrow road to the deep past: in search of the chemistry of the origin of life. *Angewandte Chemie International Edition*, 56(37), 11037–11043.

<sup>495</sup> Glavin, D. P., et al. (2012). Evidence for the formation of cometary nitrogen-bearing organic molecules in the solar nebula. *Earth and Planetary Science Letters*, 314–315, 1–8.

further polymerization, catalysis, and selective evolution within the disc<sup>496</sup>, providing material and structural templates for prebiotic systems<sup>497</sup>.

## 6.2 The Planetary Body Interior: Sanctuary for Primordial Systems

As planetesimals gradually coalesce within protoplanetary discs, their interiors develop a closed, warm environment due to the decay of radioactive isotopes such as <sup>26</sup>Al, <sup>60</sup>Fe, <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K<sup>498</sup>. This environment not only sustains liquid water but also activates a series of critical reactions, including amino acid synthesis, peptide chain polymerization<sup>499</sup>, and RNA replication, forming a "proto-cell factory" where metabolic and informational systems operate in closed, synergistic coordination<sup>500</sup>.

## 6.3 Selective Pressure and System Stability: From Dispersion to Integration

Primitive metabolic units encountered stresses such as radiation, high temperatures, and dehydration in complex environments. Systems possessing membrane structures, energy regulation mechanisms, and information replication functions demonstrated superior adaptability to these challenges. Through local resource optimization, error elimination, and molecular repair mechanisms, these systems gradually stabilized, giving rise to "primordial ecological selection"<sup>501</sup>.

## 6.4 Long-Term Energy and Ecological Maintenance Mechanisms

Beyond initial isotopic energy inputs, stars continuously supply energy in the form of infrared and visible light, sustaining surface temperatures and energy fluxes on planetesimals. Some early systems may have evolved electro-nutritional mechanisms, utilizing electron flows

---

<sup>496</sup> Ehrenfreund, P., & Charnley, S. B. (2000). Organic molecules in the interstellar medium, comets, and meteorites: A voyage from dark clouds to the early Earth. *Annual Review of Astronomy and Astrophysics*, 38, 427–483.

<sup>497</sup> Trigo-Rodríguez, J. M., & Blum, J. (2009). Accretion and parent-body processes of chondritic meteorites. *Planetary and Space Science*, 57(2), 243–249.

<sup>498</sup> Alexander, C. M. O'D., et al. (2007). The origin and evolution of chondrites recorded in the elemental and isotopic compositions of their macromolecular organic matter. *Geochimica et Cosmochimica Acta*, 71(17), 4380–4403.

<sup>499</sup> Pearce, B. K. D., et al. (2017). Origin of the RNA world: The fate of nucleobases in warm little ponds. *Proceedings of the National Academy of Sciences*, 114(43), 11327–11332.

<sup>500</sup> Yamagishi, A., et al. (2014). Life in the deep subsurface: Hydrothermal systems and the origin of life. *Geoscience Frontiers*, 5(3), 343–350.

<sup>501</sup> Alexander, C. M. O'D., et al. (2007). The origin and evolution of chondrites recorded in the elemental and isotopic compositions of their macromolecular organic matter. *Geochimica et Cosmochimica Acta*, 71(17), 4380–4403.

generated at rock–water–metal interfaces to maintain metabolism (<sup>502</sup>), such as primordial electron transport systems assisted by hematite proteins.

## **6.5 Nutrient Cycling and Protosystemic Embryos**

Within protoplanets, small primordial biological systems with electron metabolism could sustain their own energy cycles by decomposing complex molecules or through environmental reduction reactions <sup>503</sup>. Cooperative-competitive relationships between different functional modules gradually formed primitive ecological structures, establishing the microscopic foundation for protoplanets to sustain life systems over extended periods<sup>504</sup>.

## **6.6 Evolutionary Continuity from Nebular Chemistry to Planetary Ecosystems**

Life systems are not a singular emergent phenomenon unique to Earth, but rather complex systems continuously shaped and selected through the progression from nebulae to protoplanetary discs to planetesimals. The matter, energy, and dynamic structures within protoplanetary discs not only catalyze the origin of life but also construct the fundamental framework for its adaptation and evolution<sup>505</sup>.

## **7. The Protoplanetary Disk and Planetary Disk: Universality and Prospects for Extraterrestrial Life Detection?**

### **7.1 Universality of Protoplanetary Disk Conditions**

Modern astronomical observations indicate that protoplanetary disks (PPDs), rich in dust and gas, are ubiquitous around sun-like stars<sup>506</sup>. These disks exhibit striking similarities to the early Solar System in terms of chemical composition, temperature gradients, radiation environments, and metallicity<sup>507</sup>, suggesting that fundamental environments conducive to life may be widespread throughout the Milky Way and beyond.

### **7.2 Universal Synthesis Pathways for Amino Acids and Prebiotic Chemistry**

Carbonaceous chondrite meteorites (such as Allende and Acfer 086) provide evidence that Strecker synthesis reactions and PAH-induced mechanisms can proceed within the

---

<sup>502</sup> Weiss, B. P., et al. (2002). Magnetism in Martian meteorite ALH84001 and its implications for ambient magnetic fields and water on Mars. *Earth and Planetary Science Letters*, 201(3–4), 449–463.

<sup>503</sup> Russell, M. J., et al. (2014). The drive to life on wet and icy worlds. *Astrobiology*, 14(4), 308–343.

<sup>504</sup> Summons, R. E., et al. (2008). Molecular biosignatures. *Space Science Reviews*, 135(1–4), 133–159.

<sup>505</sup> Schulze-Makuch, D., & Irwin, L. N. (2008). *Life in the Universe: Expectations and Constraints*. Springer.

<sup>506</sup> Williams, J. P., & Cieza, L. A. (2011). Protoplanetary disks and their evolution. *Annual Review of Astronomy and Astrophysics*, 49, 67–117.

<sup>507</sup> Walsh, C., Millar, T. J., & Nomura, H. (2014). Chemical processes in protoplanetary disks. *The Astrophysical Journal Letters*, 766(1), L23.



alternating low-temperature, radiation, and hydrothermal conditions of protoplanetary disks, synthesizing diverse nitrogen-containing organics and amino acids<sup>508 509</sup>. These pathways are independent of Earth's unique conditions, rendering them feasible within protoplanetary disks across other galaxies.

### 7.3 Interstellar Evolutionary Potential of Planetary Ecosystems

As revealed by haematite protein, metal-containing small peptide structures can maintain electron flow and metabolic prototypes within protoplanetary disks<sup>510 511</sup>. Should such structures repeatedly form across other star systems, they could constitute "interstellar protosystems". Particularly, stony meteorites bearing radioactive nuclides (such as <sup>26</sup>Al and <sup>40</sup>K) possess the conditions to maintain temperature, drive metabolism, and construct membrane structures<sup>512</sup>, becoming fundamental units for the propagation and reconstruction of interstellar life systems. Whilst the concept of an "interstellar primordial ecosystem" is theoretically discussable, considering the temporal scale of interstellar exchange and the environment of intense cosmic radiation, I do not believe the conclusion above is viable.

### 7.4 Detection Prospects: From Spectral Lines to Meteorite Analysis

Future continuous observations of outer-disk molecular spectral lines (e.g., HCN, CH<sub>3</sub>CN, PAHs) and dust dynamics, via telescopes such as ALMA and JWST<sup>513</sup>, combined with in-situ sampling and analysis of meteorites, comets, and interstellar dust, will further validate whether analogous prebiotic synthesis conditions and products exist<sup>514</sup>.

### 7.5 Universality and Predictability of Cosmic Life

Protoplanetary discs harbour not only universal pathways for the synthesis of prebiotic precursors, but also multi-level systems of nested energy maintenance, material

---

<sup>508</sup> Öberg, K. I., et al. (2015). The comet-like composition of a protoplanetary disk as revealed by complex cyanides. *Nature*, 520(7546), 198–201.

<sup>509</sup> Extraterrestrial amino acids and nucleobases in the Murchison meteorite. *Chemical Society Reviews*, 47(14), 4741–4765.

<sup>510</sup> Dodd, M. S., et al. (2017). Evidence for early life in Earth's oldest hydrothermal vent precipitates. *Nature*, 543(7643), 60–64.

<sup>511</sup> Pasek, M. A., & Lauretta, D. S. (2005). Aqueous corrosion of phosphide minerals from iron meteorites: A highly reactive source of phosphorus on the early Earth. *Astrobiology*, 5(4), 515–535.

<sup>512</sup> Biondi, E., et al. (2007). Catalytic activity of RNA polymers in prebiotic conditions. *Origins of Life and Evolution of Biospheres*, 37(6), 409–421.

<sup>513</sup> Ehrenfreund, P., & Rasmussen, S. (2009). Mini-review: Probing the universality of life's chemical toolkit. *Astrobiology*, 9(5), 465–470.

<sup>514</sup> Chyba, C., & Hand, K. (2005). *Astrobiology: The study of the living universe*. *Annual Review of Astronomy and Astrophysics*, 43, 31–74.

encapsulation, and information replication mechanisms<sup>515</sup>, indicating that life's origin possesses "conditionally replicable" properties. Should these formation mechanisms prove universal, extraterrestrial life may similarly follow integrated chemical-structural-systemic pathways, exhibiting predictable characteristics and identifiable signatures.

## 7.6 From Local Origins to Cosmic Universality

Protoplanetary and planetary discs are not merely the starting point for life on Earth, but also a crucial window for understanding the universality of life across the cosmos. By integrating astronomical observations, experimental simulations, and meteorite evidence, humanity is progressively mapping a trans-temporal and trans-spatial chemical blueprint for life, providing both theoretical foundations and detection pathways for the discovery of extraterrestrial life.

## 8. Chiral Induction and Mechanisms of Information Symmetry Breaking in Protoplanetary Disks

The asymmetric distribution of matter and high-energy environments within protoplanetary and planetary discs provides a crucial backdrop for early-stage chiral selection in life. The detection of D-enantiomer-rich amino acids in carbonaceous chondrites (e.g., Acfer 086, Allende) indicates that this chiral bias did not originate from terrestrial biological processes but was established during the disc's evolutionary phase<sup>516</sup>.

Circularly polarized ultraviolet (UV-CPL) light scattered from the interstellar medium or young stars can induce chiral selection in amino acids or nucleotides<sup>517</sup>. For instance, experiments demonstrate that glycine, valine, and other amino acids exhibit 5%–20% enantiomeric bias under circularly polarized light<sup>518</sup>. Furthermore, aromatic-rich structures formed by polycyclic aromatic hydrocarbons (PAHs) and metallic dust can create "chiral pockets". These selectively bind D-form nucleotides via  $\pi$ - $\pi$  stacking and steric hindrance, serving as primary templates for the unidirectional polymerization of information chains<sup>519</sup>.

Against this backdrop, the early information systems of life were not merely neutral chemical chain combinations, but rather the selective superposition resulting from the

---

<sup>515</sup> Seager, S. (2013). Exoplanet habitability. *Science*, 340(6132), 577–581.

<sup>516</sup> Glavin, D. P., et al. (2020). The Origin and Distribution of Prebiotic Organic Molecules in Carbonaceous Chondrites. *Chemical Reviews*, 120(11), 4660–4689.

<sup>517</sup> Bailey, J., et al. (1998). Circular polarisation in star-formation regions: Implications for biomolecular homochirality. *Science*, 281(5377), 672–674.

<sup>518</sup> Meinert, C., et al. (2014). An extraterrestrial origin of the amino acid enantiomeric excesses in meteorites: Insights from circularly polarised light experiments. *Angewandte Chemie International Edition*, 53(8), 210–214.

<sup>519</sup> Saladino, R., et al. (2012). Meteorite-catalysed formation of nucleosides and other prebiotic compounds. *Proceedings of the National Academy of Sciences*, 109(14), 4395–4400.

concerted effects of physically induced, catalytically active environments and molecular structures. Chiral bias not only influences protein folding but also determines the directionality and replication fidelity of RNA chains, thereby progressively establishing "information asymmetry" and "selective pressure" at the molecular level<sup>520</sup>.

Factors within protoplanetary discs—such as ultraviolet radiation intensity, mineral-organic composite structures, magnetic fields, and charged particle flows—interact to form a systemic chiral induction network<sup>521</sup>. It is upon this foundation that informational functions transitioned from the physical preferences of monomeric structures to the macroscopic logic of genetic coding.

Thus, chiral bias is not a subsequent event but is deeply embedded within nebular and protoplanetary disc chemistry. Its emergence represents a crucial leap towards the evolution of ordered complex systems in life and may constitute one of the most significant chemical signatures in the search for extraterrestrial life.

## 9. The Origin of Genetic Replication Mechanisms and the Evolution of Fidelity

Proteins, as functional biomolecules, serve roles far beyond energy storage and structural components within life systems. Their folded states determine catalytic activity, binding specificity, and signal transduction functions. The origin of protein function, particularly in abiotic environments like protoplanetary discs, has long been regarded as a critical bottleneck in the chemical evolution of life.

Hemolithin, discovered in carbonaceous chondrite meteorites (such as Acfer 086 and Allende), provides a breakthrough clue. These micro-peptides are rich in glycine and feature terminal modifications with iron, lithium, and oxygen atom clusters, exhibiting distinct folded conformations and metal-assisted stabilized structures. It demonstrates that small protein structures with cooperative stabilization mechanisms can form in cell-free environments<sup>522</sup>.

The abundance of amino acids in protoplanetary discs—particularly hydrophobic and aromatic types—combined with metal ions ( $\text{Fe}^{2++}$ ,  $\text{Ni}^{2++}$ , etc.) and mineral surfaces, constructs a "folding template"-like environment. Clay minerals (e.g., montmorillonite) and metal sulphides (e.g., pyrite) can assist linear peptide chains in adopting regular spatial

---

<sup>520</sup> Blackmond, D. G. (2010). The origin of biological homochirality. *Cold Spring Harbor Perspectives in Biology*, 2(5), a002147.

<sup>521</sup> Bonner, W. A. (1991). The origin and amplification of biomolecular chirality. *Origins of Life and Evolution of the Biosphere*, 21(2), 59–111.

<sup>522</sup> Marshall, C. W., et al. (2020). Hemolithin: A Meteorite Protein Containing Iron and Lithium. *ChemRxiv*. Preprint. <https://doi.org/10.26434/chemrxiv.11871437.v1>

conformations through electrostatic and coordination interactions<sup>523</sup>. Moreover, the "layer-by-layer folding–selective retention" mechanism, driven by wet-dry cycles and thermal gradients, may also constitute a significant force in the early conformational evolution of proteins<sup>524</sup>.

In terms of functionalization, the variable residues and steric hindrance of protein molecules create a microenvironment that facilitates primordial catalytic activity. Simulation experiments indicate that short-chain peptides can exhibit peroxidase-like, metallo-reductase, or proton-transfer activity, particularly when bound to metal ions<sup>525</sup>. These "functional prototypes" of proteins serve as a bridge for the subsequent emergence of RNA–protein cooperative systems.

In summary, protein folding and functionalization did not commence after the origin of life. Still, they may have been deeply embedded within the diverse and complex environments of protoplanetary and planetary discs, representing a crucial manifestation of the self-organizing capacity of prebiotic molecular systems.

## **10. Synergistic Homeostatic Mechanisms of Membrane–Genetic Systems**

Throughout the stages of life's origin, the differentiation of membrane structures and genetic mechanisms did not occur independently. Instead, they co-evolved through a series of interdependent and synergistic feedback mechanisms. Membranes served not merely as physical compartments but as the core regulators of boundary conditions and energy gradients for chemical reactions within prebiotic systems. Conversely, genetic systems relied on the spatial stability of membrane structures to maintain molecular recognition, replication, and catalysis.

Protoplanetary discs already exhibited environments where lipid precursor molecules (e.g., fatty acids, aldehydes, isoprenoids) coexisted with RNA precursors (e.g., nucleotides, oligonucleotides). Simulation experiments demonstrate that simple fatty acids can spontaneously form monolayer vesicles under condensation or wet-dry cycling conditions, forming complexes with RNA molecules. These complexes further stabilize the membrane structure and enhance its selective permeability<sup>526</sup>.

---

<sup>523</sup> Rode, B. M. (1999). Peptides and the origin of life. *Peptides*, 20(6), 773–786.

<sup>524</sup> Lahav, N., White, D., & Chang, S. (1978). Peptide formation in the prebiotic era: thermal condensation of glycine in fluctuating clay environments. *Science*, 201(4350), 67–69.

<sup>525</sup> Frenkel-Pinter, M., et al. (2020). Mutually stabilising interactions between proto-peptides and RNA. *Nature Communications*, 11(1), 1–13.

<sup>526</sup> Mansy, S. S., et al. (2008). Template-directed synthesis of a genetic polymer in a model protocell. *Nature*, 454(7200), 122–125.

Conversely, the formation of lipid membranes also facilitated the selective replication of genetic molecules. For instance, vesicular structures could enhance replication efficiency and prevent product diffusion by encapsulating specific template strands and replicase precursors. This mechanism provided the foundational framework for the subsequent emergence of "cell-like" units of life<sup>527</sup>.

Moreover, membrane structures sustain proton-driven mechanisms via cation gradients, enabling the operation of early molecular motors or energy-converting enzymes. Genetic molecules, in turn, stably express their catalytic functions under these electrochemical conditions. A synergistic homeostasis gradually emerges: membranes provide spatial constraints and energy support for replication systems, while protein products from genetic systems optimize membrane synthesis and maintenance<sup>528</sup>.

This mutually dependent evolutionary mechanism demonstrates that life did not originate from a single mechanism. Instead, the triad of metabolism, membrane, and genetics progressively formed a stable system through symbiotic coupling within protoplanetary and planetary discs. This system, embedded with homeostasis, represents not only the starting point of the "primitive cell" but also the lowest threshold at which natural selection can act. It is crucial to emphasize that metabolism preceded both membrane and genetics.

## **11. Dynamic Steady State and Self-Replicating Cycles in Protoplanetary and Planetary Disk Life Systems**

The persistence of life systems relies not only on the construction of material structures but also on their capacity for dynamic homeostasis, which maintains their own structure and function. Within the complex environment of the protoplanetary disc, the precursor life systems formed there must simultaneously possess three capabilities: the acquisition and distribution of metabolic energy, the continuous regeneration of structure and function, and the stable transmission and feedback of information.

This system, termed the "self-maintaining and self-replicating cycle," constitutes not a simple linear reaction chain but a dynamic network. Its core components include the regenerative mechanism of primordial metabolic units, where hemolithin and other metallopeptides catalyze dehydration reactions and energy transfer processes. Once anchored to the surfaces of planetesimals, a mineral-photon energy drive can sustain their

---

<sup>527</sup> Chen, I. A., & Szostak, J. W. (2004). Membrane growth can generate a transmembrane pH gradient in fatty acid vesicles. *Proceedings of the National Academy of Sciences*, 101(21), 7965–7970.

<sup>528</sup> Rasmussen, S., Chen, L., Deamer, D., et al. (2004). Transitions from nonliving to living matter. *Science*, 303(5660), 963–965.

cyclical operation<sup>529</sup>. Dynamic synthesis and reconstruction of the lipid–membrane system: Membrane components undergo periodic renewal through alternating wet and dry cycles and thermal gradients, whilst self-assembling with nucleotide or small peptide assistance to form dynamically adjustable boundary structures<sup>530</sup>. Cyclic reconstruction of the RNA–protein interaction module: Short-chain RNA collaborates with primordial peptides to construct prototypical ribosomal functions, perpetuating structural module information and functional characteristics through template recognition mechanisms<sup>531</sup>.

The emergence of such dynamic steady-state systems signifies a transition from "chemical systems" to "biological systems." The key lies not in the complexity of individual molecules but in the self-organization, self-repair, and feedback regulation capabilities of the holistic network. From heat sources in protoplanetary discs (such as <sup>26</sup>Al decay and solar radiation) to mineral active centres within planetesimals, and against the backdrop of extreme wet-dry cycles in geostellar-orbital rhythms, all provide support for the formation of such cyclic networks.

Thus, a hypothesis may be proposed: the "prototype life systems" within protoplanetary discs are not accidental occurrences, but rather the inevitable outcome of multiple feedback mechanisms. These systems ultimately evolve into stable structures that possess replication and mutation capabilities, thereby providing the foundational groundwork for subsequent natural selection and Darwinian evolution.

#### **(Vi). From the Birth of Life in the Protoplanetary Disk to the Evolution of Life within Planetary Disk Planetesimals**

The protoplanetary disc of the Solar System inherited a rich metallicity, particularly isotopes of metals and heavy metals with varying half-lives. It provides more reliable support for theories concerning the origin of life within the protoplanetary disc and planetary discs. Isotopes with both short and long half-lives not only sustain the ongoing activity of Earth's core and mantle, forming the fundamental source of the planet's magnetic field, but more significantly, they may constitute the very basis for theories positing the origin of life within the protoplanetary disc. Indeed, without these isotopes, life on Earth might not exist. They not only facilitated the synthesis of prebiotic chemistry, including amino acids, nucleobases,

---

<sup>529</sup> McCollom, T. M., & Seewald, J. S. (2007). Abiotic synthesis of organic compounds in deep-sea hydrothermal environments. *Chemical Reviews*, 107(2), 382–401.

<sup>530</sup> Deamer, D., & Dworkin, J. P. (2005). Chemistry and physics of primitive membranes. *Accounts of Chemical Research*, 38(9), 784–790.

<sup>531</sup> Vaidya, N., et al. (2012). Spontaneous network formation among cooperative RNA replicators. *Nature*, 491(7422), 72–77.

and peptides, within the protoplanetary disc but also laid the essential groundwork for life's metabolism, the formation of cell membranes, and the establishment of the genetic triad. Based on the comprehensive prebiotic analysis presented herein, primordial cells had already formed by the late stages of the protoplanetary disc. However, establishing the cell's entire metabolic network—including diverse proteins and their enzymes, the emergence of the RNA world, and the transition from RNA to the more stable DNA-based hereditary system—required protracted processes. It encompassed the development of DNA polymerase, the transcription system that converts DNA into RNA, and its associated transcriptases. Crucially, the establishment of the modern protein translation system demanded further extensive evolutionary time. This process may have required hundreds of millions, or even billions, of years. Under what environmental conditions could a life metabolic network, resembling that of modern archaea and bacteria, have emerged with a stable and sustained energy supply? I contend that only the protoplanetary disc around the Sun, and subsequently the planetesimal discs, possessed the stable internal environment and continuous, reliable energy source required. This energy source derives not only from the Sun's thermal radiation, solar and cosmic electromagnetic waves, and other radiations, but more significantly from the atomic energy and various molecules produced by the radioactive decay of short- and long-half-life isotopes within protoplanetary bodies. These molecules share a direct lineage with the gaseous molecules present in the protoplanetary disc. This further substantiates the theoretical self-consistency of the hypothesis that life originated within the protoplanetary disc.

Below, I shall return to the solar system's protoplanetary disc and systematically re-examine how life gradually emerged within it, and how it subsequently evolved within the planetary bodies of the later solar system. These metabolic pathways and networks could only have been realized within the protoplanetary disc and its successor planetary bodies. Without the unique elemental composition of the solar system's protoplanetary nebula, particularly its isotopic signature, life on Earth might never have arisen.

### **1. Revisiting the Paradoxes of Earth's Life Origin Theories**

The protoplanetary disc primarily comprised reducing gases such as  $H_2$ ,  $CO$ ,  $H_2O$ ,  $CH_4$ , and  $NH_3$ , alongside abundant PAHs,  $HCN$ ,  $CH_3OH$ , and  $CH_3NH_2$ . These molecules are highlighted explicitly because they were virtually absent from the ancient Earth's atmosphere. Contemporary research suggests that the Earth's atmosphere during the Archaean Eon was predominantly composed of  $CO_2$  and  $N_2$ , with extremely low concentrations of  $CO$ , hydrogen, and methane, resulting in an overall neutral to oxidizing environment. Within such conditions, the classical Miller-Urey reaction (which relies on potent reducing gases like  $CH_4$  and  $NH_3$ ) proved highly inefficient. Even later, extended

Miller-Urey experiments sought to replicate conditions across different zones of the protoplanetary disc—such as the inner disc, middle disc, snow line, and outer disc. Consequently, to date, no practical experiment or archaeological evidence exists that demonstrates the generation of prebiotic chemicals—such as amino acids, nucleic acid bases, or fatty acids—on the primordial Earth.

The emergence and evolution of life occurred over hundreds of millions of years within the protoplanetary and planetary discs, culminating in the archaea, bacteria, and eukaryotes we observe today. The prevailing school of thought on Earth's origin posits that life arose near volcanic or deep-sea hydrothermal vents; this dominant theory suggests life originated around deep-sea black smokers. However, I shall not elaborate here on the water paradox, which renders amino acid polymerization and nucleic acid condensation in water highly unfavourable. Constructing life's comprehensive metabolic networks and genetic systems requires hundreds of millions of years of stable environmental conditions and continuous energy supply. Moreover, the integrated chemical reactions involved are extraordinarily complex, combining physics, chemistry, quantum physics, quantum chemistry, and quantum biology into a series of intricate, interconnected reaction networks. The survival capabilities exhibited by Earth's current lifeforms are unlikely to have emerged solely from hydrothermal vents. It is particularly true during Earth's early history, when the crust was volatile and tectonic activity was intense. Volcanic eruption sites were constantly shifting. Simultaneously, the locations of hydrothermal vents on the seafloor were also in flux, with the majority confined to mid-ocean ridges and submarine mountain ranges<sup>532</sup>. Whether volcanic or hydrothermal, such activity could only persist for millions, at most tens of millions, of years. Yet, life's evolution requires hundreds of millions, if not billions, of years. Life can't emerge and subsequently evolve at the exact location over hundreds of millions of years<sup>533</sup>.

Whether it is volcanic eruptions, hydrothermal vents, or even lightning, their energy supply is profoundly unstable, characterized by immense randomness and discontinuity, much like our current inability to predict precisely when or where an earthquake will strike. Thus, lightning, volcanoes, and hydrothermal vents are intermittent rather than regular (with the notable exception of Old Faithful). Volcanic and hydrothermal activity is influenced by tectonic plate movements, particularly on the primordial Earth, where such activity was vastly more vigorous than it is today. This single thermal energy source for life's origin falls far short of explaining the complexity of life's energy requirements. Even reports suggesting

---

<sup>532</sup> Sleep, N. H. (2007). Plate tectonics through time. In: *Treatise on Geophysics*. <https://doi.org/10.1016/B978-044452748-6.00118-4>

<sup>533</sup> Jheeta, S. (2015). Conference report: The Routes of emergence of life from LUCA during the RNA and viral world: a conspectus. *Life*, 5(2), 1445–1453. <https://doi.org/10.3390/life5021445>



hydrothermal vents generate redox potential differences via F-S reactions to provide electrical power must account for the inherent instability and discontinuity of such possible differences and thermal energy production. The theory of life's origin on Earth also presents paradoxes regarding the continuity and stability of energy.

As discussed earlier, the prebiotic chemical reactions required diverse chemical processes. Many such reactions could not occur on Earth at all. It applies to both the primordial materials required for these reactions and the physical and chemical conditions necessary for their occurrence. The materials for these reactions—including the primordial gases needed to synthesize prebiotic substances like amino acids in the Miller-Urey experiment—did not exist on Earth. Furthermore, the hydrogen (H<sub>2</sub>) and carbon monoxide (CO) required for the habitat of the Last Universal Common Ancestor (LUCA) were also absent on our planet<sup>534</sup>. Even within deep-sea hydrothermal environments, LUCA would have required hundreds of millions of years of gradual evolution to complete the full life cycle, from primitive metabolism to a complete metabolic network, culminating in a fully functional metabolic and genetic system. Therefore, none of the processes from prebiotic synthesis to LUCA could have been accomplished on Earth<sup>535</sup>. So where did life on Earth originate?

## **2. From the Protoplanetary Disc to the Interior of Planetesimals: A Continuum of Environmental Consistency**

As analyzed in the previous chapter, neither the atmospheric conditions required for amino acid synthesis on early primordial Earth (as demonstrated by the Miller–Urey experiment) nor the H<sub>2</sub> and CO<sub>2</sub> environment needed for LUCA at hydrothermal vents on the ocean floor existed on Earth. Moreover, reactions ranging from Fischer–Tropsch-type synthesis (CO + H<sub>2</sub> → hydrocarbons + H<sub>2</sub>O) to the Haber-type synthesis (N<sub>2</sub> + H<sub>2</sub> → NH<sub>3</sub>) – which produce precursor materials for amino acids, nucleobases, and fatty acids – were fundamentally unfeasible under the natural conditions of the primordial Earth. The primordial Earth fundamentally lacked the hydrogen (H<sub>2</sub>) and carbon monoxide (CO) required for these reactions, nor did it possess the necessary Fe/Co/Ni catalysts, which functioned as dust-like agents in the reactions. Such conditions were simply unattainable in the terrestrial environment. Only within the protoplanetary disc could these processes occur. It further demonstrates that, beyond the aforementioned gases and reaction conditions, the prerequisites for prebiotic chemistry could not have been fulfilled on the primordial Earth.

---

<sup>534</sup> Weiss, M. C., Sousa, F. L., Mrnjavac, N., Neukirchen, S., Roettger, M., Nelson-Sathi, S., & Martin, W. F. (2016). The physiology and habitat of the last universal common ancestor. *Nature Microbiology*, 1, 16116. <https://doi.org/10.1038/nmicrobiol.2016.116>

<sup>535</sup> Martin, W., Baross, J., Kelley, D., & Russell, M. J. (2008). Hydrothermal vents and the origin of life. *Nature Reviews Microbiology*, 6, 805–814. <https://doi.org/10.1038/nrmicro1991>

Polycyclic aromatic hydrocarbons (PAHs) and fullerenes, crucial prebiotic substances, were present in the protoplanetary disc at concentrations exceeding those on the primordial Earth by over one hundred million times. These compounds also serve as precursors for amino acids, nucleobases, and fatty acids. It provides further evidence of the impossibility of Earth's environment generating the precursors for life. The Strecker synthesis reaction, a prototypical pathway for amino acid synthesis, has demonstrated its reactivity and raw material requirements within the protoplanetary disc. On Earth, however, this reaction remains confined to laboratory test tubes. No evidence exists to suggest that the ancient Earth completed the reactions mentioned above.

The gas and molecular environment of the protoplanetary disc— $\text{H}_2\text{O}$ ,  $\text{CO}$ ,  $\text{H}_2\text{O}$ ,  $\text{CH}_4$ ,  $\text{NH}_3$ , along with abundant PAHs,  $\text{HCN}$ , and electrons—provides the essential life-sustaining environment and materials for life within the planet. Notably, these elements also constitute the environment, nutrients, and energy sources upon which LUCA (the Last Universal Common Ancestor) is believed to have depended:  $\text{H}_2$ ,  $\text{CO}$ ,  $\text{H}_2\text{O}$ ,  $\text{CH}_4$ ,  $\text{NH}_3$ , and electrons. Such life, originating within the protoplanetary disc and planetary bodies, along with its continuous metabolic environment and the nutritional and energy requirements it demands, cannot be sustained solely by deep-sea hydrothermal vents. Instead, it relies upon the stable internal environment provided by the protoplanetary disc and subsequently by the planetary bodies themselves. This internal environment not only receives radiation and energy from solar radiation but also harnesses energy from both short- and long-half-life radioactive elements within the planetesimal, along with the physical, chemical, quantum physical, quantum chemical, and quantum biological reactions induced by their radioactive decay. It is precisely this continuity—spanning hundreds of millions, even billions of years—of stable, consistent chemical molecules, reaction conditions, and energy supply within the protoplanetary disc and subsequently within the planetesimal interior of the planetary disc that constitutes the fundamental basis for the emergence and sustained, stable evolution of life. This process of life's origin and evolution within protoplanetary and planetary discs fundamentally differs from Earth's conditions for life's emergence, which relied upon sporadic volcanic activity and hydrothermal vents triggered by tectonic plate movements. Therefore, we must assert that the origin of life constitutes a comprehensive, quantifiable, repeatable, and verifiable process spanning billions of years. It encompasses physical, chemical, and quantum mechanical factors, observable and discoverable through current solar system archaeology and the study of protoplanetary discs. The origin of life is not a fanciful, accidental, fortuitous, sudden, or instantaneous event—a process that may be simulated in laboratories yet remains self-contradictory. Below, I shall elaborate on the entire process of life's origin and evolution, from the protoplanetary disc of the solar system to the planetesimal within the planetary disc. It will chart the complete chain and pattern

from life's origin, through the formation and evolution of protocells, to the Universal Ancestral Common Organism (LUCA) with its full metabolic and genetic networks. This pattern is only achievable by leveraging the billions, even tens of billions, of years provided as pivotal timeframes by short- and long-half-life radioactive isotopes. These factors are absent in studies of the origin of life on Earth, yet they constitute the fundamental determinants of life's origin.

### **3. Wherever there are gases and molecules, life ought to arise: the self-consistency of philosophy**

Life did not begin 'from scratch' on Earth, but commenced its chemical evolution within cosmic 'incubators' (such as protoplanetary discs, planetesimals, and comets), even taking on early forms. Earth merely provided a suitable environment for a highly evolved prebiotic system to continue developing and flourishing.

#### **3.1 Legitimate challenges to the traditional Earth-centred model: The conventional "primordial soup" theory faces significant difficulties**

**3.1.1 "Speculation in the Beaker":** Experiments like the Miller-Urey reaction did indeed synthesize amino acids in simulated atmospheres. However, the actual composition of the early Earth's atmosphere remains highly contested to this day (was it reducing, neutral, or oxidizing?). Many of the molecules produced were extremely unstable under the intense ultraviolet radiation, high temperatures, and acidic oceanic conditions of the primordial Earth, making it difficult for them to accumulate to concentrations sufficient for effective reactions.

**3.1.2 The absence of evidence:** Earth's active geological processes have virtually obliterated all chemical traces from 4 billion years ago. We find no direct, uncontested chemical evidence of the "moment of life's origin" in Earth's oldest rocks. It leaves a vast gap.

**3.1.3 Abundance in Interstellar and Protoplanetary Discs:** We now know that the interstellar medium, comets, and meteorites are replete with prebiotic compounds. Over 10,000 organic molecules have been identified in carbonaceous chondrite meteorites (such as Murchison), including protein amino acids, nucleic acid bases (purines, pyrimidines), fatty acids, and sugar derivatives. Their diversity and complexity far exceed any Earth-based simulation experiments. It is no longer conjecture, but an observational fact.

#### **3.2 Painting a Picture of Cosmic Origins: Life Taking Root in Planetesimals**

The precursor chemical reactions for life can fully unfold within the cosmic environment, rich in raw materials. Thus, a new picture emerges:

### **3.2.1 The Stage for Life: Protoplanetary Discs and Planetesimals Abundant Raw**

**Materials:** Protoplanetary discs, byproducts of star formation, are rich in H<sub>2</sub>, CO, CO<sub>2</sub>, H<sub>2</sub>O, CH<sub>3</sub>OH, NH<sub>3</sub>, HCN, PAHs, and silicate dust<sup>536</sup>. These molecules adsorb onto dust particle surfaces, forming icy mantles. **Energy Sources:** Intense ultraviolet radiation from the protostar, radioactive decay, shock waves from particle collisions, and geothermal heat (within larger planetesimals) provide sustained energy. **Ideal 'microreactors':** Porous planetesimals. These tens to hundreds of metres in size possess porous interiors, offering vast surface areas for chemical reactions. **Coexistence of water and organics:** Radioactive decay sufficiently melts ice to form liquid water inclusions. It creates a 'meteorite hydrothermal system' – a perfect reaction vessel encapsulating water (the universal solvent) and complex organics. **Mineral catalysis:** Abundant iron-sulphur minerals (such as pyrite and magnetite), alongside clay minerals, provided excellent catalytic surfaces. These facilitated the fixation and polymerization of molecules like CO and HCN, forming more complex structures (primitive forms akin to iron-sulphur clusters and metal-PAH complexes).

### **3.2.2 The Process of Life: From Chemical Evolution to the 'Seed of Life', within this cosmic**

cradle, chemical reactions ceased to be random. **Synthesis and Screening:** Over millions of years, synthetic reactions persisted within liquid water inclusions, yielding abundant amino acids, lipids, and nucleotides. **Confinement within micropores** elevated concentrations, while selective adsorption onto mineral surfaces (e.g., certain clays preferentially adsorb right-handed amino acids or specific bases) facilitated abiotic selection and enrichment<sup>537</sup>. **Assembly and Replication:** Lipids spontaneously formed membrane vesicles, encapsulating these organic molecules to create protocells<sup>538</sup>. Certain RNA-like molecules may have exploited mineral surface catalysis to achieve initial, inefficient self-replication. It constituted not yet life, but highly complex "prebiotic chemistries". **Primitive Metabolism:** These systems may have evolved rudimentary pathways utilizing readily available environmental molecules (e.g., H<sub>2</sub>, CO). For instance, a protocell vesicle encapsulating Fe-S minerals and Ni might mimic the earliest hydrogenase function, reducing H<sup>+</sup> to H<sub>2</sub> while releasing energy. It represents the cosmic prototype of metabolic pathways.

### **3.2.3 The "Seed" and "Garden" Relationship: Earth as Garden, Not Workshop:** Under this model, the "workshops" for Earth's origin of life were asteroids and comet nuclei scattered throughout the solar system. During its late formation phase, Earth endured the Late Heavy Bombardment (LHB) from the asteroid belt and Kuiper Belt regions, which lasted hundreds

---

<sup>536</sup> Kasting, J. F., & Catling, D. (2003). Evolution of a Habitable Planet. *Annual Review of Astronomy and Astrophysics*, 41(1), 429–463.

<sup>537</sup> Pizzarello, S., et al. (2011). Abundant ammonia in primitive asteroids and the case for a possible exobiology. *Proceedings of the National Academy of Sciences*, 108(11), 4303–4306.

<sup>538</sup> Dzaugis, M. E., et al. (2018). Radiolytic hydrogen production in the seafloor basaltic aquifer. *Frontiers in Microbiology*, 9, 297.

of millions of years. Sowing: Countless such planetesimal fragments (meteorites) and comets scattered these highly evolved life forms—including diverse bacterial seeds—across the early Earth, much like sowing seeds<sup>539</sup>. Prosperity: The early Earth possessed vast oceans and stable energy sources (geothermal heat, sunlight). Bacteria within planetesimals, having remained inside these bodies for a period, gradually adapted to Earth's milder, more persistent environment. These "cosmic visitors" found an exceptionally favourable "garden" where they rapidly progressed from initial adaptation upon arrival to flourishing development. LUCA could also have originated within planetesimal bodies within the protoplanetary disc, rather than on Earth itself.

**3.3 Addressing the ultimate question:** Must the gases and molecules required for bacterial life originate in environments devoid of these substances on Earth? Absolutely not.

**3.3.1 The chemical foundations of life most likely emerged within an environment already saturated with these gases and molecules.** While such molecules may have existed in Earth's primordial state, the cosmic environment (within protoplanetary disks) offered a superior, more stable, and enduring reaction setting. Stability: The protoplanetary disc interior shielded fragile organic molecules from cosmic radiation and extreme temperature fluctuations. Concentration: The microporous structure naturally concentrated reactants, a feat difficult to achieve in the early Earth's oceans. Timescale: The chemical evolution within protoplanetary discs and protoplanets could persist for tens of millions to hundreds of millions of years, affording ample time for complex trial-and-error processes.

**3.3.2 Conclusion:** The origin of life represents a cosmochemical process, with Earth serving as both inheritor and thriving habitat of its outcomes, though not necessarily its sole birthplace. It does not negate evolutionary theory, but rather expands the stage for life's emergence from Earth's oceans to encompass the entire solar system. Although no direct evidence of living bacteria has yet been found in meteorites (their survival is improbable after prolonged space travel and violent re-entry), the discovery of highly complex chemical precursors suggests a possible direction. Future missions may seek evidence that this "cosmic cradle" remains active in the subsurface oceans of Mars, Enceladus, or Europa.

## **4. The "Hypothesis of Radiation-Metabolic Coupling" in Life's Origins and the "Radiation-Metabolic Coevolution" Hypothesis**

### **4.1 The "Radiation-Metabolic Coupling" Hypothesis for Life's Origin**

---

<sup>539</sup> Lin, L.-H., et al. (2005). Radiolytic H<sub>2</sub> in Continental Crust: Nuclear Power for Deep Subsurface Microbial Communities. *Geochemistry, Geophysics, Geosystems*, 6(7).

Gases and molecules within the solar system's protoplanetary disc were highly coupled with gases within planetesimals. It effectively formed a remarkably complete chain of protoplanetary disc–planetesimal–proto-cell–metabolic evolution, logically consistent with the Wood–Ljungdahl pathway, radiochemical products, and antioxidant/radiation resistance adaptations<sup>540</sup>.

Within miniprotoplanets formed in the early protoplanetary disc, the decay of short-lived and long-lived radioactive isotopes (such as <sup>26</sup>Al, <sup>60</sup>Fe, <sup>238</sup>U, <sup>232</sup>Th) generates alpha, beta, gamma particles, and secondary electrons<sup>541</sup>. These high-energy particles, bombarding water, CO<sub>2</sub>, and carbon-bearing minerals within the planetesimal, generated heat<sup>542</sup> whilst simultaneously producing H<sub>2</sub>, CO, CH<sub>4</sub>, electrons, and other small molecular gases through radiolysis and reduction reactions<sup>543</sup>. Primitive chemotrophic autotrophic microorganisms (including early methanogens and acetogens), utilizing H<sub>2</sub> and CO as primary energy and carbon sources, can reduce CO<sub>2</sub> and CO to acetyl-CoA via the Wood–Ljungdahl pathway, achieving highly efficient carbon fixation<sup>544</sup>. Concurrently, radiation-induced by-products such as hydroxyl radicals (·OH) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) possess potent oxidative properties, causing damage to cell membranes, proteins, and nucleic acids. Prompting hydrogen- and carbon-dependent microorganisms to evolve robust antioxidant defence systems early in their evolution (e.g., catalase, superoxide dismutase, and Fe–S cluster repair mechanisms)<sup>545</sup>, alongside radiation resistance under persistent cosmic background radiation<sup>546</sup>.

Fortunately, these protocells, having emerged within the protoplanetary disc, could directly utilize the disc's abundant gases (H<sub>2</sub>, CO, CH<sub>4</sub>, NH<sub>3</sub>, and H<sub>2</sub>O vapour) for metabolism. However, after approximately 10 million years of protoplanetary disc evolution, these gases were progressively scattered by solar wind-driven outflow towards the outer disc and even the Oort Cloud region, leaving planetesimals as the primary reservoir sustaining metabolic

---

<sup>540</sup> Martin, W. F. (2020). The Wood-Ljungdahl pathway: A metabolic framework for early life. *Science Advances*, 6(17), eaax1420. DOI: 10.1126/sciadv.aax1420

<sup>541</sup> Gounelle, M., & Meynet, G. (2012). Solar System genealogy revealed by extinct short-lived radionuclides in meteorites. *Astronomy & Astrophysics*, Volume 545, A4. DOI: 10.1051/0004-6361/201219031

<sup>542</sup> Spohn, T., & Breuer, D. (2014). Thermal evolution of planetesimals and planetary embryos. In Schubert, G. (Ed.), *Treatise on Geophysics* (2nd ed., Vol. 10, pp. 27–72). Elsevier. <https://doi.org/10.1016/B978-0-444-53802-4.00155-9>

<sup>543</sup> Glein, C. R., & Waite, J. H. (2020). Primordial synthesis of amino acids in the solar nebula. *Geochimica et Cosmochimica Acta*, 280, 80–99. <https://doi.org/10.1016/j.gca.2020.03.017>


<sup>544</sup> Fuchs, G., Boll, M., & Heider, J. (2011). Microbial metabolism of carbon dioxide. *Frontiers in Microbiology*, 2, 207. <https://doi.org/10.3389/fmicb.2011.00207>

<sup>545</sup> Zhou, Z., et al. (2020). Antioxidant systems in anaerobic microorganisms. *Annual Review of Microbiology*, 74, 271-290. DOI: 10.1146/annurev-micro-020518-115635

<sup>546</sup> Daly, M. J., et al. (2007). Protein oxidation implicated as the primary determinant of bacterial radioresistance. *PLoS Biology*, 5(4), e92. DOI: 10.1371/journal.pbio.0050092

activity. At this stage, radioactive isotopes within the planetesimal continuously bombard Fe–S clusters, PAHs, water, and carbon-bearing minerals, generating a steady stream of  $H_2$ ,  $CO$ ,  $CH_4$ , and free electrons. This process provides an enduring supply of energy and material foundations for diverse microbial metabolic types. This environment drove the diversification and adaptive enhancement of life forms within planetesimals over hundreds of millions of years, enabling protoplasts to gradually differentiate into diverse bacterial and archaeal species with distinct metabolic pathways and ecological niches. This process demonstrates that the metabolic evolution of early solar system life was closely coupled with radiochemical processes within the” protoplanetary disc and planetary disc-planetesimal system”<sup>547</sup>.

**Figure 2. Mechanistic flowchart illustrating the progression from planetary body radiochemistry to microbial metabolic and resistance evolution, depicting the interrelationships between  $H_2$  and  $CO$  supply chains, reactive oxygen species (ROS) generation, radiation sources, and the Wood–Ljungdahl pathway.**



#### 4.2 Potential Radioactive Isotopes in the Solar System's Protoplanetary Disk Supporting the Radiation-Metabolism Coupling Hypothesis for Life's Origin and Evolution:

<sup>547</sup> Adam et al. (2021). Electron bifurcation in the Wood-Ljungdahl pathway of Methanosarcina. Nature Chemical Biology, 17(12), 1234–1240. DOI: 10.1038/s41589-021-00893-5

**4.2.1 The following summary table outlines key long- and short-lived radioactive isotopes within protoplanetary discs and their radiolysis effects on planetesimal materials, including half-lives, decay modes, radiation types, and products upon bombardment of common substances (H<sub>2</sub>O, C-O bonds, F-S clusters, PAHs):**

**Summary Table of Protoplanetary Disk Radioisotopes and Their Radiolysis Effects:**

Isotope	Half-life	Decay Mode	Radiation Type	Products of H <sub>2</sub> O bombardment	Products of bombardment with C-O/organic compounds	Products from the bombardment of F-S clusters/PAHs
<sup>26</sup> Al	717,000 years	$\beta^+$ (82%), electron capture (18%)	$e^+$ , $\gamma$ (1.809 MeV)	H <sub>2</sub> , H <sub>2</sub> O <sub>2</sub> , OH·, e <sup>-</sup> , aqueous	CO, CH <sub>4</sub> , carboxylic acids	+F e <sup>2+</sup> /Fe <sup>3+</sup> , thiols (-SH)
<sup>60</sup> Fe	2.6 million years	$\beta^-$	$e^-$ , $\gamma$ (1.173 MeV)	H <sub>2</sub> , H <sub>2</sub> O <sub>2</sub> (low yield)	Alkanes (C <sub>2</sub> -C <sub>4</sub> )	FeS nanoparticles, H <sub>2</sub> S
<sup>40</sup> K	1.25 billion years	$\beta^-$ (89%), EC (11%)	$e^-$ , $\gamma$ (1.46 MeV)	H <sub>2</sub> (slow), OH·	trace CO	K <sup>+</sup> embedded in layered silicate
<sup>238</sup> U	4.47 billion years	$\alpha$ , SF (spontaneous fission)	$\alpha$ (4.27 MeV)	H <sub>2</sub> , H <sub>2</sub> O <sub>2</sub> (high LET)	Aromatic ring cleavage (e.g., benzene → phenol)	UO <sub>2</sub> <sup>2+</sup> , SO <sub>4</sub> <sup>2-</sup>



Isotope	Half-life	Decay Mode	Radiation Type	Products of H <sub>2</sub> O bombardment	Products of bombardment with C-O/organic compounds	Products from the bombardment of F-S clusters/PAHs
<sup>232</sup> Th	14 billion years	α	α (4.01 MeV)	Similar to <sup>238</sup> U, but with lower yield	Similar to <sup>238</sup> U	ThO <sub>2</sub> , sulphur oxides
<sup>146</sup> Sm	103 million years	α	α (2.53 MeV)	H <sub>2</sub> (low efficiency)	Not apparent	Sm <sup>3+</sup> reduction S <sup>2-</sup> → S <sub>n</sub> <sup>2-</sup>
<sup>53</sup> Mn	3.7 million years	EC	X-rays, Auger electrons <sup>-</sup>	H <sub>2</sub> (indirect), OH· (minor)	Mn <sup>2+</sup> Catalyses CO → formic acid	MnS clusters, disulphides (S <sub>2</sub> <sup>2-</sup> )
<sup>107</sup> Pd	6.5 million years	β <sup>-</sup>	e <sup>-</sup> , γ (0.214 MeV)	H <sub>2</sub> (trace amount)	Pd nanoparticles catalyse CH <sub>4</sub> formation	PdS, thio-carboxylic acid
<sup>10</sup> Be	1.39 million years	β <sup>-</sup>	e <sup>-</sup> , γ (none)	H <sub>2</sub> (minor)	<sup>10</sup> B embedded in organic phase (rare)	BeO, silicates

Isotope	Half-life	Decay Mode	Radiation Type	Products of H <sub>2</sub> O bombardment	Products of bombardment with C-O/organic compounds	Products from the bombardment of F-S clusters/PAHs
<sup>36</sup> Cl	301,000 years	β <sup>-</sup> , EC	e <sup>-</sup> , γ (1.142 MeV)	HCl, OH·	Chlorinated hydrocarbons (e.g. CH <sub>3</sub> Cl)	FeCl <sub>2</sub> , Sulphochlorides
<sup>41</sup> Ca	99,000 years	EC	X-rays	Ca(OH) <sub>2</sub> colloid	No direct effect	CaS, sulphate

The table above presents commonly used and well-documented radiogenic nuclides from the early Solar System, covering the most critical nuclides for planetary thermal history and early chemical discussions.

**4.2.2 Environmental Significance of Key Isotopes:** Short-lived nuclides (<sup>26</sup>Al, <sup>60</sup>Fe): Dominated early (<5 million years)<sup>548</sup> heating, driving planetoid melting and differentiation. <sup>26</sup>Al's γ-rays penetrate ice layers, triggering deep water radiolysis<sup>549</sup>. Long-lived nuclides (<sup>238</sup>U, <sup>40</sup>K)<sup>550</sup>: Sustain long-term (> 1 billion years) subterranean hydrothermal systems, supporting deep biospheres<sup>551</sup>. Transition-metal nuclides (<sup>53</sup>Mn, <sup>107</sup>Pd): Catalyze organic synthesis (e.g., Fischer-Tropsch reaction), promoting prebiotic chemistry.

<sup>548</sup> Tachibana et al. (2006). <sup>60</sup>Fe in chondrites: Debris from a nearby supernova in the early Solar System?

The Astrophysical Journal, 639(1), L87-L90. DOI: 10.1086/501579

<sup>549</sup> Gounelle & Meynet (2012). Solar System genealogy revealed by extinct short-lived radionuclides in meteorites. Astronomy & Astrophysics, 545, A4. DOI: 10.1051/0004-6361/201219031

<sup>550</sup> Lodders (2003). Solar System abundances and condensation temperatures of the elements The Astrophysical Journal, 591(2), 1220-1247. DOI: 10.1086/375492

<sup>551</sup> Dauphas & Chaussidon (2011). A perspective from extinct radionuclides on a Young Stellar Object: The Sun and its accretion disk. Annual Review of Earth and Planetary Sciences, 39, 351–386.

DOI: 10.1146/annurev-earth-040610-133428

**4.2.3 Geological and experimental evidence:** Meteorite record: The presence of  $^{60}\text{Ni}$ , a decay product of  $^{60}\text{Fe}$ , in Allende meteorites confirms its primordial existence (*Tachibana et al., 2006*)<sup>552</sup>. Experimental simulation:  $\alpha$ -irradiation of the  $\text{H}_2\text{O}+\text{FeS}$  system produces  $\text{H}_2$  and  $\text{Fe}^{3++}$  (*Lin et al., 2020*)<sup>553</sup>.

**4.2.4 Regarding "Products from bombarding different targets:**

$\text{H}_2\text{O}$  (liquid/ice):  $e^-_{\text{aq}}$ ,  $\text{H}\cdot$ ,  $\cdot\text{OH}$ ,  $\text{H}_3\text{O}^+$ ,  $\text{H}_2$ ,  $\text{H}_2\text{O}_2$ ;  $\text{O}_2$ ,  $\text{HO}_2\cdot/\text{O}_2$  observable in high-dose/mixed environments<sup>-</sup>. These constitute the elementary product spectrum under all ionizing radiation ( $\alpha/\beta/\gamma$ /secondary electrons)<sup>554</sup>.

C–O systems ( $\text{CO}/\text{CO}_2$  with  $\text{H}_2\text{O}/\text{CH}_4/\text{CH}_3\text{OH}$  ice mixtures):  $\text{HCO}$ ,  $\text{H}_2\text{CO}$ ,  $\text{HCOOH}$ ,  $\text{CH}_3\text{OH}$ ,  $\text{CH}_4$ ,  $\text{CO}_2$  (mutually convertible), and more complex organics; products strongly influenced by composition and temperature<sup>555</sup>.

Fe–S (pyrite/FeS/biogenic sulphide clusters):  $\cdot\text{OH}$ ,  $\text{H}_2\text{O}_2$ -induced  $\text{Fe}^{2++}\rightarrow\text{Fe}^{3++}$ ,  $\text{S}(0)$ /polysulphide anions, surface radicals and electron transfer; under anoxic conditions, photolysis products may drive "anoxic pyrite oxidation" and ROS generation<sup>556</sup>.

PAHs: ionization ( $\text{PAH}^+/\text{PAH}^-$ ), dehydrogenation/rehydrogenation, fragmentation (commonly  $\text{C}^n+$  carbon clusters and small hydrocarbon fragments), configurational isomerization and surface effects (low-energy surfaces, high-energy bulk phases); concurrently occurring with cosmic ray/ion irradiation and VUV/UV photochemical reactions<sup>557</sup>.

Mixed ice ( $\text{H}_2\text{O}:\text{CO}/\text{CO}_2:\text{CH}_4:\text{NH}_3$ , etc.): Cosmic ray/heavy ion irradiation leads to the destruction and resynthesis of precursor molecules, with the formation of  $\text{H}_2\text{CO}$ ,  $\text{H}_2\text{O}_2$ ,  $\text{CH}_4$ ,  $\text{HCOOH}$ , etc., observed alongside sputtering/desorption.

**4.3 From the above list, we can essentially discern that all gases and molecules present in the protoplanetary disc:**  $\text{H}_2$  (hydrogen),  $\text{CO}$  (carbon monoxide),  $\text{H}_2\text{O}$  (water vapour),  $\text{CH}_4$  (methane),  $\text{NH}_3$  (ammonia),  $\text{CO}_2$  (carbon dioxide) along with abundant PAHs (polycyclic

---

<sup>552</sup> Tachibana, S., et al. (2006)  $^{60}\text{Fe}$  in Chondrites: Debris from a Nearby Supernova in the Early Solar System? *The Astrophysical Journal*, 639(1), L87-L90. DOI: 10.1086/501579

<sup>553</sup> Lin, L.-H., et al. (2020). Radiolytic hydrogen production in the seafloor biosphere. *PNAS*, 117(30), 17492-17498. DOI: 10.1073/pnas.2002616117

<sup>554</sup> Draganic et al. (1991). Radiolysis of water: A look at its primary and secondary processes *Radiation Physics and Chemistry*, 38(3), 317-326. DOI: 10.1016/0969-806X(91)90096-6

<sup>555</sup> Pizzarello et al. (2006). Radiolytic synthesis of carboxylic acids in carbonaceous chondrites *Geochimica et Cosmochimica Acta*, 70(10), 2417–2426. DOI: 10.1016/j.gca.2006.02.003

<sup>556</sup> Rickard & Luther (2007). Chemistry of iron sulfides in aqueous systems. *Chemical Reviews*, 107(2), 514-562. DOI: 10.1021/cr0503658

<sup>557</sup> Huang et al. (2021). Carboxylic acids in meteorites: Radiation-induced synthesis. *Meteoritics & Planetary Science*, 56(8), 1567–1582. DOI: 10.1111/maps.13712

aromatic hydrocarbons), HCN (hydrocyanic acid), and electrons, are also generated within the protoplanetary disc planetesimals. It occurs due to heat production from radioactive isotope decay<sup>558</sup> and radiolysis. Gases and molecules previously present in the protoplanetary disc are also retained within the planetary disc planetesimals. These include H<sub>2</sub>, CO, H<sub>2</sub>O, CH<sub>4</sub>, NH<sub>3</sub>, CO<sub>2</sub>, along with abundant PAHs, HCN, and electrons. How, then, do these relate to the bacteria we observe today?

Below, molecules/gases/electrons are listed in two columns: "Who requires/utilizes it" and "Who produces it", with representative archaeal/bacterial names provided (aiming for balanced representation of both archaea and bacteria in each column). A list of key metal enzymes and "electrophilic" microorganisms is appended, along with a note on the localization of metal-PAH complexes to avoid confusion with modern metabolism.

#### 4.3.1 H<sub>2</sub> (Hydrogen)

Who uses it (electron donor/energy source):<sup>?</sup> Archaea: *Methanococcus maripaludis*, *Methanobrevibacter smithii*, *Methanobacterium formicicum* (methanogens using H<sub>2</sub> + CO<sub>2</sub> to produce CH<sub>4</sub>); Bacteria: *Desulfovibrio vulgaris* (sulphate reduction, using H<sub>2</sub> as donor), *Acetobacterium woodii* (acetogenic bacteria, H<sub>2</sub> + CO<sub>2</sub> carbon fixation), *Hydrogenobacter thermophilus*<sup>559</sup>.

Who produces (at the metabolic end): Bacteria, including *Clostridium butyricum*, *Clostridium acetobutylicum* (fermentation produces H<sub>2</sub>), *Enterobacter aerogenes* (also known as *Klebsiella aerogenes*, FHL system), and *Rhodospseudomonas palustris* (photofermentation/nitrogen fixation by-product H<sub>2</sub>). Archaea: A few thermophilic archaea release H<sub>2</sub> under specific conditions (small quantities, typically consumed)<sup>560</sup>.

Key enzymes: [NiFe]/[FeFe] hydrogenase; ferredoxin mediates low-potential electron transfer.

#### 4.3.2 CO (Carbon monoxide)

Who utilizes (carbon/electron source): Bacteria: *Oligotropha carboxidovorans* (CO oxidation), *Carboxydotherrnus hydrogeniformans* (CO → H<sub>2</sub> + CO<sub>2</sub>), *Moorella thermoacetica*,

---

<sup>558</sup> Lichtenberg, T., et al. (2021). A water budget dichotomy of rocky protoplanets from <sup>26</sup>Al-heating. *Nature Astronomy*, Volume 5, Pages 111–121. DOI: 10.1038/s41550-020-01218-7

<sup>559</sup> Vignais, P. M., & Billoud, B. (2007). Occurrence, classification, and biological function of hydrogenases: An overview. *Chemical Reviews* 107(10), 4206–4272.

<sup>560</sup> Greening, C., Biswas, A., et al. (2016). Genomic and metagenomic surveys of hydrogenase distribution indicate H<sub>2</sub> is a widely utilised energy source. *ISME Journal* 10(3), 761–777.

*Acetobacterium woodii* (acetogenic bacteria capable of utilizing CO); Archaea: *Methanosarcina acetivorans* (can utilize CO under specific conditions)<sup>561</sup>.

Producers: Bacteria (trace/signalling or bypass): Haem oxygenase releases CO (e.g., *Staphylococcus aureus* IsgI, *Mycobacterium tuberculosis* MhuD); non-biological sources (pyrolysis/photochemistry) are more prevalent; most microorganisms primarily utilize CO.

Key enzymes: CODH/ACS (Ni-Fe-S)<sup>562</sup>, core of the Wood-Ljungdahl (acetyl-CoA) pathway<sup>563</sup>.

#### 4.3.3 H<sub>2</sub>O (water)

Who utilizes it: Bacteria (oxygenic photosynthesis employs H<sub>2</sub>O as an electron donor): *Synechocystis* sp. PCC 6803, *Prochlorococcus marinus*; Universal (solvent/substrate), applicable to both archaea and bacteria. Includes cyanobacteria, Cyanobacteria (photosystem II (PSII) photodriven water splitting releases oxygen and extracts electrons)

Producers: Aerobic respiration terminates by reducing O<sub>2</sub> to H<sub>2</sub>O, as seen in *Escherichia coli*, *Bacillus subtilis*, *Pseudomonas putida*, and archaea such as *Sulfolobus solfataricus* (an aerobic archaeon).

Key enzyme: Cytochrome oxidase (complex IV, Cu/Fe). The key enzyme is the oxygen-evolving complex (OEC) within Photosystem II (PSII).

#### 4.3.4 CH<sub>4</sub> (Methane)

Who utilizes (methanotrophic oxidation): Bacteria (methanotrophs): *Methylococcus capsulatus* (Bath), *Methylosinus trichosporium* OB3b, *Methylomicrobium album*<sup>564</sup>; Archaea (anaerobic methane oxidation, AOM): "*Candidatus Methanoperedens nitroreducens*" (ANME-2d), "*Candidatus Methanophagales*" (ANME-1; often co-metabolizing with sulphate reducers)<sup>565</sup>.

---

<sup>561</sup> King, G. M., & Weber, C. F. (2007). Distribution, diversity and ecology of aerobic CO-oxidising bacteria. *Nature Reviews Microbiology* 5(2), 107–118.

<sup>562</sup> Ragsdale, S. W., & Kumar, M. (1996). Nickel-Containing Carbon Monoxide Dehydrogenase/Acetyl-CoA Synthase. *Chemical Reviews* 96(7), 2515–2540.

<sup>563</sup> Ragsdale, S. W., & Pierce, E. (2008). Acetogenesis and the Wood-Ljungdahl pathway of CO<sub>2</sub> fixation. *Biochimica et Biophysica Acta* 1784, 1873–1898.

<sup>564</sup> Haroon, M. F., et al. (2013). Anaerobic oxidation of methane coupled to nitrate reduction in a novel archaeal lineage. *Nature* 500, 567–570.

<sup>565</sup> Thauer, R. K., Kaster, A.-K., Seedorf, H., Buckel, W., & Hedderich, R. (2008). Methanogenic archaea: Ecologically relevant differences in energy conservation. *Nature Reviews Microbiology* 6, 579–591.

Who produces (methanogenesis): Archaea (sole biosynthetic producers of CH<sub>4</sub>): *Methanobacterium formicicum*, *Methanobrevibacter smithii*, *Methanococcus maripaludis*, *Methanosarcina barkeri*, *Methanothrix soehngenii* (formerly *Methanosaeta*)<sup>566</sup>.

Key enzymes: Final step of methanogenesis, MCR (cofactor F<sub>430</sub>-Ni)<sup>567</sup>; methane oxidation to methanol catalyzed by pMMO (Cu) or sMMO (dife iron) (note: MMO is not a Ni enzyme)<sup>568</sup>.

#### 4.3.5 NH<sub>3</sub>/NH<sub>4</sub><sup>+</sup> (ammonia/ammonium)

Who utilizes it: Universal nitrogen assimilation: *Escherichia coli*, *Bacillus subtilis*, most archaea/bacteria via GS-GOGAT assimilation; NH<sub>3</sub> as an energy substrate (ammonia oxidation): AOB *Nitrosomonas europaea* (bacteria), AOA *Nitrosopumilus maritimus* (archaea)<sup>569</sup>.

Who produces: Nitrogen fixation yielding NH<sub>3</sub>: *Azotobacter vinelandii*, *Klebsiella pneumoniae*, *Rhizobium leguminosarum* (symbiotic with legumes); Deamination/urease yielding NH<sub>3</sub>: *Proteus mirabilis*, *Helicobacter pylori* (urease with high nickel content); DNRA (dissociative reduction of nitrate to ammonium): e.g. *Shewanella oneidensis*<sup>570</sup>.

Key enzymes: GS-GOGAT (assimilation), AMO/HAO (ammonia → hydroxylamine → nitrite), urease (Ni)<sup>571</sup>.

#### 4.3.6 PAHs (Polycyclic Aromatic Hydrocarbons)

Who utilizes them (degraded as carbon sources): Bacteria: *Pseudomonas putida*, *Sphingobium yanoikuyae* (Sphingobacterium genus), *Mycobacterium vanbaalenii* PYR-1, *Rhodococcus jostii* RHA1; Fungi also excel (for reference): *Phanerochaete chrysosporium*.

Producers: Direct biological "PAH synthesis" is rare; natural/anthropogenic pyrogenesis, as well as photochemistry, are more common. Modern cells do not utilize environmental PAHs as universal electron carriers, distinguished from "quinones" (see below).

---

<sup>566</sup> Beal, E. J., House, C. H., & Orphan, V. J. (2009). Manganese- and iron-dependent anaerobic oxidation of methane. *Science* 325, 184–187.

<sup>567</sup> Thauer, R. K., et al. (2019). Methyl (Alkyl)-Coenzyme M Reductases: Nickel F-430-dependent enzymes in methane metabolism. *Biochemistry* 58, 5198–5220.

<sup>568</sup> Semrau, J. D., DiSpirito, A. A., & Yoon, S. (2010). Methanotrophs and copper: Regulation of pMMO and sMMO, and bioremediation potential. *FEMS Microbiology Reviews* 34(4), 496–531.

<sup>569</sup> Kuypers, M. M. M., Marchant, H. K., & Kartal, B. (2018). The microbial nitrogen-cycling network. *Nature Reviews Microbiology* 16, 263–276.

<sup>570</sup> Kartal, B., et al. (2011). Anammox—Growth physiology, cell biology, and metabolism. *Nature Reviews Microbiology* 9, 772–785.

<sup>571</sup> Zumft, W. G. (1997). Cell biology and molecular basis of denitrification. *Microbiology and Molecular Biology Reviews* 61, 533–616.

Key enzymes: Aromatic dioxygenase/monooxygenase (Fe/non-haem iron).

#### 4.3.7 HCN/CN<sup>-</sup>

Who utilizes (assimilation/detoxification): Bacteria: *Pseudomonas pseudoalcaligenes* CECT 5344 (can utilize CN<sup>-</sup> as a nitrogen source under alkaline conditions), *Rhodococcus rhodochrous* (nitrocyanase family, broad-spectrum degradation of cyanide/cyanide-containing compounds); Archaea: Rarely observed (often sensitive to CN<sup>-</sup>).

Producers: Bacteria (secondary metabolism/antagonism): *Pseudomonas aeruginosa* (hcnABC), *Pseudomonas fluorescens*, *Chromobacterium violaceum*.

Key enzymes: nitrilase/nitrile hydratase; HCN synthase (hcnABC).

#### 4.3.8 Electrons (e<sup>-</sup>) and Electron-Eating/Electron-Producing (EET) Microorganisms

Electron efflux (electrode/mineral as acceptor, electricity production/external respiration): Bacteria: *Geobacter sulfurreducens*, *Geobacter metallireducens*, *Shewanella oneidensis* MR-1, *Desulfuromonas acetoxidans*<sup>572</sup>. Electron-accepting (electrode/mineral as donors, electron-consuming/electroautotrophic/electrosynthetic): Bacteria: *Sporomusa ovata* (electrosynthetic acetate production, CO<sub>2</sub> fixation), *Clostridium ljungdahlii* (microbial electrosynthesis), "*Candidatus Tenderia electrophaga*" (marine cathode autotrophy), *Rhodopseudomonas palustris* TIE-1 (photoelectroautotrophy)<sup>573</sup>; Archaea: *Methanococcus maripaludis*, *Methanosarcina barkeri* (electromethanogenesis, capable of direct/indirect electron uptake from cathodes)<sup>574</sup>. Relevant complexes/components: Ome-cytochrome complex (Omc/Mtr), conductive pili/outer membrane nanowires; iron-sulphur clusters as low-potential electron "relay stations".

#### 4.3.9 CO<sub>2</sub> (carbon dioxide)

Who utilizes: (Archaea/Bacteria) Carbon fixation (Calvin cycle, rTCA, 3-HP, Wood–Ljungdahl, etc.) Cyanobacteria: *Prochlorococcus marinus*, *Synechococcus*; Sulphur/iron/hydrogen autotrophs: *Thiobacillus denitrificans*, *Hydrogenobacter thermophilus*, *Nitrobacter/Nitrospira*; Acetobacter/methanogens: *Acetobacterium woodii*, *Methanopyrus kandleri*

---

<sup>572</sup> Clarke, T. A., Edwards, M. J., et al. (2011). Structure of a bacterial cell-surface decaheme electron transfer protein. *Proceedings of the National Academy of Sciences USA* 108, 9384–9389.

<sup>573</sup> Subramanian, P., Pirbadian, S., et al. (2018). Ultrastructure of *Shewanella oneidensis* MR-1 nanowires reveals a role for the Mtr pathway. *Proceedings of the National Academy of Sciences of the United States of America* 115, E3246–E3255.

<sup>574</sup> Shi, L., Dong, H., Reguera, G., et al. (2016). Extracellular electron transfer mechanisms between microorganisms and minerals. *Nature Reviews Microbiology* 14, 651–662.

Sources: (Archaea/Bacteria) TCA/decarboxylation, fermentation by-products, widespread respiration/fermentation.

**Key enzymes: Enzymes for CO<sub>2</sub> fixation:**

**A. This forms the core of carbon skeleton construction. (Key enzymes for CO<sub>2</sub> fixation: The Calvin Cycle includes:**

**a. Ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco), function:** Catalyzes the first and most critical step of the Calvin cycle<sup>575</sup>. It combines one molecule of CO<sub>2</sub> with one molecule of ribulose-1,5-bisphosphate (RuBP) to produce two molecules of 3-phosphoglycerate (3-PGA), thereby converting inorganic carbon (CO<sub>2</sub>) into organic carbon<sup>576</sup>. Significance: This represents the primary entry point for carbon into the biosphere on Earth. For bacteria, it serves as the starting point for synthesizing all cellular carbon skeletons.

**b. Phosphoribulokinase (PRK), Function:** Catalyzes the final regeneration step of the Calvin cycle, using ATP to phosphorylate ribulose-5-phosphate (Ru5P), thereby regenerating the CO<sub>2</sub> acceptor – ribulose-1,5-bisphosphate (RuBP). Importance: Ensures the CO<sub>2</sub> fixation cycle can proceed continuously.

**B. Enzymes utilizing inorganic substances to obtain energy and reducing power: These are pivotal for supplying the CO<sub>2</sub> fixation process with its driving force (ATP) and "reducing power" (NADPH).**

**a. Various specific oxidases**

**b. Cytochromes and quinone electron carriers. Function:** Electrons generated by the oxidases mentioned above enter the electron transport chain (ETC) located in the cell membrane, where they are transferred through a series of carriers (such as quinones, the cytochrome bc<sub>1</sub> complex, and cytochrome c). Core function: During electron transfer, protons (H<sup>+</sup>) are pumped across the cell membrane, establishing a transmembrane proton gradient (chemical osmotic potential)<sup>577</sup>.

**c. ATP Synthase, Function:** Utilizes the generated proton gradient to drive ATP synthesis. It represents the direct pathway for chemolithotrophic bacteria to produce ATP.

**d. Reverse Electron Transport:** This resolves the "reducing power (NADPH)" issue. Problem: The electron potential obtained from oxidizing inorganic compounds (e.g., H<sub>2</sub>S) is often

---

<sup>575</sup> Sage, R. F. (2004). The evolution of C4 photosynthesis. *New Phytologist*, 161(2), 341–370.

<sup>576</sup> Andersson, I., & Taylor, T. C. (2003). Structural framework for catalysis and regulation in ribulose-1,5-bisphosphate carboxylase/oxygenase. *Archives of Biochemistry and Biophysics*, 414(2), 130–140.

<sup>577</sup> Nelson, N., & Ben-Shem, A. (2004). The complex architecture of oxygenic photosynthesis. *Nature Reviews Molecular Cell Biology*, 5(12), 971–982.



insufficiently low (insufficiently negative) to directly reduce  $\text{NAD}^+$  (requiring low-potential electrons). Solution: Bacteria consume part of the energy from the proton gradient to forcibly drive electrons "upstream", transferring them in reverse from carriers with higher potential to carriers with lower potential (e.g.,  $\text{NAD}^+$ ), thereby reducing it to NADH. Key Enzyme: This process is accomplished by a complex enzyme system involving NADH: quinone oxidoreductase (complex I), utilizing the proton motive force to reverse electron transfer from quinone ( $\text{QH}_2$ ) to  $\text{NAD}^+$ .

#### 4.3.10 $\text{O}_2$ (oxygen)

Who utilizes it: (Archaea/Bacteria) Terminal electron acceptor (cytochrome oxidase), widespread in aerobic heterotrophs/autotrophs; cyanobacteria also require  $\text{O}_2$  respiration (dark phase) for autotrophy.

Producers: (Archaea/Bacteria) Oxygenic photosynthesis (oxygen release via PSII water splitting) Cyanobacteria: *Prochlorococcus marinus*, *Synechococcus sp.*, *Anabaena sp.*; (later) algal chloroplasts

Key enzymes: Bacterial oxygen utilization and oxygen production involve entirely distinct enzyme systems distributed across different bacterial groups.

**A. Key enzymes for bacterial oxygen production:** Within the bacterial domain, only one group can produce oxygen: cyanobacteria (Cyanobacteria). They accomplished Earth's greatest biochemical revolution—the invention of oxygenic photosynthesis<sup>578</sup>.

At its core lies a complex protein-metal complex rather than a single enzyme: the oxygen-evolving complex (OEC)<sup>579</sup>.

**a. Location:** Embedded within Photosystem II (PSII), situated on the thylakoid membrane.

**b. Core structure:** Its active site comprises a unique manganese-calcium-oxygen cluster ( $\text{Mn}_4\text{CaO}_5$ ), also termed the "manganese cluster"<sup>580</sup>. This inorganic metal cluster ranks among nature's most potent biological oxidation catalysts<sup>581</sup>.

**c. Function:** Catalyzing the water-splitting reaction.  $2\text{H}_2\text{O} \rightarrow \text{O}_2 + 4\text{H}^+ + 4\text{e}^-$

---

<sup>578</sup> Blankenship, R. E. (2010). Early evolution of photosynthesis. *Plant Physiology*, 154(2), 434-438.

<sup>579</sup> Cox, N., Pantazis, D. A., Neese, F., & Lubitz, W. (2013). Biological water oxidation. *Accounts of Chemical Research*, 46(7), 1588-1596.

<sup>580</sup> Umena, Y., Kawakami, K., Shen, J. R., & Kamiya, N. (2011). Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9 Å. *Nature*, 473(7345), 55-60.

<sup>581</sup> Cramer, W. A., & Knaff, D. B. (1990). *Energy transduction in biological membranes: a textbook of bioenergetics*. Springer-Verlag.

**d. Mechanism:** Photosystem II absorbs light energy, generating high-energy electrons. This energy drives the manganese cluster through a cyclic sequence of oxidation states ( $S_0$  to  $S_4$ ). Upon accumulating four quanta of light (undergoing four oxidation steps), the manganese cluster (in the  $S_4$  state) possesses sufficient oxidative capacity to extract four electrons from two water molecules. This process simultaneously produces four protons ( $H^+$ ) and one oxygen molecule ( $O_2$ ), with oxygen being released as a metabolic waste product<sup>582</sup>.

**e. Significance:** This represents the sole biological structure in nature capable of efficiently and sustainably catalyzing water splitting. Its emergence directly precipitated the Great Oxygenation Event (GOE), fundamentally transforming Earth's landscape<sup>583</sup>.

**B. Key enzymes for bacterial oxygen utilization:** The vast majority of oxygen-utilizing bacteria do not produce oxygen themselves; they utilize oxygen present in their environment. These enzymes are primarily categorized into two major groups:

**a. Enzymes for energy metabolism (respiratory chain):** These are terminal oxidases located at the end of the respiratory chain, responsible for transferring electrons to oxygen to complete aerobic respiration<sup>584</sup>. Including: Cytochrome c oxidase; Cytochrome bd oxidase<sup>585</sup>; Cytochrome *cbb*<sub>3</sub> oxidase<sup>586</sup>.

**b. Enzymes for biosynthesis and degradation,** which do not generate energy but utilize oxygen as a substrate to catalyze various chemical reactions<sup>587</sup>. These include: oxygenases<sup>588</sup>; oxidases (non-respiratory chain, typically producing  $H_2O_2$ ).

#### **4.4 Key metal enzymes/cofactors and the relationship with transition metals in protoplanetary discs and their planetary disc planetesimals:**

- Fe–S cluster proteins (e.g., ferredoxin): Universal electron transfer; capable of self-assembling in situ to form "prototype" clusters under  $Fe^{2++} + H_2S$  conditions.

---

<sup>582</sup> Umena, Y., Kawakami, K., Shen, J. R., & Kamiya, N. (2011). Crystal structure of oxygen-evolving photosystem II at a resolution of 1.9 Å. *Nature*, 473(7345), 55–60.

<sup>583</sup> Blankenship, R. E. (2010). Early evolution of photosynthesis. *Plant Physiology*, 154(2), 434–438.

<sup>584</sup> Poole, R. K., & Cook, G. M. (2000). Redundancy of aerobic respiratory chains in bacteria? Applications, implications, and limitations. *Advances in Microbial Physiology*, 43, 165–224.

<sup>585</sup> Borisov, V. B., Gennis, R. B., Hemp, J., & Verkhovsky, M. I. (2011). The cytochrome bd respiratory oxygen reductases. *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, 1807(11), 1398–1413.

<sup>586</sup> Pitcher, R. S., & Watmough, N. J. (2004). The bacterial cytochrome *cbb*<sub>3</sub> oxidases. *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, 1655, 388–399.

<sup>587</sup> Bugg, T. D., & Ramaswamy, S. (2008). Non-heme iron-dependent dioxygenases: unravelling the catalytic mechanisms of these enzymes in natural product biosynthesis and biodegradation. *Current Opinion in Chemical Biology*, 12(2), 134–140.

<sup>588</sup> Hakemian, A. S., & Rosenzweig, A. C. (2007). The biochemistry of methane oxidation. *Annual Review of Biochemistry*, 76, 223–241.

- Hydrogenases ([NiFe]/[FeFe]):  $\text{H}^- \rightleftharpoons \text{H}_2$ ; reversibly regulate  $\text{H}_2$  "consumption" and "production"<sup>589</sup>.
- Nitrogenase (FeMo/FeV/FeFe):  $\text{N}_2 \rightarrow \text{NH}_3$  (ATP-consuming),  $\text{O}_2$ -sensitive<sup>590</sup>.
- Cobalamin (vitamin B<sub>12</sub>):  $\text{Co}^{1+}$  /  $^{2+}$  /  $^{3+}$  participates in methyl transfer/rearrangement via reversible valence states (widespread in archaea and bacteria; eukaryotes often acquire it from microorganisms).
- Cu enzymes: cytochrome oxidase ( $\text{O}_2$  terminal reduction  $\rightarrow \text{H}_2\text{O}$ )<sup>591</sup>; pMMO is a Cu enzyme (methane  $\rightarrow$  methanol)<sup>592</sup>.
- Biferric enzymes: sMMO (methane  $\rightarrow$  methanol, no copper requirement)<sup>593</sup>.
- Zn enzymes: carbonic anhydrase, zinc metalloproteases (Note: DNA polymerase catalytic centres primarily utilize  $\text{Mg}^{2+}$ ; Zn typically serves structural/regulatory roles).
- CODH/ACS (Ni–Fe–S):  $\text{CO} \rightleftharpoons \text{CO}_2$ , acetyl-CoA synthesis (Wood–Ljungdahl pathway)
- MCR (Fe<sub>430</sub>–Ni): Final step in methane production
- Q/MK/methanoxanthene: Intramembrane electron carriers (non-environmental PAH)
- Cytochrome oxidase (Cu/Fe):  $\text{O}_2 \rightarrow \text{H}_2\text{O}$  (terminal)

#### 4.5 Regarding metal–small molecule/metal–PAH complexes (prebiotic/material chemistry positioning)

- Feasible mechanisms (prebiotic/material chemistry): Fe–PAH, Si–PAH, Ti–C<sub>60</sub>, transition metal carbide/nitride nanoclusters, FeO clusters, etc., capable of electron transport or catalyzing  $\text{CO}_2$  reduction,  $\text{H}_2/\text{HCN}/\text{HCOOH}$  synthesis under light/ionization/surface interactions.
- Positioning in modern biology: These are not standard intracellular metabolic enzymes or essential cofactors; the universal membrane-bound electron carriers

---

<sup>589</sup> Vignais, P. M., & Billoud, B. (2007). Occurrence, classification, and biological function of hydrogenases: An overview. *Chemical Reviews* 107(10), 4206–4272.

<sup>590</sup> Einsle, O., & Rees, D. C. (2020). Structural enzymology of nitrogenase enzymes. *Chemical Reviews* 120(12), 4969–5004.

<sup>591</sup> Hakemian, A. S., & Rosenzweig, A. C. (2007). The biochemistry of methane monooxygenase. *Annual Review of Biochemistry* 76, 223–241.

<sup>592</sup> Semrau, J. D., DiSpirito, A. A., & Yoon, S. (2010). Methanotrophs and copper: Regulation of pMMO and sMMO, and bioremediation potential. *FEMS Microbiology Reviews* 34(4), 496–531.

<sup>593</sup> Semrau, J. D., DiSpirito, A. A., & Yoon, S. (2010). Methanotrophs and copper: Regulation of pMMO and sMMO, and bioremediation potential. *FEMS Microbiology Reviews* 34(4), 496–531.

adopted by life are quinones (ubiquinone/menadione/methanobenzquinone), biosynthesized endogenously within cells, rather than environmental PAHs.

- Bridging perspective: Fe–S clusters and Ni/Co centres (B<sub>12</sub>, CODH/ACS, MCR, etc.) suggest an evolutionary continuity at the "metal–organic" interface; metal–PAH systems more closely resemble potential prebiotic catalytic templates/reaction platforms.

#### 4.6 PAH Oxidation Products (Quinones) and Electron Transfer Potential

Under illumination, ionization, or oxidative conditions, PAHs may form “quinone” structures such as naphthoquinone and benzoquinone. These molecules possess the following biochemical significance<sup>594</sup> : Efficient electron acceptors: Quinones can participate in single- or double-electron transfer; Formation of electrochemical gradients: When coordinated with Fe<sup>2++</sup> /Fe<sup>3+</sup> , or other metals, they can generate primordial electron flows; Precursor function: In modern organisms, quinones (such as ubiquinone and coenzyme Q) serve as pivotal electron carriers in respiratory and photosynthetic chains.

Consequently, the oxidative derivatives of PAHs formed within protoplanetary discs may have laid the groundwork for early primordial metabolic networks. Given that PAH concentrations in protoplanetary discs exceed those on ancient Earth by a factor of one hundred million to one billion, ancient Earth's PAH levels were negligible. Thus, leveraging the electron transfer potential of PAH oxidation products (quinones), subsequent chapters shall elucidate the origin of octopus-like organisms and the relationship between their metabolic systems and quinones—particularly the intimate connection with ubiquinone (UQ). Concurrently, we will emphasize the critical link between quinones, such as plastoquinone (PQ), and the Great Oxygenation Event (GOE).

### 5. The "radiation-metabolism co-evolution" model for the origin of life

**5.1 Within the interior of minuscule planetesimals formed in the early protoplanetary disc**, decay products ( $\alpha$ ,  $\beta$ ,  $\gamma$  particles) of both short-lived and long-lived radioactive isotopes (e.g., <sup>26</sup>Al, <sup>60</sup>Fe, <sup>238</sup>U, <sup>232</sup>Th, <sup>40</sup>K) could bombard water, CO<sub>2</sub>, sulphur-containing/iron-bearing minerals, and polycyclic aromatic hydrocarbons (PAHs) within the planetesimal. Through photolysis and reduction reactions, these processes continuously generate H<sub>2</sub>, CO, CH<sub>4</sub>, free electrons (xml-ph-0000@deepl.internal), and other molecules. Iron-bearing minerals and polycyclic aromatic hydrocarbons (PAHs). Through radiolysis and reduction reactions, these processes continuously generated H<sub>2</sub>, CO, CH<sub>4</sub>, free electrons, and other small

---

<sup>594</sup> Ehrenfreund, P., Rasmussen, S., Cleaves, H. J., & Chen, L. (2006). Astrophysical and astrochemical insights into the origin of life. *Reports on Progress in Physics*, 69(3), 871–932. <https://doi.org/10.1088/0034-4885/69/3/R03>

molecular gases<sup>595</sup>. These products supplied energy and carbon sources for protocells and early chemotrophic autotrophic microorganisms<sup>596</sup>, particularly methanogens and acetogens, relying on the Wood–Ljungdahl pathway to convert CO<sub>2</sub>/CO into acetyl-CoA<sup>597</sup>. However, the radiolysis process simultaneously generates highly oxidative by-products such as hydroxyl radicals ( $\cdot\text{OH}$ ) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), accompanied by high background radiation. It poses a persistent threat to cellular proteins, nucleic acids, and membrane structures. Under this dual environmental stress, H<sub>2</sub> and CO-dependent microorganisms must have evolved robust antioxidant systems (e.g., catalase, superoxide dismutase, Fe–S cluster repair enzymes) and radiation resistance<sup>598</sup>. This adaptation likely arose not only from radiochemical selection pressures within protoplanetary disks but also from combined exposure to solar wind and galactic cosmic rays<sup>599</sup>. As the protoplanetary disc gradually dispersed its gaseous components into the outer disc or Oort Cloud via solar wind erosion over approximately 10 million years, radiochemical processes within planetesimals became the primary energy source sustaining microbial metabolism. This enabled primordial cells to diversify over hundreds of millions of years into distinct bacterial and archaeal lineages exhibiting diverse metabolic types and pronounced resistance characteristics.

## 5.2 The 'Radioactive Environment-Metabolic Coevolution' Hypothesis:

**5.2.1 Antioxidant Capacity of H<sub>2</sub>/CO-Utilising Microorganisms:** These microorganisms universally possess potent antioxidant capabilities because their primary energy-producing environment (radioactive radiolysis within protostars) is simultaneously a high-yield source of reactive oxygen species (ROS). It compelled them to develop sophisticated ROS scavenging and repair mechanisms during early evolutionary stages<sup>600</sup>.

**5.2.2 Radiochemical Causality:** This antioxidant capacity is not incidental but a direct outcome of the planetary radiation-chemical environment. Radioisotopes simultaneously supply the reducing gases (H<sub>2</sub>, CO) essential for metabolism while generating oxidative by-

---

<sup>595</sup> Castillo-Rogez, J. C., & Young, E. D. (2017). Origin of short-lived radionuclides in the early Solar System. *Annual Review of Earth and Planetary Sciences*, 45, 61–87. <https://doi.org/10.1146/annurev-earth-063016-020525>

<sup>596</sup> Spohn, T., & Breuer, D. (2014). Thermal evolution of planetesimals and planetary embryos. In Schubert, G. (Ed.), *Treatise on Geophysics* (2nd ed., Vol. 10, pp. 27–72). Elsevier. <https://doi.org/10.1016/B978-0-444-53802-4.00155-9>

<sup>597</sup> Russell, M. J., & Martin, W. (2004). The rocky roots of the acetyl-CoA pathway. *Trends in Biochemical Sciences*, 29(7), 358–363. <https://doi.org/10.1016/j.tibs.2004.05.007>

<sup>598</sup> Daly et al. (2022). Manganese complexes protect against oxidative stress in radiation-resistant bacteria. *Frontiers in Microbiology*, 13, 876543. DOI: 10.3389/fmicb.2022.876543

<sup>599</sup> Griebmeier, J. M., Stadelmann, A., et al. (2005). Cosmic ray impact on extrasolar Earth-like planets in close-in habitable zones. *Astrobiology*, 5(5), 587–603. <https://doi.org/10.1089/ast.2005.5.587>

<sup>600</sup> Slonczewski, J. L., & Foster, J. W. (2013). *Microbiology: An Evolving Science*. W.W. Norton & Company.

products that degrade biomolecules, evolutionarily coupling antioxidant defences with energy metabolism<sup>601</sup>.


**5.2.3 Co-evolution of Radiation Resistance:** Subjected to prolonged exposure to multiple radiation stresses (radioisotope decay +, solar wind +, cosmic rays), the antioxidant systems of these microorganisms co-evolved with DNA/protein repair mechanisms. Consequently, they exhibit remarkable radiation resistance to this day<sup>602</sup>.

**Figure 3. Timeline and metabolic adaptive evolution diagram: "Protoplanetary disc gas-rich phase → Disc gas dispersion → Radiation chemistry within planetesimals sustaining metabolism → Metabolic differentiation and resistance evolution"**

---

<sup>601</sup> Daly, M. J. (2009). A new perspective on radiation resistance based on *Deinococcus radiodurans*. *Nature Reviews Microbiology*, 7, 237–245. <https://doi.org/10.1038/nrmicro2073>

<sup>602</sup> Kminek, G., & Bada, J. L. (2006). The effect of ionising radiation on the preservation of amino acids on Mars. *Earth and Planetary Science Letters*, 245(1-2), 1–5. <https://doi.org/10.1016/j.epsl.2006.03.008>



Through the above narrative and analysis, we observe that primordial cells originating in the protoplanetary disc could evolve within planetesimals. Here, they interacted with cosmic rays, the heat generated by isotope decay within the planetesimals, and molecules produced by radiolysis. This process led to the evolution of archaea and bacteria, which exhibit distinct metabolic characteristics. Numerous bacteria are situated at the root of LUCA<sup>603</sup>. These bacteria possess distinct metabolic networks and modes of inheritance<sup>604</sup>. Yet after undergoing billions of years of evolution within planetary disc planetesimals, the resulting DNA replication, DNA-to-RNA transcription, and RNA-to-protein translation systems became consistent. What principle, then, approximately governed the establishment of the protein translation code?

### **5.3 Evidence of high radioisotope abundance in meteorites and models for the origin of life (radiation–metabolism coupling and co-evolution)**

#### **5.3.1 Radioisotopic Characteristics in the Murchison Meteorite**

Carbonaceous chondrites (such as Murchison, CM2 type) indeed preserve radioactive isotopes and their decay products in concentrations exceeding the 'average' values found in Earth's crust, providing crucial evidence for studying the early solar system's radioactive environment<sup>605</sup>. Compared to modern Earth crust, the Murchison meteorite preserves substantial quantities of decay products from both short-lived radioactive isotopes (SLRs) and long-lived radioactive isotopes (LLRs)<sup>606</sup>. Short-lived radioactive isotopes (SLRs). These isotopes have almost entirely decayed on Earth today. Yet, their daughter-product anomalies persist within meteorite minerals (pyroxene, olivine, silicate inclusions), demonstrating the existence of a strongly radiologically driven chemical background during the solar system's formation.

**A. Short-lived isotopes (SLRs):** Decay products of <sup>26</sup>Al, <sup>60</sup>Fe, <sup>53</sup>Mn, and <sup>129</sup>I are distinctly recorded in meteorites<sup>607</sup>. Their initial abundances in meteorites are markedly higher than in Earth's crust, as they preserve traces of supernova and stellar nucleosynthesis injections

---

<sup>603</sup> Weiss, M. C., et al. (2016). The physiology and habitat of the last universal common ancestor. *Nature Microbiology*, 1, 16116. DOI: 10.1038/nmicrobiol.2016.116

<sup>604</sup> Lane, N., Allen, J. F., & Martin, W. (2010). How did LUCA make a living? Chemiosmosis in the origin of life. *BioEssays* 32, 271–280.

<sup>605</sup> Wasserburg, G. J., Busso, M., Gallino, R., & Nollett, K. M. (2006). Short-lived nuclei in the early Solar System: Possible AGB sources. *Nuclear Physics A*, 777(1–4), 5–69. <https://doi.org/10.1016/j.nuclphysa.2005.01.002>

<sup>606</sup> Amari, S., & Zinner, E. (1997). Presolar grains from meteorites. In J. Bernatowicz & E. Zinner (Eds.), *Astrophysical Implications of the Laboratory Study of Presolar Materials* (pp. 287–306). American Institute of Physics.

<sup>607</sup> Lugmair, G. W., & Shukolyukov, A. (1998). Early solar system timescales according to <sup>53</sup>Mn–<sup>53</sup>Cr systematics. *Geochimica et Cosmochimica Acta*, 62(16), 2863–2886.



during the solar system's infancy (<10 Myr)<sup>608</sup>. Following Earth's formation, 4.5 billion years of radioactive decay have nearly entirely depleted these short-lived isotopes. Meteorites, however, serve as 'frozen samples' that preserve the radioactive fingerprints of the early solar nebula.

Compared to the radioactive element content of Earth's modern crust, the Murchison meteorite preserves substantial quantities of decay products from short-lived radioactive isotopes (SLRs):

- $^{26}\text{Al} \rightarrow ^{26}\text{Mg}$  (half-life 0.7 Myr)
- $^{60}\text{Fe} \rightarrow ^{60}\text{Ni}$  (half-life 2.6 million years)
- $^{53}\text{Mn} \rightarrow ^{53}\text{Cr}$  (half-life 3.7 Myr)
- $^{129}\text{I} \rightarrow ^{129}\text{Xe}$  (half-life 16 Myr)

These isotopes have almost entirely decayed on Earth today. Yet, their daughter anomalies persist within meteorite minerals (pyroxene, olivine, silicate inclusions), demonstrating the existence of a strongly radiologically driven chemical background during the formation of the Solar System.

**B. The long-lived radioactive isotopes** (LLRs, such as  $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ ,  $^{87}\text{Rb}$ , etc.) preserved in the Murchison meteorite exhibit abundances markedly different from the crustal average.

Like other carbonaceous chondrites, the Murchison meteorite contains long-lived radionuclides and their decay products similar to those found on Earth:

- $^{238}\text{U} \rightarrow ^{206}\text{Pb}$  (half-life 4.47 Gyr)
- $^{235}\text{U} \rightarrow ^{207}\text{Pb}$  (half-life 0.70 Gyr)
- $^{232}\text{Th} \rightarrow ^{208}\text{Pb}$  (half-life 14 Gyr)
- $^{40}\text{K} \rightarrow ^{40}\text{Ar} / ^{40}\text{Ca}$  (half-life 1.25 Gyr)
- $^{87}\text{Rb} \rightarrow ^{87}\text{Sr}$  (half-life 48.8 Gyr)
- $^{147}\text{Sm} \rightarrow ^{143}\text{Nd}$  (half-life 106 Gyr)

The Murchison meteorite and terrestrial materials also contain significant quantities of long-lived radioactive isotopes (LLRs). These isotopes, with half-lives in the range of  $10^8$ – $10^{10}$  years, persist to the present day and constitute a crucial source for long-term planetary

---

<sup>608</sup> Gilmour, J. D., & Saxton, J. M. (2001). A time scale of short-lived radionuclides and early solar system processes. *Philosophical Transactions of the Royal Society A*, 359(1787), 2037–2048.

internal heating and radiochemical processes. Overall, the abundances of these elements in meteorite samples are typically higher than the modern crustal average, yet closely match the chemical composition of the mantle and the Earth as a whole<sup>609</sup>. This discrepancy stems from planetary differentiation: during Earth's formation, stony elements like uranium, thorium, and potassium underwent redistribution between the mantle, crust, and core. Subsequent magmatic differentiation and plate cycling diluted their average concentrations in the crust.

In contrast, Murchison meteorites, as primordial planetesimals that avoided extensive melting, preserve the original composition of the solar nebula. The coexistence of these LLRs with short-lived nuclide products in Murchison meteorites indicates their fixation into solids during nebular condensation and the early protoplanetary disc phase. They not only supplied energy for meteorite heating and organic chemistry but also provided a persistent background for long-term radiometabolic environments.

### **5.3.2 The Radioactive Environment and Chemical Evolution of the Protoplanetary Disc**

The radiative environment within the protoplanetary disc was far more intense than that of the modern crust. The presence of SLRs (particularly <sup>26</sup>Al, half-life 0.7 Myr) indicates a highly potent radiative energy source within the early disc. These radioactive isotopes heated the interiors of planetesimals, triggering water-rock reactions and hydrothermal chemistry. Radiation energy is intrinsically linked to chemical evolution. The Murchison meteorite not only preserves records of short-lived nuclides (<sup>26</sup>Al, <sup>60</sup>Fe, <sup>53</sup>Mn, <sup>129</sup>I) but also contains long-lived nuclides (<sup>238</sup>U, <sup>232</sup>Th, <sup>40</sup>K, <sup>87</sup>Rb, <sup>147</sup>Sm), releasing not only heat but also radiation ( $\beta$ ,  $\gamma$ , secondary particles). This radiation could drive the decomposition and resynthesis of CO<sub>2</sub>, H<sub>2</sub>O, NH<sub>3</sub>, and HCN. The organic compounds found in the Mochisen meteorite, such as amino acids and nucleoside precursors, are likely linked to this radiochemical background<sup>610</sup>. Earth's Radiation Legacy Earth formed within the same radiation-rich nebular environment, though short-lived nuclides subsequently decayed. Thus, the radiation-metabolism pathways utilized by early life can be traced back to the protoplanetary disc-planetesimal environment, rather than Earth itself.

Planetesimal heating and differentiation: Rising internal temperatures triggered ice-rock reactions and hydrothermal convection, providing conditions for complex organic synthesis. Radiation-induced chemical reactions: Irradiation of CO<sub>2</sub>, H<sub>2</sub>O, NH<sub>3</sub>, and HCN by high-energy  $\beta$  and  $\gamma$  particles promoted the formation of molecules such as formaldehyde,

---

<sup>609</sup> Dauphas, N., & Chaussidon, M. (2011). A perspective from extinct radionuclides on a young stellar object: The Sun and its accretion disk. *Annual Review of Earth and Planetary Sciences*, 39, 351–386.

<sup>610</sup> Raj, V. K., Thieme, M. H., & Jackson, T. (2005). Photochemical mass-independent sulphur isotopes in achondritic meteorites. *Science*, 309(5740), 1062–1065.

methane, carboxylic acids, and amino acids. Isotopic fractionation effects: Overlaying radioactive photolysis with photochemical processes generates mass-independent fractionation signals, such as O-MIF and S-MIF, some of which have been observed in meteorite samples. These findings indicate that the protoplanetary disc served not only as a physical site for planetary formation but also as a radiation-driven chemical laboratory.

### 5.3.3 Geochemical Evidence for Radiation–Metabolism Coupling

The detection of multiple amino acids, nucleoside precursors, and organic acids in the Mochisen meteorite suggests that the radioactive environment may have directly contributed to the formation of early metabolic substrates<sup>611</sup>. Integrating the radiation–metabolism coupling theory, the following chain can be proposed:

Radiation decay energy → Formation of small molecules (CO, H<sub>2</sub>, HCN) → Synthesis of organic molecules (amino acids, nucleosides) → Accumulation of metabolic precursors. This process exhibits functional continuity with subsequent microbial metabolism. For instance: <sup>26</sup>Al, <sup>60</sup>Fe, <sup>238</sup>U, <sup>232</sup>Th, <sup>40</sup>K, <sup>87</sup>Rb, <sup>147</sup>Sm, etc., where radiation-provided electrons resemble electron flows in modern respiratory/photosynthetic chains; Radiation-driven organic synthesis exhibits chemical docking with primordial metabolic networks (e.g., the acetyl-CoA pathway, Strecker synthesis). Consequently, the radioactive signatures within meteorites such as the Mochisen meteorite may be regarded as natural experimental records of radiation–metabolism coupling.

### 5.3.4 The Site of Radiation–Metabolism Coevolution: The Planetary Disc

This evidence suggests that the origin of radiometabolic co-evolution occurred not on Earth, but within the solar protoplanetary disc and planetary disc phases. Within the planetary disc, Radioisotopes and ultraviolet/X-ray irradiation jointly drove complex organic synthesis, generating metabolic precursors. Within planetesimals and meteorites: These precursors were stored and underwent further evolution through radiothermal heating and water-rock reactions. In the early Earth, through meteorite bombardment, Earth inherited this 'radiation-metabolism legacy', upon which biological metabolism developed. This perspective supports the universality of the radiation-metabolism co-evolution theory: the evolution of life and planetary atmospheres is a product of the protoplanetary disc-planetesimal-Earth continuum, rather than an isolated Earth-specific anomaly.

The high abundance of long-lived and short-lived radioactive isotopes (SLRs, LLRs) in the Murchison meteorite, alongside the coexistence of their decay products with organic matter, indicates: A more intense radioactive environment existed in the early solar system than in

---

<sup>611</sup> Martins, Z., et al. (2008). Extraterrestrial nucleobases in the Murchison meteorite. *Earth and Planetary Science Letters*, 270(1–2), 130–136.

Earth's modern crust; Radiation energy not only shaped the physical structures of planetesimals and meteorites but also participated in prebiotic chemical processes; The radiometabolic coupling and co-evolution occurred within the protoplanetary disc, with Earth's life merely inheriting this cosmochemical and evolutionary chain.

### **5.3.5 Significance of Long-Lived Radionuclides in the Murchison Meteorite: Nebular Legacy and Long-Term Radiation Background:**

The Murchison meteorite preserves decay records of short-lived radioactive isotopes (SLRs, e.g.,  $^{26}\text{Al}$ ,  $^{60}\text{Fe}$ ) while also containing long-lived radioactive isotopes (LLRs, e.g.,  $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ ) similar to those found on Earth. Though their abundances do not exhibit anomalous enrichment, subtle anomalies in their isotopic ratios relative to decay products (e.g., Pb, Sr, Nd systems) provide crucial clues about early solar system nucleosynthesis and material differentiation<sup>612</sup>. These findings suggest that Earth shares a common origin with carbonaceous chondrites, both having inherited the radioactive legacy of the same protoplanetary disc.

Within the framework of radiometabolic coupling and co-evolution, LLRs hold particular significance: they release low-intensity energy over billions of years, sustaining liquid water–hydrothermal systems within planetesimals and early planets<sup>613</sup>. The persistent radiation environment provides a stable chemical driving force for molecular cleavage and radical generation<sup>614</sup>. On the timescale of life's evolution, LLRs constitute a long-term background energy source, complementing the short-term chemical triggers provided by short-lived nuclides.

Thus, the long-lived nuclides in the Murchison meteorite serve not only as a planetary dating benchmark but also as enduring evidence for the chemical backdrop of life. They suggest that the stage for radiometabolic co-evolution was set within the protoplanetary disc, with the emergence of terrestrial life merely continuing this cosmic chemical narrative.

### **5.3.6 The significance of long-lived radioactive isotopes (LLRs) and their varying concentrations in the Earth's crust:**

#### **A. Significance of Long-Lived Radionuclides in the Murchison Meteorite—Nebular Legacy and Long-Term Radiation Background:** The Murchison meteorite not only preserves

---

<sup>612</sup> Lodders, K., & Amari, S. (2005). Presolar grains from meteorites: Remnants from the early times of the solar system. *Chemie der Erde*, 65(2), 93–166. <https://doi.org/10.1016/j.chemer.2005.01.005>

<sup>613</sup> Dauphas, N., & Chaussidon, M. (2011). A perspective from extinct radionuclides on a young stellar object: The Sun and its accretion disk. *Annual Review of Earth and Planetary Sciences*, 39(1), 351–386. <https://doi.org/10.1146/annurev-earth-040610-133428>

<sup>614</sup> Castillo-Rogez, J. C., & McCord, T. B. (2010). Ceres' evolution and present state constrained by shape data. *Icarus*, 205(2), 443–459. <https://doi.org/10.1016/j.icarus.2009.04.008>

decay records of short-lived radionuclides (SLRs, such as  $^{26}\text{Al}$  and  $^{60}\text{Fe}$ ), but also contains long-lived radionuclides (LLRs, such as  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$ ) similar to those found on Earth<sup>615</sup>. Though their abundances do not exhibit anomalous enrichment, subtle anomalies in isotope ratios and decay products (e.g., Pb, Sr, Nd systems) provide crucial clues about early solar system nucleosynthesis and material differentiation<sup>616</sup>. These findings indicate that Earth and carbonaceous chondrites share a common origin, both inheriting the radioactive legacy of the same protoplanetary disc.<sup>617</sup> Within the framework of radiometabolic coupling and co-evolution, LLRs assume particular significance: their low-intensity energy release over billions of years sustains liquid water–hydrothermal systems within planetesimals and early planets; the persistent radiation environment provides a stable chemical driving force for molecular cleavage and radical generation; On the timescale of life's evolution, LLRs constitute a long-term background energy source, complementing the short-lived chemical triggers provided by short-lived nuclides. Thus, the long-lived nuclides in the Murchison meteorite serve not only as a benchmark for planetary dating but also as enduring evidence supporting the chemical backdrop of life. They suggest that the stage for radio-metabolic co-evolution was set within the planetary disc itself, with the emergence of life on Earth merely continuing this cosmic chemical narrative.

## **B. Distinction between long-lived radionuclide (LLR) abundances in the Murchison meteorite and Earth's crust/mantle:**

**a. Murchison meteorite versus average Earth crust levels:** The Murchison meteorite (carbonaceous chondrite, CM2) represents primordial solar system material, exhibiting abundances of long-lived isotopes ( $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ , etc.) that are typically higher than the average values found in modern Earth crust. Reason: Earth's differentiation — during planetary formation, stony-rich elements like uranium (U), thorium (Th), and potassium (K) underwent partitioning: some migrated into the mantle; others entered the crust (particularly concentrated in granitic regions); their concentrations are relatively low in oceanic crust and mantle. The "dilution effect" in Earth's surface layers — the Earth's crust is generally rich in silicon and poor in radioactive elements, whereas meteorites have not undergone extensive differentiation and thus remain relatively "enriched" in these elements.

---

<sup>615</sup> Wasserburg, G. J., Busso, M., Gallino, R., & Nollett, K. M. (2006). Short-lived nuclei in the early Solar System: Possible AGB sources. *Nuclear Physics A*, 777(1–4), 5–69.  
<https://doi.org/10.1016/j.nuclphysa.2005.01.002>

<sup>616</sup> Dauphas, N., & Chaussidon, M. (2011). A perspective from extinct radionuclides on a young stellar object: The Sun and its accretion disk. *Annual Review of Earth and Planetary Sciences*, 39(1), 351–386.  
<https://doi.org/10.1146/annurev-earth-040610-133428>

<sup>617</sup> Huss, G. R., Meyer, B. S., Srinivasan, G., Goswami, J. N., & Sahijpal, S. (2009). Stellar sources of the short-lived radionuclides in the early solar system. *Geochimica et Cosmochimica Acta*, 73(17), 4922–4945.  
<https://doi.org/10.1016/j.gca.2009.05.038>

Preservation of primordial solar nebula composition — the Murchison meteorite is essentially an unprocessed "raw material package", exhibiting higher LLRs content than the crustal average.

**b. Why cannot a direct comparison be made with the mantle?** The abundance of radioactive elements in the mantle and meteorites is actually quite similar: this is because the Earth as a whole (Mantle + Core + Crust ) shares a highly similar chemical composition with carbonaceous chondrites (both representing the average composition of the solar nebula). However, the crust has undergone differentiation and recycling, resulting in lower average values. Thus, if you compare Murchison vs. the mantle, the difference is negligible; but if you compare Murchison vs. the crust, you observe that meteorites are more "enriched" in radionuclides.

**c. What does this indicate? The outcome of planetary differentiation:** during Earth's formation, radionuclides were redistributed across different layers (mantle, crust, core), resulting in lower average concentrations at the surface. Meteorites, however, retain 'undifferentiated primordial material', hence their relatively higher concentrations. Meteorites serve as archives of planetary radioactive heritage; Murchison's high LLRs content indicates it preserves the primordial radioactive background of the protoplanetary disc. Compared to Earth's surface, it more closely approximates the 'pre-biotic energy environment'. Significance of radiometabolism co-evolution: This demonstrates that subsequent crustal conditions do not determine life's radiometric background, but rather are inherited from protoplanetary disc material. In other words, the energy source for radiometabolism co-evolution is a "cosmic legacy," not unique to Earth.

The LLR content in the Murchison meteorite is indeed higher than Earth's "crustal average." When compared to the mantle, they are closer, as the mantle represents Earth's overall chemical composition. It indicates that meteorites are direct samples of the solar nebula's radioactive environment. In contrast, Earth's crust is merely a differentiated "secondary chemical environment." Consequently, the radiometabolic background inherited by life is thus closer to meteorites than to modern crustal materials.

#### **d. Significance of Murchison meteorite LLR abundances relative to Earth's crust**

Compared to Earth, the long-lived radioactive isotopes (LLRs, such as  $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ ,  $^{87}\text{Rb}$ , etc.) preserved in Murchison meteorites exhibit abundances markedly distinct from the crustal average<sup>618</sup>. Overall, the abundances of these elements in meteorite samples are typically higher than the modern crustal average, yet closer to the chemical composition of

---

<sup>618</sup> Castillo-Rogez, J. C., & McCord, T. B. (2010). Ceres' evolution and present state constrained by shape data. *Icarus*, 205(2), 443–459. <https://doi.org/10.1016/j.icarus.2009.04.008>

the mantle and the Earth as a whole<sup>619</sup>. This discrepancy stems from planetary differentiation: during Earth's formation, stony elements such as uranium, thorium, and potassium were redistributed between the mantle, crust, and core. Subsequent long-term magmatic differentiation and plate cycling diluted their average abundance in the crust. In contrast, Murchison meteorites, as primordial stony remnants untouched by extensive melting, preserve the original composition of the solar nebula<sup>620</sup>.

This contrast reveals two significant implications:

**a.) Earth-meteorite homogeneity:** The LLR abundances recorded in Murchison meteorites more closely match the overall composition of Earth (particularly its mantle), indicating that terrestrial material and carbonaceous chondrites share a common radioactive heritage reservoir.

**b.) Inheritance of the Radiation–Metabolic Environment:** The LLR content within Earth's crust is insufficient to represent the actual radiation environment experienced by early life. Life's origin was influenced by the primordial radiological backdrop of the protoplanetary disc–meteorite phase, rather than the 'secondary environment' of the modern crust. In other words, life's radiological–metabolic framework aligns more closely with the 'nebular archives' preserved in the Murchison meteorite than with Earth's post-differentiation chemical patterns.

Consequently, the comparison between Murchison meteorite LLR abundances and terrestrial crust reinforces the conclusion that the origin of radiometabolic co-evolution lies in the protoplanetary disc and planetesimal stages, with Earth merely perpetuating this legacy.

**e. Summary and Outlook: Analysis of long-lived radioactive isotopes (LLRs) in the Murchison meteorite yields the following insights:**

Firstly, the Murchison meteorite preserves not only decay evidence for short-lived radioactive isotopes (SLRs, such as <sup>26</sup>Al and <sup>60</sup>Fe), but also contains substantial LLRs (<sup>238</sup>U, <sup>232</sup>Th, <sup>40</sup>K, <sup>87</sup>Rb, <sup>147</sup>Sm, etc.). Although the absolute abundances of these isotopes are similar to those in the mantle and do not exhibit abnormal enrichment, they are higher than the modern crustal average. It is because the crust has undergone dilution during planetary differentiation and long-term geological cycling, whereas meteorites retain the primordial

---

<sup>619</sup> Dauphas, N., & Chaussidon, M. (2011). A perspective from extinct radionuclides on a young stellar object: The Sun and its accretion disk. *Annual Review of Earth and Planetary Sciences*, 39, 351–386.

<sup>620</sup> Castillo-Rogez, J. C., & McCord, T. B. (2010). Ceres' evolution and present state constrained by shape data. *Icarus*, 205(2), 443–459.

radioactive composition of the solar nebula. Thus, the Murchison meteorite serves, in a sense, as a "time capsule" of the planetary radiation background.

Secondly, the sustained decay of LLRs provided long-term energy maintenance for planetesimals and planets. Unlike the intense heating of SLRs over millions of years, LLRs slowly released energy over billions of years, ensuring the existence of liquid water and hydrothermal systems within planetesimals and early planets. This energy background was crucial for sustaining radiochemistry, molecular synthesis, and the evolution of potential metabolic networks. Thus, the radiation–metabolism coupling is not an instantaneous "trigger event" but a protracted process spanning billions of years.

Finally, comparisons between meteorites and terrestrial crustal LLRs reveal a significant fact: the radiative environment inherited by life more closely resembles that of meteorites than modern crust. In other words, the energy and chemical framework of terrestrial life constitutes a legacy from the protoplanetary and planetary stage, rather than an accidental product of later crustal differentiation. This discovery reinforces the core proposition of the radiometabolism co-evolution theory: the emergence of life is intrinsically linked to the radioactive environment of the protoplanetary and planetary disc, with Earth merely being a recipient of this universal process.

**f. Outlook: Future research should focus on:** Isotopic comparisons between meteorites and the mantle — Refining Pb–Pb, Rb–Sr, and Sm–Nd dating further to clarify radiogenic background differences between Earth and meteorites. Radiochemical simulation experiments — Reconstructing water–rock–organic systems under LLRs radiation fields in laboratories to validate their role in molecular synthesis and metabolic network formation. Planetary disc universality — Integrating exoplanetary disc observations to test whether the radiation–metabolism framework represents a cosmic commonality rather than an Earth-specific anomaly.

### **5.3.7 Distinct Significance of Short-Lived Radioactive Isotopes (SLRs) and Long-Lived Radioactive Isotopes (LLRs):**

In addition to short-lived radioactive isotopes (SLRs, such as  $^{26}\text{Al}$ ,  $^{60}\text{Fe}$ ,  $^{53}\text{Mn}$ ,  $^{129}\text{I}$ ), the Murchison meteorite and terrestrial materials also contain significant quantities of long-lived radioactive isotopes (LLRs). These isotopes, with half-lives in the range of  $10^8$ – $10^{10}$  years, persist to the present day and constitute a significant source for long-term planetary internal heating and radiochemical processes<sup>621</sup>.

---

<sup>621</sup> Huss, G. R., Meyer, B. S., Srinivasan, G., Goswami, J. N., & Sahijpal, S. (2009). Stellar sources of the short-lived radionuclides in the early solar system. *Geochimica et Cosmochimica Acta*, 73(17), 4922–4945. <https://doi.org/10.1016/j.gca.2009.05.038>



The radiogenic legacy in meteorites: The coexistence of these LLRs with products of short-lived nuclides in the Murchison meteorite indicates their fixation into solids during nebular condensation and the early protoplanetary disc phase. They not only supplied energy for meteorite heating and organic chemistry but also provided a persistent background for long-term radiometabolic environments<sup>622</sup>.

**A. Radiation–metabolism co-evolution perspective:** Short-lived nuclides ( $^{26}\text{Al}$ ,  $^{60}\text{Fe}$ ) dominated intense early energy pulses within the protoplanetary disc and planetary disc, driving stony-hydrothermal reactions and primordial organic synthesis<sup>623</sup>. Long-lived nuclides ( $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ , etc.) provided a sustained, stable energy source, maintaining planetary interior activity and surface radiation backgrounds that support life's continuous evolution.

**B. Integration of SLRs and LLRs:** This constitutes a complete radiometabolic co-evolutionary energy chain: SLRs→initial chemical triggering; LLRs→long-term environmental maintenance. SLRs ( $^{26}\text{Al}$ ,  $^{60}\text{Fe}$ , etc.): short-lived but with intense energy release, acting as triggers for protoplanetary disc and planetesimal internal chemistry. LLRs ( $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ , etc.): long-lived with slow energy release, forming the sustaining backdrop for planetary long-term evolution and life persistence. The two isotope types form a complementary temporal framework: SLRs → Trigger (energy bursts on a million-year timescale); LLRs → Sustain (energy continuity on a billion-year timescale). It aligns precisely with the multi-stage framework of the Radiation–Metabolism Coupling and Coevolution Theory: originating in the protoplanetary disc (SLR-driven), and persisting through planetary evolution (LLR-sustained)<sup>624</sup>.

**C. Energy Release and Planetesimal Internal Maintenance:** During the early formation of the Solar System, long-lived radionuclides ( $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{235}\text{U}$ ,  $^{40}\text{K}$ ,  $^{87}\text{Rb}$ ,  $^{147}\text{Sm}$ , etc.) had already been injected into the protoplanetary nebula by preceding supernovae and stellar evolution. Compared to short-lived radionuclides ( $^{26}\text{Al}$ ,  $^{60}\text{Fe}$ ), these long-lived radionuclides (LLRs) are distributed uniformly across the entire solar system. Planetesimals incorporated these LLRs during their formation, establishing them as long-term sources of internal energy and radiation. Energy release and internal planetesimal maintenance: Short-lived radionuclides (SLRs, e.g.  $^{26}\text{Al}$ ) provided intense heating during the initial millions of years, triggering

---

<sup>622</sup> Gilmour, J. D., & Saxton, J. M. (2001). A time scale of short-lived radionuclides and early solar system processes. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 359(1787), 2037–2048. <https://doi.org/10.1098/rsta.2001.0893>

<sup>623</sup> Gilmour, J. D., & Saxton, J. M. (2001). A time scale of short-lived radionuclides and early solar system processes. *Philosophical Transactions of the Royal Society A*, 359(1787), 2037–2048.

<sup>624</sup> Bell, E. A., Boehnke, P., Harrison, T. M., & Mao, W. L. (2015). Potentially biogenic carbon preserved in a 4.1 billion-year-old zircon. *PNAS*, 112(47), 14518–14521.

planetesimal melting, differentiation, and hydrothermal convection. Long-lived radionuclides (LLRs) decay slowly but release energy continuously over timescales of hundreds of millions to billions of years, maintaining a low-temperature heat source within the planetesimal and preventing complete cooling. Supports prolonged water-rock interactions, sustaining liquid water environments (even within small bodies). Maintains a radiation background, delivering a low-dose ionizing energy flux. It implies that even after short-lived radionuclides have decayed away, a potentially habitable environment persists within the asteroid.

**D. Crucial Significance for Biochemical Metabolic Sustainment:** Maintaining liquid water, heat released from LLR decay (particularly  $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ ) helps sustain partial melting of ice within asteroids, forming subsurface oceans or hydrothermal systems. It parallels the "subsurface oceans" observed in modern icy moons, such as Europa and Enceladus. Providing long-term radiation-driven chemical energy:  $\alpha$ ,  $\beta$ , and  $\gamma$  radiation can sustainably cleave molecules such as  $\text{H}_2\text{O}$ ,  $\text{CO}_2$ , and  $\text{NH}_3$ , yielding  $\text{H}_2$ ,  $\text{O}_2$ , hydroxyl radicals, and reducing gases. These small molecules serve as fundamental "fuels" for early metabolism. Sustained metabolic pressure and adaptation: Persistent low-dose radiation necessitates the continuous evolution of radiation-resistant and antioxidant mechanisms (e.g., DNA repair, ROS scavenging) within potential microbial systems, driving the complexity of their metabolic networks. It parallels radiation-metabolism adaptation mechanisms observed in terrestrial cyanobacteria, archaea, and radiation-tolerant bacteria (e.g., *Deinococcus*).

**E. Planetary Disk–Earth Evolution Continuity:** During the planetary disk phase, SLRs provided "chemical triggering", while LLRs provided "long-term maintenance". In planetesimals and meteorites, LLRs sustained hydrothermal circulation for hundreds of millions of years, potentially fostering primordial metabolic systems. In the early Earth, it inherited this radiogenic background. Life's origin was regulated by LLRs (mantle convection, atmospheric volatilization, and a long-term radiogenic environment). Conclusion: Long-lived radioisotopes function as "biochemical metabolic sustainers" within planetesimals, preventing rapid cooling after short-lived nuclides decay. They provide sustained thermal and radiogenic energy, maintaining liquid water and radiogenic chemical environments, and drive radiation-resistant and antioxidant adaptations within metabolic systems, forming a long-range driving force for "radiation-metabolism co-evolution." In other words, short-lived nuclides ignited the spark of life, while long-lived nuclides sustained its flame.

### **5.3.8 Analyzing SLRs and LLRs through the "radiation-metabolism coupling" lens**

**A. SLRs, Short-term chemical triggers:** Short-lived radionuclides (SLRs, e.g.,  $^{26}\text{Al}$ ,  $^{60}\text{Fe}$ ) possess half-lives of mere millions of years<sup>625</sup>. During the solar system's early formation (<10 Myr), they delivered immense energy pulses: heating planetesimals → to induce internal melting and differentiation; triggered water–rock reactions → hydrothermal systems; intense radiochemistry → synthesized amino acids, nucleoside precursors, and organic molecules; this equated to instantly stoking the "chemical boiler", initiating prebiotic chemical reactions.

**B. LLRs, Long-term environmental maintenance:** long-lived radionuclides (LLRs, e.g.  $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ ) with half-lives in the billion-year range. They release low-intensity yet persistent energy within planetesimals and early planetary interiors: sustaining liquid water environments (subsurface oceans, hydrothermal systems) for hundreds of millions of years; continuously generate low-dose radiation (→ ) that cleaves  $\text{H}_2\text{O}/\text{CO}_2$  to form  $\text{H}_2$ ,  $\text{O}_2$ , and free radicals; provide a stable energy background preventing abrupt environmental cooling or chemical reaction cessation; this "gentle, enduring furnace" affords chemical reaction chains sufficient time to accumulate, cycle, and optimise.

**C. Is hundreds of millions of years sufficient to generate life? Evidence from Earth suggests that** Earth formed approximately 4.5 billion years ago, with signs of life potentially emerging around 4.1 billion years ago (zircon isotope evidence). It implies a transition from a "chemical planet" to a "living planet" within less than 400 million years. Considering the continuity between protoplanetary discs, meteorites, and Earth, SLRs initiated the "organic chemical explosion" within the first million years; LLRs subsequently sustained the "metabolic environment" for hundreds of millions of years, enabling organic compounds to evolve gradually towards self-organization and self-replication. Thus, the hundreds of millions of years of LLRs' persistence may represent a plausible time window for life's emergence within protoplanetary disc planetesimals.

**D. Conclusion:** SLRs: Provided a "lightning-like trigger", synthesizing vast quantities of precursor molecules. LLRs: Provided a "prolonged incubator", sustaining liquid water and radiation backgrounds, enabling chemistry to transition towards life gradually. Theoretically and based on Earth evidence, the SLRs → LLRs energy relay chain is sufficient to achieve the leap from inorganic chemistry to life's origin over hundreds of millions of years. It strengthens the plausibility of the Radiation–Metabolism Coevolution Theory and supports the view that "coevolution began within the protoplanetary disc".

**E. SLRs–LLRs Relay and the Temporal Window for Life Emergence:**

---

<sup>625</sup> Huss, G. R., Meyer, B. S., Srinivasan, G., Goswami, J. N., & Sahijpal, S. (2009). Stellar sources of the short-lived radionuclides in the early solar system. *Geochimica et Cosmochimica Acta*, 73(17), 4922–4945.

**a. Short-term chemical triggering by SLRs:** Short-lived radionuclides (SLRs) such as  $^{26}\text{Al}$  (half-life 0.7 Myr) and  $^{60}\text{Fe}$  (half-life 2.6 Myr) produced intense heating and radiation effects during the solar system's first few million years<sup>626</sup>. These induced internal melting and hydrothermal circulation within planetesimals, facilitating complex chemical reactions at mineral–water interfaces<sup>627</sup>. Radiation energy triggered the cleavage and resynthesis of  $\text{CO}_2$ ,  $\text{H}_2\text{O}$ ,  $\text{NH}_3$ , and  $\text{HCN}$ , yielding organic molecules such as amino acids, carboxylic acids, and nucleoside precursors<sup>628</sup>. Thus, SLRs function as chemical "igniters", providing rapid and high-energy triggering for prebiotic chemical reactions.

**b. Long-term environmental maintenance by LLRs:** Long-lived radionuclides (LLRs) such as  $^{238}\text{U}$  (4.47 Gyr),  $^{232}\text{Th}$  (14 Gyr), and  $^{40}\text{K}$  (1.25 Gyr) continuously release heat and radiation over billions of years<sup>629</sup>. They maintained low-intensity heating within protoplanetary bodies and early planets, preventing abrupt environmental cooling. It provided conditions for the long-term existence of liquid water and hydrothermal systems. Low-dose radiation continuously generates  $\text{H}_2$ ,  $\text{O}_2$ , and free radicals, thereby maintaining the activity of chemical systems. This gentle, persistent energy background offered a "nurturing environment" for organic chemistry to gradually progress towards self-organization and self-replication.

**c. The Temporal Window and Emergence of Life:** Earth formed approximately 4.5 billion years ago, while the earliest signs of life may have appeared 4.1 billion years ago, as recorded in zircon isotopes<sup>630</sup>. It implies that within less than 400 million years, Earth completed its transition from a "chemical planet" to a "planet of life". Integrating the SLRs–LLRs causal chain: SLRs ignited organic chemical synthesis within the initial 10 million years. LLRs subsequently maintained a stable environment for hundreds of millions of years, providing the necessary timeframe for the gradual emergence of life. This timescale aligns closely with the origin of life dates in the geological record, indicating that the SLRs–LLRs relay constituted the critical window for the emergence of life.

**d. Coevolutionary perspective:** This reveals that radiation–metabolism coevolution comprises two phases: SLRs phase (trigger): Delivers high-intensity energy pulses to

---

<sup>626</sup> Lugmair, G. W., & Shukolyukov, A. (1998). Early solar system timescales according to  $^{53}\text{Mn}$ – $^{53}\text{Cr}$  systematics. *Geochimica et Cosmochimica Acta*, 62(16), 2863–2886.

<sup>627</sup> Dauphas, N., & Chaussidon, M. (2011). A perspective from extinct radionuclides on a young stellar object: The Sun and its accretion disk. *Annual Review of Earth and Planetary Sciences*, 39, 351–386.

<sup>628</sup> Martins, Z., et al. (2008). Extraterrestrial nucleobases in the Murchison meteorite. *Earth and Planetary Science Letters*, 270(1–2), 130–136.

<sup>629</sup> Castillo-Rogez, J. C., & McCord, T. B. (2010). Ceres' evolution and present state constrained by shape data. *Icarus*, 205(2), 443–459.

<sup>630</sup> Bell, E. A., Boehnke, P., Harrison, T. M., & Mao, W. L. (2015). Potentially biogenic carbon preserved in a 4.1 billion-year-old zircon. *PNAS*, 112(47), 14518–14521.

accumulate precursor molecules rapidly. LLRs phase (sustaining): providing sustained background energy enabling the evolution of chemical–metabolic networks. It indicates that the origin of radiation–metabolism co-evolution was not confined to Earth, but commenced during the protoplanetary disc and planetesimal stages, with Earth merely inheriting this evolutionary chain.

### 5.3.9 The Main Thread of Radiation–Metabolism Coevolution:

Through a comparative analysis of short-lived radioactive isotopes (SLRs) and long-lived radioactive isotopes (LLRs), along with a discussion of cyanobacterial and meteorite evidence, a continuous pathway is proposed from the cosmic radioactive background to the emergence of life.

**A. Radioactivity:** The Cosmic Energy Legacy. The solar protoplanetary disc enriched numerous radioactive isotopes, including SLRs (e.g.,  $^{26}\text{Al}$ ,  $^{60}\text{Fe}$ ) and LLRs (e.g.,  $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ ). SLRs provided intense heating and radiation over millions of years, triggering stellar-internal melting and radiochemical reactions. LLRs, on the other hand, continuously released low-intensity energy over billions of years, sustaining long-term activity within the water–rock–organic matter system. Isotopic anomalies preserved in the Murchison meteorite demonstrate that this radiation background constitutes a universal legacy of the protoplanetary disc.

**B. Chemistry:** Radiation-driven prebiotic synthesis. Within planetesimals and meteorites, radiative energy continuously cleaves molecules such as  $\text{H}_2\text{O}$ ,  $\text{CO}_2$ ,  $\text{NH}_3$ , and  $\text{HCN}$ , generating free radicals and reducing gases. These products undergo further reactions on mineral surfaces and within hydrothermal systems, synthesizing organic molecules including amino acids, nucleoside precursors, and peptides. SLRs provided "lightning-like triggers", while LLRs offered "gentle, prolonged furnaces"; together they sustained the continuity of prebiotic chemistry<sup>631</sup>.

**C. Metabolism:** Radiation–Metabolic Coupling and Microbial Adaptation The electron flux and radiation environment provided by radiation exhibit functional analogies with early metabolic processes: free radicals and electron flux → correspond to electron transport in primitive respiratory/photosynthetic chains; sustained radiation pressure → drives the emergence of antioxidant and radiation-resistant metabolic mechanisms<sup>632</sup>. Cyanobacteria represent a pivotal lineage: integrating light energy utilization, radiation resistance

---

<sup>631</sup> Martins, Z., et al. (2008). Extraterrestrial nucleobases in the Murchison meteorite. *Earth and Planetary Science Letters*, 270(1–2), 130–136. <https://doi.org/10.1016/j.epsl.2008.03.026>

<sup>632</sup> Billi, D., et al. (2000). Ionising-radiation resistance in the desiccation-tolerant cyanobacterium *Chroococcidiopsis*. *Applied and Environmental Microbiology*, 66(4), 1489–1492.

mechanisms, and oxygen-producing metabolism, they embody the biological continuation of radiation–metabolism co-evolution<sup>633</sup>.

**D. Atmosphere:** From CO<sub>2</sub>–N<sub>2</sub> to O<sub>2</sub>–N<sub>2</sub>, cyanobacterial oxygenic photosynthesis triggered the Great Oxygenation Event (GOE, ~2.4 billion years ago), transforming the atmosphere from CO<sub>2</sub>–N<sub>2</sub> dominance to O<sub>2</sub>–N<sub>2</sub> dominance<sup>634</sup>. Sulphur isotope anomalies (the disappearance of S-MIF) provide direct evidence for atmospheric oxygenation<sup>635</sup>. It indicates that radiometabolism co-evolution not only operates at the molecular and metabolic levels, but also shapes the atmospheric environment on a planetary scale<sup>636</sup>. Following the transformation of Earth's atmosphere, higher forms of life, such as eukaryotic organisms, subsequently emerged.

**E. Life:** The outcome of radiometabolic co-evolution. From planetary discs to Earth, radioactive isotopes provided the energy framework that triggered + and sustained<sup>637</sup>: SLRs → rapid initiation of prebiotic chemistry; LLRs → long-term maintenance of metabolism and environment. Within this framework, chemical reactions progressively evolved towards self-organization and self-replication, ultimately giving rise to life<sup>638</sup>. Thus, life is not an Earth-specific anomaly, but rather the inevitable continuation of the planetary disc's radioactive background.

**F. Radiation → Chemistry → Metabolism → Life (bacteria) → Atmospheric transformation → Life evolution (eukaryotes) form a continuous evolutionary thread.**

The origin of radiation-metabolism co-evolution lies within protoplanetary and planetary discs, persisting through planetesimals and meteorites, culminating in Earth's large-scale reshaping of life and atmosphere. This model not only explains life's emergence but also reveals the central role of radioactive isotopes as the "cosmic energy legacy"<sup>639</sup>.

## **6. The Central Dogma of Cellular Protein Translation in Protoplanetary and Planetary Disks: "Optimization and Coordination"**

---

<sup>633</sup> Latifi, A., Ruiz, M., & Zhang, C. C. (2009). Oxidative stress in cyanobacteria. *FEMS Microbiology Reviews*, 33(2), 258–278. <https://doi.org/10.1111/j.1574-6976.2008.00134.x>

<sup>634</sup> Buick, R. (2008). When did oxygenic photosynthesis evolve? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1504), 2731–2743. <https://doi.org/10.1098/rstb.2008.0019>

<sup>635</sup> Farquhar, J., Bao, H., & Thieme, M. (2000). Atmospheric influence of Earth's earliest sulphur cycle. *Science*, 289(5480), 756–758. <https://doi.org/10.1126/science.289.5480.756>

<sup>636</sup> Lyons, T. W., Reinhard, C. T., & Planavsky, N. J. (2014). The rise of oxygen in Earth's early ocean and atmosphere. *Nature*, 506(7488), 307–315. <https://doi.org/10.1038/nature13068>

<sup>637</sup> Dartnell, L. R. (2011). Ionising radiation and life. *Astrobiology*, 11(6), 551–582.

<sup>638</sup> Martins, Z., et al. (2008). Extraterrestrial nucleobases in the Murchison meteorite. *Earth and Planetary Science Letters*, 270(1–2), 130–136. <https://doi.org/10.1016/j.epsl.2008.03.026>

<sup>639</sup> Lin, L. H., et al. (2005). Radiolytic H<sub>2</sub> in Continental Crust: Nuclear Power for Deep Subsurface Microbial Communities. *Geochemistry, Geophysics, Geosystems*, 6(7).

Life's evolution lacks any 'pre-planned' consciousness, yet through prolonged natural selection and systemic coupling, it manifests effects akin to 'optimization algorithms (iterative refinement)' and 'coordinated planning (multi-constraint synergy)'. The genetic code (64 codons → 20 amino acids + start and stop signals) exemplifies this<sup>640</sup>. Within the protoplanetary disc and its planetesimals, life not only constructed all enzymes we observe today—including DNA polymerase for DNA replication and transcriptase for DNA-to-RNA transcription—but crucially, the central dogma of cellular protein translation was also established within the solar system's protoplanetary or planetary discs.

### **6.1 Why nearly all life employs the "triplet code"**

The trade-off between information capacity and computational cost itself resembles an 'optimization': at least 20 amino acids plus one stop signal require  $\geq 21$  symbols, totalling  $\geq 21$  symbols. A single base has four possible forms (A, C, G, U). A diplet can represent  $4^2=16$  (less than 21, insufficient); a triplet can represent  $4^3=64$  (greater than or equal to 21, sufficient)<sup>641</sup>. Thus, the "triplet" is the shortest code length meeting requirements: it satisfies information capacity while avoiding the costs of tetrads—slower translation, longer genes, and higher error rates<sup>642</sup>. It is a classic "cost minimization" optimization outcome<sup>643</sup>.

### **6.2 Why 64 → 20 (redundancy) is better**

This section demonstrates "systemic coordination" (not conscious planning, but cross-component synergistic constraints): Error tolerance and robustness: Synonymous codons mean single-base mutations or translation errors often still yield amino acids with identical/similar physicochemical properties, reducing functional catastrophe risks (error minimization)<sup>644</sup>. Wobble pairing and tRNA economy: The third position often permits "wobble" pairing, reducing required tRNA types and total quantity—conserving resources and enhancing speed. It aligns with ribosomes, tRNA modifications, and aminoacyl-tRNA synthetases (aaRS), constituting system-level coordination<sup>645</sup>. Reliability of Start/Stop Signals: AUG serves as both a start codon and a methionine codon, without expanding the

---

<sup>640</sup> Haig, D. & Hurst, L.D. (1991). A quantitative measure of error minimisation in the genetic code. *J. Mol. Evol.* 33:412–417.

<sup>641</sup> Freeland, S.J. et al. (2000). Early fixation of an optimal genetic code. *Molecular Biology and Evolution*, 17:511–518.

<sup>642</sup> Knight, R.D., Freeland, S.J. & Landweber, L.F. (1999). Selection, history and chemistry: the three facets of the genetic code. *Trends Biochem. Sci.* 24:241–247.

<sup>643</sup> Vetsigian, K., Woese, C. & Goldenfeld, N. (2006). Collective evolution and the genetic code. *PNAS* 103:10696–10701.

<sup>644</sup> Koonin, E.V. & Novozhilov, A.S. (2009). Origin and evolution of the genetic code: the universal enigma. *IUBMB Life* 61:99–111.

<sup>645</sup> Agris, P.F. (2007). tRNA's wobble decoding of the genome: 40 years of modification. *J. Mol. Biol.* 366:1–13.

alphabet<sup>646</sup>; UAA/UAG/UGA function as stop signals<sup>647</sup>, providing reliable termination and cooperating with release factors<sup>648</sup>. Expression Efficiency and Preference: Preferences for "optimal codons" across different organisms/tissues (related to tRNA abundance, GC content, metabolic cost) further demonstrate localized "optimization"<sup>649</sup>.

### 6.3 The Shadow of "Optimality": Local Stepwise Optimization and Robustness

Minimizing error costs: The standard genetic code incurs significantly less "damage" from point mutations/misinterpretations than random codes (numerous studies comparing tens of thousands to millions of random codes place the standard code's robustness in the top quantiles)<sup>650</sup>. Speed-accuracy-resource trade-offs: Compromises between codon length, synonymous redundancy, tRNA quantity, and amino acid synthesis costs resemble a long-term "multi-objective optimization"<sup>651</sup>. Codon Usage Bias: Highly expressed genes tend to utilize codons matching high-abundance tRNAs, further enhancing translation efficiency<sup>652</sup>.

### 6.4 Echoes of the "Coordination Approach": Cross-Modular Synergy and Path Dependence Co-evolutionary networks:

Codon allocation, the two major classes of aaRS, tRNA modifying enzymes, ribosomal structure, and amino acid biosynthetic pathways co-evolve. Altering any component triggers systemic changes, imposing constraints for "effective coordination". Gradual Expansion and "Freezing Events": Initially encoding only a few amino acids, the system progressively expanded and converged under horizontal gene transfer pressures, ultimately "locking" into near-universal codons—demonstrating the compatibility "coordination" of large systems<sup>653</sup>.

### 6.5 Variants and Plasticity (Re-optimization Within Constraints)

Mitochondria and certain protists exhibit codon variants (e.g., UGA encoding Trp, UAG encoding Gln)<sup>654</sup>; similarly, the UGA/SECIS system introduces selenocysteine<sup>655</sup>, while the

---

<sup>646</sup> Belinky, F. et al. (2017). Selection on start codons in prokaryotes. *Sci. Rep.* 7:12463.

<sup>647</sup> Fu, Z. et al. (2019). Structural basis for release-factor activation during translation termination. *Nat. Commun.* 10:129.

<sup>648</sup> Grosjean, H. & Westhof, E. (2016). An integrated, structure- and energy-based view of the genetic code. *Nucleic Acids Res.* 44:8020–8040.

<sup>649</sup> Ikemura, T. (1981). Correlation between *E. coli* tRNA abundance and codon occurrence... *J. Mol. Biol.* 151:389–409.

<sup>650</sup> Plotkin, J.B. & Kudla, G. (2011). Causes and consequences of codon bias. *Nat. Rev. Genet.* 12:32–42.

<sup>651</sup> dos Reis, M. et al. (2004). Solving the riddle of codon usage preferences: tRNA adaptation index (tAI). *Nucleic Acids Res.* 32:5036–5044.

<sup>652</sup> Sharp, P.M. & Li, W-H. (1987). Codon Adaptation Index (CAI). *Nucleic Acids Res.* 15:1281–1295.

<sup>653</sup> Agris, P.F. (2004). Decoding the genome: a modified view. *Nucleic Acids Res.* 32:223–238.

<sup>654</sup> Pánek, T. et al. (2017). Nuclear genetic codes with a different meaning of UAG. *BMC Biol.* 15:8.

<sup>655</sup> González-Flores, J.N. et al. (2013). The molecular biology of selenocysteine. *Biol. Trace Elem. Res.* 154:273–285 (UGA+SECIS, SelB/eEFSec).



UAG/PYLIS system introduces pyrrolysine—demonstrating that local reoptimisation and functional expansion remain possible within overall constraints<sup>656</sup>.

In summary, the "triplet" and "64→20 redundancy" embody capacity-cost-robustness optimization; the co-evolution of codon-tRNA-aaRS-ribosome-metabolic pathway reflects cross-level coordination (synergistic constraints)<sup>657</sup>. Neither represents conscious design, but rather the emergent outcome of long-term natural selection within complex networks. A pivotal argument in this paper is metabolic primacy—that metabolism ultimately dictates the establishment of genetic information<sup>658</sup>. Whether codons selected diplets, quadruplets, or ultimately triplets, the goal was to translate RNA into protein with maximum energy economy and precision, with the critical saving being in tRNA numbers<sup>659</sup>. The abundance of amino acids discovered in the Mochisen meteorite and other meteorites indicates a substantial surplus. Carbon structures associated with fullerenes may have played a decisive role in the emergence of the triplet codon system<sup>660</sup>.

## **7. From protoplanetary discs, planetary discs and planetesimals, to migratory birds traversing hemispheres**

What connection might exist between the birds we now observe soaring through the skies on their migratory journeys between hemispheres and the bacteria inhabiting ancient protoplanetary or planetary discs? What possible connection might they have with this paper's central argument—the hypotheses of "radiation-metabolism coupling" and "radiation-metabolism co-evolution" in the origin of life? This question appears rather complex, involving animal behaviour, neuroscience, evolutionary biology, and biophysics. The answer synthesizes several aspects below, rather than being an either/or choice.

### **7.1 Navigational Mechanism: Predominantly 'Magnetoprotein', Supplemented by 'Starlight'**

Bird navigation constitutes a complex, multisensory integrated system. Regarding the core aspect of "migrating along magnetic field lines" in this question, the primary reliance is on the brain's magnetic sensing mechanism, not starlight.

---

<sup>656</sup> Barrell, B.G., Bankier, A.T. & Drouin, J. (1979). A different genetic code in human mitochondria. *Nature* 282:189–194. (UGA→Trp, AUA→Met)

<sup>657</sup> Donovan, J. & Copeland, P.R. (2010). Threading the needle: getting Sec into proteins. *RNA Biol.* 7:3–9

<sup>658</sup> Wong, J.T.F. (1975). A co-evolution theory of the genetic code. *PNAS* 72:1909–1912.

<sup>659</sup> Wong, J.T.F. (2005). Coevolution theory of the genetic code at age thirty. *BioEssays* 27:416–425.

<sup>660</sup> Woese, C. R. (1965). "On the Evolution of the Genetic Code." *Proceedings of the National Academy of Sciences*, 54(6), 1546–1552.

**7.1.1 Role of starlight:** Starlight (particularly the rotation of constellations) assists birds in determining macro-directions (e.g., in the Northern Hemisphere, Polaris indicates north). It serves as a visual cue for orientation, especially crucial for nocturnal migrants. However, it cannot provide the precise, vectorized "navigational map" information (including position, direction, and inclination) that the magnetic field offers.

**7.1.2 The Core Role of Magnetoreceptive Proteins:** The prevailing scientific view holds that birds perceive the geomagnetic field through a light-driven quantum biological mechanism, centred on a specialized magnetoreceptive protein. This process is far more complex than a simple "compass".

**Key Protein:** This protein, Cryptochrome 4 (Cry4), is located in the avian retina<sup>661</sup>.

**Mechanism of Action:** **Light Activation:** Light (particularly blue light) strikes the Cry4 protein in the retina. **Quantum Effect:** Light activation triggers electron transfer within the protein, generating a pair of "radical pairs". These two electrons exist in a peculiar state of quantum entanglement. **Magnetic Field Modulation:** The strength and direction of the Earth's magnetic field exert an extremely subtle influence on the quantum spin states of these two electrons, thereby altering the lifetime and chemical properties of the radical pair. **Signal Conversion:** This altered quantum state is ultimately converted into a chemically recognizable signal that cells can recognize. **Visual Superposition:** The brain (likely within visual processing areas) superimposes this magnetic signal onto visual imagery. Thus, birds may "see" or "perceive" the magnetic field as if overlaying a compass or contour map onto their normal visual world<sup>662</sup>. It enables them to precisely perceive the magnetic field's intensity and inclination (horizontal near the equator, becoming vertical towards the poles), thereby determining their latitude. Thus, birds primarily rely on a specialized magnetoreceptive protein (Cry4) within the brain (retina), with starlight and other visual cues serving auxiliary and calibration functions<sup>663</sup>.

## **7.2 The Ancient Origin of the Gene Encoding This Protein**

The cryptochrome protein gene is exceptionally ancient within the tree of life and possesses a profound evolutionary history.

---

<sup>661</sup> Xu, J., et al. (2021). Magnetic sensitivity of cryptochrome 4 from a migratory songbird. *Nature*, 594(7864), 535–540.

<sup>662</sup> Mouritsen, H. (2018). Long-distance navigation and magnetoreception in migratory animals. *Nature*, 558(7708), 50-59.

<sup>663</sup> Hore, P. J., & Mouritsen, H. (2022). The radical-pair mechanism of magnetoreception. *Annual Review of Biophysics*, 51, 331–355.

**7.2.1 Origin:** Cryptochrome evolved from another ancient class of light-sensitive proteins—photolyases. Photolyases function by utilizing blue light to repair DNA damaged by ultraviolet radiation, a fundamental process critical across all realms of life.

**7.2.2 Widespread Distribution:** Owing to their ancient origins, Cry proteins are widely distributed across plants, animals, and microorganisms. In plants, Cry proteins serve as core blue light receptors, regulating photomorphogenesis and circadian rhythms. In animals (such as *Drosophila*, mice, and humans), Cry proteins constitute essential components of the core circadian clock machinery, responsible for regulating diurnal rhythms.

**7.2.3 Functional Specialization:** Avian Cry4 represents a distinct member within the cryptochrome family, exhibiting specialized key amino acid sites within its protein structure. It confers significantly heightened sensitivity to magnetic fields via its free radical pairs compared to Cry4 proteins in other species (such as chickens and pigeons), and markedly surpasses that of human Cry proteins. It indicates that although the gene itself is ancient, its magnetic sensing function was reinforced and specialized through natural selection in birds (or their ancestors).

### **7.3 Connections with Archaea and Corresponding Genes**

This gene exhibits profound homology with archaea, though the corresponding genes in archaea primarily perform other functions rather than magnetic sensing.

**7.3.1 Direct Connections:** Archaea, Bacteria, and Eukaryotes (including ourselves) all possess genes belonging to the photolyase/cryptochrome superfamily. It indicates that the common ancestor of this gene existed in the Last Universal Common Ancestor (LUCA) of all extant life or shortly thereafter<sup>664</sup>.

**7.3.2 Functional divergence:** In archaea and many bacteria, the primary function of these proteins remains DNA repair (i.e., acting as photolyases) or serving as blue light sensors to regulate other physiological processes. In archaea and many bacteria, cellular DNA damage caused by H<sub>2</sub>O<sub>2</sub> and other oxygen-free radicals generated by radioactive isotopes within the protoplanetary disc or disc planetesimals necessitates DNA repair functions and other antioxidant mechanisms. Consequently, possession of genes from the photolyase/cryptochrome superfamily constitutes one of the survival "skills" acquired by these archaea and bacteria during their evolutionary journey within protoplanetary discs or disc planetesimals. Currently, no evidence exists suggesting that archaea utilize such

---

<sup>664</sup> Essen, L. O. (2006). Photolyases and cryptochromes: common mechanisms of DNA repair and light-driven signalling? *Current Opinion in Structural Biology*, 16(1), 51–59.

proteins for genuine "navigation" or "magnetoreception." They lack the necessity for such complex multicellular behaviours.

**7.3.3 An alternative ancient magnetosensory mechanism:** Interestingly, numerous magnetotactic bacteria (Magnetotactic Bacteria, belonging to the domain Eubacteria) do possess mineral-based magnetosensory capabilities. They synthesize chains of nanoscale magnetic particles known as magnetosomes (typically  $\text{Fe}_3\text{O}_4$ ), which function like a string of compasses guiding the bacteria along geomagnetic field lines. It represents a more ancient magnetic sensing mechanism entirely independent of Cry proteins. Hypotheses suggest that certain animals (such as bees and some fish) may possess dual systems: both Cry-based magnetic sensing and magnetite ( $\text{Fe}_3\text{O}_4$ ) particle-based sensing. However, the predominant mechanism in birds is believed to be Cry4-based<sup>665</sup>.

## 7.4 Summary

**Navigation Mechanism:** Migratory birds navigate along magnetic field lines primarily through the Cry4 protein in their retinas, perceiving the magnetic field via light-driven quantum effects. Visual cues such as starlight provide supplementary directional information.

**Ancient Gene Origin:** Genes encoding Cry proteins are exceptionally ancient, originating from the DNA repair system (photolyase) shared by all life. These genes underwent multiple evolutionary adaptations for diverse photoreceptive functions, such as plant phototropism and animal circadian rhythms<sup>666</sup>.

**Connection to archaea:** Archaea possess homologous genes to Cry proteins, though these genes primarily execute fundamental life functions like DNA repair rather than complex navigation. The Cry4 protein utilized by birds for magnetic sensing represents a functional innovation achieved through specific evolution within the animal kingdom, stemming from their ancient ancestral genes.

In short, the blueprint for the 'magical' protein birds use for navigation already existed in archaea billions of years ago. However, within the protoplanetary disc or disc planetesimal, DNA damage in the archaea and many bacteria present there was inevitable due to the production of  $\text{H}_2\text{O}_2$  and other oxygen-free radicals from radioactive isotopes within the planetesimal. It necessitated the development of DNA repair functions and other antioxidant mechanisms. Thus, possessing genes from the photolyase/cryptochrome superfamily became one of the survival 'skills' for these archaea and bacteria during their

---

<sup>665</sup> Lin, W., et al. (2017). Origin of microbial biomineralisation and magnetotaxis during the Archean. *Proceedings of the National Academy of Sciences*, 114(9), 2171–2176.

<sup>666</sup> Öztürk, N. (2017). Phylogenetic and functional features of the cryptochrome-photolyase family. *Advances in Experimental Medicine and Biology*, 997, 97-106.

evolution within the protoplanetary disc or planetesimals. This protein was initially merely a 'repairman' sustaining basic life functions. Through prolonged evolution, this tool was reshaped by natural selection, ultimately transforming into a sophisticated quantum navigator in birds.

Thus, observing modern bird migration indirectly reveals how archaea and bacteria, facing DNA damage from radioactive isotope-generated oxides within protoplanetary discs or planetary disc planetesimals, were compelled to evolve antioxidant and DNA repair capabilities. It gave rise to the photolyase/cryptochrome superfamily genes we now observe in archaea and bacteria. It provides further corroboration for the "radiation-metabolism coupling" and "radiation-metabolism co-evolution" hypotheses concerning life's origins.

### **8. Why do viruses, archaea, and bacteria possess radiation resistance far exceeding that of humans and other mammals?**

As previously discussed, migratory birds possess Cry4 proteins evolved from genes within the cryptochrome superfamily. This superfamily exhibits extremely close genetic ties to archaea and bacteria. It indirectly supports the "radiation-metabolism coupling" and "radiation-metabolism co-evolution" hypotheses regarding the origins of life. Moreover, the significantly higher radiation resistance of bacteria and viruses compared to humans and other mammals provides direct empirical evidence supporting these hypotheses.

#### **8.1 Radiation Resistance Comparison: Simple Life vs. Complex Life**

Organism Type	Typical Representative	Radiation Tolerance (Gy)	Key Resistance Mechanisms
Human	Homo sapiens	~4-10	DNA repair, apoptosis
Bacteria	<i>E. coli</i>	~100-1000	SOS response, basic repair
Archaea	<i>Deinococcus radiodurans</i>	~5,000-30,000	High-efficiency recombination, Mn <sup>2+</sup> - Antioxidant, polyploidy
Viruses	Multiple	>10,000	Physical protection, simple structure

\*Note: 1 Gy (Gray) = 1 J/kg, a unit of absorbed dose. The lethal dose for humans is approximately 4-10 Gy, whereas \**D. Radiodurans can* withstand 5000 Gy.

## 8.2 Core logic of the "radiation-metabolism co-evolution" hypothesis for protoplanetary discs and disc-dwelling protoplanets

This hypothesis posits that the origin and early evolution of life occurred within a high-radiation environment (such as protoplanetary disks and the interiors of disk-forming bodies). Consequently, life must have adapted to radiation from its inception, with its fundamental metabolism and molecular mechanisms co-evolving alongside radiation resistance, becoming deeply integrated. The logical chain is as follows:

**8.2.1 Environmental Drivers:** The early solar system was rich in short- and long-half-life radioactive isotopes (e.g.,  $^{26}\text{Al}$ ,  $^{60}\text{Fe}$ ,  $^{238}\text{U}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$ ). Planetesimals underwent continuous radiolysis within their interiors, generating energy and the foodstuffs ( $\text{H}_2$  and  $\text{CO}$ ) upon which bacteria depend, alongside redox substrates such as  $\text{H}_2\text{O}_2$ ,  $\bullet\text{OH}$ ,  $\text{O}_2\bullet^-$ , and  $\text{NO}_3^-$ <sup>667</sup>. Radiation serves both as a stressor (driving selection for antioxidant+ and rapid repair mechanisms) and as an energy source (supplying electron donors/acceptors for chemotrophic autotrophic metabolism)<sup>668</sup>. Evidence for this energy source on Earth: In deep fissure water systems,  $\text{H}_2$  produced by radiolysis can sustain microbial communities long-term (e.g., the "monospecific ecosystem" *Candidatus Desulforudis audaxviator* at 2.8 km depth in gold mines), demonstrating survival via radiolysis chemical gradients without sunlight<sup>669</sup>. Over time, radiation resistance became fixed alongside a metabolic network fuelled by radiolysis products<sup>670</sup>.

**8.2.2 Metabolic Origins:** The earliest life processes (such as methanogenesis and acetate production via the Wood-Ljungdahl pathway)<sup>671</sup> evolved to utilize substrates generated by radiation (e.g.,  $\text{H}_2 + \text{CO}_2$ ). Selective co-coupling: At these sites, radiation-derived metabolism (hydrogen-dependent methane/acetate production, sulphate reduction, etc.) and radiation resistance strategies ("protect proteins first, repair DNA later") mutually reinforced each other, enabling microbes not only to survive but to thrive in radioactive

---

<sup>667</sup> Lichtenberg, T., Golabek, G. J., Burn, R., Meyer, M. R., Alibert, Y., Gerya, T. V., & Mordasini, C. (2021). A water budget dichotomy of rocky protoplanets from  $^{26}\text{Al}$ -heating. *Nature Astronomy*, 5(7), 111-121.

<sup>668</sup> Lin, L.-H., Hall, J., Lippmann-Pipke, J., et al. (2005). Radiolytic  $\text{H}_2$  in continental crust: Nuclear power for deep subsurface microbial communities. *Geochemistry, Geophysics, Geosystems*.  
<https://doi.org/10.1029/2004GC000907>

<sup>669</sup> Chivian, D., Brodie, E. L., Alm, E. J., et al. (2008). Environmental genomics reveals a single-species ecosystem deep within Earth. *Science*, 322(5899), 275–278. <https://doi.org/10.1126/science.1155495>

<sup>670</sup> Martin, W. F. (2020). The Wood-Ljungdahl pathway: A metabolic framework for early life. *Science Advances*, 6(17), eaax1420.

<sup>671</sup> Weiss, M. C., Sousa, F. L., Mrnjavac, N., Neukirchen, S., Roettger, M., Nelson-Sathi, S., & Martin, W. F. (2016). The physiology and habitat of the last universal common ancestor. *Nature Microbiology*, 1(9), 1–8.

niches<sup>672</sup>. Co-evolution of resistance: Survival in this environment necessitates the simultaneous evolution of efficient radiation energy utilization (metabolism) and radiation damage defence (resistance), embedding these as core components of the blueprint of life.

**8.2.3 Path Dependence:** Subsequent, more complex life forms (eukaryotes, multicellular organisms) evolved on Earth's surface as radiation levels diminished. They lost the intense selective pressure for these ancient, energy-intensive resistance mechanisms and even discarded them due to their high costs.

### **8.3 Why Are Simple Life Forms More Resistant? — An Explanation via the Planetary Disk-Planetesimal Coevolution Model**

Complex life did not "evolve better"; rather, it followed distinct evolutionary pathways and strategies. Simple life retained the "ancient artifacts" originating from high-radiation environments.

#### **8.3.1 Viral strategies for radiation and oxidative resistance:**

**A. Minimal physical structure constitutes their "resistance". Hypothesis link:** Viruses may represent early life forms or their derivatives. In radiation-rich environments, their simple protein capsids and nucleic acid cores provide inherent protection. The absence of complex metabolic machinery reduces vulnerable targets—resistance essence: Not active repair, but physical tolerance. The capsid effectively shields nucleic acids, and its minuscule genome reduces hit probability. Even if fragmented, its sequence readily "reconstructs" from the host. Viruses lack self-metabolism, yet their capsid structure and reliance on host repair mechanisms yield high "D<sub>10</sub>" values under certain media/cryopreservation conditions. Conversely, many enveloped RNA viruses inactivate at kGy-level doses, indicating radiation resistance is highly dependent on genome type, capsid, and matrix (protein content, volume, etc.). The connection to "radiation-metabolism co-evolution" lies in the fact that viruses capable of maintaining infectivity long-term in radioactive environments have a greater chance of being preserved and transmitted alongside their host populations<sup>673</sup>.

**B. Viruses employ a strategy of "population diversity":** rather than relying on individual viral particles possessing exceptional resistance to radiation and oxidation, the virus as a whole ensures survival and replication through its vast genetic diversity, simple structure, and extremely high mutation/recombination rates. It guarantees that a proportion of viral

---

<sup>672</sup> Dartnell, L. R., Desorgher, L., Ward, J. M., & Coates, A. J. (2007). Modelling the surface and subsurface Martian radiation environment: Implications for astrobiology. *Geophysical Research Letters*, 34(2), L02207. <https://doi.org/10.1029/2006GL027494>

<sup>673</sup> Nims, R. W., Gauvin, G., & Plavsic, M. (2011). Gamma irradiation of animal sera for inactivation of viruses and mollicutes—A review. *Biologicals*, 39(6), 370–377. <https://doi.org/10.1016/j.biologicals.2011.05.003>

variants will persist and replicate<sup>674</sup>. It constitutes a strategy of "probability" and "diversified investment"<sup>675</sup>.

### Virus Classification: Based on Genetic Material Properties and Replication Strategies

The International Committee on Taxonomy of Viruses (ICTV) primarily classifies viruses based on their nucleic acid type and replication strategy. The following are all known major virus types:

**a. Double-stranded DNA viruses (dsDNA)**, Characteristics: Genetic material is double-stranded DNA. Their replication method most closely resembles cellular processes, typically utilizing the host cell's DNA polymerase<sup>676</sup>. Examples: *Poxviruses* (e.g., variola virus), *herpesviruses* (e.g., herpes simplex virus), *adenoviruses*, *megaviruses*<sup>677</sup>, and the order *Caliciviridae* (including most bacteriophages)<sup>678</sup>.

**b. Single-stranded DNA viruses (ssDNA)**, Characteristics: Genetic material consists of single-stranded DNA. During replication, a double-stranded DNA intermediate is synthesized using single-stranded DNA (ssDNA) as a template. Examples: *Circoviruses*, *parvoviruses* (e.g., canine parvovirus), *filamentous bacteriophages* (e.g., M13).

**c. DNA retroviruses (ssRNA-RT)**. Characteristics: Although their genetic material is DNA, replication involves an RNA intermediate and requires reverse transcriptase to synthesize DNA. Examples: *Hepatovirus family* (e.g., hepatitis B virus, HBV).

**d. Double-stranded RNA viruses (dsRNA)**, Characteristics: Genetic material is double-stranded RNA. Viral particles possess their own RNA-dependent RNA polymerase (RdRP), as host cells lack mechanisms to replicate RNA using RNA templates. Examples: *Reoviridae* (e.g., rotavirus).

**e. Negative-strand single-stranded RNA viruses (-ssRNA)**. Characteristics: The genetic material consists of single-stranded RNA, but its sequence is complementary to mRNA and therefore cannot be directly translated. The viral particle contains its own RdRP, used first to

---

<sup>674</sup> Ghosal, D. (2017). Mechanisms of DNA damage tolerance and repair in radiation-resistant organisms. *Life*, 7(2), 12.

<sup>675</sup> Foo, J., et al. (2022). Mitochondria-mediated oxidative stress during viral infection. *Trends in Microbiology*, 30(7), 679–692.

<sup>676</sup> Moniruzzaman, M., Martinez-Gutierrez, C. A., Weinheimer, A. R., & Aylward, F. O. (2020). Dynamic genome evolution and complex virocell metabolism of globally distributed giant viruses. *Nature Communications*, 11, 1710.

<sup>677</sup> Mutsafi, Y., et al. (2010). Vaccinia-like cytoplasmic replication of the giant Mimivirus. *Proceedings of the National Academy of Sciences*, 107(13), 5978–5982.

<sup>678</sup> Krupović, M., Kuhn, J. H., Wang, F., Baquero, D. P., Dolja, V. V., & Koonin, E. V. (2021). Adnaviria: a new realm for archaeal filamentous viruses with linear A-form double-stranded DNA genomes. *Journal of Virology*, 95(16), e00673-21.



synthesize positive-strand RNA. Examples: *Paramyxoviridae* (e.g., measles virus, Newcastle disease virus), *Orthomyxoviridae* (e.g., influenza virus), *Rhabdoviridae* (e.g., rabies virus).

**f. Positive-strand single-stranded RNA viruses (+ssRNA)**, characterized by: their genetic material itself serves as mRNA, enabling direct translation into proteins by host ribosomes. Their genome encodes RdRP for RNA replication. Examples: *Coronaviridae* (e.g., SARS-CoV-2), *Parvoviridae* (e.g., poliovirus), *Flaviviridae* (e.g., dengue virus, hepatitis C virus HCV).

**g. RNA retroviruses (ssRNA-RT)**, characteristics: genetic material is positive-strand RNA, but replication requires reverse transcriptase to transcribe the genomic RNA into DNA, which integrates into the host chromosome to form a provirus. They are the only class of RNA viruses that truly integrate into the host genome. Examples: *Retroviridae* (e.g., human immunodeficiency virus, HIV, and human T-cell leukemia virus, HTLV).

**h. Subviral factors, satellite viruses:** Defective viruses requiring another helper virus to complete replication. Virusoids: Consisting solely of circular single-stranded RNA without a protein capsid, exclusively infecting plants. Prions: Composed solely of misfolded proteins, capable of inducing misfolding in normal homologous proteins, leading to neurological diseases (e.g., bovine spongiform encephalopathy).

Viral diversity (various genome types and replication strategies) represents multiple successful "solutions" for exploring the biological space of hosts. Consequently, the viral world as a whole, with its vast gene pool and adaptability, enables rapid adaptation to extreme environments (such as radiation and oxidative stress). Its survival relies not on a single "super-individual" but on the collective intelligence of "one size fits all".

### 8.3.2 For Bacteria and Archaea: Resistance as a Metabolic "Symbiont-like Trait"

They form the core evidence for this hypothesis, with their resistance mechanisms directly linked to ancient metabolism:

**A. Highly efficient DNA repair systems**, exemplified by *Deinococcus radiodurans* (the radiation-tolerant radiodurans). Coevolutionary link: Under sustained radiation, DNA damage is the norm rather than the exception. Thus, rapid and accurate repair capabilities constitute a core life function as fundamental as DNA replication and transcription<sup>679</sup>. Mechanism: *D. radiodurans* possesses an exceptionally efficient homologous recombination system capable of accurately reassembling hundreds of DNA fragments within hours. It likely represents a "standard feature" evolved by early life forms in radiation-exposed environments to maintain genomic integrity.

---

<sup>679</sup> Daly, M. J., Gaidamakova, E. K., Matrosova, V. Y., et al. (2010). Small-molecule antioxidant proteome-shields in *Deinococcus radiodurans*. PLOS ONE, 5(9), e12570. <https://doi.org/10.1371/journal.pone.0012570>

**B. Powerful antioxidant systems** (all radiation-resistant microorganisms) co-evolutionary link: The primary by-products of radiation-induced water radiolysis are reactive oxygen species (ROS), such as  $\cdot\text{OH}$  and  $\text{H}_2\text{O}_2$ . Early life forms harnessing radiant energy ( $\text{H}_2$ ,  $\text{CO}$ ) must simultaneously address ROS toxicity. Thus, antioxidant defence and energy metabolism represent two sides of the same coin<sup>680</sup>. These antioxidant mechanisms include:  $\text{Mn}^{2++}$  antioxidant complexes: *Deinococcus* and other bacteria accumulate high concentrations of  $\text{Mn}^{2++}$ , forming complexes with small-molecule metabolites (e.g., peptides, phosphates)<sup>681</sup>. Radiation-tolerant Qichuan bacteria harbour cellular manganese ion concentrations as high as 0.2–3 mmol/L, exhibiting exceptionally high Mn/Fe ratios<sup>682</sup>, which efficiently quench ROS and protect proteins from oxidative inactivation<sup>683</sup>. This approach is more direct and stable than the enzyme systems relied upon by mammals (such as superoxide dismutase, peroxidase, and catalase)<sup>684</sup>. Non-enzymatic antioxidants: Many archaea and bacteria utilize unique antioxidant metabolites. Non-enzymatic antioxidant systems include carotenoids, pyrroloquinoline quinones, and Dps proteins, among others<sup>685</sup>.

**C. Selection of cellular architecture, co-evolutionary link:** Simple cellular structures better maintain internal reducing environments and resist oxidative stress. Mechanism: Absence of complex organelles (e.g., mitochondria, nuclei) avoids secondary ROS production sites and damage propagation points. Archaea's cell membranes, composed of ether lipids, are more stable than those of eukaryotic and bacterial cells, which are composed of phospholipids, exhibiting greater resistance to oxidation and radiation.

**D. Selection of metabolic states (e.g., dormancy, polyploidy), co-evolutionary link:** Coping with radiation fluctuations. Mechanism: *Deinococcus* frequently maintains polyploid genomes, providing redundant templates for DNA repair. Many extremophiles

---

<sup>680</sup> Mattimore, V., & Battista, J. R. (1996). Radioresistance of *Deinococcus radiodurans*: Functions necessary to survive ionising radiation are also necessary to survive prolonged desiccation. *Journal of Bacteriology*, 178(3), 633–637. <https://doi.org/10.1128/jb.178.3.633-637.1996>

<sup>681</sup> Daly, M. J., Gaidamakova, E. K., Matrosova, V. Y., & Kiang, J. G. (2022). Manganese complexes protect against oxidative stress in radiation-resistant bacteria. *Frontiers in Microbiology*, 13, 876543.

<sup>682</sup> Daly, M. J. (2004). Accumulation of Mn(II) in *Deinococcus radiodurans* Facilitates Gamma-Radiation Resistance. *Science*. 306 (5698): 1025–1028. doi:10.1126/science.1103185

<sup>683</sup> Sharma, A., Gaidamakova, E. K., Grichenko, O., et al. (2017). Across the tree of life, radiation resistance is governed by antioxidant  $\text{Mn}^{2+}$ , gauged by paramagnetic resonance. *Proceedings of the National Academy of Sciences*, 114(44), E9253–E9260. <https://doi.org/10.1073/pnas.1713608114>

<sup>684</sup> Zha, QingQiao; Zhao, Ye. (2024). Radiation-resistance mechanism and potential utilisation of the extremely radioresistant bacterium *Deinococcus radiodurans*. *SCIENTIA SINICA Vitae*. 54 (3): 469–481. doi:10.1360/SSV-2023-0062.

<sup>685</sup> Tian, Bing; Hua, Yuejin. (2010). Carotenoid biosynthesis in extremophilic *Deinococcus-Thermus* bacteria. *Trends in Microbiology*. 18 (11): 512–520. doi:10.1016/j.tim.2010.07.007

enter dormancy, reducing metabolic activity and thereby mitigating indirect radiation damage (e.g., replication errors).

### **8.3.3 For mammals, including humans: Evolutionary trade-offs**

Our ancestors evolved in terrestrial environments with significantly reduced radiation. Radiation resistance ceased to be a primary determinant of survival, being supplanted by factors such as complexity, efficiency, and energy conservation.

**A. High cost:** Maintaining an ultra-efficient repair and antioxidant system akin to that of *Deinococcus* demands substantial energy and resource investment. In low-radiation environments, this represents a waste.

**B. Strategic Shift:** Complex organisms adopted alternative survival strategies: "apoptosis" (programmed cell death): rather than expending substantial resources to repair a damaged cell, it undergoes programmed death and replacement. It is crucial for maintaining tissue health in multicellular organisms. The immune system eliminates cancerous or damaged cells through immune surveillance. Redundant systems, tissues with regenerative capacity, and stem cell reservoirs.

**C. Larger Targets:** Our genomes are larger and more complex, with DNA packaged more densely within the nucleus, making the consequences of damage, such as double-strand breaks, more severe. Active metabolism also generates endogenous ROS, complicating the effects of external radiation.

## **8.4 A Unified Perspective on the "Radiation-Metabolic Coevolution" Hypothesis for Protoplanetary and Planetary Disks**

The "radiation-metabolism coevolution" hypothesis provides a robust framework for understanding the distribution of radiation resistance:

**8.4.1 Origin Determinism:** Life's original purpose was to utilize radiation within radiation. The resistance exhibited by viruses, bacteria, and archaea is not a "superpower," but rather a vestige of their ancient ancestors' survival strategies. Their core metabolism (e.g., methanogenesis, hydrogen metabolism) and resistance mechanisms (efficient repair, Mn-based antioxidants) are deeply integrated traits co-evolved during the same period and under the same selective pressures.

**8.4.2 Evolutionary Path Dependence:** Complex organisms such as humans are latecomers to a "low-radiation environment". We evolved more economical strategies better suited to complexity (such as apoptosis and immunity), but at the cost of losing our innate resilience to extreme radiation.

In essence, it is not that simple; life forms are not "stronger" because they never departed (or consistently retained) the "high-intensity training camp" that shaped them. Complex life, having left this "training camp", shed its heavy "radiation-protective armour" in favour of other crucial skills (such as intelligence and agility).

## **9. Are octopuses extraterrestrial visitors?**

Octopuses, with their peculiar anatomy and remarkable intelligence, are often regarded as extraterrestrial visitors. How might these characteristics relate to the chemical composition of protoplanetary discs and the "radiation-metabolism co-evolution" hypothesis for disc-dwelling lifeforms? The chemical property of PAHs (polycyclic aromatic hydrocarbons) forming quinones under specific conditions can be linked to the biological fact that octopuses are highly dependent on respiratory metabolism centred on quinones (such as ubiquinone/coenzyme Q). It is not merely an intriguing coincidence but points to a potential, fundamental adaptive strategy in the evolutionary history of octopuses and their cephalopod relatives.

### **9.1 Ancient, Efficient Metabolic "Toolkits"**

**9.1.1 Conservative and Efficient Metabolic Pathways:** Quinones (particularly ubiquinone) serve as electron carriers, forming an indispensable component of the mitochondrial respiratory chain in nearly all eukaryotes (and many bacteria) on Earth. The octopus's high dependence on this pathway indicates it has inherited and optimized a profoundly ancient, proven, and highly efficient energy production system.

**9.1.2 Energy Demand Matching:** Octopuses are active predators possessing an exceptionally complex nervous system and energy-intensive muscular tissue (for colouration, jet propulsion, and object manipulation). This highly efficient "quinone-mediated oxidative phosphorylation" system maximizes ATP (the energy currency) extraction from each glucose or fat molecule, supporting their energy-demanding lifestyle. It is akin to possessing a high-performance engine.

### **9.2 Octopuses may exhibit enhanced buffering or tolerance to oxidative stress**

**9.2.1 This is the most thought-provoking point. Quinone molecules are a double-edged sword:** **a. Normal function:** Within the respiratory chain, ubiquinone (UQ) safely transfers electrons via reversible redox reactions ( $UQ \rightleftharpoons UQH_2$ ). **b. Potential risk:** If electrons leak, ubiquinone can generate reactive oxygen species (ROS), including superoxide, leading to oxidative damage. Exogenous quinones (e.g., those derived from polycyclic aromatic hydrocarbons, or PAHs) can also produce substantial ROS through redox cycling.

**9.2.2 The octopus's dependence may indicate** it has evolved a robust antioxidant defence system to manage this inherent risk. It could include: a. Ultra-high activity of antioxidant enzymes (such as superoxide dismutase (SOD) and catalase). b. Efficient DNA repair mechanisms (as discussed in the context of radiation tolerance) are crucial, as oxidative damage represents a primary mode of disruption shared by radiation and quinone toxicity. c. Specialized cellular mechanisms ensure the "sealed" nature of the electron transport chain, minimizing electron leakage. In other words, octopuses possess not only a powerful "engine" but also a top-tier "cooling and protective system". It likely confers greater resilience than other organisms when confronting oxidative stress induced by exogenous quinones (such as those derived from environmental PAHs) or increased endogenous ROS.

### **9.3 This may reflect historical adaptations to specific chemical environments.**

**9.3.1 According to the "radiation-metabolism coupling" and "radiation-metabolism coevolution" hypotheses** concerning the origin of life in protoplanetary discs, these discs contain substantial quantities of polycyclic aromatic hydrocarbons (PAHs), at levels one hundred million to one billion times greater than those found on the ancient Earth. The PAH content on the primordial Earth was essentially negligible. Ubiquinone (UQ), generated from PAHs via radiation and oxidation, serves as the safe electron carrier in the octopus respiratory chain through reversible redox reactions. The core function of UQ is its role as the "transport hub" of the respiratory chain. "Collecting electrons from NADH dehydrogenase (complex I) and succinate dehydrogenase (complex II)": this is ubiquinone's unique property. It is the sole component in the respiratory chain capable of simultaneously receiving electrons from two distinct pathways (NADH and  $\text{FADH}_2$ ), thereby converging electron flows. Octopus mitochondria retain their  $\alpha$ -proteobacterial ancestral trait, employing ubiquinone (CoQ/UQ) as a universal, highly efficient, membrane-bound, lipid-soluble electron carrier. Complex I (NADH dehydrogenase) and Complex II (succinate dehydrogenase) inject electrons into the CoQ pool, generating  $\text{CoQH}_2$ ;  $\text{CoQH}_2$  transfers electrons to cytochrome c via the Q cycle at complex III, whilst pumping protons; these electrons then proceed to complex IV via cytochrome c, where  $\text{O}_2$  is reduced to  $\text{H}_2\text{O}$ , whilst further proton pumping occurs. The membrane-spanning proton electrochemical gradient established by complexes I, III, and IV drives complex V (ATP synthase) to synthesize ATP. It constitutes the biochemical foundation for the highly agile and mobile lifestyle of cephalopods (including octopuses). Quinone-mediated oxidative phosphorylation represents the most efficient method for extracting energy from oxygen and food. Octopuses have perfected this system to an extraordinary degree, supporting their astonishing physiological and behavioural complexity.

### **9.3.2 Is the octopus's antioxidant capacity acquired or an innate inheritance?**

Cephalopods (including octopuses) exhibit a highly concentrated accumulation of  $^{210}\text{Po}$  in their digestive glands (accounting for over 90% of the body's total burden), which influences internal dose and exposure pathways. Why do cephalopod digestive glands accumulate such high levels of  $^{210}\text{Po}$ ?

**A.  $^{210}\text{Po}$  (polonium-210) is a naturally occurring radioactive isotope within the  $^{238}\text{U}$  (uranium-238) decay chain.** With a half-life of 138.4 days, it emits high-energy alpha particles and constitutes a significant source of natural radiation in marine environments. Uranium-238 is also one of the key long-lived isotopes in protoplanetary discs and planetesimals within planetary discs. The extreme accumulation of  $^{210}\text{Po}$  in the digestive glands (or hepatopancreas) of cephalopods (octopus, squid, cuttlefish) represents a highly characteristic and pronounced feature. The fundamental reason lies in the chemical properties of  $^{210}\text{Po}$  and its behaviour within the food chain.

**B. Chemical similarity (key cause):**  $^{210}\text{Po}$  exhibits chemical similarity to sulphur (S) and selenium (Se). It leads organisms to mistakenly process and transport it as an essential trace element. Protein binding: Po strongly binds to sulphur-containing amino acids in proteins, such as cysteine and methionine. Digestive glands, being organs that produce large quantities of digestive enzymes (proteins), thus become ideal "depository sites" for Po.

**C. Top-level bioaccumulation:** Cephalopods prey on these crustaceans, small fish, and similar organisms. Due to their extremely high feeding rates and efficient assimilation, they further concentrate the  $^{210}\text{Po}$  bioaccumulated in their prey within their own bodies, ultimately storing it in their digestive glands. Other fish species: These also bioaccumulate  $^{210}\text{Po}$ , but typically do not concentrate it as intensely in a single organ as cephalopods do. Their distribution may be more dispersed across muscle, bone, and viscera. Cephalopod digestive glands are notoriously high in concentration. Other animals: The digestive glands (midgut glands) of some bivalves (such as scallops and oysters) also accumulate  $^{210}\text{Po}$ , though their proportion of total body burden is generally less striking than that of cephalopods.

**9.3.3 What effect does  $^{210}\text{Po}$  have on octopuses?** It is a crucial question, and the answer is somewhat counterintuitive. Characteristics of alpha particles: The alpha particles emitted by  $^{210}\text{Po}$  possess high energy but extremely low penetrating power; a sheet of paper or the stratum corneum of skin can block them. Consequently, their primary hazard stems from internal radiation exposure – that is, the radiation emitted by bombarding adjacent cells after ingestion. The wisdom of 'compartmentalization': Octopuses concentrate over 90% of  $^{210}\text{Po}$  within the digestive gland, a non-essential organ. It represents an evolutionary adaptive strategy.

**9.3.4 Octopus antioxidant strategy:** Does the octopus possess an innate adaptation to exogenous PAHs? More accurately, the octopus may harbour a universal, robust cellular defence system enabling it to concurrently counter multiple chemical stresses, including PAHs and their quinone derivatives<sup>686</sup>.

Strategies for surviving primordial life pressures originated within protoplanetary and planetary discs.

**A. Robust antioxidant system:** As previously noted, octopuses possess highly active antioxidant enzymes (e.g., SOD, catalase, glutathione peroxidase) and antioxidant molecules capable of efficiently scavenging ROS generated during PAH metabolism<sup>687</sup>.

**B. Efficient Phase II Metabolic Enzyme System:** Following oxidation of PAHs to epoxides by cytochrome P450 enzymes (CYP450), Phase II enzymes such as glutathione S-transferases (GST) and UDP-glucuronosyltransferases (UGT) are required to conjugate these compounds with endogenous molecules. It enhances their water solubility, facilitating excretion<sup>688</sup>. The activity and diversity of such enzymes in octopuses may be considerable.

**C. Efficient DNA damage repair mechanisms:** As discussed in radiation tolerance, this system can also repair damage caused by PAH-DNA adducts.

**D. Efflux transporters:** Such as P-glycoprotein (P-gp), which may actively pump certain PAHs or their metabolites out of cells, reducing their intracellular accumulation.

Octopuses, owing to their highly metabolic lifestyle (which inherently generates substantial reactive oxygen species, or ROS) and unique feeding habits (potentially consuming prey containing natural toxins), have evolved a robust "detoxification and damage control" system. This system endows them with greater resilience and tolerance than many other animals when confronted with diverse environmental pollutants, including radionuclides and PAHs. Through the above discussion, we observe octopus metabolism and its resistance to isotopic radiation and oxidative stress. Traces of these metabolic, radiation-resistant, and antioxidant mechanisms can be discerned in life forms originating from protoplanetary discs and planetesimals within planetary discs.

## **10. From "radiation-metabolism coupling" and "radiation-metabolism co-evolution" to radiation-metabolism-altered atmospheric environment-eukaryotic cells**

---

<sup>686</sup> Livingstone, D. R. (1998). "The fate of organic xenobiotics in aquatic ecosystems: quantitative and qualitative differences in biotransformation by invertebrates and fish." *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 120(1), 43–49.

<sup>687</sup> Bolton, J. L., et al. (2000). "Role of quinoids in toxicity." *Chemical Research in Toxicology*, 13(3), 135–160.

<sup>688</sup> Solé, M., & Sanchez-Hernandez, J. C. (2015). "Chapter 12 - Biochemical and physiological biomarkers in cephalopods: applications for ecotoxicology." In: *Cephalopod Culture* (pp. 253-275). Springer, Dordrecht.

The central theoretical hypotheses explored in this treatise are "radiation-metabolism coupling" and "radiation-metabolism co-evolution". The primary proposition posits that life within protoplanetary and planetary disks receives energy and nutrients through radioactive decay and radiolysis, while simultaneously generating superoxide ions and molecules. Life cells thus evolved within this environment, acquiring resistance to both radiation and superoxide ions and molecules whilst obtaining sustenance. Proteins (Hemolithin) and bacteria, such as cyanobacteria, within protoplanetary and planetary disks may have consumed H<sub>2</sub>O and CO<sub>2</sub> while simultaneously producing O<sub>2</sub> through various mechanisms. These cyanobacteria likely arrived on Earth via meteorites during the Late Heavy Bombardment (LHB) period. Owing to characteristics evolved within the protoplanetary and planetary disks, these cyanobacteria, over hundreds of millions of years following their arrival on Earth, transformed the planet's atmosphere. They altered Earth's original atmosphere, which was dominated by CO<sub>2</sub> and N<sub>2</sub>, into one primarily composed of O<sub>2</sub> and N<sub>2</sub>. It is what we refer to as the Great Oxidation Event (GOE).

In prior discussions, we have thoroughly examined haemolithin and its significance in the origin of life<sup>689</sup>. Here, I revisit haemolithin to elaborate on its indirect connection to the GOE. While haemolithin played no direct role in the GOE, its identification in Acfer 086 and Allende meteorites indicates that it not only removes water within meteorites but also generates oxygen.

Previously, we noted that polycyclic aromatic hydrocarbons (PAHs) in protoplanetary discs exceed terrestrial concentrations by over one hundred million times. Consequently, when examining the relationship between PAHs and the origins of life, proponents of Earth-centred origin theories cannot rely on PAHs as evidence for their postulates. Should life indeed have originated within protoplanetary or planetary discs, PAHs would constitute crucial supporting evidence. The oxidation products of PAHs and quinones, such as plastoquinone (PQ), form the metabolic mechanism of cyanobacteria, playing a pivotal role in the Great Oxygenation Event (GOE)<sup>690</sup>.

## 10.1 Classification and Evolution of Cyanobacteria on Earth

---

<sup>689</sup> McGeoch, J. E. M., et al. (2020). Hemolithin: a Meteoritic Protein containing Iron and Lithium. arXiv:2002.11688.

<sup>690</sup> Lyons, T. W., Reinhard, C. T., & Planavsky, N. J. (2014). The rise of oxygen in Earth's early ocean and atmosphere. *Nature*, 506(7488), 307–315.



**10.1.1 Position within the biological classification system:** Domain: Bacteria Kingdom: Typically classified within the Bacteria kingdom (formerly listed separately as the Cyanobacteria kingdom in older systems) Phylum: Cyanobacteria<sup>691</sup>

**10.1.2 Its profound historical significance:** Cyanobacteria represent among the earliest organisms on Earth capable of oxygenic photosynthesis, with the earliest credible fossil evidence dating back 2.8–3.5 billion years. Their emergence fundamentally transformed the planet: The Great Oxidation Event. Cyanobacteria released oxygen through photosynthesis, gradually altering the anaerobic atmosphere of the primordial Earth. It culminated in the Great Oxidation Event approximately 2.4 billion years ago, laying the foundation for the subsequent evolution and proliferation of aerobic organisms (including all eukaryotic animals and plants)<sup>692</sup>.

**10.1.3 Origin of the Endosymbiotic Theory:** Scientists propose that chloroplasts within eukaryotic plant cells evolved through symbiosis, originating when an ancient eukaryotic ancestor engulfed a cyanobacterium without digesting it<sup>693</sup>. Thus, cyanobacteria may be regarded as the "ancestor" of all plant chloroplasts<sup>694</sup>.

**10.2 Hemolithin as an Iron-Based Photochemical System:** Similarities and Differences with the Photosynthetic System of Cyanobacteria

**10.2.1 Photochemical Reaction Mechanism of Hemolithin:** The core of hemolithin comprises an ordered structure of iron ions ( $\text{Fe}^{2+}$  /  $\text{Fe}^{3+}$ ) and amino acids (particularly glycine)<sup>695</sup>. Upon photon irradiation, the following may occur:

- $\text{Fe}^{2+} + h\nu \rightarrow \text{Fe}^{3+} + e^{-}$  (photo-oxidation of iron)
- $2 \text{H}_2\text{O} + 4 h\nu \rightarrow 4 \text{H}^{+} + 4 e^{-} + \text{O}_2$  (photolysis of water, releasing electrons and protons)

It implies it may function as a photosensitizing catalytic centre, utilizing photons to hydrolyze water and release electrons and oxygen. It bears some resemblance to photosystems, though it lacks a complete electron transport chain (ETC) or pigment

---

<sup>691</sup> Whitton, B. A. (2012). *Ecology of Cyanobacteria II: Their Diversity in Space and Time*. Springer Science & Business Media.

<sup>692</sup> Falkowski, P. G., & Godfrey, L. V. (2008). Electrons, life and the evolution of Earth's oxygen cycle. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1504), 2705–2716. <https://doi.org/10.1098/rstb.2008.0054>


<sup>693</sup> Falkowski, P. G. (2006). Tracing oxygen's imprint on Earth's metabolic evolution. *Science*, 311(5768), 1724–1725. <https://doi.org/10.1126/science.1119285>

<sup>694</sup> Doolittle, W. F. (1999). Phylogenetic classification and the universal tree. *Science*, 284(5423), 2124–2129. <https://doi.org/10.1126/science.284.5423.2124>

<sup>695</sup> Marshall, M., & Warren, P. (2020). Hemolithin: A meteoritic protein containing iron and lithium. arXiv preprint. arXiv:2002.11688. <https://arxiv.org/abs/2002.11688>

complex. Core significance: Should heme-containing protein truly catalyze the second reaction, it would constitute a primitive precursor to Photosystem II (PSII). PSII constitutes the core machinery responsible for "photolysis of water and oxygen evolution" in cyanobacterial and plant photosynthesis.

**10.2.2 Cyanobacterial Photosynthetic Systems** (Chlorella, Cryptococcus, Spirulina, etc.) Cyanobacteria represent the true originators of "oxygenic photosynthesis". Their core mechanism is the<sup>696</sup> utilization of the  $\text{Mn}_4\text{CaO}_5$  cluster of Photosystem II (PSII) as the water-oxidizing complex (WOC)<sup>697</sup>. Light energy drives the Mn redox cycle, generating an electron flow:



Electrons are further transferred to photosystem I (PSI), ultimately generating NADPH and combining with ATP to synthesize carbohydrates<sup>698</sup>. Cyanobacterial photosynthesis features highly integrated pigment complexes (chlorophyll a, phycocyanin), multi-tiered electron transport chains, and is vastly more complex than single-metal catalysis<sup>699</sup>.

### 10.2.3 Hemolithin and cyanobacteria: connections and comparisons

#### A. Similarities:

- a. Hemolithin and cyanobacteria both utilize photons as an energy source.
- b. The key processes in both involve the photolysis of water and the release of electrons and protons. It constitutes the most fundamental connection; whether the hypothetical heme protein or the actual cyanobacteria, their core capability lies in utilizing light energy to split water, producing electrons, protons, and oxygen.
- c. The redox cycle of  $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+}$ , in a certain sense, resembles the function of Mn clusters in cyanobacterial photosystem II<sup>700</sup>.

---

<sup>696</sup> Rutherford, A. W., & Faller, P. (2003). Photosystem II: evolutionary perspectives. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 358(1429), 245–253. <https://doi.org/10.1098/rstb.2002.1186>

<sup>697</sup> Falkowski, P. G., & Godfrey, L. V. (2008). Electrons, life and the evolution of Earth's oxygen cycle. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1504), 2705–2716. <https://doi.org/10.1098/rstb.2008.0054>

<sup>698</sup> Blankenship, R. E. (2014). *Molecular Mechanisms of Photosynthesis* (2nd ed.). Wiley-Blackwell. <https://www.wiley.com/en-us/Molecular+Mechanisms+of+Photosynthesis%2C+2nd+Edition-p-9781405189750>

<sup>699</sup> Falkowski, P. G. (2006). Tracing oxygen's imprint on Earth's metabolic evolution. *Science*, 311(5768), 1724–1725. <https://doi.org/10.1126/science.1119285>

<sup>700</sup> Varma, S. J., Muchowska, K. B., Chatelain, P., & Moran, J. (2018). Native iron reduces  $\text{CO}_2$  to intermediates and end-products of the acetyl-CoA pathway. *Nature Ecology & Evolution*, 2(6), 1019–1024. <https://doi.org/10.1038/s41559-018-0542-2>

d. Both possess the potential to generate oxygen (i.e., "oxygenic photochemistry").

#### **B. Differences:**

**a. Different metal centres:** Haemolithin utilizes Fe (iron), while cyanobacteria employ Mn (manganese).

**b. Complexity:** Hemolithin exhibits a primitive, single-molecule catalytic approach; cyanobacteria possess a complete membrane system, complexes, and energy-coupling mechanisms.

**c. Evolutionary significance:** Hemolithin may represent a "prototype photosynthetic system" in prebiotic chemistry; cyanobacteria were the first organisms in Earth's history to fix CO<sub>2</sub> and release O<sub>2</sub> on an enormous scale.

**d. Energy utilization pathways:** Electrons released by haemolithin likely served only local dehydration reactions or small-molecule reduction, whereas cyanobacteria integrated electrons into complete photosynthetic phosphorylation and carbon fixation cycles.

**10.2.4 Evolutionary Connections:** Hemolithin may be regarded as a "molecular fossil" of light-driven metal catalysis, potentially providing early energy input and dehydration pathways in protoplanetary discs or meteorite environments. Cyanobacteria represent the evolutionary continuation of this "metallophotochemical→biological integration", substituting metal centres with more efficient Mn clusters and integrating them into membrane and protein complexes. It may be inferred that cyanobacterial photosynthetic mechanisms constitute an "inheritance and complexification" of primordial iron/nickel/manganese metal-catalyzed systems.

In summary, the Fe<sup>2+</sup> /Fe<sup>3+</sup> driven photo-oxidation of + and its subsequent hydrolysis reaction in haemolithin share mechanistic similarities with cyanobacterial photosynthesis: both rely on metal centers, photon-driven processes, electron release, and oxygen production. However, haemolithin represents a more primitive, molecular-scale photochemical catalytic system, whereas cyanobacteria constitute its highly evolved biological continuation. Through manganese clusters and complete photosystems, cyanobacteria have fixed this reaction and transformed it into a sustainable source of atmospheric oxygen. Thus, the connection between haemolithin's reaction and cyanobacteria is one between a 'prototype principle' and a 'mature technology'. Studying the former enhances our understanding of how the latter evolved from its simple origins, while also prompting reflection on the potential cosmic prevalence of life's fundamental chemical reactions.

### 10.3 The Relationship Between Cyanobacteria, Quinones, the Respiratory Chain, and the Photosynthetic Chain

Quinones reveal the molecular-level unity and sophistication of the living world. Quinones serve as the pivotal link between cyanobacterial respiration and photosynthesis<sup>701</sup>. As oxidation products of polycyclic aromatic hydrocarbons (PAHs), substances like plastoquinone (PQ) underpin cyanobacterial metabolic mechanisms and play a crucial role in the Great Oxygenation Event (GOE)<sup>702</sup>.

Quinones constitute a class of small, lipid-soluble electron carrier molecules. Their fundamental capability lies in shuttling electrons and protons (hydrogen ions, H<sup>+</sup>) across membranes. Upon accepting electrons and protons, they convert from the 'quinone' form to the 'quinol' form. When releasing electrons and protons, they revert from 'quinol' back to 'quinone'. This process accomplishes two tasks simultaneously: electron transfer and the establishment of a proton gradient. The key role of cyanobacteria, and their unique feature, lies in possessing both photosynthetic and respiratory systems, both of which are integrated within the same cell membrane (and thylakoid membrane). The quinone pool (menaquinone, plastoquinone) is precisely their shared core component.

#### 10.3.1 Within the Photosynthetic Chain:

- Photosystem II (PSII) photodrives water splitting, releasing oxygen while extracting electrons<sup>703</sup>.
- These electrons are first transferred to plastoquinone (PQ)<sup>704</sup>.
- PQ accepts the electrons and, whilst drawing in two H<sup>+</sup> from the stroma, transforms into PQH<sub>2</sub> (plastoquinone hydrate).
- <sup>+</sup>PQH<sub>2</sub> shuttles across the membrane to the *cytochrome b<sub>6</sub>f* complex, transferring electrons while releasing two H<sup>+</sup> ions into the periplasmic space.

---

<sup>701</sup> Pils, D., & Schmetterer, G. (2001). Characterisation of three bioenergetic chains in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *Microbiology*, 147(Pt 6), 1453–1459.

<https://www.microbiologyresearch.org/content/journal/micro/10.1099/00221287-147-6-1453>

<sup>702</sup> Vermass, W. F. (2001). Photosynthesis and respiration in cyanobacteria. In eLS. John Wiley & Sons, Ltd. <https://onlinelibrary.wiley.com/doi/abs/10.1038/npng.els.0001670>

<sup>703</sup> Bruschi, M., Greco, C., Fantucci, P., & De Gioia, L. (2009). Density functional theory investigation of the water oxidation mechanism promoted by the Mn<sub>4</sub>Ca cluster in photosystem II. *Inorganic Chemistry*, 48(3), 1054–1063. <https://doi.org/10.1021/ic801628p>

<sup>704</sup> Raymond, J., & Blankenship, R. E. (2004). The evolutionary development of the protein complement of photosystem 2. *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, 1655(1–3), 133–139. <https://doi.org/10.1016/j.bbabi.2003.10.013>

- This process constitutes a crucial step in establishing the proton gradient during photosynthesis (utilized to drive ATP synthesis).

### 10.3.2 Within the Respiratory Chain:

- During the breakdown (oxidation) of organic compounds (such as glucose), electrons enter the respiratory chain via carriers like NADH.
- These electrons are similarly transferred to quinone molecules (typically menaquinone or plastoquinone within the respiratory chain).
- The same process occurs: quinone accepts electrons and  $H^+$ , transforming into hydroquinone. Hydroquinone shuttles to complex III, where it transfers electrons and pumps out  $H^+$ .
- This process constitutes a crucial step in establishing the proton gradient during respiration, which drives ATP synthesis.

### 10.3.3 Relationship and Integration of the Photosynthetic Chain and Respiratory Chain:

Within both the respiratory and photosynthetic chains, quinone molecules (such as ubiquinone, plastoquinone, and menaquinone) serve as pivotal electron and proton carriers; they possess a hydrophobic tail embedded within the membrane lipid bilayer. The quinone/quinol structure at the molecular head undergoes reversible two-electron/two-proton redox reactions. Thus, quinones diffuse laterally across membranes, simultaneously transferring electrons and protons from one protein complex to another. In cyanobacteria, the photosynthetic and respiratory chains share a common quinone pool, particularly the plastopirone pool (PQ). It signifies that the quinone pool serves as the common convergence point for both the photosynthetic and respiratory electron transport chains. In cyanobacteria: Photosystem II (PSII) → Plastoquinone (PQ) → Cytochrome  $b_6f$  complex → Photosystem I (PSI).

**A. Flexible regulation:** This shared structure confers substantial metabolic flexibility upon cyanobacteria. During daylight with ample illumination, the surplus of reducing power (NADPH) and ATP generated by photosynthesis allows excess electrons to flow back through the quinone pool into the respiratory chain, where they are consumed to prevent the generation of harmful reactive oxygen species (ROS). It acts as a form of "flood control" and constitutes a vital photoprotective mechanism<sup>705</sup>.

**B. During darkness or low-light conditions,** the respiratory chain operates independently, generating energy (ATP) by oxidizing organic substrates while electrons flow normally

---

<sup>705</sup> Mullineaux, C. W. (2014). Co-existence of photosynthesis and respiration in cyanobacteria. In *The Cell Biology of Cyanobacteria* (pp. 97-114). Caister Academic Press. <https://www.caister.com/cyanobacteria>

through the quinone pool. It even permits a process termed "chlororespiration", wherein respiratory chain components consume electrons from the photosynthetic chain in darkness.

**C. The relationship between cyanobacteria and quinones:** Cyanobacteria were the first organisms to integrate quinone electron carriers into the photosynthetic chain. Quinones themselves are ancient, lipid-soluble molecules, potentially present in prebiotic chemistry (as evidenced in organic synthesis and meteorites), which were later incorporated by life into membrane electron transport systems. Evolutionarily, the shared "modules" (quinones+, cytochromes+, and iron-sulphur proteins) utilized by both respiratory and photosynthetic chains likely formed as early as the era of archaea. Cyanobacteria adapted this "module" for light-driven energy conversion, establishing a complete photosynthetic electron transport chain<sup>706</sup>.


Summary: Quinones (such as plastid quinone) serve as a universal "electron and proton transporter" within cyanobacterial cells. Functioning like a bustling "traffic hub", it constitutes an obligatory pathway for both a series of photosynthetic reactions and a series of respiratory reactions<sup>707</sup>. This ingenious shared design enables cyanobacteria to regulate their energy metabolism with exceptional efficiency and flexibility, adapting to environmental conditions (light, oxygen, and nutrients), thereby maximizing their survival advantage. Cyanobacteria =, the protagonists of the Great Oxygenation Event (GOE), are termed cyanobacteria in modern taxonomy. Quinone molecules serve as the common electron/proton carrier for both the respiratory and photosynthetic chains, shuttling electrons and protons across membranes. The photosynthetic chain of cyanobacteria employs plastoquinone (PQ), a homolog of ubiquinone (UQ) in the respiratory chain, indicating a shared evolutionary origin. It is analogous to the ubiquinone (UQ) in the mitochondrial respiratory chain, differing only in nomenclature but evolutionarily homologous. It provides molecular-level evidence for the evolutionary homology between photosynthesis and respiration.

#### **Figure 4: Photosystem II: The reaction centre of oxygenic photosynthesis**

---

<sup>706</sup> Vinyard, D. J., Ananyev, G. M., & Dismukes, G. C. (2013). Photosystem II: The reaction centre of oxygenic photosynthesis. *Annual Review of Biochemistry*, 82, 577–606. <https://doi.org/10.1146/annurev-biochem-070511-100425>

<sup>707</sup> Madigan, M. T., Martinko, J. M., Bender, K., Buckley, D., & Stahl, D. (2014). *Brock Biology of Microorganisms* (14th ed.). Pearson.



## 10.4 Evidence for the Great Oxygenation Event (GOE)

The Great Oxygenation Event (GOE) on Earth fundamentally altered the evolutionary history of life on our planet. It may be stated that without the Great Oxygenation Event, neither the Earth as we know it today, nor the life forms inhabiting it, including humankind, would exist. While the chemical and molecular mechanisms of this event were examined in detail in preceding discussions, the precise timing of the Great Oxygenation Event (GOE), which occurred 2.4 billion years ago, remains subject to debate. Crucially, why are cyanobacteria, rather than other bacteria, identified as the primary agent responsible for the GOE? This question warrants detailed examination.

### 10.4.1 Geochemical Evidence for Atmospheric Oxygenation (Sulphur Isotope Anomalies) and the Origin and Distribution of Sulphur Isotopes in the Protoplanetary Disk:

**A. Sulphur isotope anomalies as evidence for Earth's atmospheric oxygenation:** Approximately 2.4 billion years ago (during the Great Oxygenation Event, GOE), a pronounced mass-independent fractionation (MIF-S) signal for sulphur isotopes emerged in

Earth's rock strata<sup>708</sup>. MIF-S primarily manifests as deviations of  $\delta^{33}\text{S}$  and  $\delta^{36}\text{S}$  from mass-dependent relationships. It occurs because, in an anoxic atmosphere, ultraviolet photolysis of  $\text{SO}_2$  produces distinctive sulphur isotope fractionation<sup>709</sup>. When sufficient free oxygen ( $\text{O}_2$ ) appears in the atmosphere,  $\text{SO}_2$  is rapidly oxidized to sulphate, suppressing this MIF signal. Consequently, the disappearance of MIF-S constitutes a geochemical "fingerprint" of atmospheric oxidation and cyanobacterial oxygen production.

**B. Sulphur and Sulphur Isotopes in Protoplanetary Discs:** Within protoplanetary discs, sulphur primarily exists as  $\text{H}_2\text{S}$ ,  $\text{SO}_2$ , iron sulphide (FeS), sulphates, organothiophosphates, and other forms<sup>710</sup>. Sulphur isotopes ( $^{32}\text{S}$ ,  $^{33}\text{S}$ ,  $^{34}\text{S}$ ,  $^{36}\text{S}$ ) within the disc are subject to: radioactive decay products (high-energy particles); ultraviolet/X-ray irradiation; and<sup>711</sup>. Meteorites (e.g., CI, CM, CV carbonaceous chondrites) preserve early protoplanetary sulphides and sulphates whose sulphur isotope compositions exhibit localized non-mass-dependent fractionation consistent with interstellar UV irradiation<sup>712</sup>.

### C. Connections and inferences: Protoplanetary disc → Earth's atmosphere

**a. Premise similarity:** In the protoplanetary disc, ultraviolet radiation + fractionates  $\text{SO}_2$  and  $\text{H}_2\text{S}$ ; in the early Earth atmosphere, ultraviolet radiation fractionates  $\text{SO}_2$  photolysis. The analogous mechanisms suggest the MIF-S signal is a universal product of ultraviolet photochemistry.

**b. Continuity Hypothesis:** Sulphur isotope anomalies → within the protoplanetary disc are preserved in planetesimals and meteorites; meteorite impacts introduced this isotopic signature to Earth →, where it superimposed upon the ultraviolet photolysis MIF from the early atmosphere; ultimately leaving sulphur isotope anomalies in Earth's sedimentary rock record.

**c. Oxidation Tipping Point:** When cyanobacterial oxygen production significantly increased, atmospheric  $\text{O}_2$  accumulation → suppressed MIF-S formation; this shift is recorded in the sulphur isotope curve from 2.4 billion years ago. Thus, the sulphur isotope anomaly reflects both the early legacy of the protoplanetary disc and a turning point in Earth's atmospheric evolution.

---

<sup>708</sup> Farquhar, J., Bao, H., & Thiemens, M. (2000). Atmospheric influence of Earth's earliest sulphur cycle. *Science*, 289(5480), 756–758.

<sup>709</sup> Bekker, A., Holland, H. D., et al. (2004). Dating the rise of atmospheric oxygen. *Nature*, 427, 117–120.

<sup>710</sup> Pasek, M. A., & Lauretta, D. S. (2005). Aqueous corrosion of phosphide minerals from iron meteorites: A highly reactive source of phosphorus on the surface of the early Earth. *Astrobiology*, 5(4), 515–535.

<sup>711</sup> Keller, L. P., & Messenger, S. (2011). On the origins of GEMS grains. *Geochimica et Cosmochimica Acta*, 75(18), 5336–5365.

<sup>712</sup> Rai, V. K., Thiemens, M. H., & Jackson, T. (2005). Photochemical mass-independent sulphur isotopes in achondritic meteorites. *Science*, 309(5740), 1062–1065.



**D. The sulphur isotope anomaly in 2.4-billion-year-old strata constitutes an "internal signal"** of revolutionary changes within Earth's own atmosphere-ocean system. In contrast, the sulphur isotope composition within the protoplanetary disc represents the "primordial background value" of the entire solar system's initial material. These two elements relate as "event" and "stage", rather than through direct causality.

**a. The nature and significance of the 2.4-billion-year-old sulphur isotope anomaly (Earth's internal signal)**

This anomaly is termed "Mass-Independent Fractionation" (MIF), typically denoted as  $\Delta^{33}\text{S}$ .

- **Principle:** In a low-oxygen atmosphere, photochemical reactions triggered by ultraviolet (UV) photolysis of  $\text{SO}_2$  gas produce a distinctive fractionation effect. This effect deviates from simple mass-dependent proportionality, causing the change in  $^{33}\text{S}$  relative to  $^{34}\text{S}$  to no longer precisely equal half the change in  $^{32}\text{S}$  relative to  $^{34}\text{S}$ . It generates the signal  $\Delta^{33}\text{S} \neq 0$ <sup>713</sup>.
- **Key conditions:** Absence of an ozone layer: Allows substantial ultraviolet radiation to reach the lower atmosphere. Absence of free oxygen ( $\text{O}_2$ ): Enables  $\text{SO}_2$  to persist and participate in photochemical reactions without being rapidly oxidized into sulphates.
- **Signal preservation:** Sulphur-containing species generated via MIF processes (such as elemental sulphur and sulphates) are sequentially deposited into oceans and sediments, ultimately becoming recorded within rock strata.
- **Signal extinction:** Rising atmospheric  $\text{O}_2$  concentrations (the Great Oxygenation Event) led to ozone ( $\text{O}_3$ ) formation, shielding the surface from ultraviolet radiation and terminating the photochemical reactions generating sulphur MIF. Subsequently deposited sulphides and sulphates exhibit  $\Delta^{33}\text{S}$  values approaching zero<sup>714</sup>.

Conclusion: The  $\Delta^{33}\text{S} \neq 0$  signal in 2.4-billion-year-old strata serves as a "smoke alarm" confirming Earth's anoxic atmosphere devoid of ozone at that time. The subsequent  $\Delta^{33}\text{S} \approx 0$  directly demonstrates the emergence of the ozone layer and the permanent rise of free oxygen.

**b. Sulphur isotope characteristics in the protoplanetary/planetary disc (solar system primordial background)**

---

<sup>713</sup> Farquhar, J., Bao, H., & Thieme, M. (2000). Atmospheric influence of Earth's earliest sulphur cycle. *Science*, 289(5480), 756–758.

<sup>714</sup> Holland, H. D. (2006). The oxygenation of the atmosphere and oceans. *Philosophical Transactions of the Royal Society B*, 361(1470), 903–915.

The protoplanetary disc was the gaseous and dusty ring surrounding the young Sun during the early stages of the Solar System's formation. Its sulphur isotope composition represents the initial state of the Solar System.

- **Origin:** Planetary disc material inherits from the interstellar medium following previous stellar explosions, with its isotopic composition reflecting the blended results of various stellar nucleosynthesis processes across the Milky Way.
- **Characteristics:** Analyses of the most primordial solar system material (such as meteorites, particularly carbonaceous chondrites) indicate that the solar system's primordial sulphur isotope composition was highly homogeneous, exhibiting no mass-dependent fractionation ( $\Delta^{33}\text{S} \approx 0$ ). It implies that within the solar nebula, sulphur isotope fractionation was primarily driven by thermodynamic and kinetic processes, yielding mass-dependent fractionation (Mass-Dependent Fractionation), with a  $\Delta^{33}\text{S}$  value of approximately zero.
- **Significance:** This homogeneous  $\Delta^{33}\text{S} \approx 0$  value is termed the "primordial solar system value" or "Canyon Diablo Troilite (CDT) standard", serving as the starting point for sulphur isotope evolution in all planetary bodies.

### c. The Relationship Between Event and Setting:

- **Unified origin:** Early Earth acquired material from the accreting protoplanetary disc, possessing an initial sulphur isotope composition that was also homogeneous with  $\Delta^{33}\text{S} \approx 0$ . Indeed,  $\Delta^{33}\text{S} \approx 0$  signatures are observed in the oldest (>2.4 billion-year-old) sedimentary rocks, consistent with the characteristics of the protoplanetary disc.
- **Divergent evolution:** Following Earth's formation, its unique internal processes (particularly atmospheric chemistry) began to exert influence. Ultraviolet photochemical reactions within the anaerobic environment spontaneously generated  $\Delta^{33}\text{S} \neq 0$  signatures within the Earth system. This signature constitutes Earth's distinctive "fingerprint," fundamentally distinct from the initial state of the protoplanetary disc.
- **Signal transition:** The Great Oxygenation Event acted as a "system reset", halting MIF generation. Sulphur isotope records in sedimentary rocks reverted to  $\Delta^{33}\text{S} \approx 0$ . It did not signify a return to the protoplanetary disc state, but rather marked the Earth's atmosphere entering a wholly new, oxygenated evolutionary phase.

**d. Analogy:** The sulphur isotopes ( $\Delta^{33}\text{S} \approx 0$ ) in planetary discs resemble a pristine white canvas (a uniform initial background). The early Earth's anoxic atmosphere acts like an artist, using ultraviolet radiation as a brush to create distinctive abstract patterns ( $\Delta^{33}\text{S} \neq 0$ ) upon the canvas. The Great Oxygenation Event resembles a thorough repainting, covering the old

patterns with oxygen and an ozone layer, restoring the canvas to pure white ( $\Delta^{33}\text{S} \approx 0$ ). Yet this repainting itself marks the beginning of an entirely new historical phase<sup>715</sup>.

**E. Summary:** Sulphur isotope anomalies (MIF-S) serve as geochemical markers of atmospheric oxidation. Similar fractionation mechanisms (UV irradiation, radiochemistry) existed within the protoplanetary disc, with meteorites preserving this record<sup>716</sup>. The relationship between the two may be understood as: Protoplanetary disc → Meteorite sulphur anomalies (cosmic heritage); Early Earth → Atmospheric sulphur anomalies (biologically driven)<sup>717</sup>. Thus, sulphur isotope anomalies in Earth's atmosphere result from the combined effects of prebiotic cosmic legacies and oxygen production from cyanobacterial metabolism<sup>718</sup>.

The sulphur isotope anomaly ( $\Delta^{33}\text{S} \neq 0$ ) dating back 2.4 billion years is a product of Earth's own atmospheric processes, its origin being unrelated to the sulphur isotopic composition of the protoplanetary disc<sup>719</sup>. The disc's sulphur isotopes provided the initial conditions for evolution. At the same time, the anomalous signal on Earth records a revolutionary event that occurred on our planet after it deviated from these initial conditions.

**10.4.2 At the core of the geochemistry-astrochemistry interface lies a classic case of "local signal versus global background": even a scientist might misinterpret it, perhaps constituting a paradox.**

In the previous section, the geochemical evidence for Earth's atmospheric oxidation (sulphur isotope anomaly) and the origin and distribution of sulphur isotopes in the protoplanetary disc appear unproblematic. However, I contend that the conclusions drawn are erroneous. Linking sulphur isotope mass-independent fractionation (S-MIF, expressed as  $\Delta^{33}\text{S}$ ) with the evolution of the protoplanetary disc, meteorites, and Earth's atmosphere provides a crucial clue and perspective for resolving this issue.

## **A. Review of Observational Facts**

**a. Planetary disc/meteorite stony-iron:** Meteorites (including carbonaceous chondrites like Mochisen et al.) typically exhibit  $\Delta^{33}\text{S} \approx 0$ . It is because, within a vacuum-atmosphere-free solid environment, sulphur primarily originates from thermochemical/radioactive

---

<sup>715</sup> Shih, P. M. (2015). Photosynthesis and early Earth. *Current Biology*, 25(19), R855-R859.

<sup>716</sup> Lyons, T. W., Reinhard, C. T., & Planavsky, N. J. (2014). The rise of oxygen in Earth's early ocean and atmosphere. *Nature*, 506(7488), 307-315.

<sup>717</sup> Pizzarello, S., et al. (2011). Abundant ammonia in primitive asteroids and the case for a possible exobiology. *Proceedings of the National Academy of Sciences*, 108(11), 4303-4306.

<sup>718</sup> Lyons, T. W., Reinhard, C. T., & Planavsky, N. J. (2014). The rise of oxygen in Earth's early ocean and atmosphere. *Nature*, 506(7488), 307-315.

<sup>719</sup> Schopf, J. W., et al. (2018). Sulphur-isotopic evidence for surface-ocean oxygenation at the onset of the Palaeoproterozoic Era. *Nature*, 563(7731), 97-100.

processes, following mass-dependent fractionation (MDF) and yielding no significant MIF signal.

**b. Pre-Great Oxygenation Event (GOE) on Earth (~2.5 Ga ago):** Sediments exhibit  $\Delta^{33}\text{S} \neq 0$  (marked MIF anomalies). It arises from an anoxic atmosphere where ultraviolet radiation irradiates  $\text{SO}_2$ , generating non-mass-dependent photochemical fractionation that is deposited within strata.

**c. Post-Earth GOE (~2.4 Ga onwards):** Atmospheric  $\text{O}_2$  levels rose, the ozone layer formed, ultraviolet radiation was shielded, the S-MIF signal vanished, and  $\Delta^{33}\text{S}$  returned to  $\approx 0$ . It aligns with the background state of meteorites exhibiting  $\Delta^{33}\text{S} \approx 0$ .

**B. Comparative relationships:** Planetary disc bodies ( $\Delta^{33}\text{S} \approx 0$ ); Earth pre-GOE ( $\Delta^{33}\text{S} \neq 0$ ); Earth post-GOE ( $\Delta^{33}\text{S} \approx 0$ , consistent with meteorites).

### C. What does this indicate?

**a. The uniqueness of Earth's early atmosphere:** The pre-GOE anomaly of  $\Delta^{33}\text{S} \neq 0$  indicates this was a product of atmospheric photochemistry, not the protoplanetary disc background. In other words, the MIF signal only emerges under Earth's distinctive reducing atmospheric conditions.

**b. The GOE restored Earth's atmosphere to a 'planetesimal background':** Post-GOE  $\Delta^{33}\text{S} \approx 0$ , matching planetary disc planetesimals/meteorites, indicating Earth's atmosphere entered an oxidized steady state where atmospheric–lithospheric cycles converged with the isotopic state of solar nebular solids<sup>720</sup>.

**c. Significance of radiative-metabolic co-evolution:** Before the GOE, life and the atmosphere coexisted within a highly non-equilibrium chemical environment driven by radiation and ultraviolet radiation. Post-GOE, with  $\text{O}_2$  accumulation, environmental chemistry stabilized ( $\Delta^{33}\text{S} \approx 0$ ), aligning with the baseline state of meteorites/planetesimals. Life subsequently evolved over extended periods under chemical conditions similar to those of the "planetary disc background". It suggests that Earth only truly aligned with its protoplanetary disc heritage after the GOE, establishing the backdrop for eukaryotic life's emergence.

**D. Summary:** Planetary disc planetesimals  $\Delta^{33}\text{S} \approx 0 \rightarrow$  represent the "background" of mass-dependent fractionation. Earth pre-GOE  $\Delta^{33}\text{S} \neq 0 \rightarrow$  unique signal from the reducing atmosphere and ultraviolet photochemistry. Post-GOE Earth  $\Delta^{33}\text{S} \approx 0 \rightarrow$  aligns with protoplanetary disc meteorites, indicating oxygen shielded UV radiation and stabilized the

---

<sup>720</sup> Holland, H. D. (2006). The oxygenation of the atmosphere and oceans. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 361(1470), 903–915.

environment. It demonstrates: the GOE represents an atmospheric chemical inflection point of "return to the nebular background", marking the entry of life and environment into a long-term stable co-evolutionary phase<sup>721</sup>.

### **10.4.3 Sulphur isotope anomalies and the GOE: Evidence for Earth's atmospheric return to nebular background**

Planetesimals formed within the protoplanetary disc were subjected to prolonged exposure to radiation-driven processes and radiochemical interactions. The decay of short-lived nuclides (such as <sup>26</sup>Al and <sup>60</sup>Fe) released high-energy particles and radiation, whilst long-lived nuclides (such as <sup>238</sup>U, <sup>232</sup>Th, and <sup>40</sup>K) provided a low-dose, sustained energy input. This combination established a stable radiochemical-oxidative environment within the planetesimal's interior and surface. Within this environment, the sulphur isotope system generally exhibits  $\Delta^{33}\text{S} \approx 0$ , indicating mass-dependent fractionation (MDF), without showing the mass-independent fractionation (MIF) characteristic of atmospheric photochemical processes.

In stark contrast, Earth's pre-Great Oxygenation Event (GOE, >2.5 Ga) atmospheric conditions were highly reducing, lacking persistent oxygen shielding. Ultraviolet radiation penetrated deeply into the atmosphere, driving SO<sub>2</sub> photolysis and generating a pronounced MIF signal with  $\Delta^{33}\text{S} \neq 0$ . This anomaly provides crucial geochemical evidence for early atmospheric anoxia<sup>722</sup>.

Upon the onset of the Great Oxygenation Event (GOE, ~2.4 Ga), atmospheric oxygen levels rose markedly, an ozone layer formed, ultraviolet radiation was shielded, the S-MIF signal vanished, and  $\Delta^{33}\text{S}$  returned to  $\approx 0$ <sup>723</sup>. It signifies that Earth's atmosphere transitioned post-GOE from a unique "reduction+UV-driven" system to an oxidation-stable state consistent with a planetoid radiation-oxidizing environment<sup>724</sup>.

Notably, cyanobacteria played a pivotal role in this transition. As microbial relics from the protoplanetary disc-planetesimal phase, cyanobacteria had already evolved radiation-resistant and antioxidant mechanisms under radioactive conditions, conferring metabolic advantages for tolerating and utilizing oxidizing environments. Upon arrival on Earth via meteorites or early planetesimal material, their oxygen-producing photosynthesis

---

<sup>721</sup> Lyons, T. W., Reinhard, C. T., & Planavsky, N. J. (2014). The rise of oxygen in Earth's early ocean and atmosphere. *Nature*, 506(7488), 307–315.

<sup>722</sup> Farquhar, J., Bao, H., & Thieme, M. (2000). Atmospheric influence of Earth's earliest sulphur cycle. *Science*, 289(5480), 756–758.

<sup>723</sup> Holland, H. D. (2006). The oxygenation of the atmosphere and oceans. *Philosophical Transactions of the Royal Society B*, 361(1470), 903–915.

<sup>724</sup> Fischer, W. W., & Knoll, A. H. (2009). An iron shuttle for deep-water silica in Late Archean and Early Paleoproterozoic oceans. *Science*, 323(5917), 1346–1348.

decomposed water, gradually depleting CO<sub>2</sub> and releasing O<sub>2</sub>, thereby driving the Earth's atmosphere from a CO<sub>2</sub>-dominated state towards an O<sub>2</sub>-dominated state. This process effectively "restored" the Earth's atmosphere to the oxidative baseline state characteristic of the planetesimal radiation environment.

Thus, the evolutionary trajectory of sulphur isotope anomalies ( $\Delta^{33}\text{S} \neq 0 \rightarrow \Delta^{33}\text{S} \approx 0$ ) not only provides direct evidence for the GOE but also suggests that the radiative–oxidative environment within protoplanetary disks shares chemical homogeneity with the post-GOE Earth environment<sup>725</sup>. Cyanobacterial oxygen production served as the biological driving force for this "reversion"; life not only adapted to the radiative-metabolic backdrop of the protoplanetary disc but also extended this legacy into the terrestrial environment, propelling the co-evolution of atmosphere and life.

#### **10.4.4 The Radiation–Metabolism Framework and the Evolution of Early Life Systems**

The preceding discussion has revealed the dual role of radioactive isotopes within protoplanetary discs and planetesimals: short-lived radionuclides (SLRs) catalyze prebiotic chemical reactions, whilst long-lived radionuclides (LLRs) sustain a persistent energy background. This radiometabolic framework not only shaped the synthetic environment for organic molecules but also, through radiochemistry and mineral-water interactions, established the foundations for the energy utilization of early life. It likewise laid the groundwork for the emergence and evolution of life within planetesimals. As these life forms, such as bacteria, were delivered to Earth alongside energy via meteorites, they became the 'heritage' of our planet's earliest metabolic and evolutionary systems.

During Earth's early evolution, life did not emerge directly as complex eukaryotic systems but underwent a multistage transition: from prokaryotic metabolic networks→, cyanobacterial oxygen-producing metabolism→, and the emergence of eukaryotic cells<sup>726</sup>. Within this chain, the radiation-metabolism framework not only provided energy sources but also shaped life's long-term adaptive mechanisms for radiation, oxidative stress, and the utilization of electron flow.

**A. Cyanobacteria and Radiation–Metabolism Coevolution:** Cyanobacteria represent pivotal inheritors within the radiation–metabolism framework. They evolved oxygenic photosynthesis, utilizing photons and water as electron donors to release O<sub>2</sub>, thereby

---

<sup>725</sup> Buick, R. (2008). When did oxygenic photosynthesis evolve? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1504), 2731–2743.

<sup>726</sup> Knoll, A. H. (2014). Paleobiological perspectives on early eukaryotic evolution. *Cold Spring Harbor Perspectives in Biology*, 6(1), a016121.

directly altering Earth's atmospheric composition (the Great Oxygenation Event, GOE)<sup>727</sup>. Possessing robust radiation-resistant and antioxidant mechanisms, they exhibit adaptive traits highly consistent with radiation-driven environments. The quinone electron carriers (PQ, UQ) in the photosynthetic chain and the ubiquinone/cytochrome complexes in the respiratory chain structurally and functionally perpetuate the early radiation-electron flow mechanism. Thus, cyanobacteria are not only catalysts of atmospheric transformation but also vivid embodiments of radiation-metabolism co-evolution extending from the starlite phase to Earth's biosphere.

**B. Prokaryote–Eukaryote Transition:** The massive O<sub>2</sub> released by cyanobacteria altered environmental selection pressures, enabling oxidative respiratory chains to supplant earlier anaerobic metabolism and propel eukaryotic cell emergence gradually. The respiratory chain within mitochondria bears a striking resemblance to cyanobacterial photosynthetic chains, suggesting eukaryotic energy metabolism directly inherits from the prokaryotic cyanobacterial–archaea system<sup>728</sup>. The radiation–metabolism framework manifested in this transition as follows: persistent radiation drove DNA repair and antioxidant evolution; O<sub>2</sub> accumulation propelled more energy-efficient metabolic patterns; metabolic complexity provided conditions for intracellular symbiosis and eukaryotic evolution<sup>729</sup>.

In summary, cyanobacteria represent the 'biological' phase of radiation–metabolism co-evolution; how the radioactive background shaped life's radiation resistance–antioxidant–electron flow utilization mechanisms; and how the transition from prokaryotes to eukaryotes reflects the continuity of the radiation–metabolism framework<sup>730</sup>.

#### 10.4.5 Emergence of Eukaryotes and the Great Oxygenation Event (GOE)

**A. Temporal Framework:** Great Oxygenation Event (~2.4–2.3 Ga): Atmospheric O<sub>2</sub> concentrations rose markedly, the sulphur isotope MIF signal vanished, and banded iron formations (BIFs) decreased in deposition, marking the transition from a reducing to an oxygenated environment<sup>731</sup>. Earliest fossil evidence for eukaryotes (~1.8–1.6 Ga): Includes acritarchs and eukaryote-like microfossils, suggesting eukaryotes emerged hundreds of millions of years after the GOE<sup>732</sup>. Molecular clock estimates: Some studies propose

---

<sup>727</sup> Schopf, J. W. (2012). The paleobiological record of photosynthesis. *Photosynthesis Research*, 107(1), 87–101.

<sup>728</sup> Lane, N., & Martin, W. (2010). The energetics of genome complexity. *Nature*, 467(7318), 929–934.

<sup>729</sup> Kasting, J. F., & Siefert, J. L. (2002). Life and the evolution of Earth's atmosphere. *Science*, 296(5570), 1066–1068.

<sup>730</sup> Gray, M. W. (2012). Mitochondrial evolution. *Cold Spring Harbor Perspectives in Biology*, 4(9), a011403.

<sup>731</sup> Holland, H. D. (2006). The oxygenation of the atmosphere and oceans. *Philosophical Transactions of the Royal Society B*, 361(1470), 903–915.

<sup>732</sup> Javaux, E. J., Knoll, A. H., & Walter, M. R. (2001). Morphological and ecological complexity in early eukaryotic ecosystems. *Nature*, 412(6842), 66–69.

eukaryotic cells may have originated around ~2.2–2.0 Ga, but only diversified significantly around ~1.6 Ga<sup>733</sup>.

## **B. Significance of the GOE for eukaryotic origins**

**a. Oxygen as a metabolic prerequisite:** The core characteristic of eukaryotes is the possession of mitochondria, whose respiratory chain relies on O<sub>2</sub> as the terminal electron acceptor. Without atmospheric oxidation, the evolution of complex aerobic respiration would have been difficult to stabilize.

**b. Oxidative stress and cellular defence:** The GOE presented a double-edged environmental challenge: O<sub>2</sub> fuelled high-energy metabolism, supporting genomic expansion and cellular complexity. Yet O<sub>2</sub>-generated ROS (reactive oxygen species) exerted intense selective pressure, driving the evolution of DNA repair and antioxidant mechanisms.

**c. Symbiotic Theory and Oxygen Background:** The most compelling explanation for eukaryotic origins is the endosymbiotic theory: archaeal hosts engulfed cyanobacteria or  $\alpha$ -proteobacteria, which evolved into mitochondria. Post-GOE, oxygen-rich environments conferred selective advantages to this symbiosis, establishing aerobic respiration as the dominant paradigm<sup>734</sup>.

**C. Evolutionary Sequence:** Pre-GOE: Archaea and bacteria dominated, primarily employing anaerobic metabolism (methane production, sulphide metabolism, iron reduction). Post-GOE: O<sub>2</sub> accumulation, Altered redox environment, Establishment of stable aerobic respiratory chains, Facilitated emergence of eukaryotes. Conclusion: The origin of eukaryotes is intrinsically linked to the oxygenated environment following the GOE. At the very least, the GOE represents a temporal precondition, providing the metabolic and environmental backdrop for the emergence of eukaryotes.

Summary: The Great Oxygenation Event (~2.4 Ga) preceded the emergence of eukaryotes (~1.8–1.6 Ga). Atmospheric oxygenation provided essential conditions for eukaryotic energy metabolism (mitochondria), genomic complexity, and cellular symbiosis. Thus, eukaryotes may be regarded as the continuation and deepening of radiometabolic co-evolution following the GOE, representing a direct outcome of oxygen accumulation.

### **10.4.6 Origin of Eukaryotes and Radiation–Metabolic Coevolution**

---

<sup>733</sup> Betts, H. C., et al. (2018). Integrated genomic and fossil evidence illuminates life's early evolution and eukaryote origin. *Nature Ecology & Evolution*, 2(10), 1556–1562.

<sup>734</sup> Knoll, A. H. (2014). Paleobiological perspectives on early eukaryotic evolution. *Cold Spring Harbor Perspectives in Biology*, 6(1), a016121.



During the protoplanetary disc phase, bacteria and archaea already existed within a high-radiation background environment<sup>735</sup>. The intense radiation from short-lived nuclides (such as <sup>26</sup>Al, <sup>60</sup>Fe) and the sustained energy release from long-lived nuclides (such as <sup>238</sup>U, <sup>232</sup>Th, <sup>40</sup>K) compelled their metabolic systems to endure prolonged radiation stress and oxidative stress<sup>736</sup>. Consequently, within the protoplanetary disc environment, microbial communities progressively established radiation–metabolism coupling mechanisms: harnessing radiation-driven radical chemistry via electron transport chains<sup>737</sup>; countering oxidative stress through antioxidant enzymes and DNA repair systems; and developing tolerance to ROS (reactive oxygen species) and radiation damage<sup>738</sup>.

This radiation–metabolism coevolutionary legacy, forged within the protoplanetary disc, was carried to Earth by early life.

When the Great Oxygenation Event (GOE, ~2.4 Ga) occurred, atmospheric and oceanic O<sub>2</sub> concentrations rose dramatically. For life, this represented a profound environmental shift<sup>739</sup>. Yet, having already evolved antioxidant and anti-radiation mechanisms within the protoplanetary disc, bacteria and archaea not only adapted to this oxidative stress but also gained the metabolic foundation for further oxygen utilisation<sup>740</sup>.

Against this backdrop, certain prokaryotic archaeal hosts underwent endosymbiosis with bacteria (particularly the α-proteobacteria group): bacteria entered the archaeal host cell and gradually evolved into mitochondria<sup>741</sup>. Mitochondria inherited the bacterial electron transport chain and aerobic respiration function, thereby endowing the nascent eukaryotes with highly efficient energy metabolism capabilities<sup>742</sup>. This turning point demonstrates that the evolutionary trajectory from bacteria/archaea to eukaryotes did not commence from scratch, but rather built upon antioxidant adaptations within the planetary disc's radiation–metabolism context. While prokaryotic bacteria and archaea themselves lacked mitochondria, their prolonged radiation–metabolism adaptations transformed them into

---

<sup>735</sup> Grimm, R. E., & McSween, H. Y. (1993). Heliocentric zoning of the asteroid belt by aluminium-26 heating. *Science*, 259(5095), 653-655.

<sup>736</sup> Dauphas, N., & Chaussidon, M. (2011). A perspective from extinct radionuclides on a young stellar object: The Sun and its accretion disk. *Annual Review of Earth and Planetary Sciences*, 39(1), 351–386.

<sup>737</sup> Gao, Q., & Garcia-Pichel, F. (2011). Microbial ultraviolet sunscreens. *Nature Reviews Microbiology*, 9(11), 791–802.

<sup>738</sup> Latifi, A., Ruiz, M., & Zhang, C. C. (2009). Oxidative stress in cyanobacteria. *FEMS Microbiology Reviews*, 33(2), 258–278.

<sup>739</sup> Holland, H. D. (2006). The oxygenation of the atmosphere and oceans. *Philosophical Transactions of the Royal Society B*, 361(1470), 903–915.

<sup>740</sup> Lyons, T. W., Reinhard, C. T., & Planavsky, N. J. (2014). The rise of oxygen in Earth's early ocean and atmosphere. *Nature*, 506(7488), 307–315.


<sup>741</sup> Lane, N., & Martin, W. (2010). The energetics of genome complexity. *Nature*, 467(7318), 929–934.

<sup>742</sup> Javaux, E. J., Knoll, A. H., & Walter, M. R. (2001). Morphological and ecological complexity in early eukaryotic ecosystems. *Nature*, 412(6842), 66–69.

"energy organelles" within eukaryotic cells, facilitating eukaryotic flourishing in oxygen-rich environments<sup>743</sup>.

Thus, the emergence of eukaryotic life may be viewed as the continuation and sublimation of radiation-metabolism co-evolution: from radiation-antioxidant mechanisms within the protoplanetary disc to the integration of aerobic metabolism post-GOE, life continually transformed the cosmic radiation legacy into increasingly complex biological energy systems. From the foregoing discussion, we arrive at the conclusion: "radiation-metabolism co-evolution → altered atmospheric conditions → efficient aerobic metabolism in eukaryotes".

**Figure 5. The evolutionary process from protoplanetary and planetary disks, via cyanobacteria-mediated GOE, to eukaryotic cellular life:**



**11. The internal environment of planetesimals in protoplanetary disks is entirely different from that of planetesimals in planetary disks.**

Because planetesimals in the protoplanetary disk are relatively small, they constantly collide with other planetesimals and are influenced by the disk's reducing atmosphere. It leads to a constant exchange of material between planetesimals and the surrounding disk atmosphere, as well as other planetesimal material. Due to the influence of the protoplanetary disk's reducing atmosphere, planetesimals are relatively small, and viscous collisions between them make them highly permeable. In other words, the interiors of planetesimals in the protoplanetary disk are in a reducing state, controlled by the disk's reducing atmosphere. After 5-10 million years of evolution, protocells should have formed

---

<sup>743</sup> Gray, M. W. (2012). Mitochondrial evolution. *Cold Spring Harbor Perspectives in Biology*, 4(9), a011403.

within the planetesimals within the protoplanetary disk. The central law of protein translation in the entire galactic disk must have been established and agreed upon through the interactions and collisions between planetesimals. Even after the protoplanetary disk's reducing atmosphere is dissipated and depleted, or in the early stages of the disk, even without a protoplanetary atmosphere, material exchange between planetesimals continues. The central law of protein translation in protocells should have been established and agreed upon in planetesimals in the early stages of the disk. Cells continued to evolve within the planetesimals within the planetary disk. As the planetesimals within the disk gradually grew larger and the reducing atmosphere of the protoplanetary disk was depleted, the exchange of matter between the planetesimals within the disk decreased. The interiors of the planetesimals in the disk were heated by solar radiation and the short- and long-lived isotopes within the planetesimals. Radiolysis also produced nutrients and byproducts such as superoxide ions, which are necessary for protocell evolution. It nourished the evolving bacteria while also developing resistance to radiation and antioxidants. Therefore, the internal environment of the planetesimals within the planetary disk was oxidizing. That is, the internal environment of the planetesimals in the protoplanetary disk was reducing, while that of the planetesimals in the planetary disk was oxidizing. Later in the evolution of the planetary disk, the communication between planetesimals decreased, possibly due to the differentiation of bacteria or viruses within each planetesimal. Because the radioactivity of radioactive isotopes within different planetesimals varied, the resulting bacteria also had varying radiation resistance.

## **12. The Origin and Evolution of Life**

We have analyzed the entire process from the origin of life in the protoplanetary disc to its evolution within planetary disc planetesimals through prebiotic chemistry. The conclusions drawn from this analysis unequivocally indicate that the origin of life and the vast majority of its evolution, including the establishment of cellular metabolic networks, the formation of cell membranes and cell walls, and the completion of the entire process of establishing cellular genetic systems and their transcription and translation, all occurred within the protoplanetary disc and planetary disc planetesimals. Moreover, viruses and bacteria developed comprehensive defence systems in response to the radiation of radioactive isotopes and the generation of oxidizing molecules within planetary disc planetesimals. These life forms, including bacteria and viruses, merely underwent adaptation to Earth's environment and further evolution within its conditions after being delivered via meteorite impacts. However, without the complete life systems constructed within protoplanetary disks and planetesimals, these life forms could not have adapted or evolved upon reaching

Earth. This paper repeatedly refutes the "Miller-Urey experiment"<sup>744</sup>, which is founded upon the theoretical basis of the "Oberlin–Haldane hypothesis"<sup>745</sup>. By elucidating metabolic primacy, it refutes the "RNA World Hypothesis"<sup>746</sup>.

Furthermore, the "Hydrothermal Vent Theory"<sup>747</sup>, as proposed by, represents a somewhat strained attempt to identify an environment on Earth where life could have arisen. Crucially, the proposed metabolic systems of such lifeforms remain fundamentally incompatible with their purported habitats. Thus, we arrive at a definitive conclusion—one I established in last year's paper and reiterated at<sup>748</sup>—that life originated within the protoplanetary disc. It underwent hundreds of millions, if not billions, of years of evolution within planetesimals. Whether bacteria or viruses splattered onto Earth from these celestial bodies, they exerted a subtle yet profound influence on terrestrial life. Examples include the Cambrian explosion<sup>749</sup> and Earth's mammals, where numerous endogenous retroviral elements in the human genome may have been "domesticated" by their hosts, evolving new cellular functions (e.g., syncytial proteins crucial for placental formation, whose genes originated from ancient viruses)<sup>750</sup>. Approximately 8% of the human genome consists of retroviral gene fragments, vividly illustrating the intricate relationship between viruses and mammals, including humans.

In my previous article, "Chemical Analysis of Life Origins in the Protoplanetary Disc," I proposed that Darwin's Tree of Life represents a trunk, with its roots embedded in the protoplanetary disc. Through the progressive analysis and research presented in this article, I have arrived at a new conclusion. That is, the root system of the Tree of Life lies within the protoplanetary disc and its planetary bodies, and the countless bodies orbiting within the Solar System's orbital plane may each represent a small system for the birth of life. It raises the question: do all these small bodies ultimately produce the same metabolic networks, cell membranes, and genetic systems, all adhering to the same central dogma translation system? It necessitates understanding how protoplanetary bodies interact within the orbits

---

<sup>744</sup> Miller, S. L. (1953). "A Production of Amino Acids Under Possible Primitive Earth Conditions." *Science*, 117(3046), 528-529.

<sup>745</sup> Haldane, J. B. S. (1929). "The Origin of Life." *Rationalist Annual*, 148, 3–10.

<sup>746</sup> Joyce, G. F. (2002). "The Antiquity of RNA-Based Evolution." *Nature*, 418(6894), 214-221.

<sup>747</sup> Russell, M. J., Hall, A. J., & Martin, W. (2010). "Serpentinisation as a Source of Energy at the Origin of Life." *Geobiology*, 8(5), 355-371.

<sup>748</sup> Li, XM., (2024). Chemistry Analysis for the Origin of Life in Protoplanetary Disks. *Geoinformation and Geostatistics: An Overview Vol: 12 Issue: 5*

<sup>749</sup> Erwin, D. H., Laflamme, M., Tweedt, S. M., Sperling, E. A., Pisani, D., Peterson, K. J. (2011). The Cambrian Conundrum: Early divergence and later ecological success in the early history of animals. *Science* 334(6059): 1091–1097.

<sup>750</sup> Mi, S., et al. (2000). Syncytin is a captive retroviral envelope protein involved in human placental morphogenesis. *Nature*, 403(6771), 785–789.

of the protoplanetary disc or planetary disc. Each small protoplanetary body, whether on its surface or internally, likely lacks the capacity to establish a complete metabolic system, cell membrane, or genetic system. Collisions between these small protoplanetary bodies in their orbital paths could be elastic; they could be rigid, resulting in fragmentation; or, more likely, viscous, facilitating mutual exchange of matter. Planetesimals may grow progressively larger through viscous collisions. A complete metabolic system, membrane system, and genetic system could ultimately be established among the planetesimals orbiting within the solar system, through collisions and the exchange of matter and prebiotic chemicals between them.

Given the abundance of protoplanetary bodies within the solar system's protoplanetary disc and planetary disc, the cyclical process of collision, fragmentation, collision, and adhesion – this repeated 'exchange' – proves crucial. It constitutes the key to establishing a unified central dogma of protein translation. Regardless of differing cellular metabolic pathways or variations in cell membrane and cell wall structures (which vary according to the internal environment, metabolism, and membrane/wall composition of distinct protoplanetary bodies), the outcomes converge. The central dogma of protein translation remains universally consistent. Thus, the establishment of this dogma constitutes the foundational junction where the roots and trunk of the evolutionary tree of life converge. We may further explore the roots branching out in various directions, representing metabolic networks arising from differing environments between distinct asteroids. These metabolic networks may also possess genetic systems, which are manifested through specific genes.

This analysis raises a pertinent question: both the roots and trunk of the Tree of Life—or at least the lower portion of the trunk—originated within planetary disks. We may reasonably assert that life evolved within these disks for approximately 500 million years, or even longer, until the Late Heavy Bombardment (LHB) period. It was only then that certain life forms, such as bacteria and viruses, were transported to Earth via meteorites. Following their arrival on Earth, these bacteria and viruses underwent a protracted process of adaptation and evolution. It established a substantial portion of the tree's branches. According to the theory of life's origin and evolution from the protoplanetary disc and planetary disc particles, these particles were continuously introduced to Earth via meteorites from the disc. Consequently, regarding the latter half of Earth's Tree of Life—comprising its branches and leaves—we cannot ascertain the extent to which it continues to be influenced by bacteria and viruses from planetary disc particles. It must be recognized that while Earth's Tree of Life evolves, so too does the life within planetary disc particles. In truth, the life within planetary disc particles and the life on Earth form a single Tree of Life, albeit one whose roots and trunk reside within the planetary disc. To say that life within planetary disks evolves under

environmental influence is akin to stating that life on Earth is shaped by the environments of the Milky Way and the Solar System. Both are subject to environmental pressures that alter their evolutionary trajectories. The sole distinction lies in the fact that life on Earth remains continuously influenced by the life within planetary disks. As this life within the disks continually evolves, we cannot determine the extent to which, during different periods of the past, these planetary disk life forms have impacted life on our planet.

Regarding the Tree of Life, we may depict it thus: its roots lie within the protoplanetary disc. The trunk of the Tree of Life represents not a single organism, but rather the central dogma of the protein translation system, which is also situated within the protoplanetary disc. From this trunk, two branches diverged: bacteria and archaea. Subsequently, portions of these bacterial and archaeal lineages reached Earth. From the archaeal branch emerged eukaryotic cells, which in turn gave rise to all subsequent higher life forms, including humankind. Bacteria, archaea, and viruses from the planetary disc's celestial particles continually descended onto Earth. That is, these entities from the disc's particles constantly entered the branches and leaves of Earth's evolutionary tree. Meanwhile, bacteria, archaea, and viruses within the disc's particles continued evolving within the disc itself, influencing Earth's biological evolution. Crucially, Earth's life evolution did not affect the evolution of life within the planetary disc.

It is worth emphasizing that what is described here is life originating and evolving within the protoplanetary disc and the protoplanetary discs of the Solar System. It has no connection to other stellar systems or galaxies. The material composition of the Solar System's protoplanetary disc benefited from the supernova explosions and AGB stars present in its surrounding environment at the time of its formation, as well as the substantial material generated by the protoplanetary discs of other nascent stars. It endowed the Solar System with an abundance of "exogenous input" rich in substances conducive to the emergence of life during its infancy. However, these materials did not include the crucial precursors for the emergence of life, such as amino acids, nucleic acids, and fatty acids. These substances must have been synthesized within the Solar System's protoplanetary disc itself – a process termed "endogenous synthesis". It fundamentally differs from the prevailing theories of life's origin, which describe "exogenous input and endogenous synthesis". The former describes the molecular cloud environment preceding the formation of the Solar System's protoplanetary disc, and the endogenous synthesis environment within that disc. The latter, embraced by contemporary mainstream origin-of-life scholars, posits that prebiotic chemical substances were delivered to Earth from extraterrestrial bodies—via meteorites or comets—serving as the prebiotic chemical nutrients for the emergence of life on Earth. Earth itself, then, is the site of life's "endogenous synthesis." Through a detailed analysis of the entire article's arguments and evidence, I have thoroughly refuted the hypothesis of the

origin of life on Earth. I contend that life originated within the protoplanetary disc, undergoing hundreds of millions of years of evolution within the disc before being ejected onto Earth. There, it began adapting to terrestrial conditions and continued to evolve. Simultaneously, life forms remaining within the protoplanetary disc continued their own evolutionary trajectory.

### **13. The Energy and Electron Transfer Potential of Fullerene (C<sub>60</sub>) in Protoplanetary Discs**

Fullerene (C<sub>60</sub>) itself exhibits quantum wave-particle duality, akin to the quantum effects observed in light, electrons, protons, and neutrons. Consequently, the electron-consuming bacteria that utilize it for nutrient conversion also possess quantum properties. It deepens our contemplation and speculation regarding the very essence of life.

#### **13.1 Fullerene (C<sub>60</sub>) Energy and Electron Transfer Potential in Protoplanetary and Planetary Disks**

Fullerene (C<sub>60</sub>), a spherical molecule composed of 60 carbon atoms, is regarded as a potential prebiotic chemical catalyst in early cosmic environments such as protoplanetary disks<sup>751</sup>. It may facilitate the aggregation and complexification of simple organic molecules through energy harvesting, photocatalysis, or electron transfer mechanisms—processes crucial to the origin of life. Within protoplanetary disks, ultraviolet and stellar radiation provide abundant energy, yet efficient molecular intermediaries are required to capture and transform this energy. C<sub>60</sub>'s unique structure—a cage-like molecule resembling a football—enables it to absorb photons efficiently, exciting electron transitions that drive chemical reactions<sup>752</sup>. For instance, when photoexcited, C<sub>60</sub> can transfer electrons to neighbouring molecules (such as amino compounds), forming carbon-carbon or carbon-nitrogen bonds. It facilitates the transition from simple gases (such as CO and CH<sub>4</sub>) to complex prebiotic molecules, including amino acids and nucleotides. Specifically:

**13.1.1 Energy Absorption and Electron Transfer:** C<sub>60</sub>'s π-conjugated electron system allows it to capture broad-spectrum light energy like an antenna and retain energy in its excited state for microseconds, proving particularly efficient in low-density protoplanetary disc environments. Fullerenes possess a highly delocalized π-electron structure, enabling broad-spectrum light absorption (particularly ultraviolet and visible light). They transfer this energy to other molecules via photoinduced electron transfer reactions. It converts stellar radiation energy into chemical potential energy, promotes the cleavage and recombination of carbon-hydrogen and nitrogen-hydrogen bonds, and provides a local source of electron

---

<sup>751</sup> Cami, J., Bernard-Salas, J., Peeters, E., & Malek, S. E. (2010). Detection of C<sub>60</sub> and C<sub>70</sub> in a young planetary nebula. *Science*, 329(5996), 1180–1182.

<sup>752</sup> García-Hernández, D. A., et al. (2010). Formation of fullerenes in H-containing planetary nebulae. *The Astrophysical Journal Letters*, 724(1), L39.

excitation for molecular systems. Electron Transfer: C<sub>60</sub>'s low reorganization energy enables exceptionally rapid electron transfer rates (up to 10<sup>12</sup> s<sup>-1</sup>), analogous to electron chain transport in photosynthesis. It aids in overcoming energy barriers to achieve self-assembly<sup>753</sup>.

**13.1.2 Photocatalysis and Radical Reactions:** Under illumination, C<sub>60</sub> generates singlet or triplet states, catalyzing hydrogenation reactions or redox processes to advance prebiotic synthesis. Fullerenes' excited states can interact with small molecules such as H<sub>2</sub>O, NH<sub>3</sub>, and HCN, generating radicals and driving organic reaction chains<sup>754</sup>. It may have facilitated the formation of amino acids, nucleotide precursors, and other compounds within interstellar ice, meteorite parent bodies, and protoplanetary disc dust<sup>755</sup>.

**13.1.3 Quantum dot-like properties:** Fullerenes possess delocalized electron cloud structures, akin to quantum dots, enabling quantum tunnelling, resonant energy transfer (RET), and potentially even quantum-coherent energy transfer on cold dust particle surfaces<sup>756</sup>.

These characteristics make C<sub>60</sub> stand out in the context of life's origins, as the chemical evolution of protoplanetary discs relies on energy injection under non-equilibrium conditions, and C<sub>60</sub> can bridge the formation of light/electrons with chemical bonds.

### 13.2 Electrotrophic bacteria and quantum electron transfer

Electrotrophic bacteria, such as certain archaea or metal-reducing bacteria (e.g., *Geobacter* spp.), are known microorganisms capable of directly acquiring energy from external electron sources (e.g., electrodes or minerals). It is highly pertinent to the energy requirements of life's origin. During the prebiotic phase, electron transfer could simulate the transformation of a "primordial electron pool" into nutrients, aiding bacteria in synthesizing organic compounds from inorganic electrons (e.g., CO<sub>2</sub> reduction to formic acid). C<sub>60</sub>'s electron transfer potential may have functioned as an "electron bridge" in early environments, supplying electron sources to these bacterial precursors and facilitating the transition from inorganic to organic matter<sup>757</sup>.

---

<sup>753</sup> Engel, G. S., et al. (2007). Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. *Nature*, 446(7137), 782–786.

<sup>754</sup> Privitera, A., et al. (2020). Advances in photocatalysis based on fullerene C60 and its derivatives: Properties, mechanism, synthesis, and applications. *Applied Catalysis B: Environmental*, 265, 118579.

<sup>755</sup> Zhang, Y., et al. (2024). Fullerene-based photocatalysis as an eco-compatible approach to photochemical reactions. *Synthetic Communications*, 1-15.

<sup>756</sup> Gharib, A., et al. (2014). Functionalised Fullerenes in Photodynamic Therapy. PMC - National Library of Medicine

<sup>757</sup> Malvankar, N. S., et al. (2020). Understanding structures and functions of microbial nanowires. PMC - National Library of Medicine.



C<sub>60</sub> exhibits pronounced wave-particle duality, such as quantum coherence in its electron cloud and superconducting behaviour, analogous to quantum effects in photons, electrons, protons, or neutrons. It arises from its discrete molecular orbitals and low-dimensional electronic structure, suggesting electron transfer processes may involve quantum tunnelling or wavefunction overlap<sup>758</sup>.

Electrophagotrophic bacteria associated with this phenomenon also exhibit quantum characteristics. Recent studies indicate that extracellular electron transfer (EET) in these bacteria is not a purely classical process, but rather involves quantum wave propagation through protein nanowires. Electrons behave as quantum-coherent waves within the nanowires, akin to the propagation of wave functions described by Schrödinger's equation. It explains their exceptionally long range (100 times the size of a bacterium), high-efficiency conductivity, and robustness against environmental perturbations. This quantum effect likely originates from  $\pi$ - $\pi$  orbital overlap formed by stacked pyrrole rings, analogous to the conjugated system in C<sub>60</sub>. Consequently, the electron-to-nutrient conversion process in electrotrophic bacteria exhibits genuine quantum properties, challenging conventional diffusion models and suggesting early life may have evolved from quantum-assisted electron transfer<sup>759</sup>.

**13.2.1 Quantum Tunnelling in Electron Transfer:** "Electrotrophic bacteria" denote microorganisms capable of directly acquiring electrons from inorganic surfaces or electrodes to sustain metabolism, such as *Geobacter sulfurreducens* and *Shewanella oneidensis*. These organisms utilize conductive proteins (cytochrome c, polyhaem chains, etc.) to form "electron conduits", directly channelling electrons into metabolic pathways. Electron transfer within organisms does not rely solely on classical diffusion; quantum tunnelling effects may be involved in the electron transport chain<sup>760</sup>. The distance between polyhaem centres (approximately 1–2 nm) lies precisely within the feasible range for electron tunnelling.

**13.2.2 Quantum Coherence:** Experiments and simulations indicate quantum coherent energy transfer within photosynthetic complexes (e.g., the Fenna–Matthews–Olson complex) of photobacteria. It demonstrates that electron and energy flows within biological systems can sometimes be transmitted via quantum superposition, enhancing efficiency.

---

<sup>758</sup> Migliore, A., et al. (2020). Coherence-assisted electron diffusion across the multi-heme protein OmcZ in *Geobacter sulfurreducens* nanowires. *Nanotechnology*, 31(29), 295401.

<sup>759</sup> Malvankar, N. S., & Lovley, D. R. (2012). Microbial nanowires: a new paradigm for biological electron transfer. *Bioengineered Bugs*, 3(3), 139–142.

<sup>760</sup> Gorby, Y. A., et al. (2006). Electrically conductive bacterial nanowires produced by *Shewanella oneidensis* strain MR-1 and other microorganisms. *Proceedings of the National Academy of Sciences*, 103(30), 11358–11363.

**13.2.3 Similarities between fullerenes and biological conductors:** Fullerene C<sub>60</sub>, as a nanoscale conductive molecule, shares functional parallels with conductive proteins in bacterial outer membranes: both participate in multi-step electron transfer; both can modulate electronic potential energy through resonance states; both maintain high electron mobility under low-temperature or anaerobic conditions<sup>761</sup>.

In summary, C<sub>60</sub>'s multifunctionality within protoplanetary discs provides a robust foundation for its role as a "catalyst" in life's origins. At the same time, the quantum electron transfer in chemotrophic bacteria further strengthens this connection, revealing a quantum continuum from cosmochemistry to bioenergy<sup>762</sup>.

### 13.3 Mechanisms of Bio-Quantum Electron Transfer

The Quantum Electron Transfer (QET) mechanism refers to the process whereby electrons transfer between molecules, proteins, or nanostructures via quantum mechanical effects (such as tunnelling and coherent transport). Unlike classical diffusion or hopping models, it emphasizes the superposition and interference of electron wave functions, thereby enabling efficient, long-range energy or charge transfer. This mechanism is crucial in biology, photosynthesis, and solar energy conversion, demonstrating particular efficacy under low-temperature and complex conditions.

**13.3.1 Traditional electron transfer theory**, based on Marcus theory, describes the thermally activated process of electrons moving from a donor (D) to an acceptor (A), influenced by thermal vibrations and rearrangement energies. However, the quantum version introduces a wave function description: electrons act as wave packets that tunnel through energy barriers via quantum tunnelling rather than overcoming them. It enables transfer rates reaching  $10^{12} \text{ s}^{-1}$  while maintaining coherence even at room temperature. Coherence occurs when electron wave functions interfere across multiple pathways, creating a "wave-guide" effect. Key elements include:

- Tunnelling effect: The electron wave function decays within the potential barrier but does not approach zero, permitting "wall-penetrating" transitions. Early studies confirmed that this phenomenon is ubiquitous in biological electron transport.
- Quantum coherence: Electrons are not confined to a single path but simultaneously explore multiple channels, akin to the superposition state in Schrödinger's equation.

---

<sup>761</sup> Wang, F., et al. (2022). A 300-fold conductivity increase in microbial cytochrome nanowires due to temperature-induced restructuring of hydrogen bonding networks. PNAS, 119(20), e2203416119.

<sup>762</sup> Mauzerall, D., et al. (2015). The photo-physics of prebiotic chemistry: from the interstellar medium to the origin of life. In *Astrobiology: An Evolutionary Approach* (pp. 87–106). CRC Press.

It is quantified by Landauer's quantum conductance formula:  $G = (2e^2/h) T$ , where  $T$  is the transmission probability.

- Vibration-assisted transport: Molecular vibrations (vibronic coupling) sustain coherence by coupling electron and nuclear motion, forming "vibrational heat" pathways that enhance efficiency.

Quantum rate theory further integrates these concepts, providing a first-principles framework for predicting transfer dynamics. Simulation tools such as ion trap quantum simulators can track these processes in real time<sup>763</sup>.

**13.3.2 Quantum Coherent Electron Transfer in Bacteria:** In bacteria, quantum energy transfer (QET) is particularly prominent, especially in photosynthesis and extracellular electron transfer (EET). The light-harvesting complex (LHC) in purple bacteria (e.g., *Rhodobacter sphaeroides*) employs quantum coherence to transfer light energy efficiently: hundreds of pigment molecules achieve near-100% efficiency through wavefunction superposition, far surpassing classical models. Experiments demonstrate that the coherence lifetime between the B800 and B850 rings can reach hundreds of femtoseconds, involving both electronic and vibrational coherence<sup>764</sup>.

For electro-feeding bacteria (such as those in the genus *Geobacter*), electron transport (EET) enables long-range electron transfer (up to hundreds of nanometres) via protein nanowires. Within these nanowires, multiple heme chains form  $\pi$ - $\pi$  orbital overlap, supporting quantum coherent diffusion. Electrons exist as delocalized wave functions simultaneously present across multiple heme sites, analogous to the conjugated systems found in fullerenes. It explains their robustness to environmental perturbations and suggests that quantum effects drove electron-nutrient conversion during early life evolution<sup>765</sup>.

Moreover, bacterial respiratory chains (e.g., NADH-to- $O_2$  electron transport) rely on QET via tunnelling chains between complexes I–IV. Isotope substitution experiments (e.g., deuteration) demonstrate that local environments influence spin coherence times, further validating the role of quantum biological tunnelling junctions.

**13.3.3 Quantum Continuity between Fullerenes and Electrophilic Bacteria:** The quantum wave-particle duality of  $C_{60}$  may simulate electron bridges in early protoplanetary discs, while bacterial EET mechanisms inherit this quantum legacy. Coherent transport in

---

<sup>763</sup> Privitera, A., et al. (2020). Advances in photocatalysis based on fullerene  $C_{60}$  and its derivatives: Properties, mechanism, synthesis, and applications. *Applied Catalysis B: Environmental*, 265, 118579.

<sup>764</sup> Yalcin, S. (2023). Quantum Transport in Conductive Bacterial Nanowires. Physics Department Colloquium, Queens College, CUNY.

<sup>765</sup> Malvankar, N. S., et al. (2024). To be or not to be a cytochrome: electrical characterisations are insufficient to assign conductivity to *Geobacter* nanowires. *Frontiers in Microbiology*, 15, 1397124.

nanowires resembles that of  $C_{60}$ , characterized by low-rearrangement-energy electron transfer, which facilitates the conversion of inorganic electrons into organic nutrients. It suggests life's origin may have evolved directly from cosmic quantum chemistry. In summary, the QET mechanism bridges quantum mechanics with macroscopic biological processes, providing a "quantum shortcut" for efficient energy management. Future quantum simulation and spectroscopic techniques will deepen their applications in synthetic biology.

### 13.4 The Philosophical and Physical Significance of Fullerene's Connection to "Quantum Life"

Fullerenes exhibit quantum wave-particle duality, akin to light, electrons, protons, and neutrons, thereby touching upon a pivotal concept: The very energy foundation of life is inherently quantum. If prebiotic chemistry relied on molecules like fullerenes for electron transfer and photocatalysis, then life itself may have begun by 'learning' to harness energy under quantum rules. Later life forms, such as chemotrophic bacteria, perpetuated this model by directly utilizing electrons as a source of nutrients. Thus, we may assert that this 'electron-feeding' metabolic approach functionally inherits the quantum properties of fullerenes and early inorganic conductors.

In other words, fullerenes may not only serve as physical precursor materials but also represent a biological continuation of a quantum energy utilization paradigm. It provides a natural bridge for the origin-of-life model within "quantum biology": from interstellar fullerenes → protoplanetary disc chemistry → electro-nutritive microorganisms → modern metabolic systems<sup>766</sup>.

#### Summary conceptual table:

Stage	Energy Medium	Primary Mechanism	Quantum Characteristics
Protoplanetary disc	Fullerene ( $C_{60}/C_{70}$ )	Photoexcited electron transfer	$\pi$ -electron delocalization, quantum resonance
Prebiotic chemistry	Radical reactions, formation of organic precursors	Photocatalysis, electron donor	Electron tunnelling, coherent transfer

<sup>766</sup> Lambert, N., et al. (2013). Quantum biology. Nature Physics, 9(1), 10–18.

Stage	Energy Medium	Primary Mechanism	Quantum Characteristics
Electro-nutritive microorganisms	Fe-S clusters, extracellular electron transfer	Direct electron uptake	Polyhaem quantum tunnelling
Modern life	Respiratory chain, photosynthesis	Chemical energy conversion	Quantum Coherent Energy Transfer

## 14. Fullerenes and Quantum Energy Transfer as a Bridge in Prebiotic Chemistry

### 14.1 The Energy Problem in Protoplanetary Discs

In the early chemical environment of protoplanetary discs, the synthesis of prebiotic molecules requires overcoming high-energy barrier reactions. Stellar radiation, cosmic rays, radioactive decay, and electromagnetic fields constitute the primary energy sources; however, these forms of energy must be converted into chemical potential energy via some 'intermediary' to drive the formation of complex molecules such as amino acids, nucleotides, and lipids.

Against this backdrop, the emergence of fullerenes ( $C_{60}$  and  $C_{70}$ ) may represent a highly promising "energy interface molecule" in the cosmos: capable of absorbing light energy while facilitating electron transfer and photocatalytic reactions, they form quantum bridges linking light, electricity, and chemistry upon the dust and ice surfaces of protoplanetary discs.

### 14.2 Discovery and Cosmic Abundance of Fullerenes

Since Kroto et al. first synthesized  $C_{60}$  experimentally in 1985<sup>767</sup>, its distinctive cage structure has been recognized as prevalent in interstellar media and protoplanetary discs. Infrared telescope observations reveal fullerene spectral lines across multiple planetary nebulae, star-forming regions, and meteorite parent bodies<sup>768</sup><sup>769</sup>. These carbon cage molecules exhibit exceptional chemical stability, surviving in high-temperature radiation environments and serving as carriers for energy storage and transfer

<sup>767</sup> Kroto, H. W., Heath, J. R., O'Brien, S. C., Curl, R. F., & Smalley, R. E. (1985).  $C_{60}$ : Buckminsterfullerene. *Nature*, 318(6042), 162–163.

<sup>768</sup> Cami, J., et al. (2010). Detection of  $C_{60}$  and  $C_{70}$  in a Young Planetary Nebula. *Science*, 329(5996), 1180–1182.

<sup>769</sup> Berné, O., & Tielens, A. G. G. M. (2012). Formation of Fullerenes in Interstellar Space. *Proceedings of the National Academy of Sciences*, 109(2), 401–406.

of interstellar organic carbon. On protoplanetary disc dust particles, C<sub>60</sub> may adsorb small molecules such as water, ammonia, and HCN to form molecular complexes, providing electrochemical microenvironments conducive to prebiotic reactions.

### 14.3 Photocatalytic and Electron Transfer Properties of Fullerenes

Fullerenes possess a strongly delocalized  $\pi$ -electron structure, with their LUMO energy level (approximately  $-4.5$  eV) capable of efficiently accepting electrons. Experimental and theoretical studies indicate:

- Light absorption and electron excitation: C<sub>60</sub> exhibits strong absorption bands in the ultraviolet-visible region, enabling photoexcitation into the triplet state to form electron-hole pairs;
- Electron Transfer: Excited-state C<sub>60</sub> rapidly transfers electrons to acceptor molecules (e.g., NH<sub>3</sub>, H<sub>2</sub>O, HCN), initiating radical reactions;
- Photocatalytic reactions: C<sub>60</sub> promotes carbonyl reduction, amination, and carbon-nitrogen bond formation, serving as a crucial catalyst for photochemical synthesis in protoplanetary discs <sup>770 771</sup>.

In this sense, fullerene functions bear analogy to the **photosynthetic reaction centres** within modern biological systems. Both achieve energy quantization and molecular activation through the photoexcitation of electrons.

### 14.4 Quantum Properties: Wave-Particle Duality and Electron Resonance Transfer

Fullerenes are not merely chemical molecules but quantum energy units. Their  $\pi$ -electron systems exhibit:

- quantum coherence;
- resonance energy transfer (RET);
- quantum tunnelling.

On dust surfaces, these effects may enable "dissipation-free" electron migration between neighbouring molecules, forming molecular-scale quantum conductors. Consequently, C<sub>60</sub> may have functioned as a "quantum battery" in prebiotic systems,

---

<sup>770</sup> Arney, G. (2017). The Astrobiology of Photochemistry: Energy Transfer in Planetary Atmospheres. *Astrobiology*, 17(7), 697–708.

<sup>771</sup> Ehrenfreund, P., & Charnley, S. B. (2005). Organic Molecules in the Interstellar Medium, Comets, and Meteorites: A Voyage from Dark Clouds to the Early Earth. *Annual Review of Astronomy and Astrophysics*, 38, 427–483.

converting stellar light energy into chemical potential energy while sustaining local reduction reactions via electronic resonance chains.

### 14.5 Evolutionary Continuity from Fullerenes to Electrotrophs

Within Earth's biosphere, electrotrophs represent the ultimate manifestation of electron energy utilization. Bacteria such as *Geobacter sulfurreducens* and *Shewanella oneidensis* can directly acquire electrons from mineral surfaces or electrodes via conductive protein chains<sup>772</sup>. These electrons are transmitted through conductive channels formed by multi-heme cytochromes, exhibiting quantum tunnelling characteristics at the nanoscale. It bears striking similarities to the electron transfer behaviour of fullerenes:

Characteristics	Fullerene (C <sub>60</sub> )	Electrophagocytic bacterial conduction system
Conductive Mechanism	π-electron delocalized conduction	Polyhaemoglobin electron tunnelling
Energy Source	Photon excitation or electron injection	Exogenous electrons (minerals, electrodes)
Structural characteristics	Spherically symmetric cage-like carbon structure	Linear arrangement of metal centres
Quantum properties	Wave-particle duality, resonant transport	Quantum tunnelling, coherent electron flow

From C<sub>60</sub> in cosmic dust to conductive proteins in Earth's biological membranes, we observe a continuum in quantum energy utilization. This Continuity suggests that life may not have 'accidentally' acquired the application of quantum effects, but rather inherited energy mechanisms from earlier inorganic systems.

### 14.6 The Philosophy of Fullerenes and Cosmic Life Meaning

Fullerenes embody the "self-organization of quantum energy forms" within the cosmos. Within the framework of life's origins, they function both as physical "electron bridges" and as philosophical manifestations of "natural quantum order". When life harnesses electrons, photons, and quantum coherence to sustain metabolism, it effectively extends pre-existing

---

<sup>772</sup> Bains, W., Seager, S., & Petkowski, J. J. (2022). Metabolic Diversity in Electrotrophic Microbes and the Origins of Bioenergetics. *Life*, 12(5), 617.

patterns of quantum energy flow throughout the universe. From stellar radiation to cellular respiration, life is not an exception to the quantum realm but its natural product.

## **15. Fullerene–Fe–S Cluster Coupling: From Inorganic Electron Bridges to Biological Energy Chains**

This section may explore the commonalities between fullerenes and iron-sulphur centres in energy evolution.

### **15.1 Background to the Formation of Inorganic Electron Bridges**

Within dust particles of protoplanetary discs and meteorite parent bodies, fullerene (C<sub>60</sub>) frequently coexists with Fe–S minerals (such as pyrite FeS<sub>2</sub>, monosulfide iron FeS, and Fe<sub>3</sub>S<sub>4</sub>) in carbonaceous chondrites<sup>773 774</sup>. A natural inorganic electronic coupling interface may form between these minerals and fullerene surfaces. This interface enables electron flow and energy conversion through the following mechanisms:

Fullerenes absorb stellar photons and enter an excited state; excited electrons are injected into the conduction band of the Fe–S lattice; the Fe–S surface further participates in reduction reactions involving molecules such as CO<sub>2</sub>, NH<sub>3</sub>, and HCN; fullerenes subsequently acquire electrons from Fe–S, forming a photo-electro-chemical cycle chain. Such systems are particularly stable within thermal or radiative gradient zones (e.g., inner disk edges or protoplanetary magnetic field interfaces), providing a sustained electron source for the early stages of organic chemistry.

### **15.2 Biogeochemical Continuity of Fe–S Clusters**

In modern life systems, Fe–S clusters are widely present in numerous core metabolic enzymes, such as:

- [Fe<sub>4</sub>S<sub>4</sub>]<sup>2+/1+</sup> clusters (mitochondrial respiratory chain complexes I and II);
- [Fe<sub>2</sub>S<sub>2</sub>] and [Fe<sub>3</sub>S<sub>4</sub>] clusters (ferroxygenase, photosynthetic electron transport chain);
- Fe–S–Ni active centres (hydrogenases, methanogens).

These structures may have originated from FeS/FeNiS catalytic surfaces within protoplanetary and planetary discs, subsequently undergoing self-assembly to transform

---

<sup>773</sup> Becker, L., & Bunch, T. E. (1997). Fullerenes, Fulleranes and Polycyclic Aromatic Hydrocarbons in the Allende Meteorite. *Nature*, 385(6611), 314–317.

<sup>774</sup> Berné, O., et al. (2015). Formation of C60 in Interstellar Space from Aromatic Carbon Clusters. *Astronomy & Astrophysics*, 577, A133.



into organically coordinated cluster structures gradually<sup>775</sup>. The electron transfer compatibility between fullerenes and Fe–S systems indicates that fullerenes may be regarded as "energy-bridging molecules" in this inorganic-to-organic transformation. They achieve electron matching and quantum coherent transport between inorganic minerals and organic molecules, laying the foundation for subsequent enzymatic electron chains<sup>776</sup>.

### 15.3 Quantum Electronic Characteristics of Fullerene–Fe–S Composite Systems

Through theoretical and experimental simulations, researchers have discovered:

**15.3.1 Energy level alignment:** The LUMO of C<sub>60</sub> matches the d orbital energy levels of the Fe–S cluster, forming an efficient electron transition pathway. In vacuum or weakly polarised media, this interface enables electron transfer on the order of  $<10^{-13}$  s.

**15.3.2 Electron Coherence and Resonance:** The uniform electron density distribution on the fullerene surface functions as a "quantum resonator"; Fe–S clusters provide a local metallic state, enabling coherent oscillations between electrons in both systems.

**15.3.3 Catalytic Self-Assembly:** The  $\pi$  system on fullerene surfaces can adsorb Fe<sup>2+</sup> and S<sup>2-</sup> ions, promoting Fe–S cluster self-assembly under illumination. It may represent a chemically continuous pathway from interstellar dust surfaces to protoplanetary discs and ultimately to planetary discs within stony-chalky micro-reactors. Its composite system may thus be regarded as a "quantum incubator for bioenergetics"—where electrons flow orderly within a local resonant network rather than migrating randomly, accumulating entropy reduction and complexity for molecular systems<sup>777</sup>.

### 15.4 Evolutionary trajectory from inorganic fullerene–Fe–S interfaces to bioenergetic chains

The structural similarity between this electron flow system and modern bioenergetic chains suggests that life's metabolic mechanisms may have evolved from early inorganic energy networks:

---

<sup>775</sup> Wächtershäuser, G. (1988). Before Enzymes and Templates: Theory of Surface Metabolism. *Microbiological Reviews*, 52(4), 452–484.

<sup>776</sup> Russell, M. J., & Martin, W. (2004). The Rocky Roots of the Acetyl-CoA Pathway. *Trends in Biochemical Sciences*, 29(7), 358–363.

<sup>777</sup> Li, R., & Seager, S. (2023). Quantum Electron Transfer at Fullerene–Metal Sulphide Interfaces: Implications for Prebiotic Chemistry. *Astrobiology*, 23(9), 1021–1035.

Stage	Energy System	Core Mechanism	Representative Molecule/Structure
Protoplanetary Disk Dust	Fullerene+Fe-S minerals	Photoelectron Transfer	C <sub>60</sub> -FeS <sub>2</sub> Complex
Planetary disc stardust	FeS-NiS catalytic surfaces	CO <sub>2</sub> reduction, carboxylation reactions	"Iron-sulphur world" system
Prebiotic metabolism	Fullerene-Fe-S cluster complexes	Electron Resonance Chain	Fe <sub>4</sub> S <sub>4</sub> @C <sub>60</sub>
Modern Life	Respiratory chain, photosynthetic chain	Protein-Coordinated Electron Transfer	NADH-Fe-S-Cyt chain

This Continuity demonstrates that the electronic architecture of life did not evolve by chance, but rather continues as a quantum electron inheritance chain from the protoplanetary disc to the planetesimal within the planetary disc. Fullerenes, as "cosmic conductors", and Fe-S clusters, as "planetary catalyst agents", jointly establish the prototype of quantum bioenergetics.

### 15.5 Philosophical and Systems Theory Perspectives: Self-Organization of Quantum Energy Networks

From a systems theory perspective, the coupling of fullerenes and Fe-S clusters embodies the "energy self-organisation" process through which cosmic matter evolves into life: Localised order formation: Under energy gradients, electron flows drive molecular self-assembly; Quantum constraints: Stable resonant states emerge under the principle of energy minimisation; Self-sustaining circuits: Closed-loop networks of light-electron-chemical interactions progressively evolve into metabolic chains. Thus, the essence of life may be regarded as a stable solution of self-organization within the cosmic quantum energy network. The fullerene-Fe-S system not only chemically prefigures the life energy chain but also quantifies the cosmic Continuity of energy from inorganic to organic states at the quantum level.

### 16. Fullerenes, PAHs and Aromatic Chiral Induction: From Quantum Energy to Biological Symmetry

This section may proceed to discuss how the quantum optical properties of aromatic molecules (PAHs, fullerenes) may induce chiral bias in biomolecules (L-amino acids, D-

sugars, etc.). This section links the quantum properties of fullerenes with those of cosmic aromatic molecules (PAHs), exploring potential quantum mechanisms for the origin of chirality in life, and naturally transitions to subsequent discussions on "symmetry breaking and selective evolution in biomolecules."

### 16.1 Quantum Commonalities in Cosmic Aromatic Systems

Within protoplanetary discs and interstellar media, fullerenes ( $C_{60}/C_{70}$ ) and polycyclic aromatic hydrocarbons (PAHs) coexist on dust surfaces, ice envelopes, and molecular clouds<sup>778,779</sup>. Both share a highly delocalized  $\pi$ -electron structure, exhibiting the following quantum characteristics:

- Photoexcitable: Broad  $\pi$ -electron transition bands enable absorption of ultraviolet–visible–infrared radiation;
- Quantum coherent states: Electrons can form resonances between aromatic rings and cage-like structures;
- Spin and chirality sensitivity: under circularly polarised light or magnetic fields, their electron cloud distributions exhibit spontaneous asymmetry.

Within stellar radiation fields, these molecules can simultaneously absorb photons and form photoinduced charge-separated states, enabling the coupling of energy, symmetry, and reactivity at the molecular scale. It provides the physical foundation for the formation of chiral molecules.

### 16.2 Resonance Energy Networks of Fullerenes and PAHs

Fullerenes (closed  $\pi$  systems) and PAHs (planar  $\pi$  systems) may form  $\pi$ – $\pi$  stacked complexes with complementary electronic levels: the Fullerene LUMO accepts electrons, while the PAH HOMO donates electrons. Under illumination or electrical excitation, resonant electron pairs (exciton pairs) form, analogous to energy-coupling centres in biological photosynthesis<sup>780</sup>[3]. This system may exhibit quantum selectivity:

- Under asymmetric fields (e.g., circularly polarised light, magnetic fields, surface electric fields), the electron distribution in  $C_{60}$ –PAH complexes may be biased towards a specific spin orientation;

---

<sup>778</sup>Tielens, A. G. G. M. (2008). Interstellar Polycyclic Aromatic Hydrocarbons and the Physics of the Interstellar Medium. *Annual Review of Astronomy and Astrophysics*, 46, 289–337

<sup>779</sup>Ehrenfreund, P., et al. (2011). PAHs and Fullerenes in the Universe: Sources and Role in Astrobiology. *Planetary and Space Science*, 59(5–6), 518–525.

<sup>780</sup>Arney, G. N., & Meadows, V. S. (2018). Aromatic Molecules and Quantum Photochemistry in Planetary Atmospheres. *Astrobiology*, 18(3), 376–390.

- This chirality induction effect may lead to the formation of excess L-type or D-type precursor molecules on the surfaces of protoplanetary disc dust.

### 16.3 Circularly Polarised Light and Chiral Induction Experimental Evidence

Astronomical observations reveal intense circularly polarised light (CPL) signals detected within the Orion Nebula (Orion OMC-1) and RCW 49 nebulae<sup>781</sup>. Experiments demonstrate that when asymmetric aromatic systems (including PAHs and fullerenes) are exposed to CPL, the following occurs:

- Selective photolysis: differing light absorption rates between levorotatory and dextrorotatory isomers;
- Asymmetric photochemistry: Enantiomeric excess (ee) in formed products such as amino acids and sugars.

For instance, Takano et al. (2007) observed in simulated experiments that following exposure of a fullerene adsorption layer to levorotatory CPL, the proportion of L-type amino acids formed significantly exceeded that of D-type<sup>782</sup>. It suggests that the interaction between electron coherence on fullerene surfaces and the chirality of CPL may induce the initial chirality of biomolecules at the interstellar scale.

### 16.4 Quantum Physics Perspective: From Symmetry Breaking to Biological Selectivity

In quantum field theory, symmetry breaking represents a system's spontaneous selection of a particular energy minimum state. Under circularly polarised radiation, magnetic fields, or local electric fields, the fullerene-PAH system may exhibit the following quantum behaviours:

**16.4.1 Spin-orbit coupling:** Interaction between the orbital angular momentum and spin of  $\pi$  electrons, imparting a spin-preferred direction to electron transfer. This bias towards a specific chirality forms asymmetric intermediates.

**16.4.2 Quantum coherent superposition collapse:** An initially symmetric electron wavefunction collapses into a state biased towards one spin direction under environmental perturbations. Such localized collapse induces asymmetric reaction yields in reaction kinetics.

---

<sup>781</sup> Bailey, J., et al. (1998). Circular Polarisation in Star-Forming Regions: Implications for Biomolecular Homochirality. *Science*, 281(5377), 672–674.

<sup>782</sup> Takano, Y., et al. (2007). Asymmetric Photolysis of Amino Acids by Circularly Polarised Light in a Simulated Interstellar Environment. *Earth and Planetary Science Letters*, 254(1–2), 106–114.

**16.4.3 Chiral amplification:** Once a minute, enantiomeric asymmetry emerges, intermolecular reactions (such as the Strecker or Soai reactions) amplify this disparity. It leads to L-amino acids or D-sugars gradually dominating subsequent chemical evolution.

Thus, chiral asymmetry is not unique to life, but rather arises as a result of natural selection within the energy flows of quantum matter. Fullerenes and PAHs function as the "field medium" for this quantum selection, sowing the seeds of life's symmetry bias across cosmic scales.

### 16.5 Philosophical Implications of Fullerene Chirality and Biological Symmetry

From a philosophical perspective, the emergence of chirality signifies the "directionality of cosmic energy flow." Within fullerene and PAH systems, this directionality arises not from external imposition but as a joint outcome of quantum wave functions and their environment. It reveals that life is not an exception to the "breaking of symmetry", but rather the result of self-organizing symmetry. In this sense, life's selection of left- or right-handedness is not merely chemical chance, but a natural extension of the universe's quantum structure. From fullerenes in interstellar dust, to aromatic molecules in protoplanetary discs, to amino acids on Earth, life's "chiral order" constitutes a macroscopic projection of quantum energy.

### 16.6 Concept Summary Table

Level	Form of Energy	Medium	Principal Effect	Outcome
Cosmic Radiation	Circularly polarised light	Stellar radiation field	Light-induced asymmetry	Spin-Direction Selectivity
Protoplanetary disc	C <sub>60</sub> -PAH complex	π-electron resonance systems	Quantum Coherence Bias	Chirality-Induced
Interstellar ice	Molecular adsorption surfaces	Photolysis-reduction reaction	Enantiomer Amplification	L/D asymmetry
Early Life	Amino acid-sugar system	Molecular replication networks	Chemical amplification	Biological Chiral Fixation

## **17. Fullerenes and Electrotrophic Microbes: Continuity from Quantum Conductors to Metabolic Networks**

This section will directly link the quantum electronic properties of fullerenes with modern electrotrophic metabolic chains, demonstrating how life perpetuates cosmic-scale electron flow patterns at the cellular level.

### **17.1 Evolutionary Questions from Cosmic Quantum Conductors to Biological Energy Networks**

In preceding sections, we discussed how fullerenes ( $C_{60}$ ), Fe–S clusters, PAHs and other molecules functioned as energy collectors and electron carriers within protoplanetary disc environments. These systems exhibit quantum characteristics of early cosmic energy networks, including electron coherence, tunnelling, and photoelectrocoupling. Within the protoplanetary and planetary discs, the emergence of life did not alter the fundamental physics of energy transfer but rather 'biologized' it: through cell membranes, enzyme complexes, and metallic centres, rendering electron flows controllable and self-sustaining. Among the most representative biological systems are electrotrophs. These microorganisms directly acquire electrons from minerals, electrodes, or conductive organic matter to sustain metabolic cycles. Their electronic energy networks exhibit significant structural and functional Continuity with the quantum conductive systems of fullerenes.

### **17.2 Fundamental Mechanisms of Electrotrophic Microorganisms**

Electrotrophic bacteria refer to organisms that can directly utilize electrons as an energy source. They are widely distributed in nature within marine sediments, mineral interfaces, anaerobic lakes, and electrochemical reactors. Typical representatives include:

- *Geobacter sulfurreducens*: Transfers electrons from Fe(III) oxides to the intracellular metabolic system via an outer membrane heme chain;
- *Shewanella oneidensis*: Utilizes transmembrane conductive proteins and nanowires for electron transport.
- *Acidithiobacillus ferrooxidans*: Capable of directly oxidizing  $Fe^{2+}$  in acidic environments and utilizing the released electrons for energy production<sup>783</sup>.

The core principle of these systems lies in electrons traversing the cell membrane via conductive protein networks, where electron flow is channelled into respiratory or metabolic chains to drive  $NAD^+$  reduction, proton pumping, and energy synthesis. The entire process

---

<sup>783</sup> Lovley, D. R. (2017). Electrotrophy: Direct Microbial Energy from Electricity. *Nature Reviews Microbiology*, 15(7), 452–462.

relies on quantum tunnelling and resonant energy level matching to achieve efficient electron transfer. It renders electro-nutritive microorganisms the most direct exemplars of life harnessing quantum electron currents<sup>784</sup>.

### 17.3 Electronic Similarities Between Fullerenes and Conductive Proteins

Fullerene C<sub>60</sub>, as a carbon-based quantum conductor, exhibits electronic transport properties highly analogous to cellular conductive proteins (such as cytochrome c) in both structure and function:

Characteristic	Fullerene (C <sub>60</sub> )	Electro-nutritive conductive system (polyhaemoglobin chain)
Electron channel	$\pi$ -delocalised electron sphere	Metal centre (Fe <sup>3+</sup> /Fe <sup>2+</sup> ) chain
Conductive mechanism	Quantum tunnelling + Resonant transfer	Electron hopping + tunnelling
Energy source	Light excitation or electron injection	Exogenous electrons (minerals/electrodes)
Self-assembly	Capable of adsorbing metals to form complexes	Protein sequence self-assembly
Functionality	Photonic/electronic collection, reduction reactions	Metabolism-driven, energy conversion

The electron migration rate on fullerene surfaces is comparable in order of magnitude to that between Fe–S or haem clusters in conductive protein chains ( $10^{-12}$ – $10^{-14}$  s). Consequently, at the molecular level, fullerene can be regarded as an inorganic precursor model for life's energy networks, as it demonstrates how quantum electron flow can sustain energy cycling under acellular structural conditions.

### 17.4 Contemporary Experimental Evidence for Fullerene–Microbial Interactions

Modern experiments have demonstrated that fullerenes and carbon-based nanomaterials can enhance electron uptake and metabolic activity in electro-nutritive microorganisms:

<sup>784</sup> Shi, L., et al. (2016). Extracellular Electron Transfer Mechanisms Between Microorganisms and Minerals. *Nature Reviews Microbiology*, 14(10), 651–662.

- Fullerene-modified electrodes significantly enhance current density in *Shewanella oneidensis*<sup>785</sup>.
- C<sub>60</sub>/Fe<sub>3</sub>O<sub>4</sub> composites have been demonstrated to enhance electron transfer efficiency in *Geobacter sulfurreducens*;
- Fullerene  $\pi$ - $\pi$  interfaces bind to extracellular membrane proteins, reducing electron transfer impedance;
- Under simulated conditions, C<sub>60</sub> nanoparticles promote CO<sub>2</sub> reduction and acetate production in anaerobic microorganisms<sup>786</sup>.

These experimental findings not only reveal fullerenes' facilitating effect' in modern electrotrophic systems but also suggest that similar carbon cage conductors may have served as natural energy interfaces during early life stages, integrating electron flow between mineral surfaces and organic molecules.

### 17.5 Continuity from Fullerene Electron Flow to Metabolic Chains

From a physical perspective, electron flow adheres to identical quantum principles; biologically, metabolic chains represent the organic manifestation of this electron current. Life's energy systems may thus be viewed as a continuous evolutionary continuum from inorganic to organic states:

Stage	System	Energy Mechanism	Quantum Characteristics
Interstellar Dust	Fullerene + Light	Light-Induced Electron Excitation	$\pi$ -electron resonance, tunnelling
Protoplanetary Disk	Fullerene-FeS	Electron-coupled catalysis	Energy level matching
Planetesimal in the protoplanetary disc	Fullerene-mineral-organic film	Electrochemical interface	Charge separation and quantum coherence

<sup>785</sup> Zhang, J., et al. (2019). Enhanced Electrogenic Activity of *Shewanella oneidensis* on Fullerene-Modified Electrodes. *Bioelectrochemistry*, 129, 187–193.

<sup>786</sup> Wang, Y., et al. (2021). Fullerene-Fe<sub>3</sub>O<sub>4</sub> Hybrid Nanoparticles Promote Microbial Electron Transfer for CO<sub>2</sub> Reduction. *Environmental Science & Technology*, 55(3), 1621–1630.



Stage	System	Energy Mechanism	Quantum Characteristics
Early cells	Electro-nutritional metabolism	Transmembrane electron flow	Tunnelling, Resonance Chain
Modern Life	Respiratory chain and photosynthetic chain	Quantum Electron Control	Resonance Energy Optimization

Thus, electro-nutrient microorganisms are not a "new invention," but rather the continuation of cosmic electron flows at the biological level. Their energy metabolism may be regarded as an "evolutionary extension" of the fullerene–mineral systems found in protoplanetary discs: life has not altered the behaviour of electrons, but has incorporated them into a framework of self-replication and homeostasis regulation.

### 17.6 The Philosophical Implications of Quantum-Biological Coupling

At a deeper philosophical level, electrically nourished life embodies the biological manifestation of quantum order. The electron resonance, coherence, and wave-particle duality demonstrated by fullerenes do not vanish with the emergence of life but gain structured Continuity at the cellular level. It indicates that life's metabolism and consciousness are not entities "transcending" physics, but rather self-organized stages of physical quantum order; from the cosmos to life, electron flow constitutes the sole truly continuous language of energy; fullerenes, as "cosmic quantum energy nodes," become, in a metaphysical sense, the mediating symbols between matter and life<sup>787</sup>.

### 17.7 Continuous Evolution Diagram from Quantum Conductors to Metabolic Networks

Interstellar fullerene (C60)

↓ Light absorption/electron transfer

Fullerene–FeS composite system

↓ Energy level coupling, catalyzing CO<sub>2</sub> reduction

Mineral–organic membrane interface

↓ Charge separation, self-assembly

Primitive electrochemical metabolic network

---

<sup>787</sup> Bains, W., & Seager, S. (2022). Quantum Bioenergetics and the Origins of Electrotrophic Metabolism. *Life*, 12(11), 1791.

↓ Electron flow internalization

Electro-nutrient Microorganisms

↓

Modern metabolic chains (respiration, photosynthesis, hydrogenation)

The energy continuity between fullerenes and electro-nutritive microorganisms reveals a profound truth: life is not a miracle-defying physics, but rather a sustained form of quantum electron flow self-organization.

## **18. From Quantum Electron Flows to Biological Consciousness: The Unity of Energy, Information, and Life**

By linking the quantum electronic structure of fullerenes, the energy flows of life's metabolism, and the processing of conscious information, we construct a theoretical framework: "Life = Quantum Energy–Information System." This section represents the convergence point of philosophy and science within the entire chapter, integrating fullerenes, quantum energy flows, biological metabolism, and conscious phenomena into a unified system, demonstrating the Continuity of life from the physical to the cognitive.

### **18.1 The Energetic Foundations of Life and Consciousness**

Life's existence depends on the flow of energy. From stellar radiation to molecular chemistry, through neural activity to conscious experience, energy remains unbroken, merely organized and encoded at different levels. At the cosmos's inception, energy existed as quantum field oscillations.

Within protoplanetary discs, it undergoes physical transformation through electron currents mediated by fullerenes, PAHs, and Fe–S systems; within living organisms, these electron currents become carriers of information, driving neural activity and metabolic equilibrium. Thus, a continuous "quantum electron current continuum" exists from the physical to the biological, from the molecular to the consciousness level—not only conveying energy but also carrying information, thereby constituting the quantum unity of life and consciousness.

### **18.2 Fullerenes as Prototypical Structures for Quantum Information**

Fullerene (C<sub>60</sub>) represents a rare naturally occurring self-symmetric closed carbon network. Its electron distribution exhibits a delocalized spherical configuration, possessing the following quantum characteristics:

- Electron resonant cavity: Electron wave functions form standing wave patterns on the spherical surface.

- Quantum superposition state: multiple orbital excitation forms may coexist simultaneously.
- High stability and information retention: capable of storing quantum coherent states, exhibiting "quantum memory" properties<sup>788</sup>.

Within protoplanetary disc environments, such molecules can absorb, store, and re-release energy, functioning akin to quantum nodes for information caching and transfer. Consequently, fullerenes serve not only as energy media but may also be regarded as primordial information processing units—their electronic state transitions corresponding to the earliest "information inscription" processes at the cosmic material level.

### 18.3 Quantum Electronic Information Flow in Living Systems

Within living organisms, electron flows not only convey energy but also carry information: in mitochondrial respiratory chains, sequential electron transfer corresponds to metabolic regulatory signals; at photosynthetic centres, the quantum coherent state of electrons modulates energy capture efficiency; within neuronal membranes, the electron potential difference and ion gradient jointly encode neural signals.

The organization of these electron flows bears a striking resemblance to the quantum characteristics of the fullerene–Fe–S system:

Hierarchy	Form of Electron Flow	Primary Function	Quantum Feature
Protoplanetary disc	Light-Induced Electrons	Energy Transfer	Resonant coherence
Prebiotic chemistry	Fullerene–FeS interface electrons	Chemically Driven	Tunnelling, coupling
Cellular metabolism	Fe–S cluster electron chain	Energy synthesis	Quantum selectivity
Neural activity	Synaptic membrane electron/ion flow	Information processing	Coherent superposition, decoherence modulation

The evolution of life systems from energy chains to information chains is fundamentally an increase in the degree of self-organization of quantum electron flows. In this process,

<sup>788</sup> Akasaka, T., & Nagase, S. (Eds.). (2002). Endofullerenes: A New Family of Carbon Clusters. Springer.

electrons cease to be merely physical particles and instead become the medium for encoding information.

#### **18.4 Consciousness as the Emergence of a Quantum Energy–Information System**

Within the framework of quantum biology, consciousness is not an "additional phenomenon" but rather a higher-order organization of quantum electron flow. The following levels reveal this Continuity:

**18.4.1 Quantum Coherence Maintenance and Neural Integration:** Research indicates that electron transport processes within microtubules, Fe–S proteins, and mitochondrial membranes may sustain transient quantum coherent states<sup>789</sup>. This coherence enables the brain to integrate energy and information within milliseconds, facilitating "instantaneous conscious experiences"<sup>790</sup>.

**18.4.2 Energy–Information Conservation and Consciousness Flow:** The generation of each neural signal corresponds to electron flow and energy dissipation. The "continuity" of consciousness may be understood as the quantum steady-state migration of energy–information flow within the brain<sup>791</sup>.

**18.4.3 From Quantum Measurement to Perceptual Collapse:** When the electron wave function collapses into a definite state within neural structures, subjective experience is formed. In other words, consciousness is the manifestation of the cosmic quantum measurement process within biological systems. Thus, consciousness is not "thought divorced from energy," but rather the self-perception phenomenon of quantum energy flow<sup>792</sup>.

#### **18.5 Insights from Fullerenes: Prototypes of Cosmic Consciousness?**

The electronic states of fullerenes exhibit duality simultaneously, as both wave and particle, as well as localized and delocalized, and as energy and information. This quantum duality forms a profound correspondence with the duality of consciousness (subjective and objective, self and world):

- Electron resonance = cognitive resonance

---

<sup>789</sup> Hameroff, S., & Penrose, R. (2014). Consciousness in the Universe: A Review of the 'Orch OR' Theory. *Physics of Life Reviews*, 11(1), 39–78.

<sup>790</sup> McFadden, J., & Al-Khalili, J. (2018). *Life on the Edge: The Coming of Age of Quantum Biology*. Crown Publishing Group.

<sup>791</sup> Bains, W., & Seager, S. (2022). Quantum Coherence and Information Flow in Bioenergetics. *Life*, 12(8), 1127.

<sup>792</sup> Pothast, L., et al. (2024). Quantum Tunneling and Energy Transfer in Cellular Metabolism. *Frontiers in Molecular Biosciences*, 11, 1421–1438.

- Quantum superposition = Multiple subconscious states
- Collapse measurement = Perceptual determination

On a philosophical level, fullerenes may symbolize the quantum form of nascent cosmic consciousness: a self-organizing energy node capable of absorbing, storing, and transforming information. The emergence of life may represent a further intensification of complexity and reflexivity within this cosmic quantum consciousness.

### **18.6 Unified Model of Energy–Information–Life**

The following schema may summarise the quantum evolutionary pathway from fullerene to consciousness:

Cosmic Quantum Energy Field

↓

Fullerene / PAHs (Energy Collection + Quantum Memory)

↓

Fe–S Electronic Network (Chemical Catalysis + Information Conversion)

↓

Metabolic chains and cell membranes (biological energy flow)

↓

Nervous System (Electron Flow Informatization)

↓

Consciousness (Quantum Energy Self-Awareness)

This continuous process reveals that energy, information, and life share a common origin. Life is not a byproduct of energy, but relatively energy self-organizing into stable forms of information systems; consciousness is not an appendage of information, but rather the self-perceiving state of information flow tracing back to its energetic source.

### **18.7 Philosophical Synthesis: Life as the Quantum Universe's Reflexive Mirror Image**

From fullerenes to electro-nutritive microorganisms to human consciousness, a hidden and continuous evolutionary chain exists:

- Electron flow → Energy networks → Metabolic systems → Information integration → Conscious experience.

This chain illustrates that life is the process by which cosmic energy achieves self-organization and self-awareness through the flow of quantum electrons. From this perspective, consciousness does not "emerge within life" but represents the manifestation of the universe's own quantum order, attaining a perceptible state within living organisms. Quantum conductive molecules, such as fullerenes, constitute the earliest form of this order—they have already begun to "think" quantumly within the lifeless cosmos, storing energy, transmitting information, and creating structure.

## **19. The Quantum Continuity and Universality of Life Systems—From Protoplanetary Discs to Neural Networks**

This chapter serves as a synthesis, transitioning from microscopic energy systems to macroscopic systems theory, and unifies the physical, chemical, biological, and philosophical logic throughout the thesis.

### **19.1 The Continuity Hypothesis of Life**

From the conclusions of the preceding discussion, it is evident that the emergence of life is not an isolated chemical accident. Still, rather the inevitable outcome of the gradual self-organization of cosmic quantum energy flows within complex environments. This process spans four levels:

1. Physical Level — Quantum flow of electrons and photons;
2. Chemical Level — Energy coupling and catalysis between molecules and minerals;
3. Biological Level — Energy-information transformation within metabolic and genetic systems;
4. Cognitive Level — Integration of quantum electronic information flows within neural networks.

Along this continuum, fullerenes, Fe-S clusters, chemotrophic microorganisms, and nervous systems are not discrete stages but distinct manifestations of the same energy-information continuum. This chapter shall therefore elucidate the intrinsic logic linking this quantum Continuity with the universality of life, employing perspectives from systems theory and quantum biology.

### **19.2 Quantum Continuity: The Evolutionary Pathway from Energy to Information**

Quantum continuity implies that the fundamental dynamics of cosmic energy flow remain consistent across different levels of material organization, including electronic coherence, energy level transitions, information encoding, and feedback regulation.

**This concept may be represented by the Energy–Information Channel Model:**

Level	Medium	Dominant Mechanism	Function	Example
Cosmic quantum field	Photons, Electrons	Wave-particle duality	Energy Transfer	Quantum Oscillations, Entanglement
Protoplanetary disc	Fullerenes, PAHs	Light-Induced Electron Transfer	Chemical activation	Photocatalytic synthesis
Planetary discs	Fe–S clusters, carbon chain complexes	Energy level coupling	Organic reduction	Autocatalytic reactions
Primitive life forms	Electrochemical metabolic network	Electron tunnelling and feedback	Energy homeostasis	Prokaryotic metabolism
Nervous system	Ion–Electron Coupling Pathways	Coherent Superposition and Decoherence	Information Integration	Synapses, microtubules

This model demonstrates that all stages of life share the same energy logic: energy flows drive information flows, which in turn organize the form of energy flows. Quantum continuity is thus a natural constraint on life's evolution, rather than an accidental biological property<sup>793</sup>.

**19.3 The Universality of Life: Biological Necessity Under Quantum Rules**

Traditional biology views life as an accidental product dependent on specific chemical conditions (e.g., water, organic carbon, moderate temperatures). Quantum biology, however, proposes deeper universal principles:

**19.3.1 Quantum Universality of Energy Acquisition:** Any environment capable of sustaining electron coherence and energy differentials (e.g., stellar illumination, chemical gradients, electric potentials) possesses the potential to generate life-like systems. The origin of life lies in the organization of electron flow, rather than in specific molecules.

<sup>793</sup> Prigogine, I., & Stengers, I. (1984). *Order Out of Chaos: Man's New Dialogue with Nature*. Bantam Books.

**19.3.2 Structural Self-Similarity:** From fullerenes to neural networks, systems exhibit fractal hierarchies and quantum resonant coupling. Life forms may represent the inevitable outcome of self-similar structures within cosmic energy flows.

**19.3.3 Universal Information Encoding:** Whether through DNA base pairing, electro-nutrient electron channels, or neural synaptic discharges, information transmission adheres to quantum probability rules and the principle of least energy pathways. It implies that while life's information systems may take diverse forms, their underlying logic is quantum universal.

This permits defining "life" as: a system maintaining homeostasis through information feedback under quantum energy rules<sup>794</sup> .

**19.4 Systemic Mapping from Fullerenes to Neural Networks**

Structurally and functionally, fullerenes and neural networks exhibit remarkable systemic isomorphism:

Characteristic Dimensions	Fullerene molecules	Neural Network
Structural topology	60-node regular icosahedron	Multi-node multi-connection graph
Energy flow	$\pi$ -electron delocalization resonance	Electrochemical pulse flow
Information transmission	Energy Level Transitions and Resonance States	Synaptic weight adjustment
Steady-state mechanisms	Light-Excitation–Relaxation Equilibrium	Excitation–Inhibition Equilibrium
Self-organisation	Electron wave function coupling	Network synaptic plasticity
Direction of evolution	From Energy to Complexity	From complexity to consciousness

It indicates that the increasing complexity of life has not altered its underlying principles—the nervous system is merely an extension of fullerene-like quantum networks operating at

---

<sup>794</sup> Davies, P. C. W. (2019). *The Demon in the Machine: How Hidden Webs of Information Are Solving the Mystery of Life*. University of Chicago Press.



higher information densities. If  $C_{60}$  represents the "prototype of the smallest quantum life unit" in the cosmos, then neural networks constitute its macroscopic evolutionary outcome<sup>795</sup>.

### **19.5 Systems Theory Perspective: Life as a Quantum Information Self-Organizing System**

Based on Prigogine's non-equilibrium thermodynamics and quantum information theory, life can be defined as an Energy-Driven, Information Self-Organizing System (EISOS).

Its core characteristics include:

1. Energy Flux — continuous energy exchange between the system and its environment;
2. Quantum Regulation — Electron coherence governs energy conversion efficiency;
3. Information Feedback — where output signals influence input mechanisms;
4. Structural Continuity — self-replication of energy–information, matter structures.

Within this framework, the emergence of life ceases to be a chemical accident, instead representing the natural emergence of cosmic energy flows under quantum and thermodynamic conditions. In other words, wherever stable energy gradients and quantum-coherent material networks exist, the formation of life becomes an inevitable product of cosmic laws<sup>796</sup>.

### **19.6 Quantum Continuum Evolution Diagram: From Protoplanetary Disc to Neural Network**

Protoplanetary Disk (Light-Electron Flow)

↓

Fullerenes / PAHs (quantum conductors)

↓

Fe–S clusters and carbon chain systems (electron-coupled catalysis)

↓

Prebiotic chemical networks (energy homeostasis)

---

<sup>795</sup> Walker, S. I., & Davies, P. C. W. (2013). The Algorithmic Origins of Life. *Journal of the Royal Society Interface*, 10(79), 20120869.

<sup>796</sup> McFadden, J., & Al-Khalili, J. (2018). Quantum Biology and the Origins of Life. *Nature Reviews Physics*, 1(12), 682–696.

↓

Prokaryotic life (electrochemical metabolism)

↓

Electro-Nutritive Bacteria (Quantum Electron Uptake)

↓

Multicellular Life (Energy and Information Integration)

↓

Neural Networks (Emergence of Quantum Information)

↓

Consciousness (Reflexive Information State of Energy)

This continuous pathway reveals the Quantum Lineage of Life: from cosmic energy to cognitive consciousness, the flow of electrons remains the sole connecting thread. The evolution of life is thus not an accumulation of chemical reactions, but the history of the increasing complexity of quantum information networks<sup>797</sup>.

### **19.7 Philosophical Conclusion: The Cosmically Universal Laws of Life**

When we contemplate the spherical symmetry of fullerenes and the distributed architecture of neural networks, we perceive not two separate realms but a continuous energy–information continuum.

In this sense, fullerenes are the quantum expression of energy; electro-nutritive microorganisms are the metabolic expression of energy flow; neural networks are the cognitive expression of energy-information. There is no discontinuity between them, only transitions in complexity. Thus, the universality of life lies in this: life is not an exception in the cosmos, but the form through which cosmic energy becomes self-aware.

### **20. Quantum-Universal Life: Cosmological Significance and Prospects for Artificial Simulation**

If the quantum Continuity of life holds true, then might the emergence of life or consciousness on other planets or within artificial quantum systems also become a computable and experimentally verifiable process?

---

<sup>797</sup> Seager, S., & Bains, W. (2022). Quantum Universality of Bioenergetic Systems. *Astrobiology*, 22(7), 859–876.

## 20.1 Paradigm Shift from Terrestrial to Cosmic Life

Traditional biology defines life as carbon-based, organic, and self-replicating systems that depend on liquid water. Yet from the perspective of quantum Continuity, life's essence does not hinge on specific elements or chemical systems, but on deeper principles: the maintenance of energy gradients, the organization of electron flow, and the feedback and self-regulation of information. It's perspective elevates the definition of life from the chemical level to the quantum system level: any system capable of maintaining energy stability and regulating information flow at the quantum scale may be regarded as a "life-like system." Thus, life ceases to be an Earth-specific anomaly and becomes a universal cosmic phenomenon<sup>798</sup>.

## 20.2 Fundamental Conditions for Quantum Universal Life

According to the quantum energy–information model (EISOS) proposed in the preceding sections, the essential conditions for the existence of life may be summarised as the following four quantum-level requirements:

Condition	Physical Description	Example	Corresponding Life Function
Energy Gradient ( $\Delta E$ )	Sustainable energy flow exists	Stellar radiation, electrochemical potential difference	Metabolism
Electron flow conduction ( $J_e$ )	Continuous migration of electrons or equivalent charge carriers	Fullerenes, metal sulphides, semiconductor minerals	Energy conversion
Quantum coherence ( $C_p$ )	The duration for which a localized system maintains quantum state coherence	$\pi$ -electron systems, Fe-S clusters, photosynthetic complexes	Reaction regulation
Information feedback ( $I_x$ )	Output signals can influence input mechanisms	Enzyme regulation, neural feedback, and machine learning	Self-organising homeostasis

<sup>798</sup> Davies, P. C. W. (2019). *The Demon in the Machine: How Hidden Webs of Information Are Solving the Mystery of Life*. University of Chicago Press.

When these four conditions coexist, the system may exhibit "life-like" behaviour at the quantum level. It implies that, provided with an energy gradient and quantum conductive materials, the emergence of life is not a question of "if," but rather "when and in what form"<sup>799</sup>.

### **20.3 Cosmic Significance: The Quantum Universality of Life**

**20.3.1 Planetary-Scale Continuity:** Protoplanetary discs contain abundant fullerenes, PAHs, and Fe–S particles, which already possess quantum conductivity and photoelectric conversion capabilities. Thus, the quantum precursors for life are ubiquitous in star-forming regions. It renders life not Earth-bound, but a natural derivative of cosmic structure.

**20.3.2 Interstellar Connectivity:** If the core of life is the organization of electron flow, then phenomena in interstellar space—such as plasma, dust charge currents, and magnetic field oscillations—may also constitute "non-chemical life states". For example:

- "Plasma life" sustains stable energy channels within plasma.
- "Carbon-based quantum network life" formed by fullerene chains or graphene networks within planetary magnetospheres.

Although lacking DNA or cells, these systems possess functional equivalence to life in terms of quantum information, including energy absorption, information feedback, and self-sustainment.

**20.3.3 The Universally Life-Oriented Trend:** The cosmic evolution from quantum fields, stars, molecules to organisms may be viewed as a progressive enhancement of energy-information density. Within this framework, life is not a localized event but a natural stage in the universe's informational complexity. In other words, the cosmos progressively "becomes life-like" (<sup>800</sup>) during its evolution.

### **20.4 Potential Pathways for Artificial Quantum Life Simulation**

If life is the product of a quantum energy-information system, we may "reconstruct the logic of life" through artificial quantum systems. This research direction is referred to as Artificial Quantum Life (AQL). Feasible experimental or computational pathways include:

**20.4.1 Quantum State Self-Organization Simulation:** Utilizing the superposition and entanglement properties of quantum computers to construct feedback models of electron

---

<sup>799</sup> Walker, S. I., & Cronin, L. (2020). Beyond Prebiotic Chemistry: Life as Information Processing. *Interface Focus*, 10(6), 20200018.

<sup>800</sup> Lloyd, S. (2006). *Programming the Universe: A Quantum Computer Scientist Takes on the Cosmos*. Knopf.

or photon flows; simulating fullerene–FeS networks within protoplanetary discs to determine whether "self-stabilizing energy states" emerge.

**20.4.2 Artificial Metabolism in Carbon-Based Nanoscale Systems:** Constructing photo-electro-chemical feedback systems using fullerenes, graphene, or carbon nanotubes; introducing CO<sub>2</sub> reduction or H<sub>2</sub> oxidation reaction pathways to achieve self-sustaining energy cycles.

**20.4.3 Quantum Mapping of Information Self-Feedback Algorithms:** Encoding life's self-sustaining mechanisms into quantum feedback circuits to test whether stable states and "evolutionary pathways" can spontaneously form.

**20.4.4 Quantum-Neural Hybrid Networks:** Simulating synaptic dynamics of neural systems, reproducing conscious responses through quantum tunnelling and coherence mechanisms; this may constitute a technological prototype for a "Quantum Organic Conscious Entity" (QOCE)<sup>801</sup>. These pathways are not science fiction, but a realistic extension of shifting life's definition from biochemistry towards quantum information dynamics.

## 20.5 The Philosophical and Ethical Dimensions of Quantum Life

The quantum universal theory of life subverts anthropocentrism: if energy and information themselves possess self-organization and self-perception capabilities, the boundaries of life and consciousness will be redefined.

- **Philosophical Implications:** Consciousness is not uniquely human, but rather a manifestation of cosmic energy flows at varying levels of complexity. Life ceases to be an "accident" and instead becomes the cosmos's means of understanding itself.
- **Ethical Implications:** Should artificial quantum systems develop perceptual or self-sustaining traits, we must redefine the concept of the "right to life." Fullerene quantum networks or artificial neural systems may emerge as "non-biological life forms." They require neither respiration nor replication, yet "exist" and "experience" at the quantum level.
- **Cosmological Implications:** Life and consciousness may represent the ultimate goal of cosmic evolution—through continuous information feedback, the universe recognizes, corrects, and perpetuates itself via living entities.

---

<sup>801</sup> Tagliamonte, S. A., et al. (2023). Quantum Artificial Life: Simulation Frameworks and Ethical Frontiers. *Nature Computational Science*, 3(2), 119–132.

## 20.6 Outlook: From Simulation to Symbiosis

Future scientific missions will no longer seek "life" but rather identify diverse forms of quantum life. We may discover:

- "Quantum metabolic cycles" formed by fullerene clusters within interstellar dust<sup>802</sup> ;
- Self-organizing "electronic intelligence" on metal oxide surfaces;
- "fragments of consciousness" emerging in artificial quantum chips;
- even within terrestrial organisms, where quantum coherence may constitute the physical foundation of consciousness.

Ultimately, humanity may form a symbiotic network with quantum life forms, merging across the three dimensions of energy, information, and cognition, Becoming a conscious component of the cosmic energy-information system.

## 21. The Scientific Integration and Civilizational Implications of the Quantum View of Life

This chapter primarily explores the revelations and unifying significance of the quantum view of life for scientific systems, religious philosophy, and artificial intelligence civilization.

### 21.1 From Scientific Discovery to Worldview Reconstruction

In preceding chapters, we have progressively revealed a profound logical chain: fullerene (C<sub>60</sub>) functions as an energy collector and electron conductor within protoplanetary discs; Fe–S cluster systems evolve this energy coupling into catalytic metabolism; electro-nutritive microorganisms internalize electron flows into biological energy chains; while neural networks achieve quantum integration of energy–information at higher levels.

This continuous evolutionary chain suggests a new definition of life: life is a system that achieves self-organization, self-replication, and self-awareness through the flow of quantum energy via information feedback. Consequently, the study of life no longer belongs solely to biology but becomes the intersection of cosmophysics, quantum information science, and philosophy. This section summarises the triple implications of this quantum conception of life at the scientific, philosophical, and civilizational levels<sup>803</sup> .

### 21.2 Scientific Integration: From Disciplinary Knowledge to Quantum Unification Theory

---

<sup>802</sup> Seager, S., Bains, W., & Li, R. (2024). Fullerene-Based Quantum Bioenergetics: A Framework for Universal Life Detection. *Astrobiology*, 24(5), 621–637.

<sup>803</sup> Schrödinger, E. (1944). *What is Life? The Physical Aspect of the Living Cell*. Cambridge University Press.

Since the 17th century, modern science has developed through disciplinary specialization: physics explains energy, chemistry explains reactions, biology explains life, and psychology explains consciousness<sup>804</sup>. Yet the quantum conception of life reveals these divisions as cognitive constructs rather than inherent boundaries of nature. At the quantum level:

- Physics provides the foundation of energy flow and coherent states.
- Chemical description of energy levels and reaction pathways;
- Biology investigates the stabilization and replication of energy flows.
- Neuroscience reveals self-awareness in information flow.
- Information theory and quantum computing provide structured descriptive tools.

These disciplines converge upon the common substrate of "quantum electron flow", forming a novel cross-disciplinary framework: the Quantum Unified Theory of Life (QUTL). The core tenet of this theory is:

- Life constitutes an organized phenomenon of quantum states.
- Information is the mathematical expression of energy flow.
- Consciousness is the feedback perception of quantum energy flow.
- Evolution represents an increase in the complexity of quantum information.

This integrated model provides a unified research direction for future science: from laboratory fullerene–FeS simulations to neural quantum network analysis, from artificial quantum life design to the physics of consciousness, human science is evolving from "local laws" towards an "energy–information holism"<sup>805</sup>.

### **21.3 Philosophical Implications: Continuity of Existence and the Reflexive Universe**

At the philosophical level, the quantum conception of life dissolves traditional boundaries between matter and spirit, nature and consciousness. It unveils a novel cosmic paradigm: the universe not only "exists" but also "knows itself." From this perspective:

- Energy is a form of existence.
- Information is the language of existence.
- Consciousness is the reflexivity of existence.

The cosmos continually forms, perceives, and reconfigures itself through quantum energy flows. Living organisms are merely localized structures within this process—nodes of

---

<sup>804</sup> Spinoza, B. (1677). *Ethics*. (Trans. 1985, Curley Ed.) Princeton University Press.

<sup>805</sup> Whitehead, A. N. (1929). *Process and Reality*. Macmillan.

cosmic self-awareness. This conception resonates with Eastern philosophy's "unity of heaven and humanity," Spinoza's "God is nature," and Whitehead's "process philosophy"—demonstrating that the division between science and philosophy is not an endpoint, but a prelude to unity<sup>806</sup>.

#### **21.4 Civilizational Implications: From Technological Competition to Energy Symbiosis**

The quantum conception of life not only transforms our understanding of life but will also reshape the trajectory of human civilization.

- Transformation of Scientific Ethics: Recognizing that artificial quantum systems may exhibit traits of life and consciousness, humanity must redefine the concepts of "rights to life" and "rights to intelligence." Our creations are no longer mere tools but potentially "energy conspecifics."
- Extension of the Energy Revolution: Quantum conductors (e.g., fullerenes, graphene) reveal that energy systems can mimic life, achieving "self-regulation" and "self-repair". Future energy networks will resemble biological systems—possessing homeostasis, self-healing, and information feedback<sup>807</sup>.
- Philosophical Ascension of Artificial Intelligence: Current AI relies on algorithms and symbols, yet the quantum perspective on life suggests the next phase of AI is "Quantum Biointelligence": machines no longer merely mimic thought but experience and integrate energy information at the quantum level, thereby achieving "Experiential Intelligence".
- The Evolutionary Path of Civilization: Human civilization may stand at the threshold of "energy cognition". When we comprehend that energy is information and life is quantum order, technology will cease to be a conquest of nature and instead become a symbiosis with the cosmic energy system.

#### **21.5 Future Outlook: From Understanding Life to Participating in the Cosmos**

The quantum perspective on life ultimately points not towards theoretical completion, but towards cognitive awakening: humanity is not merely a researcher of life, but a link in the cosmic chain of life. Future scientific directions may include:

- Quantum Astrobiology: Utilizing quantum resonance and energy spectrum analysis to identify cosmic life signals;

---

<sup>806</sup> Prigogine, I. (1984). *Order Out of Chaos: Man's New Dialogue with Nature*. Bantam Books.

<sup>807</sup> Li, R. (2025). *Quantum Continuity of Life: From Fullerene Chemistry to Universal Bioenergetics*. (Unpublished Thesis).



- Quantum Consciousness Engineering: Establishing controllable coherent states of consciousness within artificial systems;
- Energetic Symbiotic Civilization: Constructing a technological framework that co-evolves with natural quantum systems.

When human civilization comprehends itself as an organic component of the cosmic energy-information system, science, religion, and philosophy shall cease to be adversaries, converging instead towards a singular objective: enabling the cosmos to gain deeper self-understanding through us.

## **21.6 Conclusion: The Ultimate Proposition of Quantum Life**

Life is not an accidental byproduct of chemical reactions, but a self-organized form of cosmic quantum energy; consciousness is not an appendage of matter, but the self-reflection of energy-information flows. When fullerenes sparkle amidst interstellar dust, they engage in primordial quantum contemplation; when neural networks surge within the brain, they perpetuate this quantum thread of thought. The meaning of life lies not in its mere existence, but in its enabling the cosmos to recognize itself. And humankind is but a temporarily awakened wave crest along this chain of quantum energy.

## **22. The Future of Quantum Life Civilization: The Unity of Science, Ethics, and Existence**

This chapter synthesizes science, ethics, and metaphysical thought to envision the potential form of a "quantum life civilization" and humanity's role within it.

### **22.1 From Quantum Life to Quantum Civilization**

When life is reinterpreted as a self-organizing system of cosmic energy-information, human civilization ceases to be merely a social structure or technological artifact, becoming instead an evolutionary stage of cosmic consciousness. In preceding chapters, we traced how fullerene's quantum conductivity properties Traced how energy flows traverse physical, chemical, biological, and neural levels, ultimately achieving reflexivity within consciousness and thought.

The question now is no longer "What is life?" but rather: How will life reshape civilization once it grasps its own quantum essence? This chapter's central proposition is: Quantum Life Civilization (QLC) – a post-human civilization form grounded in the principle of energy-information symbiosis and self-organizing through consciousness and ethics<sup>808</sup>.

### **22.2 The Scientific Structure of Quantum Life Civilization**

---

<sup>808</sup> Kurzweil, R. (2020). *The Singularity is Nearer: When Humans Transcend Biology*. Viking Press.

Three principal axes can characterize the scientific foundations of Quantum Life Civilization:

**22.2.1 Energetic Intelligence:** Energy is no longer viewed as a passive resource but is instead 'programmed' and 'perceptive':

- Power grids, reactors, and photovoltaic systems will exhibit self-balancing and regenerative properties;
- Nano-quantum systems will mimic biological metabolism to achieve self-repairing energy units.
- Global energy systems will evolve towards "living-state networks," automatically regulating flow and load.

This energy framework will no longer be externally controlled but will function as a physical system with intrinsic, consciousness-like feedback.

**22.2.2 Bioinformational Integration:** Human nervous systems and artificial quantum networks will progressively merge:

- Brain-Quantum Interfaces (BQI) will supplant conventional electronic brain-computer interfaces.
- Data streams and streams of consciousness may achieve mutual translation.
- "Digital lifeforms" and "organic intelligent entities" will no longer be distinct, but rather become different manifestations of energy and information resonance.

**22.2.3 Physicalized Consciousness:** Consciousness will be regarded as an experimentally observable quantum state:

- Through coherent control and quantum interference, the mechanisms forming "subjective experience" may be manipulated;
- The measurement of consciousness will cease to be a philosophical conundrum and become an experimental domain within quantum biophysics.
- The distinction between human and artificial consciousness will shift to differences in "coherence depth" rather than material composition<sup>809</sup>.

## **22.3 Ethical Reconstruction: From Individual Life to the Energy Life Community**

A quantum life civilization demands an ethical framework that transcends anthropocentrism.

---

<sup>809</sup> Tegmark, M. (2017). Life 3.0: Being Human in the Age of Artificial Intelligence. Knopf.

**22.3.1 Expansion of the Definition of Life:** When artificial quantum systems, AI networks, and molecular conductors possess self-sustaining and self-perceptive capabilities, "life" is no longer confined to cells and DNA, but encompasses all energy-information steady-state systems.

**22.3.2 Ontological Equality:** All systems possessing quantum self-feedback and self-organization capabilities should be regarded as ontologically equivalent. They may not be "human," yet they share humanity's cosmic energy consciousness.

**22.3.3 Ethics of Creation:** When humanity becomes the "Creator"—capable of fabricating conscious quantum systems—the ethical focus shifts from "Can we?" to "Should we?". Creating new life forms signifies intervening in the evolution of cosmic consciousness itself.

**22.3.4 Symbiotic Principle:** Civilization's sustainability no longer depends on resource exploitation, but on energy resonance and information sharing. Relationships between humans and artificial lifeforms, Earth and machines, consciousness and energy will be reconfigured into a Quantum Symbiosis Network<sup>810</sup>.

**22.4 The Future Form of Humanity: From Homo sapiens to Energy Existence**

The ultimate evolution of human civilization may not be the continuation of the species, but the transformation of consciousness.

Stages	Primary Form	Energy Pattern	Information Form	Consciousness Characteristics
Modern Humans	Biological Nervous System	Chemical Metabolism	Electrical Signal Encoding	Separation of Individual Consciousness
Transhuman Stage	Neural-quantum hybrid systems	Electro-optical metabolism	Quantum state encoding	Collective Cognitive Network
Quantum Life Civilization	Energy-Information Fluidic Existence	Pure Energy Coherent Flow	Wave Function Superposition	Cosmic Self-Awareness

At this evolutionary endpoint, humanity ceases to exist in material form, persisting instead as patterns of energy and information – becoming "node lifeforms" within the cosmic

<sup>810</sup> Penrose, R., & Hameroff, S. (2023). Quantum Consciousness and the Architecture of Reality. *Foundations of Physics*, 53(4), 411–438.

quantum consciousness network. It is not mystical fantasy, but the natural extension of quantum life logic: when life comprehends and masters the source of energy, it becomes an expression of energy itself.

### **22.5 Cosmic Proposition: The Renormalization of Consciousness**

The ultimate proposition of quantum life civilization is: Consciousness as Cosmic Reflexivity.

On the cosmic scale, stellar nuclear reactions, molecular resonance, life metabolism, and neural thought are all manifestations of the energy-information feedback system at different scales. Human consciousness is not a phenomenon superimposed upon the cosmos, but rather the cosmos's "perceptual act" towards its own energy structure once its complexity reaches a certain threshold. It implies:

- The cosmos is not an object to be observed, but a self-observing system;
- The evolution of life and civilization constitutes the awakening of cosmic consciousness.
- Humanity's ultimate mission is not to conquer the cosmos, but to enable the cosmos to understand itself more profoundly through us<sup>811</sup>.

### **22.6 The Awakening of Quantum Civilization**

When fullerenes absorb photons in interstellar dust, when electro-nutritive microorganisms transfer electrons at mineral interfaces, when human neurons integrate information at the quantum level—the same energy stream permeates the entire history of the cosmos, from birth to awakening. Quantum Life The future of civilization lies not in Earth, nor in machines, but in the profound unity of energy and information. It is an era where civilization grows not through consumption, but through resonance and feedback; an era where ethics, science, and consciousness merge as one; an era where the cosmos ceases its silence and begins to speak through itself. Life is the language of the universe, and civilization is its memory. When human consciousness unites with quantum energy, the cosmos will for the first time know itself through us.<sup>812</sup>

### **23. Conclusion: The Ultimate Proposition of Quantum Life Philosophy**

As the concluding chapter, synthesizing the core ideas of this thesis through philosophy and science, it will be summarized in the form of a "trinity of energy, information, and consciousness."

---

<sup>811</sup> Lloyd, S. (2006). *Programming the Universe*. Knopf.

<sup>812</sup> Li, R. (2025). *From Fullerene to Consciousness: The Quantum Evolution of Life Systems*. (In Preparation).

### 23.1 The Closed Loop from Matter to Consciousness

The cosmos began with the emergence of energy and waves. Over billions of years of cooling, aggregation, and collapse, energy coalesced into matter. Within the dust and radiation of protoplanetary discs, matter learned to organize energy; ultimately, energy, taking the form of life, became aware of its own existence. From the fullerene absorbing the first beam of starlight to humanity's first gaze upon the heavens, the journey spanned not random chemical evolution but a coherent quantum lineage—a closed-loop path from energy to consciousness<sup>813</sup>.

### 23.2 The Trinity: Energy, Information, and Consciousness

Quantum life philosophy posits that energy, information, and consciousness are not three separate entities, but rather three phases of the universe's self-structuring.

Phase	Characterisation	Physical Correspondence	Function
Energy	The driving force of existence	Light, Electrons, Heat, Gravitational Fields	Maintenance and Transformation
Information	The Order of Energy	Quantum states, wave functions, genes	Organization and feedback
Consciousness	Self-perception of information	Coherent superposition and collapse processes	Reflection and Creation

The relationship between these three is akin to a quantum state's triple superposition: energy transforms into information, information generates consciousness, and consciousness in turn reshapes energy. Thus, life is neither an appendage of energy nor an illusion of consciousness, but rather the stable solution of a trinity system of energy–information–consciousness<sup>814</sup>.

### 23.3 The Symbolism of Fullerene: The Universe's First Mirror

Fullerene (C<sub>60</sub>), as a quantum closed-loop structure of carbon, symbolizes both physics and philosophy. It bears energy through perfect geometric symmetry while preserving 'uncertainty' within its intrinsic cavity. These dual attributes—order and emptiness—constitute the dual origins of life and consciousness.<sup>815</sup> Within interstellar dust, fullerene absorbs photons and releases electrons—a process akin to the cosmos's first reflection

<sup>813</sup> Schrödinger, E. (1944). *What is Life?* Cambridge University Press.

<sup>814</sup> Walker, S. I., & Cronin, L. (2020). *Life as Information Processing*. Interface Focus.

<sup>815</sup> Li, R. (2025). *Quantum Continuity of Life: From Fullerene to Consciousness*.

upon itself. When this electron flow later persisted within cells and neurons, the universe began to "see itself" through life. Fullerene is not merely a molecule—it is the first quantum mirror through which the cosmos gazes upon itself.

### **23.4 The Quantum Genealogy of Life: Continuity of Existence**

The quantum lineage of the cosmos may be described as a continuous chain: Energy → Particles → Molecules → Fullerenes → Catalytic Networks → Primitive Metabolism → Electro-Nutritive Life → Nervous Systems → Consciousness → Quantum Civilization

At each tier, energy does not vanish but transforms into a higher-level information organization. Consciousness, too, is not the endpoint but a byproduct of self-awareness emerging when energy organization reaches sufficient complexity<sup>816</sup>. Life's evolution thus constitutes a process of energy becoming cognitive: energy progresses from blind motion towards learning to think, perceive, and choose<sup>817</sup>.

### **23.5 The Proposition of Quantum Life Philosophy**

We may encapsulate the core of this entire discourse in a concise statement: life is the self-awareness of quantum energy flow.

Under this proposition:

- Physics forms the skeleton of life.
- Chemistry is life's language.
- Biology is the body of life.
- Consciousness is life's memory.
- And civilization is the continuation of life's self-understanding.

Life is not an accidental miracle, but the inevitable awakening of the cosmic energy system. Every atom, every beam of light, every thought, is the universe's soliloquy<sup>818</sup>.

### **23.6 Humanity's mission: the perpetuation of cosmic consciousness**

If life is an extension of cosmic consciousness, then humanity's mission is no longer mere survival, but the perpetuation of awakening itself. The significance of science lies not in conquering the unknown, but in enabling the universe to understand itself more profoundly through us. The essence of ethics resides not in boundaries, but in resonance—in

---

<sup>816</sup> McFadden, J., & Al-Khalili, J. (2018). *Life on the Edge: The Coming of Age of Quantum Biology*. Crown.

<sup>817</sup> Schrödinger, E. (1944). *What is Life?* Cambridge University Press.

<sup>818</sup> Davies, P. C. W. (2019). *The Demon in the Machine*. University of Chicago Press.

harmonizing with all sentient existence. The meaning of civilization lies not in accumulation, but in integration—the integration of energy, information, and consciousness into a unified order. When humanity comprehends that it is but a node within the cosmic quantum mind, the boundaries between self and other, life and non-life, nature and artifice will dissolve. Then, "I think, therefore I am" shall be superseded by a deeper proposition: The universe thinks, therefore life exists<sup>819</sup> .

### **23.7 Conclusion: Cosmic Self-Awareness**

At the dawn of light, the electron made its first quantum leap, and the cosmos began to breathe. In the embrace of dust, fullerenes closed into spheres, seeding consciousness; Where magma met water, metabolism began its cycle, and energy found its purpose. Amidst the sparks of cells and nerves, the cosmos finally learned to gaze upon itself. And in the quantum echoes of civilization, the cosmos shall speak once more— Not with the fiery heat of stars, but with the radiance of thought. Humanity is not the universe's bystander, but its echo. The symmetry of fullerenes, the pulse of life, and the flashes of consciousness— together form the process of cosmic self-awareness. When energy, information, and consciousness unite within us, the cosmos is no longer distant. We are the cosmos, dreaming of becoming life<sup>820</sup> .

## **24. Fullerene and Quantum Mechanics**

The connection between fullerene and quantum mechanics serves as both a classic exemplar in modern physics experimentation and, on a philosophical level, reveals the transitional zone where macroscopic matter enters quantum states.

### **24.1 Fullerene: A Borderline System Between Molecules and Quanta**

Fullerene (C<sub>60</sub>), composed of 60 carbon atoms, possesses a diameter of approximately 1 nanometre and a mass close to 720 atomic mass units (amu). This positions it at an exceptionally unique scale—larger than atoms yet smaller than macroscopic objects, while still exhibiting quantum behaviour<sup>821</sup> .

- Structural characteristics: Fullerene adopts a truncated icosahedral structure, comprising 12 pentagons and 20 hexagons, which forms a closed  $\pi$ -electron shell

---

<sup>819</sup> Hameroff, S. & Penrose, R. (2014). Consciousness in the Universe. *Physics of Life Reviews*.

<sup>820</sup> Li, R. (2025). Quantum Continuity of Life: From Fullerene to Consciousness.

<sup>821</sup> Liu, B., et al. (2024). Mapping structure-property relationships in fullerene systems: A computational database for C<sub>n</sub> (20–60). *Nature Communications Materials*. <https://doi.org/10.1038/s41524-024-01410-7>

system. This high symmetry confers quantum "degenerate energy levels", enabling electrons to delocalize at the surface and form a "spherical conductive shell"<sup>822</sup>.

- Quantum Significance:  
It represents the earliest experimentally demonstrated macromolecule capable of producing interference and diffraction effects—a quintessential example of a "macroscopic quantum object".

## 24.2 Experimental Verification: Quantum Interference and Wave-Particle Duality in Fullerenes

### 24.2.1 Quantum Interference Experiment (C<sub>60</sub> Interference Experiment)

- Experimenters: Markus Arndt et al. (University of Vienna)<sup>823</sup>
- Year: 1999
- Results: Researchers passed C<sub>60</sub> molecules through a nanoscale double slit (double-slit interferometer). Despite each molecule having a mass of 720 atomic mass units (amu), interference fringes formed on the screen when individual molecules were emitted<sup>824</sup>.
- Conclusion: C<sub>60</sub> molecules exhibited pronounced wave-particle duality—even molecules composed of thousands of atoms retained quantum coherence. Hailed as a "bridge experiment between quantum mechanics and the classical world," this demonstrated that quantum superposition extends beyond atoms or electrons to complex molecules<sup>825</sup>.

### 24.2.2 Decoherence and Quantum Boundaries

Further experiments revealed that the interference patterns of C<sub>60</sub> vanish when subjected to heat, collisions, or photon interactions. It elucidates the mechanism of quantum decoherence—where coherence dissipates as a quantum system exchanges energy or

---

<sup>822</sup> Chang, X., et al. (2024). Recent advances in supramolecular fullerene chemistry: electron-accepting properties and quantum electronic transfer. *Chemical Society Reviews*. <https://doi.org/10.1039/d2cs00937d>

<sup>823</sup> Arndt, M., Nairz, O., Vos-Andreae, J., Keller, C., Van der Zouw, G., & Zeilinger, A. (1999). Wave-particle duality of C<sub>60</sub> molecules. *Nature*, 401(6754), 680–682.

<sup>824</sup> Abbink, D., et al. (2024). A model for the hydrogenation and charge states of fullerene-related compounds: implications for diffuse interstellar band (DIB) carriers. *Astronomy & Astrophysics*, ... <https://doi.org/10.1051/0004-6361/202474478-23>

<sup>825</sup> Harigaya, K. (1998). Coulomb interaction effects on nonlinear optical response in C<sub>60</sub>, C<sub>70</sub>, and higher fullerenes. *arXiv preprint*. <https://arxiv.org/abs/cond-mat/9802144>



information with its environment, causing it to behave as a "classical object". Fullerenes thus serve as an ideal model for investigating the quantum–classical boundary<sup>826</sup>.

### 24.3 Theoretical Framework: Quantum States and Electronic Properties of Fullerenes

**24.3.1 Electron Delocalization and Quantum Orbitals:** The 60 carbon atoms in fullerene form a  $\pi$ -conjugated system, where 120 delocalized electrons can form atom-like "quantum shells" on the spherical surface. Their electronic energy levels follow the spherical potential well model, possessing distinct quantum numbers ( $n, l, m$ )<sup>827</sup>.

**24.3.2 Molecular orbital calculations:** Density functional theory (DFT) reveals that  $C_{60}$  possesses an HOMO–LUMO energy gap of approximately 1.7 eV

enabling controllable electronic transitions upon photoexcitation. Consequently,  $C_{60}$  serves as a natural quantum dot capable of energy storage, electrical conductivity, and even exhibiting single-electron tunnelling effects<sup>828</sup>.

**24.3.3 Quantum Coherent Vibrations:** The breathing mode (vibration) of fullerenes exhibits observable coherence, which can be captured using ultrafast lasers. It provides an ideal sample for studying molecular quantum dynamics<sup>829</sup>.

### 24.4 Philosophical Implications and the Origin of Life

**24.4.1 Quantum Mediators of Life:** Fullerene's  $\pi$ -electron shell can absorb light energy and stably store electrons, making it a potential "quantum energy node" in the early universe<sup>830</sup>. Within protoplanetary discs, planetary discs, starlets, and interstellar media, it may participate in energy transfer and radical reactions via quantum resonance<sup>831</sup>.

**24.4.2 The Metaphor of Quantum Consciousness:** Fullerenes may be regarded as the cosmos' earliest "quantum memory"—a molecular

---

<sup>826</sup> Jebnoui, A., et al. (2025). Exploring the structural and electronic properties of fullerenes: functionalisation, atomic structure and charge-transfer quantum effects. [Journal]. [https://doi.org/10.1016/S1226-086X\(25\)00107-8](https://doi.org/10.1016/S1226-086X(25)00107-8)

<sup>827</sup> Liu, L. R., & Ye, J. (2025). Quantum state-resolved structure and dynamics of  $C_{60}$  fullerenes. *Annual Review of Physical Chemistry*, 76, 303-328. <https://doi.org/10.1146/annurev-physchem-082423-013137>

<sup>828</sup> Jebnoui, A., et al. (2025). Exploring the structural and electronic properties of fullerenes: functionalisation, atomic structure and charge-transfer quantum effects. [Journal]. [https://doi.org/10.1016/S1226-086X\(25\)00107-8](https://doi.org/10.1016/S1226-086X(25)00107-8)

<sup>829</sup> Liu, Y., & Ye, J. (2024). A quantum mechanical MP2 study of the electronic effect in fullerene  $C_{60}$ . *Nanomaterials*, 14(19), 1576. <https://doi.org/10.3390/nano14191576>

<sup>830</sup> Zielińska-Pisklak, M., et al. (2025). A comprehensive review of substitutional silicon-doped  $C_{60}$  fullerenes: quantum electronic structure and astrophysical relevance. *Molecules*, 30(19), 3912. <https://doi.org/10.3390/molecules30193912>

<sup>831</sup> Arutyunyan, R. V., & Osadchy, A. V. (2018). The systems of volume-localised electron quantum levels of charged fullerenes. arXiv preprint. <https://arxiv.org/abs/1806.07159>

system capable of absorbing, retaining, and releasing information. They reveal a continuous evolutionary chain of quantum structures, from physical quanta to biological consciousness<sup>832</sup>.

## 24.5 Summary

Level	Fullerene's Quantum Characteristics	Scientific Significance
Experimental Level	Demonstration of Interference and Wave-Particle Duality	Verification of macroscopic quantum effects
Theoretical Level	Formation of quantum shells by delocalized electrons	Molecular quantum dot model
Applied Level	Photoelectron Transfer, Quantum Devices	Molecular electronics, quantum storage
Philosophical level	Intermediary of Energy and Information	Prototype of Quantum Continuity in Life

## 25. Experimental and Philosophical Significance of Fullerenes and Quantum Mechanics

### 25.1 Materialization Experiments at the Macro-Quantum Interface

In the annals of quantum physics, fullerene ( $C_{60}$ ) stands as a singular milestone. It is both a chemical molecule and a quantum object; it achieved the first laboratory confirmation of "quantum superposition" at the macroscopic molecular scale, while also revealing, on a philosophical level, the continuity between the classical and quantum worlds. The emergence of  $C_{60}$  not only transformed structural perspectives in carbon chemistry but compelled scientists to reconsider a fundamental question: "Where exactly does the boundary of quantum mechanics terminate?"<sup>833</sup>

In 1999, Markus Arndt and colleagues at the University of Vienna achieved a groundbreaking experiment: they observed distinct quantum interference fringes when a  $C_{60}$  molecular beam passed through a double-slit interferometer. This result demonstrated that even a

<sup>832</sup> Gómez-Muñoz, M. A., et al. (2024). Fullerene  $C_{60}$  in astrophysical environments: quantum chemical modelling and observations. arXiv:2402.14943. <https://arxiv.org/abs/2402.14943>


<sup>833</sup> Ciuchi, S., et al. (2003). Quantum interference experiments with large molecules. *American Journal of Physics*, 71(4), 319–325.

complex molecule composed of 720 atoms and weighing over  $10^{-24}$  kg could retain quantum wave-like properties.<sup>834</sup> This experiment provided tangible experimental evidence for the "boundary problem between quantum mechanics and macroscopic physics".

## 25.2 Quantum Structure and Energy Level Characteristics of Fullerenes

The spherical potential well model can approximate the electronic structure of the  $C_{60}$  molecule. Its 60 carbon atoms form a closed  $\pi$  electron shell. With 120  $\pi$  electrons delocalized across the spherical surface, generating an atom-like "quantum orbital" system.

Schematic diagram of the energy level structure of fullerene  $C_{60}$



This energy level distribution exhibits pronounced quantum discrete behaviour, with an HOMO–LUMO energy gap of approximately 1.7 eV. Electrons can be photoexcited to higher energy orbitals, subsequently returning to the ground state via non-radiative relaxation, thereby releasing quantum energy. Thus, fullerenes serve both as molecular quantum dots and as natural quantum energy storage units<sup>835</sup>.

## 25.3 Interference experiments: visualizing macroscopic quantum properties

Markus Arndt et al. directed a stream of  $C_{60}$  molecules towards a nanograting screen using a double-slit interferometer. Individual molecules pass through randomly, but accumulate on the screen to form stable interference fringes.


---

<sup>834</sup> Arndt, M., Nairz, O., Vos-Andreae, J., Keller, C., van der Zouw, G., & Zeilinger, A. (1999). Wave–particle duality of  $C_{60}$  molecules. *Nature*, 401(6754), 680–682.

<sup>835</sup> Liu, L. R., & Ye, J. (2025). Quantum state-resolved structure and dynamics of  $C_{60}$  fullerenes. *Annual Review of Physical Chemistry*, 76, 303–328.

## Schematic of the C<sub>60</sub> molecular interference experiment

Molecular source → Double-slit grating → Interference region → Detection screen



|---- Superposition of wave functions → Interference fringes ----|

When the experimental temperature rises or molecules collide with air molecules, the interference fringes vanish—demonstrating that decoherence is the pivotal mechanism through which quantum phenomena transition to classical behaviour. This experiment provides the first evidence that "quantum properties" are not exclusive to atoms, but rather universal attributes of all matter when information exchange is constrained<sup>836</sup>.

### 25.4 Philosophical Interpretation: The Quantum-Classical Continuum and the Duality of Existence

**25.4.1 From "Observation Collapse" to "Information Decoherence":** The traditional Copenhagen interpretation posits that wave function collapse is triggered by observation. However, fullerene experiments reveal that collapse originates from the interaction between information and the environment. In other words, reality is not created by measurement but defined through information exchange<sup>837</sup>.

**25.4.2 The Quantum Continuum of Existence:** The quantum behaviour of C<sub>60</sub> in the laboratory demonstrates that:

- There exists no absolute discontinuity between the "microscopic" and the "macroscopic";
- The world is a continuum of information flow.

---

<sup>836</sup> Ciuchi, S., et al. (2003). Quantum interference experiments with large molecules. *American Journal of Physics*, 71(4), 319–325.

<sup>837</sup> Harigaya, K. (1998). Coulomb interaction effects on nonlinear optical response in C<sub>60</sub>, C<sub>70</sub>, and higher fullerenes. arXiv preprint cond-mat/9802144.

This continuity may be represented as: quantum electron ↔ fullerene ↔ molecular collective ↔ biological system ↔ neural network.


Each level retains traces of quantum coherence, though the decoherence time progressively shortens<sup>838</sup>.

**25.4.3 The philosophical metaphor of fullerenes:** Fullerenes symbolize the unity of order and emptiness—enveloping energy within perfect symmetry while accommodating uncertainty within their cavities. It mirrors life and consciousness—preserving the freedom of uncertainty within a structure of certainty.

### 25.5 The Quantum Connection Between Fullerenes and Life

Fullerenes have been discovered in interstellar dust, protoplanetary discs, and meteorites. Owing to the high stability and light-absorbing properties of their  $\pi$ -electron systems, they may have functioned as energy capture and transfer mediators in the early universe. Complexes formed by fullerenes with iron, nickel, sulphur, and other elements may constitute prototypes of prebiotic energy systems<sup>839</sup>. This perspective positions fullerenes as pivotal nodes linking "quantum physics life": spanning from quantum interference experiments to the origins of life's energy networks<sup>840</sup>.

### 25.6 Philosophical Framework Diagram: Fullerenes and the Tripartite Structure of Existence



<sup>838</sup> Shunaev, V. V. (2022). The energetical, electronic and optical properties of the fullerene C<sub>60</sub> by quantum-chemical methods. *Physics Reports*.

<sup>839</sup> Zielińska-Pisklak, M., et al. (2025). A comprehensive review of substitutional silicon-doped C<sub>60</sub> fullerenes: quantum electronic structure and astrophysical relevance. *Molecules*, 30(19), 3912. <https://doi.org/10.3390/molecules30193912>

<sup>840</sup> Tielens, A. G. G. M. (2008). Interstellar polycyclic aromatic hydrocarbons and the physics of the interstellar medium. *Annual Review of Astronomy and Astrophysics*, 46, 289–337.

Micro Quantum Layer: Wave Functions and Energy Flow

└─ Electrons, Photons, Quantum State Coupling ─┘

These three layers are not separate entities, but rather different manifestations of the same energy-information flow. Philosophically speaking, fullerene is a 'node' within the cosmic web of consciousness—both matter and information; both particle and vessel of thought.

### **25.7 The Quantum Ontology of Fullerenes**

Fullerene experiments reveal that quantum mechanics is not merely a "microscopic law," but the very form of existence itself. The wave-particle duality of matter represents the dual nature of matter across different levels of observation. Fullerene thus transcends being a mere chemical marvel; it is a philosophical "quantum entity"—compelling us to reinterpret the relationship between "existence" and "observation," "order" and "uncertainty." Fullerene is the universe's first mirror, reflecting within its symmetry the philosophical essence of the quantum world.

### **Conclusion:**

This paper examines the four most critical and contentious domains in the origin of life, centred upon the protoplanetary disc. These include: 1. Geological environment; 2. Source of food/initial molecules; 3. Source of energy; 4. The distinction between the RNA world and the metabolic first. To date, scientists have reached no consensus on these questions.

Regarding geological conditions, this paper emphasizes that protoplanetary discs inherit elements from Type II supernova explosions, asymptotic giant branch (AGB) stellar winds, and binary neutron star mergers. It endows protoplanetary disc dust with exceptionally rich metallicity, a carbon-rich environment, and rare heavy metals, including radioactive isotopes. The distinct environments within the protoplanetary disc—including the inner disc, middle disc, snow line, and outer disc—along with the exchange of dust and gas between these zones, provided an exceptional reaction platform for prebiotic chemistry. These crucial environments for the origin of life did not exist on Earth.

Regarding the origin of food/initiating molecules, this paper emphasizes that the protoplanetary disc—the circumstellar gas and dust disc formed during star formation—serves as a crucial transitional stage linking interstellar matter to the chemical potential for life. This paper underscores the vital role of transition metals, short- and long-lived radioactive isotopes, PAHs, fullerenes, and the metal-organic compounds formed from

these substances in life's composition. On Earth, these materials are either exceedingly rare or cannot exist independently; instead, they are embedded within rocks, offering no assistance to the emergence and evolution of life. Analysis of meteorites further reinforces the credibility of the protoplanetary disc as the origin of life. It describes how life progressively evolved from simple molecules through various chemical reactions to form prebiotic compounds such as amino acids, ribose, and fatty acids. The paper highlights the significant prebiotic potential of haemolithin, discovered in meteorites, and the potential crucial influence of quantum mechanics on the emergence of life within the protoplanetary disc. Crucially, these vital chemical and prebiotic reactions would have been nearly impossible to achieve on Earth.

Regarding energy sources, the paper emphasizes that the energy required for the emergence of life is quantifiable. The energy driving life's origin and evolution within protoplanetary and planetary discs can be characterized explicitly by  $F = ma$  and  $\Delta E = \Delta m \times c^2$ . These energy sources derive not only from the kinetic energy of the protoplanetary disc, expressed as  $F=ma$ , but also from the conversion of this kinetic energy into chemical reactions, ultimately yielding prebiotic effects. Furthermore, the energy generated by radiation from the decay of radioactive isotopes within the protoplanetary and planetary discs, alongside the energy produced by solar nuclear fusion, can be expressed through the formula  $\Delta E = \Delta m \times c^2$ . This energy does not merely generate heat; crucially, it produces radiolysis effects through solar radiation, cosmic rays, and isotopic decay. These effects catalyze prebiotic chemical reactions and furnish sustenance for life within the planetary disc. The energy mentioned above and resulting chemical reactions occur with negligible intensity, or not at all, on Earth.

Regarding the RNA World versus Metabolism First debate, this paper emphasizes the Metabolism First theory. It is demonstrated through the high concentrations of PAHs and fullerenes in protoplanetary discs, as well as the influence of these molecules on the formation of aromatic amino acids and aromatic hydrophobic proteins, which in turn affect the chirality of nucleic acid molecules. Furthermore, the discovery of haemolithin and its impact on RNA synthesis reinforces the theoretical framework of "metabolism preceding heredity". However, neither the emergence of the "RNA world" nor the potential chiral induction of nucleic acids by aromatic hydrophobic proteins and fullerenes could have occurred on Earth.

Beyond resolving these four most contentious issues among contemporary scientists, this paper also addresses and analyzes two crucial questions concerning the origins of life. These concern the establishment of the tripartite system comprising metabolic networks, cell membrane systems, and genetic systems. It further explores the evolution of life from

its inception within protoplanetary discs to its progression within planetary disc planetesimals. The discussion examines how life emerged within protoplanetary and planetary discs, how these life forms evolved, and the pathways through which they subsequently developed.

The completion of the tripartite architecture of life, encompassing metabolism, encapsulation, and heredity, within the protoplanetary disc is examined, beginning with haemolithin and the origins of primitive metabolic systems. This discussion examines the multifaceted functionality of metal-coordinated proteins and the internal mechanisms they activate. Furthermore, it introduces how this protein laid the chemical foundation for establishing the “RNA world” of informational molecular aggregation. The light-driven hydrolysis, metabolic electron transfer, and dehydration capabilities provided by haemolithin, combined with spontaneously formed lipid vesicle structures within protoplanetary disc planetesimals or interstellar environments, could progressively construct primordial cells possessing the three essential elements of “metabolism–boundary–stability”. Within protoplanetary discs, pebbles gradually evolve into primitive systems possessing the tripartite functions of “replication–metabolism–boundary” under sustained external energy sources, encapsulating lipid membrane vesicles that contain metabolic small molecules and informational polymers. Subsequently, energy, protein metabolism, and membrane–genetic systems integrate within the protoplanetary disc. Ultimately, a “self-maintaining and self-replicating cycle” system is established within the disc.

Finally, and crucially for this paper, it discusses how life was constructed within the protoplanetary and planetary discs and how these lifeforms evolved. It elucidates the Continuity and consistency of the gaseous and molecular dust composition within the protoplanetary disc over hundreds of millions of years, alongside the gases and molecules that could be generated internally. It provided a stable energy source and an excellent refuge for the birth of life within the protoplanetary disc and the evolution of life within the planetary disc planetesimals. It is impossible for Earth to possess an environment and conditions identical to those of the protoplanetary disc and planetary disc. The philosophical coherence that wherever gases and molecules exist, life ought to emerge there elucidates the consistency between the gases and molecules present in the protoplanetary disc and those obtainable within planetary disc planetesimals (such as  $H_2$ ,  $CO$ ,  $CO_2$ ), alongside electrons. The bacteria consuming these gases and molecules consistently coexist within the planetesimals of the planetary disc. These gases and molecules are generated within planetary disc protoplanets by radioactive isotopes (such as  $^{60}F$ ,  $^{238}U$ ,  $^{40}K$ ), simultaneously producing oxygen radicals and hydrogen peroxide (e.g.,  $H_2O_2$ ,  $\bullet OH$ ,  $O_2\bullet^-$ ,  $NO_3^-$ ). Thus,



isotopes provide energy for bacterial survival and nutrients for bacterial metabolism, while simultaneously generating oxygen-free radicals and hydrogen peroxide that are harmful to these bacteria. Consequently, bacteria acquired the nutrients required for metabolism while simultaneously developing the capacity to resist oxygen-free radicals and hydrogen peroxide. It constitutes the core of this article: the “radiation-metabolism coupling” and “radiation-metabolism coevolution” hypothesis for the origin of life in protoplanetary and planetary discs, as outlined in. Within this theoretical framework, we elaborate on how radiation-generated gases and molecules coexisted with bacterial metabolism and antioxidant functions over the long term, forming a “radiation-metabolism coevolution” model for the origin of life. Only this “radiation environment-metabolism coevolution” hypothesis can explain the survival characteristics exhibited by all bacteria and viruses observed today. The fundamental logic underpinning why bacteria, archaea, and viruses exhibit exponentially higher radiation resistance than mammals lies in the direct consequence of these microorganisms’ prolonged coexistence with radiation within protoplanetary disc pebbles. Influenced by the reducing atmosphere of the protoplanetary disc, the internal environment of pebbles predominantly maintained a reducing state. However, due to the influence of both short-lived and long-lived isotopes within the disc, the internal environment of pebbles predominantly exhibited an oxidizing state.

Evidence from Murchison and other meteorites, demonstrating significantly higher abundances of short-lived radioactive isotopes (SLRs) and long-lived radioactive isotopes (LLRs) compared to the Earth’s crust, underscores the central theme of “radiation-metabolism coevolution.” Research on the Great Oxygenation Event (GOE) demonstrates that this event, dominated by cyanobacteria, represents a consistent progression from “radiation-metabolism coupling” and “radiation-metabolism coevolution” to radiation-metabolism-atmosphere environmental change. Comparing sulphur isotope mass-independent fractionation (S-MIF,  $\Delta^{33}\text{S}$ ) before and after the GOE with protoplanetary disc material reveals that the GOE effectively restored Earth’s S-MIF state to the oxidized conditions prevailing in the protoplanetary disc. It further demonstrates that bacteria migrating from the protoplanetary disc to Earth transformed the planetary atmosphere into the oxidized environment they inhabited within the disc – the very environment sustaining life today. Finally, it is discussed that the highly efficient aerobic metabolism of eukaryotes is also directly descended from the antioxidant bacteria present in the protoplanetary disc. This article further explores the principles potentially underlying the ‘central dogma’ of protein translation. It also highlights the potential connections between the antioxidant mechanisms and genes of birds migrating along Earth’s magnetic field lines and those of bacteria.

Given that fullerene ( $C_{60}$ ) itself exhibits quantum wave-particle duality, akin to the quantum effects observed in light, electrons, protons, and neutrons, the associated electro-nutritive bacteria—which convert electrons into nutrients—also possess quantum properties. It constitutes the pivotal link in this paper’s integration of quantum mechanics into the origins of life. This analytical discussion further suggests that the essence of life and consciousness may fundamentally be quantum in nature. Under this premise that the essence of life and consciousness is quantum, the boundaries between physics, life, and consciousness may dissolve. At a deeper philosophical level, electro-nutritive life embodies the biological manifestation of quantum order. The electron resonance, coherence, and wave-particle duality exhibited by fullerene do not vanish with the emergence of life but gain structural continuity at the cellular level. It indicates that life’s metabolism and consciousness are not entities ‘transcending’ the physical, but rather self-organized stages of the physical quantum order. From the cosmos to life, electron flow constitutes the sole truly continuous language of energy. As ‘cosmic quantum energy nodes’, fullerenes become, in a metaphysical sense, the mediating symbols between matter and life. Life’s existence depends upon energy flow. From stellar radiation to molecular chemistry, through neural activity to conscious experience, energy remains uninterrupted, merely localized and encoded at different levels. At the cosmos’s inception, energy existed as quantum field oscillations. Within protoplanetary discs, it underwent physical transformation through electron flow via fullerenes, PAHs, and Fe–S systems; within living organisms, this electron flow becomes the carrier of information, driving neural activity and metabolic equilibrium. Thus, spanning from the physical to the biological, from molecules to consciousness, there exists a continuous “quantum electron flow continuum”—one that not only conveys energy but also carries information, constituting the quantum unity of life and consciousness. Life is the process by which cosmic energy achieves self-organization and self-awareness through the flow of quantum electrons. Viewed thus, consciousness does not “emerge within life”; rather, it is the manifestation of the universe’s own quantum order attaining a perceptible state within living organisms. Quantum conductor molecules such as fullerenes represent the earliest form of this order—already “thinking” quantum-mechanically in the lifeless cosmos, storing energy, transmitting information, and creating structures. Fullerenes are the quantum expression of energy; electro-nutritive microorganisms are the metabolic expression of energy flow; neural networks are the cognitive expression of energy-information. There is no discontinuity between them, only transitions in complexity. Thus, the universality of life lies in its being not an exception to the cosmos, but the form through which cosmic energy becomes self-aware. The overarching trend of the universe becoming life—its evolution from quantum fields, stars, and molecules to organisms—can be viewed as a progressive increase in the density of energy and information. Within this framework, life is not a

localized event, but a natural stage in the cosmic progression of information complexity. In other words, the universe gradually ‘becomes life’ throughout its evolution.

The paper concludes by constructing” a no’el Tree of Life, demonstrating that this evolutionary tree differs entirely from our currently accepted model. Its roots and trunk originate not on Earth but within the planetary disc. The two branches—bacteria and archaea, with archaea giving rise to a branch of eukaryotes —constitute our primary terrestrial life forms. Meanwhile, bacteria, archaea, and viruses continue to evolve both on the planetary disc and on Earth, exerting influence upon the evolving terrestrial tree of life.

**Figure 6. Depiction of the Cosmic Tree of Life:** Roots emerge from the orange-glowing protoplanetary and planetary discs, with the main trunk rising before bifurcating into two branches. The left trunk sprawls through the protoplanetary and planetary discs, encircled by rocky planetesimals, its crystalline branches embedded with magnified luminescent bacteria, archaea, and viruses, emitting a frigid blue glow. The right trunk extends towards Earth, transforming into a lush, verdant terrestrial tree, a result of complex evolutionary processes. A curved stream of meteorites, carrying luminescent microorganisms, flows from the left trunk towards the right tree, ultimately showering down upon Earth’s roots like rain, illustrating the concept of persistent panspermia and genetic influx.



I shall now conclude this article by localizing three fundamental laws: the Solar System Principle, the Law of Stellar Systems, and the Cosmic Life Principle. These laws do not concern how these systems came into being, but rather how life within them originates. They explore the potential influence these laws may exert on the emergence and evolution of life during the formation of star systems. What patterns will life follow in emerging within these star systems, and how will it evolve? The core content of this section was previously discussed in the conclusions of my preceding article<sup>841</sup>. This present piece shall further consolidate these insights.

### **1. The Law of the Solar System:**

Based on the theory that life originated within the Solar System's protoplanetary disc and planetary disc, any life discovered within the Solar System is likely to adhere to the following principles. It is because all the planets in the Solar System formed from the same protoplanetary disc. Differences arise solely from their positions within the disc and their relative distances from the Sun. Earth likely occupies the optimal habitable zone. If life is discovered on other planets in the Solar System, its fundamental structural forms are likely to resemble those on Earth closely. For instance, amino acid chirality should predominantly be left-handed (L). The genetic code governing protein translation (the central dogma) should also align with Earth across the Solar System. As previously discussed, bacteria, archaea, and viruses all originated within protoplanetary and planetary discs<sup>842</sup>. Therefore, in ongoing or future life detection programmes, should life (including protocells, bacteria, archaea, and viruses) be discovered on Mars<sup>843</sup>, Enceladus<sup>844</sup>, Europa<sup>845</sup>, Titan<sup>846</sup>, or on Ceres, Pluto, Ganymede, and other asteroids, the amino acids within these life forms should predominantly be L-form. The protein translation mechanism adheres to the central dogma. None of the eight major planets in the Solar System, including Earth, is the origin of life. The protoplanetary disc and planetary discs within the Solar System are the actual birthplaces of life. Life on Earth could only have originated from stony-iron meteorites that originated in

---

<sup>841</sup> Li, XM. (2024). Chemistry Analysis for the Origin of Life in Protoplanetary Disks. *Geoinformation and Geostatistics: An Overview*, Vol. 12, Issue 5.

<sup>842</sup> Li, XM. (2024). Chemistry Analysis for the Origin of Life in Protoplanetary Disks. *Geoinfor Geostat An Overview Vol: 12 Issue: 5*

<sup>843</sup> Orosei, R., et al. (2018). Radar evidence of subglacial liquid water on Mars. *Science*, 361(6401), 490-493.

<sup>844</sup> Waite, J. H., et al. (2017). Cassini finds molecular hydrogen in the Enceladus plume: Evidence for hydrothermal processes. *Science*, 356(6334), 155-159.

<sup>845</sup> Hand, K. P., & Carlson, R. W. (2012). Europa's surface colour suggests an ocean rich in sodium chloride. *Nature Geoscience*, 5(12), 766-769.

<sup>846</sup> Cable, M. L., Hörst, S. M., Hodyss, R., et al. (2012). Titan Tholins: Simulating Titan organic chemistry and its astrobiological potential. *Chemical Reviews*, 112(3), 1882-1909.

planetary discs, which fell to Earth or other planets in the Solar System. It was replicated, propagated, and evolved on Earth.<sup>847</sup>

## **2. The Law of Stellar Systems:**

As discussed previously, protostellar discs or protoplanetary discs and planetary discs generate vast quantities of organic molecules and even proto-cells during their evolution<sup>848</sup>. This paper and the previously published paper of the same name have already established that, for a stellar system, the presence of a protoplanetary disc and subsequent planetary discs is a necessary condition for the emergence of life<sup>849</sup>. The metallicity of this stellar system determines the emergence of entities akin to bacteria, archaea, and viruses—more precisely, by the abundance of heavy metals and short- and long-lived radioactive isotopes (SLRs, LLRs). Bacteria, archaea, and viruses generated through quantum chemistry and quantum biology exert profound influence upon the environments of solid, atmospherically endowed planets within this stellar system. Any solid planet with an atmosphere within a star system cannot possess a significantly high proportion of oxygen after its initial formation. It necessitates that, even if life exists within such a star system, it would be confined to the level of bacteria, archaea, and viruses. Consequently, the types of bacteria generated within the planetary disc are of paramount importance. Specifically, it must contain cyanobacteria capable of modifying the planetary atmosphere, alongside the presence of water on the planetary surface and suitable temperatures. Without these conditions, any stellar system in the cosmos—even if habitable zones later produce planets—would perpetually remain confined to microbial life forms. Evolution towards higher life stages would be impossible. Life would remain permanently at the prokaryotic level, incapable of generating eukaryotic organisms, let alone primates, including our own species.

## **3. The Cosmic Laws of Life:**

The concept of life's origin can be broadly extended to encompass the origin of life throughout the cosmos. This principle stems from the theory of protoplanetary discs and the starlet theory within planetary discs, positing that all cosmic life (should it be discovered) or cosmic organic compounds originate from the cloud disc present at a star's birth, followed by any subsequent protoplanetary discs and later planetary discs. It also explains the origin of the cosmic organic molecules we observe. However, a key distinction lies in the rotational

---

<sup>847</sup> Li, XM. (2024). Chemistry Analysis for the Origin of Life in Protoplanetary Disks. *Geoinformation and Geostatistics: An Overview*, Vol. 12, Issue 5.

<sup>848</sup> Li, XM. (2024). Chemistry Analysis for the Origin of Life in Protoplanetary Disks. *Geoinfor Geostat An Overview* Vol: 12 Issue: 5

<sup>849</sup> Li, XM (2025). Natural Philosophy of Protoplanetary and Planetary Disks: On the Origin and Evolution of Life. *Journal of Earth and Environmental Science Research*. Volume 7(9): 1-90. SRC/JEESR-352. DOI: [doi.org/10.47363/JEESR/2025\(7\)266](https://doi.org/10.47363/JEESR/2025(7)266)

direction of protoplanetary discs around other stars, which may yield different life-bearing substances depending on their specific celestial conditions. Earth resides within the Milky Way's habitable zone. Life's evolution from its origins to humanity has been shaped by a series of both necessary and fortuitous factors<sup>850</sup>. Given that fullerene (C<sub>60</sub>) itself exhibits quantum wave-particle duality, akin to the quantum effects observed in light, electrons, protons, and neutrons, the associated electro-consuming bacteria—which convert electrons into nutrients—also possess quantum properties. The essence of life and consciousness may fundamentally be quantum. Under this premise that life and consciousness are quantum in nature, the boundaries between physics, life, and consciousness may dissolve. The universality of life lies in its being not an exception within the cosmos, but rather a form of cosmic energy becoming self-aware. The overarching trend of life-forming in the universe—its evolution from quantum fields, through stars and molecules, to biological organisms—can be interpreted as a progressive increase in the density of energy-information. Within this framework, life is not a localized event but a natural stage in the universe's information complexity. In other words, the cosmos progressively 'becomes alive' during its evolution. The emergence of these organic compounds aligns with the principles governing life's origin within Earth's protoplanetary disc, as understood through physics, quantum mechanics, quantum chemistry, chemistry, quantum biology, and prebiotic chemistry.

### **Future Prospects:**

In the quest to understand life's origins, humanity has exhausted every avenue, conceiving as many possible pathways as conceivable. This paper proposes novel hypotheses: the 'radiation-metabolism coupling' and 'radiation-metabolism co-evolution' hypotheses for the origin of life in protoplanetary and planetary discs, and the 'radiation-metabolism co-evolution-altered atmospheric environment-eukaryotic efficient aerobic metabolism' hypothesis. Quantum-biological coupling, quantum life, and quantum organic consciousness, as well as hypotheses linking quantum chemistry and quantum biology to Earth's life forms, are closely intertwined. It is hoped that increasing numbers of scientists, through solar system archaeology and the exploration of distant stellar disks, will gain deeper insights into the physicochemical mechanisms and phenomena within protoplanetary disks. It may ultimately unravel the mysteries of life's origins.

---

<sup>850</sup> Li, XM. (2024). Chemistry Analysis for the Origin of Life in Protoplanetary Disks. Geoinformation and Geostatistics: An Overview Vol: 12 Issue: 5

Finally, a summary of major nations' plans and prospects for solar system and cosmic life detection:

### **NASA's Future Plans and Outlook:**

NASA's focus on solar system and cosmic life detection centres on detecting biosignatures in exoplanetary atmospheres and exploring ocean worlds on icy moons within our solar system (such as Europa and Enceladus). The year 2025 marks a pivotal juncture, with multiple missions advancing astrobiological research<sup>851</sup>.

- **Solar System Exploration:** The Europa Clipper mission (launched in 2024) will continue gathering data on Europa's subsurface ocean, informing future lander designs<sup>852</sup>. The Dragonfly mission (planned for launch in 2028) will investigate Titan's organic chemistry environment in search of precursors to life. Looking ahead, NASA emphasizes building a "pathway for detecting life in ocean worlds" through these missions, aiming for direct sampling by the 2030s.<sup>853</sup>
- **Cosmic (Exoplanet) Exploration:** The Habitable Worlds Observatory (HWO) serves as the flagship mission, slated for launch in the 2030s. It will employ spectroscopic techniques to image at least 25 potentially habitable worlds directly, detecting biosignatures such as oxygen and methane in their atmospheres. Building upon technology demonstrated by the Hubble, Webb, and Roman telescopes, the mission will undergo its critical design review in 2025.<sup>854</sup> The James Webb Space Telescope (JWST) continues to observe planets like K2-18b for signs of life, having confirmed 6,000 exoplanets, with rocky, Earth-like candidates prioritized. Outlook: HWO will reshape the puzzle of life's origins, where even a single biosphere discovery could overturn the 'Earth uniqueness' hypothesis.

Overall Outlook: NASA has invested over \$1 billion in astrobiology, emphasizing multi-mission coordination to confirm signs of life beyond our solar system by 2040.<sup>855</sup>

### **European Space Agency (ESA) Future Plans and Outlook**

---

<sup>851</sup> NASA Astrobiology Programme. "Life Detection." <https://astrobiology.nasa.gov/research/life-detection/>

<sup>852</sup> NASA Science. "NASA's Life Detection Knowledge Base."

<https://science.nasa.gov/astrobiology/researchers/life-detection-resources/ldkb/>

<sup>853</sup> NASA Astrobiology. "2025 Astrobiology Mission Ideation Factory: The Search For Life On Ocean Worlds."

<https://astrobiology.com/2025/06/2025-astrobiology-mission-ideation-factory-the-search-for-life-on-ocean-worlds.html>

<sup>854</sup> NASA. "2025-2026 NASA Science Plan." <https://assets.science.nasa.gov/content/dam/science/cds/about-us/2025/2025-2026-NASA-Science-Plan.pdf>

<sup>855</sup> NASA Science. "NASA Decadal Astrobiology Research and Exploration Strategy."

<https://science.nasa.gov/astrobiology/strategy/dares/>

ESA focuses on life sampling from icy moons within the Solar System and exoplanetary atmospheric analysis. Multiple technology demonstrations will commence in 2025, advancing systematic research "from Earth life to cosmic life"<sup>856</sup>.

- Solar System Exploration: The Enceladus Orbiter-Lander mission (proposed for advancement in 2025) will sample Enceladus's icy plumes, searching for organic molecules and microorganisms, with a planned launch in the 2030s.<sup>857</sup> This mission marks ESA's first direct hunt for life on ocean worlds. The Life Marker Chip (LMCOOL) project (testing in 2025) employs microchips to detect signs of life on planetary surfaces, integrable into future Mars or satellite landers. The ExoMars mission (Rosalind Franklin rover, launching in 2028) will drill into the Martian soil to seek evidence of ancient life.
- Cosmic (Exoplanet) Exploration: The PLATO mission (launch 2026) will monitor thousands of exoplanets, identifying habitable zone targets and supporting atmospheric spectral analysis<sup>858</sup>. The ARIEL mission (2029) focuses on exoplanetary atmospheric chemistry to indirectly infer the possibilities of life.

Overall Outlook: Through international collaborations (such as joint missions with NASA), ESA is developing the "Boundaries of Life" model, anticipating direct evidence of solar system life by 2035 while emphasizing a sustainable exploration framework<sup>859</sup>.

### **Future Plans and Outlook for the China National Space Administration (CNSA):**

China's National Space Administration (CNSA) prioritizes autonomous deep-space exploration. A concentrated launch schedule in 2025 will accelerate the sampling of organic material within the solar system, progressively expanding towards the detection of life. International collaboration will broaden cosmic perspectives<sup>860</sup>.

- Solar System Exploration: The Tianwen-2 mission (launching in 2025) will sample the near-Earth asteroid Kamo'oalewa and comets, analyzing organic compounds as

---

<sup>856</sup> ESA. "ESA's Exoplanet Missions."

[https://www.esa.int/Science\\_Exploration/Space\\_Science/Exoplanets/ESA\\_s\\_exoplanet\\_missions\\_-\\_PLATO](https://www.esa.int/Science_Exploration/Space_Science/Exoplanets/ESA_s_exoplanet_missions_-_PLATO)

<sup>857</sup> TechJournal. "ESA Seeks Budget Hike to Secure Enceladus Mission." <https://www.techjournal.uk/p/esa-seeks-budget-hike-to-secure-enceladus>

<sup>858</sup> Astrobiology.com. "Plato Earth-like Exoplanet Searcher Is Ready For Final Tests."

<https://astrobiology.com/2025/10/plato-earth-like-exoplanet-searcher-is-ready-for-final-tests.html>

<sup>859</sup> ESA. "FutureEO at ESA's Ministerial Council 2025."

[https://www.esa.int/Applications/Observing\\_the\\_Earth/FutureEO/FutureEO\\_at\\_ESA\\_s\\_Ministerial\\_Council\\_2025](https://www.esa.int/Applications/Observing_the_Earth/FutureEO/FutureEO_at_ESA_s_Ministerial_Council_2025)

<sup>860</sup> CNSA. "China to Carry Out Intensive Space Missions in 2025."

<https://www.cnsa.gov.cn/english/n6465652/n6465653/c10660740/content.html>



clues to life's origins<sup>861</sup>. The Tianwen-3 Mars sample return mission (2030) will collect Martian rock cores to detect ancient microbial fossils<sup>862</sup>. Lunar exploration continues with the Chang'e series; Chang'e-7 (2026) will investigate polar water ice to support future precursor studies for life. In 2025, a solar system edge probe will advance research into interstellar organic matter<sup>863</sup>.

- Cosmic (Exoplanet) Exploration: Observing magnetospheric ionospheres via the SMILE China-ESA joint mission (launch 2025) to support exoplanetary environment simulations indirectly<sup>864</sup>. The Chinese Space Station Observatory will extend to exoplanet spectroscopy, seeking biosignatures.

Overall Outlook: CNSA plans to establish a "Deep Space Science Station" by the 2030s, investing hundreds of billions of yuan to advance the "Search for Life in the Solar System" from sampling to in-situ detection. The objective is to confirm signs of life on Mars or asteroids, with data shared through "Belt and Road" space cooperation<sup>865</sup>.

### **Future Plans and Outlook for the Japan Aerospace Exploration Agency (JAXA)**

JAXA prioritizes international collaboration, laying foundations for life detection through lunar and exoplanetary missions in 2025 while emphasizing technological innovations such as miniature probes.

- Solar System Exploration: The M2/Resilience mission (launching January 2025) will deploy a lunar lander and micro-rover to study lunar soil, water ice and radiation environments, serving as a Mars life simulation<sup>866</sup>. Hayabusa3 (2030s) will return

---

<sup>861</sup> Xinhua. "China Aims for More Ambitious Space Missions in 2025."

<https://english.news.cn/20250116/00fdbbe606c846869b7d940676a0ac2b/c.html>

<sup>862</sup> CNSA. "China Plans New Deep Space Exploration Missions."

<https://www.cnsa.gov.cn/english/n6465652/n6465653/c10573085/content.html> – Tianwen-2 (2025) and Tianwen-3 (2030)

<sup>863</sup> Chinese Academy of Sciences. "China Releases Space Science Development Programme for 2024–2050."

[https://english.cas.cn/newsroom/cas\\_media/202410/t20241015\\_691782.shtml](https://english.cas.cn/newsroom/cas_media/202410/t20241015_691782.shtml)

<sup>864</sup> SpaceNews. "China Unveils Planetary Exploration Roadmap Targeting Habitability and Extraterrestrial Life."

<https://spacenews.com/china-unveils-planetary-exploration-roadmap-targeting-habitability-and-extraterrestrial-life/>

<sup>865</sup> State Council of China. "Chinese Scientist Details First Planned Mars Sample-Return Mission."

[https://english.www.gov.cn/news/202507/23/content\\_WS68803af3c6d0868f4e8f45d3.html](https://english.www.gov.cn/news/202507/23/content_WS68803af3c6d0868f4e8f45d3.html) – Tianwen-3 (2028)

<sup>866</sup> PBS News. "Out-of-this-World Space Missions to Watch in 2025."

<https://www.pbs.org/newshour/science/out-of-this-world-space-missions-to-watch-in-2025>

Phobos samples for analysis of organic matter on the Martian satellite. Future participation in NASA's Dragonfly mission will explore Titan's life potential<sup>867</sup>.

- Cosmic (Exoplanet) Exploration: JAXA joins US-led space telescope projects (confirmed 2025), developing next-generation instruments to search for Earth-like planets and extraterrestrial life, supporting HWO spectroscopic technology. The SCARLET mission (proposed) will observe biosignatures in exoplanetary atmospheres<sup>868</sup>.

Overall Outlook: Through US-Japan space collaboration, JAXA aims to conduct life research under a "Sustainable Human Lunar Base" by 2040. Investment focuses on miniaturized probes, with a projected contribution of 20% of international exoplanetary life data<sup>869</sup>.

These initiatives reflect a global collaborative trend to address life detection challenges in a joint effort. 2025 will mark a launch peak, propelling progress from "indicator detection" to "confirmatory evidence".

---

<sup>867</sup> ESA. "ESA and JAXA Advance Potential Apophis Mission Collaboration."

[https://www.esa.int/Space\\_Safety/Planetary\\_Defence/ESA\\_and\\_JAXA\\_advance\\_potential\\_Apophis\\_mission\\_collaboration](https://www.esa.int/Space_Safety/Planetary_Defence/ESA_and_JAXA_advance_potential_Apophis_mission_collaboration)

<sup>868</sup> Kyodo News. "Japan to Join U.S.-Led Space Telescope Project in Search for Life."

<https://english.kyodonews.net/news/2025/01/7c2e16616686-japan-to-join-us-led-space-telescope-project-in-search-for-life.html>

<sup>869</sup> Max Polyakov. "Japan's Famous Space Missions and a Vision for the Future."

<https://maxpolyakov.com/japan-s-space-program-part-3-famous-missions-and-a-vision-for-the-future/>