

This is a non-peer-reviewed preprint submitted to EarthArXiv.

This manuscript has been submitted for publication in Nature Sustainability. Please note the manuscript has yet to be formally accepted for publication. Subsequent versions of this manuscript may have slightly different content. If accepted, the final version of this manuscript will be available via the 'Peer-reviewed Publication DOI' link on the right-hand side of this webpage. Please feel free to contact any of the authors; we welcome feedback.

Two-thirds of new fossil fuel infrastructure targets critical ecosystems, elevating health risks for millions

Ginni Braich^{1,*}, Katie Fankhauser¹, Laura Castrejon Violante¹, Naia Ormaza Zulueta¹, Stella Cooper², Candy Ofime², Isaiah Lyons-Galante¹, Varnitha Kurli¹, and Zia Mehrabi¹

¹Better Planet Laboratory, University of Colorado Boulder, Colorado, USA

²Amnesty International,11 W 43rd Street 7th Floor New York, NY 10036

*Corresponding author: Ginni Braich, gubr4186@colorado.edu

Abstract

10

11

12

13

14

16

17

18

19

20

21

23

24

25

26

27

28

29

30

31

32

33

While the fossil fuel industry's role in driving climate change is well established, the elevated risks to fenceline communities and critical ecosystems remain underexplored at a global scale, limited to a handful of prior coarse assessments. Here we map over 18,000 operating fossil fuel facilities across 170 countries, and assess their placement in surrounding populations and ecosystems, relative to empirical data on known elevated risk zones. We find that $\sim 32\%$ of facilities are located within critical ecosystems, including areas important to nature's contributions to people, key biodiversity areas, and critical irrecoverable carbon reserves, and that almost 463 million people, including over 120 million children, live within 1 km of these sites, placing them at heightened risk of health impacts. Looking forward, we assess 3,500 planned facilities and find that \sim 69% are slated for, or are already under construction in, critical ecosystems, disproportionately in Asia's biodiversity hotspots, and where an additional 106 million people in then-to-be fenceline communities will live. Together, these findings build on earlier studies and establish higher spatial resolution global estimates of communities and ecosystems at elevated risk from fossil fuel infrastructure. They underscore the urgent need for stronger environmental and health impact assessments, community-engaged health monitoring, and accountability frameworks to prevent further entrenchment of unfolding climate, ecological, and environmental justice crises on the frontlines of the fossil fuel industry globally.

1 Introduction

Fossil fuels account for 79% of global greenhouse gas emissions—an estimated \$14 trillion climate liability—and are responsible for one in five deaths worldwide

through air pollution exposure (1; 2; 3; 4). Yet, while fossil fuel contributions to climate change are well-documented, the immediate environmental and health catastrophes borne by surrounding fenceline communities remain systematically 38 under-examined. These local harms constitute ongoing public health emergencies. For instance, children living near refineries experience 50% higher asthma rates (5); communities near coal mines face toxic exposures linked to kidney failure and encephalopathy (6); and catastrophic accidents, such as the Texas City refinery explosion, have released thousands of pounds of carcinogens into nearby neighborhoods (7). These impacts are not isolated to one facility type. For example, oil shale processing in Estonia's Ida-Viru County drives elevated lung cancer rates (8), while routine spills along Trinidad's coast contaminate marine ecosystems with carcinogenic hydrocarbons, threatening food security 47 and human health (9). These are just some of the stories of a larger and more widespread set of risks posed to people and ecosystems around the world, the 49 scale of which has to date largely remained under-documented. 50

The harms posed to fenceline communities by the fossil fuel industry are not 51 only environmental, but also legal and political. The recognition of the human 52 right to a clean, healthy, and sustainable environment by the UN Human Rights 53 Council (2021) and General Assembly (2022) (10; 11) followed by landmark ad-54 visory opinions (12; 13) underscores a global obligation to respect, protect, and fulfill the right under penalty of what "may constitute an internationally wrongful act which is attributable to that State" (13). Yet, implementation is grossly inadequate: fewer than 2% of countries have binding human rights and en-58 vironmental due diligence laws (hereinafter "due diligence laws"), and violence against environmental defenders alone rose 150% between 2013-2023 (14). Weak 60 protections are particularly devastating in ecologically critical areas, where disasters cascade across ecosystems and livelihoods. For example, the Deepwater 62 Horizon spill contaminated an estimated 111,000 or km² of ocean (15), while recurrent oil spills in Nigeria have destroyed 45,000 hectares of mangroves and jeopardized the livelihoods of 150,000 fishers (16; 17). Indigenous peoples—who comprise less than 5% of the global population but steward 37% of Earth's remaining natural lands (18)—face especially acute risks: in Canada, half of all reserve-based Indigenous communities are at risk of pipeline spills, yet retain little control over decisions regarding extraction on their territories (19; 20).

Despite global calls for decarbonization, energy transitions are slow, uneven, and contested. Fossil fuel interests, disinformation campaigns, and deregulation continue to delay progress (21; 22; 23), and thousands of current and planned facilities will remain operational for decades. This reality emphasizes the need for systematic assessments of the ongoing risks that these facilities pose, and to identify not only which people and ecosystems are at risk today, but also those likely to be exposed in the near future through ongoing developments.

From an epidemiological perspective, a significant body of evidence documents the human health risks of living nearby facilities. This data exists across facility types (from extraction sites and mines, to pipelines, terminals, and plants), and

documents a vast variety of evidence on distance-based elevated risks in cognitive defects (24), neuro developmental delays (25; 26; 27; 28), respiratory illnesses (29), sleep disturbance (30), pre-term birth and low birth weight (25: 26: 27: 82 28), congenital anomalies (25; 26; 27; 28), and cancer risks (29; 31; 32; 33) for populations living within the vicinity of facilities, mediated through exposure 84 pathways such as polluted light, noise, soil, water, and air. That health risks are posed by this infrastructure is unequivocal, but the extent of adverse effects can vary depending on the pathway, with health impacts identified up to 10 km 87 from the infrastructure itself in some cases. Consistently, across studies, extreme 88 health risks have been documented for people living in close vicinity, with the highest risk for multiple facility types and health outcomes documented within 1 km (Table S1). A key question therefore, is how many people, children, or other 91 demographic groups reside in these high risk zones globally? Fundamentally, 92 another way to think about this, is "what is the estimated size of fenceline 93 communities linked to the fossil fuel industry?" Because the length of pipelines globally are so extensive we assess these exposure numbers with, and without, 95 pipelines.

From an ecological perspective, we currently have two sources of information on the potential risks fossil fuel facilities pose. The first are global, but spatially coarse (e.g. ~50 km resolution), assessments of potential risks from current and future infrastructure or concessions to biodiversity and ecosystem services (34; 35; 36; 37; 38; 39). These studies provide important first order assessments but are limited in their ability to identify localized risks and siting at fine resolution within landscapes. The second is a body of literature assessing impacts of different kinds of fossil fuel facilities on species and communities through multiple pathways, e.g. through light, noise, and air pollution or through transport based disruptions. Direct documented impacts include degradation of marine mammal communication and behavior from noise pollution (40), increased mortality for whales from ship strikes (41), reduced habitat usage, foraging, and reproduction in birds and ungulates from onshore drilling (42), disruption of animal movement from pipelines (43), reduced freshwater biodiversity from mine leaching (44), foliar damage to plants and reduced growth from air pollution (45; 46), disruption of species interactions and predator-prev relationships from artificial light (47; 48), and more. The reported distances over which these ecosystem impacts occur vary depending on facility and species: from the 25 m immediate disturbance footprints of pipeline right of way allocations (43) up to 2 km for marine disruptions to marine mammals (40) and large scale environmental disasters such as explosions or spills which can stretch for >100 km (49). Because of these differences, we localize our sampling strategy using a facility type specific approach (see Methods). The key question here is, "how many of the current and planned fossil fuel facilities are sited in ecosystems that could be classed as critical?"

100

101

102

103

104

105

106

107

108

110

112

113

114

115

116

117

118

119

120

121

123

Here, we build on and extend infrastructure monitoring efforts, including the Global Energy Monitor (GEM) (50), to map approximately 18,000 unique fossil fuel facilities worldwide. We assess their siting across over 170 countries, and im-

portantly, examine the ecological criticality of the landscapes in which they are located—not only for biodiversity conservation importance, but also in terms of zones of irrecoverable carbon storage and nature's contributions to people 127 which includes area of global importance for fisheries and agriculture, freshwater quality and quantity regulation, and coastal and inland resilience alongside 129 linked cultural values. We apply conservative estimates in our sampling strat-130 egy for siting, and only assess the most critical ecosystem components—those 131 locations that fall in the 80th percentile globally of the outcomes we examine. 132 We also perform sensitivity checks that relax our spatial sampling to encompass 133 larger buffers around sites. We then quantify the human populations residing in 134 high risk zones of this infrastructure, with special attention to children, Indige-135 nous communities, residents in states with weak environmental regulation, and 136 communities living near foreign-owned facilities where due diligence laws are 137 lacking (laws that, if in place, should protect those communities by requiring 138 businesses to respect human rights in their operations, subsidiaries, and value 139 chains). Finally, we project future risks from more than 3,500 planned facili-140 ties, evaluating whether siting and impacts follow or diverge from past evidence. 141 Taken together, our study establishes a global baseline of the location and size 142 of fenceline communities associated with fossil fuel infrastructure, as evidence for scientists, advocates, and policymakers seeking to address the intertwined 144 crises of energy, equity, and environmental justice.

146 2 Results

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

163

165

Disparity in fossil fuel production and consumption emissions

We consider more than 18,000 unique currently operating fossil fuel facility locations globally, which include coal mines, coal plants, coal terminals, oil and gas extraction sites, oil and gas plants, liquefied natural gas (LNG) terminals, and pipelines for both oil and gas (Fig 1a), of which oil and gas plants are the most numerous (Fig 1b). While exploring the emissions associated with sites is not the purpose of this study (such attribution is its own complex challenge) we use daily gridded CO₂ data for 2024 (51) to understand some of the spatial disparity in carbon emissions in terms of where fossil fuel extraction or processing occurs and where the highest consumption-based emissions patterns are (Fig S1). While this may seem obvious, the surrounding communities of these sites have the burden of high localized emissions compared to baselines (Fig 1c) although the consumption of these energy sources (for transport, residential heating, etc.) largely takes place elsewhere (Fig S1). These spatial differences are important and offer initial insight into locations around the world where communities experience localized pollution, in order for other communities, in disparate locations to benefit from meeting their energy needs. There are important differences by facility: coal plants, followed by coal terminals and oil and gas plants, are the largest source of CO₂ operating emissions (Fig 1c).

Disproportionate siting in critical ecosystems




Figure 1: Fossil fuel infrastructure is widespread globally and is associated with highly localized CO_2 emissions compared to background. (a) Locations and type of currently operating fossil fuel infrastructure; (b) The number of operating fossil fuel facilities by infrastructure type; (c) Estimated 2024 mean daily CO_2 emissions (tonnes/day) in the 10 km vicinity of each fossil fuel facility type (excluding pipelines) and emissions source compared to global average terrestrial emissions (baseline = $0.67 \, \text{TC/day}$).

Of operating fossil fuel infrastructure, we find 14.7% to be within proximity to key biodiversity areas, compared to a baseline expectation (i.e. random global distribution given locations of KBAs) of 6.8% across terrestrial land and territorial seas (Fig 2a). Similarly, 14.9% of all fossil fuel infrastructure is in proximity to critical areas for nature's contributions to people (baseline: 3.0%) and 11.1% are in or nearby zones of critical irrecoverable terrestrial carbon (baseline: 4.4%) (Fig 2a). All of these represent indicators for critical ecosystems, defined here as those in which acute disasters or ongoing pollution would have disproportionate impacts on ecosystems and communities. Combined, we identify fossil fuel infrastructure disproportionately occurs in critical ecosystems with 32% of facilities sited in any of the critical ecosystem categories compared to a baseline of 11.9%, including high occurrence in critical biomes such as temperate and broadleaf forests in Eastern US and Central Europe and mangroves in Latin America (Fig 2b). The number of operations in critical ecosystems remains high when including protected areas as an alternative indicator, at 31% compared to a baseline of 18% (Figs S2, S3); although we do not include protected areas in our main results due to low coverage of biomes in the global south compared to the global north.

We also assessed facilities in proximity to critical ecosystems using a more conservative sampling strategy that ignores the footprints of facilities where no data are available. We found that even when using 1 km buffers around all facility types other than pipelines and oil and gas extraction sites, the infrastructure still disproportionately occurs in these critical ecosystems at double the rate of baseline values (Fig S4).

191 Millions live near fossil fuel infrastructure

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

189

203

204

205

207

209

We find overall ~ 463 million people, of which ~ 124 million are children, live 192 within the 1 km high risk zone of fossil fuel infrastructure (Fig 3). Of this fence-193 line population, a subset of ~ 310 million live in critical ecosystems and ~ 107 194 million live in countries with weak environmental regulations, and therefore could be endangered due to environmental degradation or regulatory failures 196 (Fig 3a). Thirty-one million live in proximity to an internationally owned facil-197 ity which could further increase their risk because of the lack of protection and 198 accountability measures in the absence of due diligence laws. And 14 million 199 people live within 1 km of fossil fuel facilities on Indigenous land, risking already 200 historically marginalized populations (Fig 3a). 201

While fenceline populations within 1 km have higher health risks due to their close proximity to these facilities, there are also documented health risks from exposure as far as 5 km (Table S2). Almost 2 billion people globally, or a quarter of the world's population, live within 5 km of fossil fuel infrastructure, which includes over 520 million children (Fig S5). As for all our estimates, this aggregate size of fenceline communities does not indicate the total number of people that will experience health impacts, but the size of the base population with elevated health risks. At the same time, it is important to note that negative health impacts for any one individual can have large and compounding




Figure 2: Fossil fuel infrastructure is disproportionately sited in critical ecosystems, risking high priority areas for nature and people. (a) Fossil fuel infrastructure situated in critical coastal and terrestrial ecosystems compared to baselines (*indicates estimates calculated in terrestrial ecosystems only); (b) Fossil fuel infrastructure and the level of ecological risk based on critical ecosystem indicators.

negative impacts on the wider community. The subsets of populations that live within 5 km and are situated in critical ecosystems, have weak environmental regulations, are in proximity to a internationally owned location, or are located on Indigenous land are also roughly 4x larger than for 1 km estimates (Fig S5).

Excluding pipelines, ~43 million people live within 1 km of fossil fuel facili-216 ties, 19 million of them are located in already critical ecosystems, 8 million in countries with weak environmental regulations, over 1 million near an interna-218 tionally owned facility and another 1+ million on Indigenous land (Fig 3b, Table 219 S2). This difference is roughly a factor of 10 from the total—in other words, 220 most fenceline communities represent people living near pipelines. This adds complexity for public health programs aiming to assess absolute health impacts 222 because while pipeline impacts on human health are documented and can be 223 severe, exposure rates per km per year can vary based on whether oil or gas 224 is being transported, the size and age of the pipeline, maintenance schedules, 225 spillage, and leakage rates (52). Notably, at 5 km the differences between in-226 cluding and excluding pipelines is reduced, by a factor of 3-4 (Table S2). 227

As population datasets have biases depending on the underlying models and data sources (53: 54), we also consider an additional population dataset (Land-229 Scan Global) for estimating people exposed. This data attempts to estimate ambient population numbers (e.g. over a 24 hour period) rather than WorldPop, 231 which estimates residence. Notably, we find that our total population estimates at 1 km with LandScan are within approximately 2% of our WorldPop-based 233 numbers and overall within 4% for other subsets of exposed populations (sit-234 uated in ecologically critical locations, foreign ownership, weak environmental 235 regulation, or on Indigenous Land) (Table S3). It should be noted however that 236 both of these datasets may underestimate rural populations (53). 237

Future expansion threatens additional communities

238

239

241

242

243

245

247

249

251

253

As of 2025, a further 3,507 fossil fuel facilities are either proposed, in development, or currently under construction in over 140 countries globally (Fig 4a). These facilities are predominately oil and gas plants (n=1,124), gas pipelines (n=801), and coal mines (n=630); striking examples include a large number of new coal mines and coal plants as well as gas pipelines in China and oil and gas plants in the United States (Fig S6). These planned facilities continue to endanger ecologically critical locations, with 59.4% anticipated to be in proximity to key biodiversity areas compared to a baseline of 6.8%, which is a prevalence 4x higher than currently operating facilities (14.7%) and suggests these environmental designations are not deterring development on these lands and that there is preferential siting in the world's most important areas for nature conservation (Fig 4b). Similar to operating facilities, 13.5% of planned facilities are in proximity to nature's contribution to people (baseline: 3.0%, operating facilities: 14.9%) and 15.0% are near areas of critical irrecoverable terrestrial carbon (baseline: 4.0%, operating facilities: 11.1%). Combining all these indicators, we elucidate that this planned infrastructure is even more disproportionately occur-




Figure 3: An estimated hundreds of millions of people, and over a hundred million children, live within 1 km of fossil fuel infrastructure globally. (a) The number of people living within 1 km of fossil fuel infrastructure, including pipelines. (b) The number of people living within 1 km of fossil fuel infrastructure, excluding pipelines.

ring in critical ecosystems, with 69% in any of the critical ecosystem indicators compared to a baseline of 11.9% (Fig S7).

The planned infrastructure also continues to expose millions more to potential 257 risks and hazards: 106 million additional people will live within the high risk 258 zone of 1 km of the proposed locations, more than 60% of whom already reside in 259 critical ecosystems (Fig 4c). Of the people exposed, 7 million will be in countries 260 with weak environmental regulations, 8 million in proximity to a location with foreign ownership, and 7 million on Indigenous land (Fig 4c). Looking further 262 afield, 339 million additional people will be within 5 km proximity of planned 263 facilities including over 100 million children. 217 million of these 5 km fenceline 264 communities will be in existing critical ecosystems, 24 million in countries with lax environmental regulation, 21 million in proximity to a foreign-held facility, 266 and 29 million located on Indigenous land (Fig S8).



Figure 4: Planned fossil fuel infrastructure will continue to have an outsized impact on ecosystems and people. (a) Locations of planned fossil fuel infrastructure (b) Planned fossil fuel infrastructure situated in critical coastal and terrestrial ecosystems compared to baselines (*estimates calculated in terrestrial ecosystems only) (c) Number of additional people currently living within 1 km of planned fossil fuel infrastructure.

3 Discussion

277

28:

283

284

285

287

288

289

291

292

293

294

295 296

Despite widespread calls for an energy transition, fossil fuels remain the dom-269 inant energy source worldwide. Our analysis demonstrates that the 18,000 270 unique facilities documented in the GEM database are not only globally ex-271 tensive and diverse in type, but are also disproportionately sited in ecologically 272 critical areas at more than triple the baseline expectations. This includes ecosys-273 tems already under severe strain, with facilities disproportionately situated in 274 close proximity to key biodiversity areas, in regions critical to nature's contributions to people, and lands with high levels of irrecoverable carbon. This 276 threatens to erode biodiversity, undermine ecological resilience, and accelerate 278 ecosystem decline. Previous assessments of fossil fuel impacts on ecosystems have largely been conducted at much coarser scales, whether through uniform 50 km footprint assumptions (34; 38) or by mapping reserves and concession 280 blocks that may never be developed (35; 36). Our study shows that when assessed at higher resolution the risks are widespread. 282

Equally important, this study establishes the first global estimates of the size of fenceline communities living in proximity to fossil fuel infrastructure. Building on decades of evidence linking exposure to increased risks of asthma, cancer, cardiovascular disease, and other chronic conditions, we quantify the number of people living in high risk zones. Prior work has assessed the global risk to ambient populations for limited infrastructure (e.g. refineries) and likely overexaggerated the number of people at risk (55). Our estimates provide a more detailed view and further disaggregate vulnerable populations, such as children, revealing the extent of people that are likely facing the highest risks. These risks are likely compounded where facilities overlap with ecologically critical areas, weak regulatory environments, foreign ownership structures, or Indigenous territories. When these communities are impacted they can face insurmountable barriers in accessing legal remedy, given both regulatory failures and the lack of accountability frameworks.

The trajectory of planned infrastructure raises even greater concern. The 3,500+ facilities currently proposed or under construction replicate and even intensify 298 the inequities documented in existing infrastructure. We find that almost 60% of these projects are slated for, or are being constructed, near key biodiversity 300 areas, with similarly disproportionate risks to nature's contributions to people. 301 These projects will put at risk millions more people to the irreversible harms of 302 fossil fuel operations. These findings underscore how new developments not only 303 contravene international agreements and climate commitments but also deepen 304 the burden on already-stressed ecosystems and fenceline populations. 305

Geographically, some of the planned expansion is concentrated in Asia's biodiversity hotspots, where the consequences for both people and nature could 307 be devastating. China's 782 proposed coal facilities run directly counter to its net-zero pledges, while India's 156 planned coal plants, mines, and terminals 309 undermine its commitment to derive half of installed electricity from non-fossilfuel sources by 2030 (56; 57). These projects, alongside actions in other major world powers, including the United States' new "Unleashing American Energy" presidential order signal a dangerous backslide: rather than accelerating renewable energy investments, governments and corporations are choosing to lock in decades of reliance on the most greenhouse-gas-intensive and polluting fuel source. The cumulative result is not only billions of tonnes of additional CO_2 annually but also the expansion of toxic exposure, ecological destruction, and negative impacts on local communities. Impacts that are both predictable and preventable.

Moving forward, four critical actions are needed to address the global fossil fuel 320 infrastructure crisis facing fenceline communities and critical ecosystems. First, comprehensive epidemiological research and associated public health systems 322 must use our understanding of the number of people in fenceline communi-323 ties to develop improved sampling programs to monitor, evaluate, and respond 324 to what are likely millions of unreported and undocumented health impacts and needs globally. Second, researchers should support grassroots networks in 326 fenceline communities with platforms for integrating these data into advocacy 327 frameworks and legal action to realize environmental rights on the frontline of 328 fossil fuel development. Third, equal efforts on both these fronts should be made 329 but with respect to assessing and responding to ecosystem health—to ensure the 330 health and rights of nature are equally documented at scale, and legal actions 331 taken to ensure they are met. Fourth, in terms of planned and in construction 332 developments, research-informed policy and continued pressure on government 333 and industry are needed to push regulation on fossil fuel permitting processes 334 beyond narrow project-level assessments to require evaluation criteria that incor-335 porate climate commitment compatibility, renewable alternatives prioritization, 336 and the rising and compounding costs to nature and ecosystems. 337

A just energy transition must center equity. This means not only phasing out 338 fossil fuel infrastructure but also prioritizing the health, safety, and rights of the 339 hundreds of millions living at the fenceline. It means integrating robust health 340 and environmental impact assessments into decision-making, aligning national 341 and global climate commitments with on-the-ground realities, and investing in 342 renewable alternatives that reduce reliance on fossil fuels. Scientists, activists, and communities together have a vital role to play: exposing the hidden costs of 344 fossil fuels, communicating risks with clarity, and demanding urgent, enforceable 345 346

Ultimately, dismantling fossil fuel dependence is not just a climate imperative—it is a moral one. The choice is stark: continue on a path of loss of ecosystem and human health, or build an energy future that prioritizes justice, resilience, and the flourishing of both people and planet.

4 Methods

352 Data Development

353 Global Energy Monitor Database

We use data from Global Energy Monitor (GEM), an organization that devel-354 ops and analyzes data on energy infrastructure, resources, and ownership (50). 355 We aggregated data from GEM on locations, capacity, throughput, operational 356 status, and ownership levels for coal mines, coal plants, coal terminals, oil and gas extraction sites, oil and gas plants, liquefied natural gas (LNG) terminals, 358 and oil and gas pipelines from their resource and ownership trackers, most re-359 cently updated between December 2024 and June 2025. Oil and gas wells are 360 not included as part of the GEM facility databases. Fossil fuel infrastructure was also filtered to include only those that had unique location coordinates and 362 had a status of operational for operating facility analysis or either proposed, announced, in development, or under construction for our planned facility analysis. 364 Both terrestrial and offshore infrastructure located within a nation's exclusive economic zone (EEZ) was retained. A total of 18,273 unique operating fos-366 sil fuel facilities as well as 3,507 planned locations were included in the final 367 dataset.

To help determine facility extents for developing ecological buffers, we also considered the oil and gas infrastructure from the Office of Fossil Energy's National Energy Technology Laboratory (NETL) which includes features on operating assets around the world (58). While this dataset did not have as many facility types and regular updates as the GEM, it did include shapefiles for undertaking sensitivity analysis of extents for some assets of interest.

CO₂ Emissions Data

Emissions data was obtained from the Global Gridded Daily CO_2 Emissions Dataset (GRACED) which has tracked daily power, industry, residential, and transportation CO_2 emissions since 2019 at a global resolution of 0.1° by 0.1° (51).

Ecological Data

All ecological data was spatially restricted to terrestrial and EEZ boundaries, excluding Antarctica, with terrestrial biomes defined using the RESOLVE Ecore-382 gions dataset (59). Analyses focused on four key ecological dimensions: 1) Key 383 Biodiversity Areas (KBAs): Spatial data provided by the KBA Partnership via 384 BirdLife International (60) 2) Nature's Contributions to People (NCPs): Spatial raster data (~2 km resolution) representing top 20th percentile of global ecosys-386 tem service values across 12 NCP types (61) 3) Irrecoverable Carbon: Raster data (~300 m resolution) of high-density biomass and soil carbon stores that 388 would not regenerate by 2050 if disturbed, considering only the top 20th percentile values (62) 4) World Database of Protected Areas (WDPAs): Developed 390 by the United Nation Environment Programme (UNEP) and the International Union for the Conservation of Nature (IUCN). Though included in supplementary analysis, WDPAs were excluded from main figures due to regional designation biases (63).

395 Population Data

Population exposure was assessed using the WorldPop 2020 gridded population dataset, which provides estimates at \sim 1 km spatial resolution globally (64), including by age and gender for year 2020. For our analysis of population counts, children were defined as under 15 years of age. For sensitivity analysis, we also considered the 2020 population dataset from LandScan Global (65) which is an ambient population estimate at \sim 1 km spatial resolution.

402 Regulation Data

The Environmental Regulatory Risk (ERR) Index was used to indicate countrylevel environmental regulation strength (66). Four countries (France, Germany, Netherlands, Norway) have due diligence laws and were omitted when considering locations that had international ownership.

To compile our due diligence law data, we employed a multi-step research pro-407 cess current as of May 1, 2025. Our primary data was compiled using the 2024 UN report "Business, planetary boundaries, and the right to a clean, healthy, 409 and sustainable environment" (67) and Boyd and Keene 2022 Policy Brief No. 410 "Essential elements of effective and equitable human rights and environ-411 mental due diligence legislation" (68). To update our primary data, a search 412 was conducted using the keywords "due diligence", "vigilance", "supply chain", 413 "responsible business", and "responsible corporation" within FAOLEX (69), a 414 legislative and policy worldwide database managed by the FAO and ECOLEX, 415 a global environmental law repository jointly maintained by the IUCN, UNEP, 416 and FAO (70), and HeinOnline which offers access to legal resources across the 417 globe (71). The research also involved a targeted review of recent news devel-418 opments through the webpages of the European Coalition for Corporate Justice and the Canadian Network on Corporate Accountability—two civil society net-420 works dedicated to business accountability (72; 73). This was complemented by Google search using the same keywords to capture additional relevant mate-422 rials. After identifying relevant due diligence laws, we consulted governmental 423 gazettes to access official legislative versions. 424

425 Indigenous Lands Data

Indigenous lands spatial data was obtained from Garnett et al. 2018 (18) under a specific use agreement for this manuscript and research project.

428 Analytical Approach

429 Facility-Associated Emissions

Buffers for emissions estimates for spatial analysis of production vs. consumption comparisons were calculated at a standard 10 km around each facility, as

airborne emissions are more dispersed compared to the effect of physical footprints on land-based impacts. Emissions baselines were calculated as the average emissions for the proportion of global terrestrial and EEZ surface area covered 434 by the gridded emissions data. Fossil fuel infrastructure proximity was then assessed by calculating the proportion of infrastructure footprints intersecting this 436 layer. Pipelines were excluded from emissions calculations due to the difficulty 437 in identifying localized emissions, many being buried underground, emissions 438 largely arising from spills, leakage, or compressions stations which would be 439 challenging to observe at the current resolution of the data, and their density 440 near population centers which could be conflated with consumption-based emis-442

443 Infrastructure Footprints and Ecological Risk

We assessed the probability of siting in critical ecosystems by developing a facility specific sampling strategy that accounted for the direct footprint of facilities 445 plus moderate 'spillover' effects outside the facility boundaries. Such spillover 446 effects include routine noise pollution, air pollution (including particulate mat-447 ter and debris), light pollution, water pollution, and disturbance from traffic 448 and transport in land, waterways, and sea, all of which can disrupt non-human species health, interactions, and movement as well as human activities which 450 depend on nature in landscapes (25; 26; 27; 28; 29; 30; 31; 32; 33; 40; 41; 49; 451 42; 43; 44; 45; 46; 47; 48). 452

For our main analysis, we took a conservative approach that varied by facility 453 type. For coal mines, existing spatial facility level footprint data exist and we 454 simply use a 1 km buffer around those facility areas; for oil, gas, and coal plants 455 spatial footprint data is not available, and so for these locations we use a 2 km radius from the plant centroid to account for the added spatial footprint 457 of these plants globally (notably this choice was taken to account for even the 458 world's largest coal, oil, and gas plants whose spatial footprints translate to 459 \sim 850 m, \sim 500 m, \sim 1.2 km, respectively; we relax this assumption in sensitivity analysis and reduce the radius to 1 km for plants, see Fig S4); for oil and gas 461 extraction fields we considered 5 km around the centroid of the field, a choice 462 validated against actual field boundaries data where it exists, showing a con-463 servative approximation of prevalence in the risks for these extensive operations often highly interspersed in working lands (Fig S9); for pipelines, we consider 465 a 25 m buffer around the width of the pipeline to represent local disturbance; 466 and for LNG terminals, which generally have a smaller footprint, we consider 467 a 1 km buffer. Notably, these buffers are simply to estimate siting issues in 468 critical ecosystems, and are not intended to indicate the full extent of potential 469 damage, for example from explosions, major spillage, pipeline breakage, or other 470 ecological disasters, which we know to have much wider footprints that those 471 ones we estimate here. 472

Ecological baselines are calculated as the proportion of global terrestrial and EEZ surface area covered by each ecological layer; that is, what percentage of global terrestrial and EEZ surfaces are in any of the ecological layers. Fossil

fuel infrastructure proximity was then assessed by calculating the proportion of infrastructure footprints intersecting these layers. A normalized ecological risk index (0-3) is computed for each facility based on the number of overlapping 478 ecological dimensions. Terrestrial sites are evaluated against three dimensions (KBA, NCP, irrecoverable carbon) while offshore sites are evaluated against 480 two (KBA, NCP). For NCP and irrecoverable carbon we only assess those sites 481 globally that fall in the 80th percentile of the distribution in order to represent 482 the most critical ecosystems globally with respect to these indicators. 483

Fossil Fuel Infrastructure Ownership 484

Ownership of fossil fuel infrastructure is classified using a hierarchical approach, 485 drawing on four levels of corporate ownership: Level 0 (official headquarters), 486 Level 1 (Owner1), Level 2 (Owner2), and Level 3 (Owner3). Ownership type 487 is based on aggregate shares. Facilities are designated as nationally owned if 488 >50% of aggregate ownership is from companies headquartered in the host coun-489 try or internationally owned if >50% of ownership is held by foreign entities. 490 In cases where ownership data were missing at a given level the classification 491 considers the next available level. For facilities spanning multiple countries, a 492 conservative approach is used: if any of the countries listed matches the owner's headquarters, the infrastructure is classified as nationally owned. For popula-494 tion exposure analysis, foreign-owned facilities where the headquarters is not in France, Germany, Netherlands, or Norway are used in the analysis as other 496 countries do not have any due diligence laws in place. 497

Fenceline Populations 498

509

To calculate the size of the fenceline communities around fossil fuel infrastruc-499 ture buffer zones were created around facility centroids or lines (for pipelines) and resident population numbers extracted using zonal statistics. Two buffer 501 distances were defined: 1 km to represent the highest risks and 5 km to capture 502 lower but still elevated risks. These distances were based on the literature where 503 studies assess health impacts for residential areas over various distances (Table 504 S1). Additional analysis on population exposure also calculated people exposed 505 at 1 and 5 km without the inclusion of pipelines, given their large extent and 506 reduced risks at 5 km, as a sensitivity analysis (Table S2).

Subsets of the fenceline communities were estimated including 1) a total num-508 ber of people within a facility's buffer, 2) the number within critical ecosystems, or all people within the buffer of a facility which also intersects an eco-510 logical risk layer, 3) those with a low environmental regulatory regime index 511 (ERRI)—people in the facility buffer where the country has an ERRI below 512 zero, 4) the number of people close to facilities that are international owned 513 and without due diligence, counted as the number of people in the facility buffer where the facility is foreign-owned and the headquarter country does not 515 have due diligence laws, and, finally, 5) those that are proximate to Indigenous 516 lands—all people in the facility buffer which also intersects with the Indigenous 517 lands layer. 518

To ensure there was no double counting of populations for current and future facilities, operating buffers were differenced from the buffers of the planned facilities. As a result all calculated populations for planned facilities are additional individuals at risk, on top of those already living nearby operating facilities.

As an additional sensitivity layer, we also assessed populations using the Land-524 scan Global population layer, which is a population dataset with the same resolution as WorldPop but constructed through ambient population mapping (65). 526 Estimating population has many challenges and global datasets have different 527 levels of sophistication in terms of how they model populations from disaggrega-528 tion of census counts to more complex approaches that utilize multiple auxiliary sources and dasymetric mapping. There are known challenges with population 530 layers, including potential for under representation of rural populations (53) and how well they are fit for different use cases (54); however, LandScan and 532 WorldPop have comparably high fidelity. To address the potential uncertainty between data products, we include an estimate of the percentage differences be-534 tween WorldPop and Landscan exposed population estimates (Table S3).

5 Contributions

```
    Project Ideation: GB, KF, LC, NO, SC, CO, ILG, VK, ZM
    Analysis (Formal methodology): GB, KF, LC, NO, ZM
```

539 Analysis (Implementation): GB, KF, LC, NO

Results (Interpretation): GB, KF, LC, NO, ZM

Writing: GB, KF, LC, NO, ZM

⁵⁴² Comments/review: GB, KF, LC, NO, SC, CO, ILG, VK, ZM

₅₄₃ 6 Data and Code Availability

The data sets and code used and created in this study are available at https://github.com/Better-Planet-Laboratory/fossil-fuel-risk.

$_{\scriptscriptstyle 46}$ 7 References

47 References

[1] IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland. July 2023.

- [2] C. W. Callahan and J. S. Mankin, "Carbon majors and the scientific case for climate liability," *Nature*, vol. 640, p. 893–901, Apr. 2025.
- 554 [3] D. O'Rourke and S. Connolly, "Just oil? the distribution of environmental and social impacts of oil production and consumption," *Annual Review of Environment and Resources*, vol. 28, p. 587–617, Nov. 2003.
- K. Vohra, A. Vodonos, J. Schwartz, E. A. Marais, M. P. Sulprizio, and
 L. J. Mickley, "Global mortality from outdoor fine particle pollution generated by fossil fuel combustion: Results from geos-chem," *Environmental Research*, vol. 195, p. 110754, 2021.
- [5] F. A. Wichmann, A. Müller, L. E. Busi, N. Cianni, L. Massolo, U. Schlink,
 A. Porta, and P. D. Sly, "Increased asthma and respiratory symptoms in children exposed to petrochemical pollution," *Journal of Allergy and Clinical Immunology*, vol. 123, p. 632–638, Mar. 2009.
- ⁵⁶⁵ [6] S. Mondal, G. Singh, and M. K. Jain, "Spatio-temporal variation of air pollutants around the coal mining areas of Jharia Coalfield, India," *Environmental Monitoring and Assessment*, vol. 192, p. 405, May 2020.
- D. Holmstrom, F. Altamirano, J. Banks, G. Joseph, M. Kaszniak,
 C. Mackenzie, R. Shroff, H. Cohen, and S. Wallace, "Csb investigation of the explosions and fire at the bp texas city refinery on march 23, 2005,"
 Process Safety Progress, vol. 25, p. 345–349, Sept. 2006.
- J. Idavain, K. Lang, J. Tomasova, A. Lang, and H. Orru, "Cancer Incidence
 Trends in the Oil Shale Industrial Region in Estonia," *International Journal* of Environmental Research and Public Health, vol. 17, p. 3833, June 2020.
- [9] A. Balgobin and N. Ramroop Singh, "Source apportionment and seasonal cancer risk of polycyclic aromatic hydrocarbons of sediments in a multi-use coastal environment containing a Ramsar wetland, for a Caribbean island,"

 Science of The Total Environment, vol. 664, pp. 474–486, May 2019.
- [10] United Nations (Human Rights Council), A/HRC/RES/48/13 Resolution adopted by the Human Rights Council on 8 October 2021. Oct. 2021.
- [11] United Nations (General Assembly), A/RES/76/300 Resolution adopted by the General Assembly on 28 July 2022. July 2022.
- [12] AO-32/25 Climate Emergency and Human Rights. July 2025.
- [13] International Court of Justice AO on Obligation of States in respect of Climate Change. July 2025.
- ⁵⁸⁶ [14] "Standing Firm: The land and environmental defenders on the frontlines of the climate crisis," tech. rep., Global Witness, 2023.
- ⁵⁸⁸ [15] NOAA Fisheries, "Deepwater horizon 10 years later: 10 questions," Apr ⁵⁸⁹ 2020.

- [16] J. O'Farrell, D. O'Fionnagáin, A. O. Babatunde, M. Geever, P. Codyre,
 P. C. Murphy, C. Spillane, and A. Golden, "Quantifying the Impact of
 Crude Oil Spills on the Mangrove Ecosystem in the Niger Delta Using AI
 and Earth Observation," Remote Sensing, vol. 17, p. 358, Jan 22 2025.
- P. S. zu Ermgassen, N. Mukherjee, T. A. Worthington, A. Acosta, A. R. d.
 Rocha Araujo, C. M. Beitl, G. A. Castellanos-Galindo, M. Cunha-Lignon,
 F. Dahdouh-Guebas, K. Diele, C. L. Parrett, P. G. Dwyer, J. R. Gair,
 A. F. Johnson, B. Kuguru, A. Savio Lobo, N. R. Loneragan, K. Longley Wood, J. T. Mendonça, J.-O. Meynecke, R. N. Mandal, C. N. Munga, B. G.
 Reguero, P. Rönnbäck, J. Thorley, M. Wolff, and M. Spalding, "Reprint
 if: Fishers who rely on mangroves: Modelling and mapping the global
 intensity of mangrove-associated fisheries," Estuarine, Coastal and Shelf
 Science, vol. 248, p. 107159, 1 2021.
- [18] S. T. Garnett, N. D. Burgess, J. E. Fa, Á. Fernández-Llamazares,
 Z. Molnár, C. J. Robinson, J. E. M. Watson, K. K. Zander, B. Austin,
 E. S. Brondizio, N. F. Collier, T. Duncan, E. Ellis, H. Geyle, M. V. Jackson, H. Jonas, P. Malmer, B. McGowan, A. Sivongxay, and I. Leiper, "A spatial overview of the global importance of Indigenous lands for conservation," Nature Sustainability, vol. 1, pp. 369–374, Jul 16 2018.
- [19] M. A. Hurlbert and R. Datta, "When the environment is destroyed, you're destroyed: Achieving Indigenous led pipeline justice," *Energy Research amp; Social Science*, vol. 91, p. 102711, 9 2022.
- [20] L. Wright and J. P. White, "Developing oil and gas resources on or near indigenous lands in canada: An overview of laws, treaties, regulations and agreements," *International Indigenous Policy Journal*, vol. 3, no. 2, pp. pp. 1–18, 2012.
- [21] J. Farrell, "Network structure and influence of the climate change countermovement," *Nat. Clim. Chang.*, vol. 6, pp. 370–374, Apr. 2016.
- [22] J. Farrell, "Corporate funding and ideological polarization about climate change," Proceedings of the National Academy of Sciences, vol. 113, no. 1, pp. 92–97, 2016.
- [23] S. Wolf, R. Bullard, J. J. Buonocore, N. Donley, T. Farrelly, J. Fleming,
 D. J. X. González, N. Oreskes, W. Ripple, R. Saha, and M. D. Willis,
 "Scientists' warning on fossil fuels," Oxford Open Climate Change, vol. 5,
 Jan. 2025.
- [24] L. M. McKenzie, R. Z. Witter, L. S. Newman, and J. L. Adgate, "Human health risk assessment of air emissions from development of unconventional natural gas resources," Sci. Total Environ., vol. 424, pp. 79–87, May 2012.

- [25] R. D. Soares, M. Dos Santos, F. R. de Moura, A. L. Muccillo-Baisch,
 P. R. M. Baisch, M. C. F. Soares, and F. M. R. da Silva Júnior, "Gestational and neonatal outcomes in cities in the largest coal mining region in brazil," Int. J. Environ. Res. Public Health, vol. 19, p. 12107, Sept. 2022.
- [26] J. A. Casey, D. A. Savitz, S. G. Rasmussen, E. L. Ogburn, J. Pollak, D. G.
 Mercer, and B. S. Schwartz, "Unconventional natural gas development and
 birth outcomes in pennsylvania, USA," *Epidemiology*, vol. 27, pp. 163–172,
 Mar. 2016.
- [27] S. Ha, H. Hu, J. Roth, H. Kan, and X. Xu, "Associations between residential proximity to power plants and adverse birth outcomes," Am. J. Epidemiol., vol. 182, pp. 215–224, Aug. 2015.
- [28] L. M. McKenzie, R. Guo, R. Z. Witter, D. A. Savitz, L. S. Newman, and
 J. L. Adgate, "Birth outcomes and maternal residential proximity to natural gas development in rural colorado," *Environ. Health Perspect.*, vol. 122,
 pp. 412–417, Apr. 2014.
- [29] H. Guo, S. H. Kota, K. Chen, S. K. Sahu, J. Hu, Q. Ying, Y. Wang, and
 H. Zhang, "Source contributions and potential reductions to health effects
 of particulate matter in india," *Atmos. Chem. Phys.*, vol. 18, pp. 15219–15229, Oct. 2018.
- [30] M. Lavoie, D. Risk, and D. Rainham, "Sociodemographic and population exposure to upstream oil and gas operations in canada," Int. J. Environ.
 Res. Public Health, vol. 21, Dec. 2024.
- [31] J. Cortes-Ramirez, S. Naish, P. D. Sly, and P. Jagals, "Mortality and morbidity in populations in the vicinity of coal mining: a systematic review,"
 BMC Public Health, vol. 18, Dec. 2018.
- [32] A.-K. Hurtig and M. San Sebastián, "Geographical differences in cancer incidence in the amazon basin of ecuador in relation to residence near oil fields," *Int. J. Epidemiol.*, vol. 31, pp. 1021–1027, Oct. 2002.
- [33] D. Roy, G. Singh, and Y.-C. Seo, "Coal mine fire effects on carcinogenicity and non-carcinogenicity human health risks," *Environ. Pollut.*, vol. 254, p. 113091, Nov. 2019.
- [34] M. B. J. Harfoot, D. P. Tittensor, S. Knight, A. P. Arnell, S. Blyth,
 S. Brooks, S. H. M. Butchart, J. Hutton, M. I. Jones, V. Kapos, J. P.
 Scharlemann, and N. D. Burgess, "Present and future biodiversity risks
 from fossil fuel exploitation," Conservation Letters, vol. 11, Apr. 2018.
- [35] N. Butt, H. L. Beyer, J. R. Bennett, D. Biggs, R. Maggini, M. Mills,
 A. R. Renwick, L. M. Seabrook, and H. P. Possingham, "Conservation.
 biodiversity risks from fossil fuel extraction," *Science*, vol. 342, pp. 425–426, Oct. 2013.

- [36] H. S. Grantham, P. Tibaldeschi, P. Izquierdo, K. Mo, D. J. Patterson,
 H. Rainey, J. E. M. Watson, and K. R. Jones, "The emerging threat of
 extractives sector to intact forest landscapes," Front. For. Glob. Chang.,
 vol. 4, July 2021.
- [37] B. W. Allred, W. K. Smith, D. Twidwell, J. H. Haggerty, S. W. Running,
 D. E. Naugle, and S. D. Fuhlendorf, "Sustainability. ecosystem services lost
 to oil and gas in north america," *Science*, vol. 348, pp. 401–402, Apr. 2015.
- [38] L. J. Sonter, M. C. Dade, J. E. M. Watson, and R. K. Valenta, "Renewable energy production will exacerbate mining threats to biodiversity," *Nat. Commun.*, vol. 11, p. 4174, Sept. 2020.
- [39] K. Kühne, N. Bartsch, R. D. Tate, J. Higson, and A. Habet, ""carbon bombs" mapping key fossil fuel projects," *Energy Policy*, vol. 166, p. 112950, 2022.
- [40] C. Erbe, S. A. Marley, R. P. Schoeman, J. N. Smith, L. E. Trigg, and C. B.
 Embling, "The effects of ship noise on marine mammals—a review," Front.
 Mar. Sci., vol. 6, Oct. 2019.
- [41] J. M. Van der Hoop, M. J. Moore, S. G. Barco, T. V. N. Cole, P.-Y.
 Daoust, A. G. Henry, D. F. McAlpine, W. A. McLellan, T. Wimmer, and
 A. R. Solow, "Assessment of management to mitigate anthropogenic effects
 on large whales," Conserv. Biol., vol. 27, pp. 121–133, Feb. 2013.
- [42] T. Rutherford, L. Maxwell, N. Kleist, E. Teige, R. Lehrter, M. Gilbert,
 D. Wood, A. Johnston, C. Mengelt, J. Tull, T. Haby, and S. Carter, "Effects of noise from oil and gas development on ungulates and small mammals—a science synthesis to inform national environmental policy act analyses,"
 tech. rep., U.S. Geological Survey, 2023.
- [43] M. L. Richardson, B. A. Wilson, D. A. S. Aiuto, J. E. Crosby, A. Alonso,
 F. Dallmeier, and G. K. Golinski, "A review of the impact of pipelines and power lines on biodiversity and strategies for mitigation," *Biodivers. Conserv.*, vol. 26, pp. 1801–1815, July 2017.
- [44] R. G. Verb and M. L. Vis, "Periphyton assemblages as bioindicators of
 mine-drainage in unglaciated western allegheny plateau lotic systems," Water Air Soil Pollut., vol. 161, pp. 227–265, Feb. 2005.
- [45] T. M. Thompson, D. Shepherd, A. Stacy, M. G. Barna, and B. A. Schichtel,
 "Modeling to evaluate contribution of oil and gas emissions to air pollution," *J. Air Waste Manag. Assoc.*, vol. 67, pp. 445–461, Apr. 2017.
- [46] R. Bobbink, K. Hicks, J. Galloway, T. Spranger, R. Alkemade, M. Ashmore, M. Bustamante, S. Cinderby, E. Davidson, F. Dentener, B. Emmett,
 J.-W. Erisman, M. Fenn, F. Gilliam, A. Nordin, L. Pardo, and W. De Vries,
 "Global assessment of nitrogen deposition effects on terrestrial plant diversity: a synthesis," *Ecol. Appl.*, vol. 20, pp. 30–59, Jan. 2010.

- [47] K. J. Gaston, T. W. Davies, J. Bennie, and J. Hopkins, "Reducing the ecological consequences of night-time light pollution: options and developments," J. Appl. Ecol., vol. 49, pp. 1256–1266, Dec. 2012.
- [48] R. Sordello, O. Ratel, F. Flamerie De Lachapelle, C. Leger, A. Dambry, and
 S. Vanpeene, "Evidence of the impact of noise pollution on biodiversity: a systematic map," *Environ. Evid.*, vol. 9, Sept. 2020.
- [49] B. R. Silliman, J. van de Koppel, M. W. McCoy, J. Diller, G. N. Kasozi,
 K. Earl, P. N. Adams, and A. R. Zimmerman, "Degradation and resilience in louisiana salt marshes after the BP-Deepwater horizon oil spill," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 109, pp. 11234–11239, July 2012.
- [50] Global Energy Monitor, "Global energy trackers," Jul 2025.
- [51] X. Dou, Y. Wang, P. Ciais, F. Chevallier, S. J. Davis, M. Crippa,
 G. Janssens-Maenhout, D. Guizzardi, E. Solazzo, F. Yan, D. Huo, B. Zheng,
 B. Zhu, D. Cui, P. Ke, T. Sun, H. Wang, Q. Zhang, P. Gentine, Z. Deng,
 and Z. Liu, "Near-real-time global gridded daily CO2 emissions," The Innovation, vol. 3, p. 100182, 1 2022.
- [52] H. Wang and I. J. Duncan, "Likelihood, causes, and consequences of focused leakage and rupture of U.S. natural gas transmission pipelines," J. Loss Prev. Process Ind., vol. 30, pp. 177–187, July 2014.
- [53] J. Láng-Ritter, M. Keskinen, and H. Tenkanen, "Global gridded population datasets systematically underrepresent rural population," Nat. Commun., vol. 16, p. 2170, Mar. 2025.
- [54] S. Leyk, A. E. Gaughan, S. B. Adamo, A. de Sherbinin, D. Balk,
 S. Freire, A. Rose, F. R. Stevens, B. Blankespoor, C. Frye, J. Comenetz,
 A. Sorichetta, K. MacManus, L. Pistolesi, M. Levy, A. J. Tatem, and M. Pesaresi, "The spatial allocation of population: a review of large-scale gridded population data products and their fitness for use," *Earth Syst. Sci. Data*,
 vol. 11, pp. 1385–1409, Sept. 2019.
- [55] K. S. Dorman, Maxwell, "Fossil fuel atlas: Illustrating the threats of fossil fuel production a rapid threat identification methodology and platform,"
 tech. rep., Stockholm Environment Institute, Global Energy Monitor, Institute for Governance Sustainable Development, 2022.
- 739 [56] Government of China, "Chinas's achievement, new goals and new measures for nationally determined contributions (unofficial translation)." https://unfccc.int/sites/default/files/NDC/2022-06/China's Accessed: 2025-8-26.
- Government of India, "India's updated nation-[57]first 743 determined contribution agreement." allv under paris 744 https://unfccc.int/sites/default/files/NDC/2022-08/India Accessed: 745 2025-8-26. 746

- ⁷⁴⁷ [58] M. Sabbatino, "Global oil & gas features database," 2018.
- [59] E. Dinerstein, D. Olson, A. Joshi, C. Vynne, N. D. Burgess, E. Wikra-748 manayake, N. Hahn, S. Palminteri, P. Hedao, R. Noss, M. Hansen, 749 H. Locke, E. C. Ellis, B. Jones, C. V. Barber, R. Hayes, C. Kormos, 750 V. Martin, E. Crist, W. Sechrest, L. Price, J. E. M. Baillie, D. Wee-751 den, K. Suckling, C. Davis, N. Sizer, R. Moore, D. Thau, T. Birch, P. Potapov, S. Turubanova, A. Tyukavina, N. de Souza, L. Pintea, J. C. 753 Brito, O. A. Llewellyn, A. G. Miller, A. Patzelt, S. A. Ghazanfar, J. Tim-754 berlake, H. Klöser, Y. Shennan-Farpón, R. Kindt, J.-P. B. Lillesø, P. van 755 Breugel, L. Graudal, M. Voge, K. F. Al-Shammari, and M. Saleem, "An 756 ecoregion-based approach to protecting half the terrestrial realm," Bio-757 Science, vol. 67, p. 534–545, Apr. 2017. 758
- 759 [60] "Key biodiversity areas 2024," Sep 2024.
- [61] R. Chaplin-Kramer, R. A. Neugarten, R. P. Sharp, P. M. Collins, S. Polasky, D. Hole, R. Schuster, M. Strimas-Mackey, M. Mulligan, C. Brandon, S. Diaz, E. Fluet-Chouinard, L. J. Gorenflo, J. A. Johnson, C. M. Kennedy, P. W. Keys, K. Longley-Wood, P. B. McIntyre, M. Noon, U. Pascual, C. Reidy Liermann, P. R. Roehrdanz, G. Schmidt-Traub, M. R. Shaw, M. Spalding, W. R. Turner, A. van Soesbergen, and R. A. Watson, "Mapping the planet's critical natural assets," Nature Ecology amp; Evolution, vol. 7, p. 51–61, Nov. 2022.
- [62] M. L. Noon, A. Goldstein, J. C. Ledezma, P. R. Roehrdanz, S. C. Cook-Patton, S. A. Spawn-Lee, T. M. Wright, M. Gonzalez-Roglich, D. G. Hole,
 J. Rockström, and W. R. Turner, "Mapping the irrecoverable carbon in earth's ecosystems," Nature Sustainability, vol. 5, p. 37–46, Nov. 2021.
- ⁷⁷² [63] "Protected planet-world protected areas," Feb 2025.
- ⁷⁷³ [64] "World pop 1 km 2020 population density," May 2025.
- V. Lebakula, J. Epting, J. Moehl, C. Stipek, D. Adams, A. Reith, J. Kaufman, J. Gonzales, B. Reynolds, S. Basford, A. Martin, W. Buck, A. Faxon,
 A. Cunningham, A. Roy, Z. Barbose, J. Massaro, S. Walters, C. Woody,
 A. Iman, A. Wilkins, E. Powell, and M. Urban, "LandScan silver edition,"
 2024.
- [66] D. Etsy and M. E. Porter, Ranking National Environmental Regulation and
 Performance: A Leading Indicator of Future Competitiveness?", p. 78–101.
 Oxford University Press, 2002.
- Total [67] United Nations (Human Rights Council), A/HRC/55/43 Business, planetary boundaries, and the right to a clean, healthy and sustainable environment. Jan. 2024.

- [68] S. Keene and D. R. Boyd, "Essential elements of effective and equitable human rights and environmental due diligence legislation," policy Brief
 No.3, UN OHCHR, 2022.
- [69] Food and Agriculture Organization of the United Nations, "Faolex," May
 2025.
- [70] International Union for Conservation of Nature, United Nations Environment Programme and Food and Agriculture Organization of the United
 Nations, "Ecolex," May 2025.
- ⁷⁹³ [71] I. William S. Hein Co., "Heinonline," May 2025.
- ⁷⁹⁴ [72] European Coalition for Corporate Justice, "European coalition for corporate justice," May 2025.
- [73] Canadian Network on Corporate Accountability, "Canadian network on corporate accountability," May 2025.

8 Supplementary Material

Table S1: Review of health impact studies, including location, distance, affected populations, and observed health impacts.

Study (Location)	Distance	Population / Sample	Observed Health Impacts	Notes	Facility Type
Casey et al., 2016 (USA, PA)	0–2 km	Pregnant women	Preterm birth, low birth weight, anomalies	Strongest effects <1 km	Gas Wells
McKenzie et al., 2012 (USA, CO & PA)	>1 km, <1 km	Residents near wells	VOCs, NOx, PM _{2.5} , asthma, bronchitis	Air quality declines with distance	Gas Wells
McKenzie et al., 2014 (USA, CO)	0–20 km	Pregnant women	Congenital defects, preterm birth	Higher prevalence near wells	Gas Wells
Osborn et al., 2011 (USA, PA)	0–1 km	Households w/ private wells	Methane migration, water contamination	Waterborne exposure	Gas Wells
Stacy et al., 2015 (USA, PA)	0–2 km	Pregnant women	Adverse perinatal outcomes	Distance- dependent risk	Gas Wells
Steinzor et al., 2013 (USA, Marcellus Shale)	0–3 km	Residents	Noise, light, stress, sleep disturbance	Indirect impacts	Gas Wells
Chen et al., 2021 (China)	0-50 km	Communities	Mortality, COPD, cardiovascular issues	Spillover >50 km	Coal Plants
Clemons et al., 2021 (USA, KY)	0–20 km	Children	Asthma hospital visits	Reduced after decommis- sioning	Coal Plants
Gasparotto & Martinello, 2021 (Global)	0–10 km	Residents	Respiratory, cardiovascular, heavy metals	Downwind effects extend 10–20 km	Coal Plants
Guo et al., 2018 (India)	0–50 km	Residents	PM _{2.5} , CEVD, IHD, COPD, cancer	Major contributor to premature mortality	Coal Plants

Study / Location	Distance	Population / Sample	Observed Health Impacts	Notes	Facility Type
Han et al., 2024 (South Korea)	2 km	General population near mines	Higher cancer incidence (all types)	Long-term $PM_{2.5}$ and coal dust exposure	Coal Plants
Henneman et al., 2023 (USA)	0–50 km	Residents near plants	Mortality risk (PM _{2.5} , SO ₂)	460,000 attributable deaths	Coal Plants
Zhang et al., 2021 (USA)	<5 km	Children	Neurobehavioral problems	Airborne exposure, coal ash	Coal Plants
Graber et al., 2013 (USA)	<5 km	Coal miners	Bronchitis, asthma, reduced lung function	Particulate exposure focus	Coal Mines
Palmer et al., 2010 (Appalachia, USA)	1–10 km	Coal mining communities	Anxiety, depression, cardiovascular issues	Mountaintop removal; psychosocial stress	Coal Mines
Ramirez et al., 2018 (Global)	0–20 km	Communities near mines	Neoplasms, circulatory, respiratory, metabolic, congenital, cancer	Broad proximity risks	Coal Mines
Roy et al., 2019 (Jharkhand, India)	1–5 km	Villages near coal mines	Air pollution, cancer rates	Airborne exposure focus	Coal Mines
Soares et al., 2022 (Brazil)	1–10 km	Pregnant women	Low birth weight, preterm birth	PM _{2.5} and toxic gases in pregnancy	Coal Mines
Dey et al., 2015 (India)	<5 sq km	Settlements near oil drilling	SO ₂ , NO ₂ , liver damage	Long-term abnormalities	Oil Fields
Hurtig & San Sebastian, 2002/2013 (Ecuador)	0–10 km	Local communities	Childhood leukemia, cancer	Higher incidence near oil fields	Oil Fields
Kponee et al., 2015 (Nigeria)	0–10 km	Local communities	Neurological, hematological effects	Benzene, hydrocarbons in water	Oil Fields

Study / Location	Distance	Population / Sample	Observed Health Impacts	Notes	Facility Type
Kudabayeva et al., 2014 (Kazakhstan)	0–10 km	School children	High thyroid volume (goitre)	More goitre proximal	Oil Fields
Di Ciaula, 2012 (Europe)	<3 km	Communities near plants	Air pollution, hospital admissions	Higher with PM_{10}	Oil & Gas Plants
Ha et al., 2015 (USA, FL)	0–10 km	Pregnant women	Preterm delivery, low birth weight	Increased risk near plants	Oil & Gas Plants
Liu et al., 2012 (USA, NY)	0-50 km	Residents	Hospitalization: asthma, ARI, COPD	Respiratory burden	Oil & Gas Plants
Solutions for Our Climate, 2021 (S. Korea)	0–100 km	Residents	COPD, diabetes, asthma, leukemia	Transboundary PM _{2.5} impacts	Oil & Gas Plants
Davis et al., 2023 (USA)	0–2 km	Residents near corridors	Soil/water contamination, air risks	Chronic exposure possible	Pipelines
Emanuel et al., 2021 / Strube et al., 2021 (USA)	0–100 km	Counties with pipelines	Environmental justice disparities	Higher risk for vulnerable populations	Pipelines
Jonasson et al., 2019 (Canada)	0–5 km	Communities near terminus	Bioaccumulating toxins	Risks to food supply	Pipelines
Shi et al., 2014 (China)	0–1 km	Local populations	Heavy metal exposure, emissions	Impacts >50m ROW	Pipelines
Greenpeace, 2024 (USA)	0–100 km	Residents near terminal	Premature deaths, asthma, lost days	Air quality monitoring, strong correlation	LNG Terminals
Michanowicz et al., 2023 (Global)	<10 km	Communities near gas leaks	Headaches, nausea, respiratory complaints	Odorant exposure, acute unclear chronic	LNG Terminals
NFPA 59A, 2021 (USA/global)	0.5–2 km	General population	Fire/explosion risk	Regulatory guidance	LNG Terminals

Study / Location	Distance	Population / Sample	Observed Health Impacts	Notes	Facility Type
US DoE, 2016 (USA)	0–5 km	Surrounding communities	NOx, VOCs, odor, noise	Acute + chronic exposure pathways	LNG Terminals
Lavoie et al., 2024 (Canada, AB)	1–3 km	Communities near wells	Cardio- respiratory issues	9% of AB pop. within 1 km of wells	Conventional Oil & Gas Wells




Figure S1: Consumption based emissions as a percentage of total 2024 CO_2 emissions.



Figure S2: Ecological risk prevalence of fossil fuel facilities near locations in the World Database of Protected Areas.




Figure S3: Risk map of fossil fuel facilities in ecologically sensitive areas (including World Database of Protected Areas).




Figure S4: Ecological risk prevalence of fossil fuel facilities with 1 km footprint buffers for coal and oil and gas plants.

Table S2: Populations in proximity to fossil fuel infrastructure at different distances, including or excluding pipelines.

Type of Exposure	1 Km	1 Km No Pipelines	5 Km	5 Km No Pipelines
Total	462,794,372	42,229,033	1,947,553,808	621,547,837
Critical Ecosystems	310,500,206	19,783,405	1,368,728,496	308,852,375
Low Environment Regulatory Regime Index	106,887,136	7,600,357	468,093,799	139,546,272
International Ownership with No Due Diligence	31,018,846	1,386,160	156,635,531	26,727,910
Indigenous Land	13,884,394	1,306,433	66,182,091	18,183,870

803




Figure S5: Number of people living within 5 km of fossil fuel infrastructure.

Table S3: Comparison of populations exposed at 1 km buffer including pipelines using WorldPop or LandScan population layers.

Type of Exposure	WorldPop	LandScan Global	Percent Difference
Total	462,794,372	452,910,802	2.16
Critical Ecosystems	310,500,206	309,088,807	0.46
Low Environment Regulatory Regime Index	106,887,136	104,297,980	2.45
International Ownership with No Due Diligence	31,018,846	29,978,095	3.41
Indigenous Land	13,884,394	13,299,079	3.56

806




Figure S6: The number of planned fossil fuel facilities by infrastructure type.




Figure S7: Fossil fuel infrastructure and the level of ecological risk of planned locations based on critical ecosystem indicators.




Figure S8: Number of additional people living within 5 km of planned fossil fuel infrastructure.




Figure S9: Comparison of Global Energy Monitor (GEM) oil and gas infrastructure 5 kilometer buffered extents with the Office of Fossil Energy's National Energy Technology Laboratory (NETL) oil and gas fields and wells extents for the United States of America.