

*This article is a non-reviewed preprint published at EarthArXiv.*

## **Supraglacial pond evolution in the Everest region, central Himalaya, 2015-2018.**

**Caroline J. Taylor<sup>1\*</sup>, J. Rachel Carr<sup>1</sup>**

<sup>1</sup>School of Geography, Politics and Sociology, Newcastle University, Newcastle Upon Tyne, NE1 7RU, UK.

**\* Correspondence:**

Caroline J. Taylor

[C.Taylor11@newcastle.ac.uk](mailto:C.Taylor11@newcastle.ac.uk)

# 1 **Supraglacial pond evolution in the Everest region, central Himalaya, 2015-2018.**

2 **Caroline J. Taylor<sup>1</sup>, J. Rachel Carr<sup>1</sup>**

3 <sup>1</sup>School of Geography, Politics and Sociology, Newcastle University, Newcastle Upon Tyne, NE1 7RU, UK.

4

5 **Correspondence:** Caroline J. Taylor ([C.Taylor11@newcastle.ac.uk](mailto:C.Taylor11@newcastle.ac.uk))

## 6 **Abstract**

7 Supraglacial ponds are characteristic of debris-covered glaciers and can greatly enhance local melt rates. They  
8 can grow rapidly and coalesce to form proglacial lakes, presenting a major hazard. Here, we use Sentinel-2A  
9 satellite imagery (10 m) to quantify the spatiotemporal changes of 6,425 supraglacial ponds for 10 glaciers in  
10 Everest region of Nepal between 2015 and 2018. During the study period, ponded area increased on all glaciers,  
11 but showed substantial temporal and spatial variation. The rate of pond growth accelerated compared to 2000-  
12 2015 (Watson et al., 2016). Both Imja and Spillway Lake expanded and Khumbu Glacier continued to develop a  
13 chain of connected ponds. 54% of ponds were associated with an ice-cliff, but the proportion of ponds with cliffs  
14 decreased during the study period. Pond location showed limited correspondence to slope, but favoured areas  
15 of lower surface velocity. Ideal conditions for pond formations have advanced up-glacier, and are now  
16 predominantly found at mid-elevations. Results indicate high-resolution imagery (< 10 m) is essential, as using  
17 Landsat data would miss 55–86 % of the total ponds found. Finally, glaciers were classified by stage of  
18 development (Komori, 2008; Robertson, 2012), with two transitioning between 2015 and 2018, suggesting lakes  
19 in the region are evolving rapidly. Furthermore, some glaciers displayed characteristics of multiple classes, so  
20 we propose an adapted classification system. Overall, our results demonstrate a trend of pond expansion in the  
21 Everest region and highlight the need for continued monitoring for hazard assessment.

22 **Key Words:** *Supraglacial ponds, remote sensing, outburst floods, hazards.*

## 23 **1. Introduction**

24 Mass loss from glaciers in the Himalayas has increased rapidly over the past 30 years, in response to climate  
25 change (e.g. Bolch et al., 2012; Kääb et al., 2012; Quincey et al., 2009; Gardelle et al., 2013). Here, ‘summer-  
26 accumulation type’ glaciers rely on summer-monsoon snowfall for mass gain (Bolch et al., 2012), which is  
27 thought to be reducing as temperatures rise (Fujita, 2008). The shrinkage of these freshwater reservoirs will  
28 have significant regional- and local-scale impacts (Immerzeel et al., 2010; Bolch et al., 2012; Dehecq et al.,  
29 2018). Seasonal monsoonal rainfall is the dominant source of water in the Himalaya, but glacial meltwater  
30 provides up to 40% of water supplies during the dry season (Immerzeel et al., 2010). Thus, the consistent  
31 negative mass balance of Himalayan glaciers may lead to long-term reductions in perennial flow supplied to  
32 major rivers outside of monsoon season (Xu et al., 2009), and could threaten the water and food security of an  
33 estimated 70 million people in the densely populated downstream catchments (Immerzeel et al., 2010).  
34 Additionally, increased supraglacial meltwater storage will likely increase the frequency of glacier related  
35 hazards in the region, particularly from Glacier Lake Outburst Floods (GLOFs) (Thompson et al., 2010). GLOFs are  
36 highly destructive and Nepal (along with Bhutan), has been identified as the most economically vulnerable to

37 these hazards (Carrivick and Tweed, 2016), making it vital to assess how the GLOF risk will evolve with climate  
38 warming.

39 The Everest region comprises three catchments spanning the Nepal/Tibet border in Eastern Nepal (Fig.  
40 1; King et al., 2017). These catchments have experienced atmospheric warming since the mid-1970's (Shrestha  
41 et al., 1999; Shrestha and Aryal, 2011) and weakened summer monsoons, which has reduced glacier  
42 accumulation (Salerno et al., 2015). Consequently, glacial mass balance in the region has been strongly negative  
43 since the 1970s (Bolch et al., 2008; Benn et al., 2012; Kääb et al., 2012). For example mass balance in the Everest  
44 Region of Nepal was  $-0.22 \pm 0.12$  m w.e.a<sup>-1</sup> between 1999 and 2011 (Gardelle et al., 2013). Glaciers in the Everest  
45 region are characterised by ice-surface rock debris, which is sourced from the surrounding hillslopes and covers  
46 approximately ~80% of the glaciated area in the region (Fushimi et al., 1980; Sakai et al., 2000; Watson et al.,  
47 2016). The presence of this surface debris substantially alters the glacier mass balance gradient, in comparison  
48 to clean-ice glaciers: melt is suppressed close to the termini, due to the presence of thick debris which insulates  
49 the underlying ice, and enhanced up glacier, where debris is thinner and increases melt (e.g. Quincey et al., 2009;  
50 Nicholson and Benn, 2013). As a result, the glaciers are characterised by high elevation accumulation areas, and  
51 lower-elevation ice tongues, which have low surface slopes and are near-stagnant, due to the low driving  
52 stresses (e.g. Bolch and Kamp, 2006; Quincey et al., 2009; Bolch et al., 2011; Dehecq et al., 2015). Consequently,  
53 glaciers in the Everest region lose mass by widespread surface lowering (i.e. down-wasting), rather than through  
54 terminus retreat (Quincey et al., 2007; King et al., 2017).

55 The low surface slopes and slow ice velocities that characterise debris-covered glacier tongues in the  
56 Everest region (Quincey et al., 2007; King et al., 2017), and the Himalaya more broadly (e.g. Kääb, 2005) facilitate  
57 supraglacial pond formation, by allowing glacial meltwater and rainfall to accumulate in depressions on the  
58 glacier surface (Reynolds, 2000; Sakai et al., 2000; Miles et al., 2016 and 2017). These ice surface lakes can then  
59 coalesce to form a larger proglacial lake, which carries the risk of producing GLOFs (e.g. Richardson and Reynolds,  
60 2000; Quincey et al., 2007; Thompson et al., 2012; Mertes et al., 2017). Supraglacial ponds are highly variable in  
61 character (e.g. in shape, size, turbidity and ice-cliffs), dynamic in nature, and are expected to become  
62 increasingly prevalent in a warming climate (Thompson et al., 2016; Miles et al., 2017; Watson et al., 2016; 2017a  
63 and 2017b). Understanding the spatial and temporal patterns of pond growth is therefore vital for accurately  
64 forecasting proglacial lake growth and the associated hazard of GLOFs (Richardson and Reynolds, 2000; Quincey  
65 et al., 2007; Benn et al., 2012).

66 As well as posing a hazard through the formation of proglacial lakes, supraglacial ponds represent  
67 hotspots for ablation on low-gradient debris-covered tongues (Miles et al., 2017; Watson et al., 2017a and  
68 2017b). This is because they have a comparatively low albedo and therefore absorb more insolation, which they  
69 transmit to surrounding ice, resulting in higher melt rates (e.g. Reynolds, 2000; Benn et al., 2001; Röhl, 2008;  
70 Miles et al., 2016; Watson et al., 2016; Mertes et al., 2017; Salerno et al., 2017). These enhanced melt rates  
71 cause the ponds to expand both horizontally, through subaerial and sub-aqueous melting at the margins, and  
72 vertically, via basal melting (e.g. Sakai et al., 2000; Röhl, 2008; Mertes et al., 2017). Variations in the spatial  
73 distribution of ponds across a glacier is thought to be governed by surface slope and velocity (e.g. Reynolds,

74 2000; Bolch et al., 2008; Quincey et al., 2009), and results in differential surface melt rates (Sakai et al., 2000;  
75 Benn et al., 2001; Miles et al., 2016; Watson et al., 2016; Mertes et al., 2017). Once formed, ponds can persist  
76 for months to years, or drain via englacial pathways (e.g. Immerzeel et al., 2014; Miles et al., 2017; Watson et  
77 al., 2017a). Patterns of pond drainage are generally governed by their interaction with crevasses and englacial  
78 features, which provide efficient drainage outlets (Benn and Lehmkuhl, 2000; Miles et al., 2017). Determining  
79 whether ponds drain regularly or persist is important, because long-duration ponding can cause substantial ice-  
80 surface melt, whereas repeated pond drainage events can convey energy into the glacier's interior (Miles et al.,  
81 2017), resulting in quite different ice loss patterns. Additionally, recent studies (e.g. Brun et al., 2017; Buri et al.,  
82 2016; Mertes et al., 2017; Watson et al., 2017b) suggest that ice-cliffs play a significant role in pond formation,  
83 by enhancing marginal pond melt and subaerial calving. Furthermore, as ponds expand, ice and debris influx into  
84 the pond from retreating cliff-tops increases, causing pond turbidity to increase and subsequently reducing  
85 albedo, initiating a positive feedback of melt (Mertes et al., 2017). These factors complicate predicting future  
86 supraglacial pond formation and evolution and make them highly dynamic features.

87 Previous studies have documented changes in supraglacial water storage across the Himalaya (Table 1;  
88 e.g. Wessels et al., 2002; Kattlemann, 2003; Bajracharya and Mool, 2007; Gardelle et al., 2011). However, the  
89 use of comparatively coarse resolution imagery (e.g. 30m resolution Landsat imagery and 15m resolution ASTER  
90 imagery) means substantial water volumes may be missed and the low repeat frequency makes differentiating  
91 between pond persistence and regular drainage difficult, which is important for quantifying the impact of the  
92 ponds on mass loss (Immerzeel et al., 2014; Miles et al., 2017). Watson et al. (2016), presented the first high  
93 temporal and spatial resolution study of supraglacial pond evolution in the Everest region, for the period 2000-  
94 2015, where a total of 9340 ponds were identified. Here, we extend this previous work to quantify supraglacial  
95 pond evolution between 2015 and 2018. This will provide an up-to-date picture of pond coverage within the  
96 region. Our main objectives are; (1) characterise the spatial and temporal evolution of supraglacial ponds, to  
97 determine the magnitude and extent of change since 2015, (2) assess the impacts of using higher resolution  
98 imagery versus lower resolution imagery on the ability to identify and assess supraglacial ponds (3) examine the  
99 impact of local glacier characteristics (glacier surface slope, ice velocities and the presence of ice cliffs) on pond  
100 formation location, area and number and (4) classify the stage of proglacial lake development across the region.

## 101 **2. Methods**

### 102 2.1 Study Site and Water-body Definitions

103 The study focuses on ten debris-covered glaciers that drain the Dudh Koshi basin, within Sagarmatha National  
104 Park, Eastern Nepal (Fig. 1). The glaciers flow predominantly in a southerly direction, with the exception of Ama  
105 Dablam (north flowing) and Imja (west flowing) glaciers (Fig. 1). All of the study glaciers have extensive debris-  
106 covered tongues and high accumulation areas. Ngozumpa, Pangbung and Khumbu glaciers have the greatest  
107 length (~15 km, ~13 km and ~11 km respectively) and Imja Glacier the shortest at ~2 km (Table 2). Of the 10  
108 glaciers in this study, 8 were also included in Watson et al. (2016) study for the period 2000-2015 (Table 2). We  
109 add Pangbung Glacier and Sumna Glacier to our study, as they are important for downstream water resources

110 and could potentially pose a threat to downstream communities due to GLOFs (Immerzeel et al., 2010). The  
111 distinction between glacial 'pond' and 'lake' remains poorly defined in the Himalayan literature, with Watson et  
112 al. (2016) referring to all surface water as ponds unless specifically named otherwise, and other authors  
113 switching between the terms 'lake' and 'pond' (e.g. Gardelle et al., 2011; Nie et al., 2013). Here, we use the term  
114 pond to refer to all bodies of water on the glacier surface. Proglacial lakes are discussed separately from ponds,  
115 and are defined as all water bodies that are outside of the glacier margin, but are in contact with it. Following  
116 this classification, 'Spillway Lake', located at the terminus of Ngozumpa Glacier, is classified as a pond as it  
117 remains bound by glacier ice on all sides, but here is discussed separately, as its very large area would skew  
118 results. Imja Lake, located at the front of the Imja/Lhotse Shar glacier complex is classed as a proglacial lake.

## 119 2.2 Data Sources

120 This study used true-colour orthorectified Sentinel 2A imagery (<10 m resolution) (available from USGS at  
121 <http://earthexplorer.usgs.gov/>) for the time period December 2015 to April 2018 (SI. Table 1). Images outside  
122 of the monsoon season (June-September) were chosen, to minimise cloud cover and where possible, images  
123 were selected from the same month to avoid inadvertently including seasonal differences into our analysis (SI.  
124 Table 1). True colour images were derived from combining the blue (490 nm), green (560 nm) and red (665 nm)  
125 bands. For each of the study years, two true-colour Sentinel-2A images from the same date were mosaicked to  
126 produce one spatially continuous dataset of the entire region. Glacier outlines were obtained from the Randolph  
127 Glacier Inventory 5.0 (available from GLIMS at <http://www.glims.org/maps/glims>) and modified manually to  
128 reflect the debris-covered area of each glacier. These were then used as glacier / land masks and only ponds  
129 located within this mask were included in the study.

## 130 2.3 Maximum Likelihood Classification and Manual Editing

131 A supervised classification technique was used to automatically delineate supraglacial ponds. First, we manually  
132 selected training sites that contained the primary land cover classes (e.g. clean ice, water, debris covered ice  
133 etc.) from the true-colour Sentinel 2A image. We then performed the Maximum Likelihood Classification (MLC)  
134 on the true-colour image, plus bands 5 and 7 (near infrared and thermal wavelengths respectively): both infrared  
135 and thermal wavelengths are absorbed by water bodies so their addition aided the classification substantially.  
136 Whilst other classifications can be used (e.g. the Hierarchical Knowledge Based Classifier (HKBC) method),  
137 previous studies suggest that MLC is the most accurate classification method for delineating water stores on  
138 glaciers (Tiwari et al., 2016). This method was repeated for all image dates. The classification results were  
139 assessed manually, by comparing automatically detected pond margins to the underlying imagery. We then  
140 manually edited any ponds where the classification had failed to accurately detect the pond margins. In total,  
141 we identified 6,533 ponds, and then extracted key statistics for analysis, specifically ponded area and number  
142 of ponds.

## 143 2.4 Controls on Pond Location

144 We assessed controls on supraglacial pond formation and growth patterns, specifically: glacier surface slope, ice  
145 velocities and the presence of ice cliffs. We derived a slope map and glacier elevation profiles from the ASTER

146 DEM (<http://earthexplorer.usgs.gov/>) to identify glaciers with particularly low slope gradients and thus potential  
147 areas for future pond development. Glacier velocities were derived from repeat-image feature tracking of  
148 Landsat images by Dehecq et al. (2015). The data used in this study are derived from average velocities for 2013-  
149 2015 and have a 120 m spatial resolution (Dehecq et al., 2015). Each of the study glaciers were divided into 10  
150 bands, representing 10% glacier surface area, to facilitate comparison of pond locations with surface slope and  
151 ice velocities.

### 152 **3. Results**

#### 153 3.1. Supraglacial Pond Change

##### 154 3.1.1. Regional Poned Area Change

155 Across the study region, total poned area increased on all 10 glaciers between 2015 and 2018 (Fig. 2). The most  
156 prominent changes in both pond number and area were observed on the three largest glaciers (SI. Table 2;  
157 Ngozumpa, Pangbung and Khumbu glaciers), which contain ~58% of the total 6,533 ponds identified in this study  
158 (Fig. 2). Specifically, poned area increased by 255,849 m<sup>2</sup> (33.3 %) on Ngozumpa Glacier, 191,386 m<sup>2</sup> (36.2 %) on  
159 Pangbung Glacier and 134,299 m<sup>2</sup> (43.1 %) on Khumbu Glacier between December 2015 and April 2018 (Fig.  
160 2). This increase mainly resulted from pond coalescence on Ngozumpa and Khumbu glaciers, as the number of  
161 ponds decreased but the poned area increased (SI. Fig. 1): the number of ponds decreased by 60 (13.5%  
162 decrease) on Ngozumpa and 3 (1%) on Khumbu (Fig. 2.). The remaining seven smaller glaciers showed an  
163 increase in poned area between 2015 and 2018, ranging from 9,664 m<sup>2</sup> on Imja Glacier (30 % increase) to  
164 120,162 m<sup>2</sup> on Lhotse Glacier (68 % increase). This increase in area was a result of both the establishment of  
165 new ponds and the coalescence of existing smaller area ponds (Fig. 2), for instance poned area on Ama Dablam  
166 Glacier increased 38,958 m<sup>2</sup> and pond number increased by 29.

167 Whilst total poned area on all glaciers increased from 2015-2018 (SI. Table 2), the number of ponds  
168 found on each glacier varied (Fig. 2). Only 2 of the 10 glaciers showed an increase in pond number: on Pangbung  
169 Glacier the number of ponds increased by 4, to a total of 200 ponds, and on Ama Dablam Glacier the number of  
170 ponds increased by 29, to a total of 67 ponds (December 2015- April 2018). The remaining 8 glaciers exhibited  
171 an overall decrease in the number of ponds, ranging from a reduction of 2 (Lhotse Glacier) to 60 (Ngozumpa  
172 Glacier). Glaciers with decreasing pond numbers generally showed an increase in poned area in 2015, 2017 and  
173 2018 (SI. Fig. 1). The exception to this inverse relationship between pond number and poned area was in 2016  
174 (SI. Fig. 1). During this year, a large decrease in pond number was observed on glaciers in the eastern part of the  
175 study area, particularly Lhotse, Lhotse Shar and Lhotse Nup glaciers (Fig. 2b). At the same time, there was a large  
176 increase in pond number on glaciers in the western part of the study area (Fig. 2c). For example Pangbung Glacier  
177 increased in pond number from 196 (2015) to 468 (2016), and Sumna Glacier increased from 78 (2015) to 182  
178 (2016). Overall, our results show that poned area increased on all glaciers during the study period 2015-2018,  
179 whereas the number of ponds generally declined, despite the considerable spatial variability observed in 2016  
180 (Fig. 2c).

181 Two water bodies in the study area were assessed separately, due to their large size; supraglacial lake  
182 'Spillway Lake' on the terminus of Ngozumpa Glacier, and proglacial lake 'Imja Lake' fronting the Imja/Lhotse  
183 Shar Glacier complex (Fig. 4). Spillway Lake, located at the terminus of Ngozumpa Glacier, underwent a net gain  
184 of 40,565 m<sup>2</sup> during the study period (December 2015 - April 2018) and reached a size of 286,367 m<sup>2</sup>. This  
185 remains the largest surface water store in the study region. The only proglacial lake identified in this study was  
186 Imja Lake, which expanded up-glacier by 455.6 m to reach an area of size of 1,493,142.68 m<sup>2</sup> by 2018. Whilst  
187 these two water bodies are currently the only very large water bodies in the study area, our data show  
188 substantial growth and coalescence of surface ponds on Pangbung and Khumbu glaciers (Fig. 3b, c).

### 189 3.1.2 Glacier-scale Pond Changes

190 Despite the overarching trends across the region, changes in supraglacial pond number and area varied from  
191 glacier to glacier and across individual glaciers (Fig. 2; Table 2). For the largest three glaciers, most of the ponded  
192 area was located near the terminus, and this persisted throughout the study period (Fig. 3a-c). The number of  
193 ponds, although demonstrated an overall decrease, remained relatively similar over the four year period, but  
194 ponded area increased (Fig. 2), which was primarily as a result of coalescence on the lower glacier tongues. (Fig.  
195 3a-c). For example, on the terminus of Ngozumpa Glacier, two smaller ponds (Fig. 3a, i and ii) formed new  
196 branches of Spillway Lake in March 2017. This reduced the total number of ponds by 2, whilst increasing the  
197 total ponded area of Spillway Lake. Similarly, along the eastern margins of Khumbu Glacier, lateral pond  
198 expansion between March 2016 and March 2017 (Fig. 3b, i and ii) resulted in coalescence of two major ponds  
199 and a reduction in the number of surface ponds by 2. Our data therefore suggests that supraglacial pond  
200 expansion on the larger glaciers in our study area results from pre-existing ponds coalescing, rather than by the  
201 formation and growth of new ponds.

202 On the seven smaller glaciers, there was substantial spatial and temporal variability in the number and  
203 area of surface ponds, both between the glaciers and across individual glaciers (Fig. 3d and Fig. 4). For example,  
204 Lhotse Shar and Imja Glacier neighbour each other, in the east of the study region (Fig. 1). However in April 2018,  
205 Lhotse Shar Glacier had over three times the ponded area (135,420 m<sup>2</sup>) of Imja Glacier (42,603 m<sup>2</sup>) as well as  
206 almost three and a half times the number of ponds (96 and 28 respectively). Sumna Glacier in the west of the  
207 region showed major variations in ponded area and number over time, as its percentage pond cover ranged  
208 from 2.16 % to 4.87 % during the four year study period. For most of the smaller glaciers, ponds occupied similar  
209 locations at each time step (e.g. Ama Dablam Glacier SI. Fig. 2a and Lhotse Glacier SI. Fig. 2b), in addition to the  
210 growth of new ponds (e.g. Lhotse Glacier SI. Fig. 2b). Sumna Glacier was the exception to this and showed a  
211 distinctive change in the spatial pattern of the pond locations during the study period: at the start of the study  
212 (2015), it had more ponds at higher elevations, but by the end (2018), ponds were most concentrated near the  
213 terminus (Fig. 3d). Overall, our results show an increase in both ponded area and number of ponds on the  
214 smaller glaciers, but this showed substantial spatial and temporal variation, even between neighbouring glaciers  
215 and on the same glacier.

## 216 3.2 Controls on Pond Location

### 217 3.2.1 Glacier Elevation Profile

218 The study glaciers have an average slope  $> 18^\circ$  at the tongue and  $< 25^\circ$  at higher elevations. In general, areas  
219 where the mean overall slope is lower ( $>10^\circ$ ) contain more ponds Overall, slope angle tends to decrease closer  
220 to the glacier termini, where slopes are generally between  $2^\circ$  and  $4^\circ$  (Fig. 5). However, with the exception of  
221 Sumna Glacier (see section 3.1.2.) there is no apparent relationship between elevation and pond number/ area  
222 on the study glaciers (Figs. 5 and 6). We assessed this in further detail by dividing each glacier into 10 equal  
223 elevation bands (to account for differences in total length and to facilitate direct comparison between glaciers)  
224 and calculating the number and area of ponds in each elevation band (Figs. 5 and 6). Bands are numbered from  
225 1 (the glacier head wall) to 10 (terminus). On six of the ten study glaciers, the number of ponds decreased from  
226 the head of the glacier to terminus, whilst the pattern was reversed on the remaining 4, so that pond numbers  
227 increased with distance up glacier (Fig. 5 and 6). Furthermore, all the study glaciers showed large variations in  
228 ponded area and number between individual elevation bands. For all study glaciers, the largest number of ponds  
229 were usually found in the central elevation bands (bands 5-6), as exemplified by Pangbung (32.2 % of ponds),  
230 Khumbu (23.1 %) and Lhotse Shar (47.8 %) glaciers (Fig. 5). The exception to this trend can be seen on Imja,  
231 Lhotse and Sumna glaciers, where the highest numbers of ponds can be found nearer to the terminus (i.e.  
232 elevation bands 9 and 10), and Ama Dablam Glacier where most ponds are located nearer the high accumulation  
233 zone (bands 1-2; Fig. 5). Furthermore, the number of ponds was much higher on Imja Glacier's terminus (band  
234 10; 6 ponds; 21.4 %) than on any of the other study glaciers. No ponds were identified in the high accumulation  
235 zone (band 1) for four of the ten study glaciers; Imja, Nuptse, Lhotse Shar and Sumna (Fig. 5 and 6). Sumna  
236 Glacier also displayed distinctive areas of higher frequency ponding (bands 6-8) and lower frequency ponding  
237 (bands 2-4) which showed no clear relationship with elevation (Fig. 3d; Fig. 5). A number of breaks in slope were  
238 identified on six of the ten study glaciers: Ama Dablam, Imja, Lhotse, Lhotse Shar, Lhotse Nup and Sumna (Fig.  
239 5). In the band immediately down glacier of the break in slope, there was an increase in pond number on four  
240 of the glaciers (Imja, Lhotse, Lhotse Shar, and Sumna; Fig. 5). This was most notable on Lhotse Glacier, where  
241 there was a change in slope in band 1 and the number of ponds increased from 1 (band 1) to 20 (Band 2) (Fig.  
242 5). Ama Dablam and Lhotse Nup were the exception to this trend, where on Ama Dablam Glacier there was a  
243 decrease in pond number from 8 to 4, following the break in slope in band 3 (Fig. 5) whilst on Lhotse Nup a break  
244 in slope in band 6 preceded a decrease in pond number from 7 to 1 (Fig. 5).

### 245 3.2.2. *Glacier Velocity*

246 Generally, glacier velocity decreased from source to terminus, being highest in bands 1-3 and lowest in the bands  
247 8-10 (Fig. 7). Where velocities were higher, the number of surface ponds was generally lower (Fig. 7). For  
248 instance, in bands 1 and 2 on Lhotse Shar Glacier there were no ponds recorded (velocity  $>20 \text{ m a}^{-1}$ ). However,  
249 further down glacier from band 3, velocities were lower ( $<20 \text{ m a}^{-1}$ ) and the number of ponds increased by 14  
250 (14.6 %) (Fig. 7). The main exception to this trend is Nuptse Glacier, where velocity is low in bands 1-2 ( $> 2 \text{ m a}^{-1}$ )  
251 and there are no ponds, but the area of higher velocity in bands 4 to 5 ( $>14 \text{ m a}^{-1}$ ) contains a total of 34 ponds  
252 (Fig. 7). In general, where velocities are lowest, for instance nearer the glacier terminus, pond number increases,  
253 following a similar trend to that of glacier slope (section 3.2.1.). For example, from band 5 onwards on Lhotse  
254 Shar Glacier, there is almost no recordable velocity and pond number reaches its highest (23 ponds) (Fig. 7). The



255 relationship between total pond area and glacier velocity is similar to that for pond number (Fig. 8): higher  
256 velocities coincide with lower ponded area, whereas lower velocities have a higher ponded area (Fig. 8). Sumna  
257 Glacier displays this relationship clearly: it has 23 % of its ponds in Bands 2-4 and 70 % in bands 6-8. This  
258 corresponds to velocities of 3 to 7  $\text{m a}^{-1}$  and  $< 3 \text{ m a}^{-1}$  respectively. Overall, our results indicate that lower  
259 velocities correspond with higher ponded area and pond number, and higher velocities generally relate to fewer  
260 ponds.

### 261 3.2.3. Ice Cliffs

262 Although ice cliffs were identified on all 10 glaciers 2015-2018, the number of cliffs changed markedly during  
263 this period, with cliffs seen to increase and decrease from year to year, and between glaciers (Fig. 9). Ngozumpa  
264 and Khumbu glaciers had the highest percentage area of the glacier covered by ice cliffs, with 4.3% and 3.92%  
265 respectively, and Sumna glacier the least (1.1%; Fig. 9). The greatest temporal variability was observed on Ama  
266 Dablam Glacier, where ice-cliff coverage decreased from 2.59 % in 2015 to 1.3 % in 2016, and then rapidly  
267 increased to 3.3 % in 2018 (Fig. 9). The number of supraglacial ponds with a corresponding ice-cliff exceeded the  
268 number without: on average, across all of the study glaciers, 54 % of ponds had a coincident ice-cliff (Fig. 9).  
269 During the study, the number of ponds without an ice cliff increased on average by 1.6 % of the total glacier  
270 surface area and this was most noticeable on the seven smaller glaciers (Fig. 9). For instance, on Sumna Glacier,  
271 the area of the ponds with a cliff remained relatively stable ( $\sim 0.8 \%$ ), whereas the area of ponds without a cliff  
272 increased from 1 % in 2015 to 1.75 % in 2018 (Fig. 9). In comparison, the number of ponds with an ice cliff  
273 increased by just 0.9 %, observed most notably on the three larger glaciers (Fig. 9). For example, on Ngozumpa  
274 Glacier, ponds with cliffs average 4.3 % compared to just 1.25 % for ponds without cliffs. This indicates pond  
275 growth can occur irrespective of ice cliff presence. 2018 saw a marked increase in ponded area without an  
276 adjacent ice cliff, increased on average by 1.4 % (average of 9 glaciers) with the exception of Sumna Glacier  
277 where there was a decrease (0.32 %).

### 278 3.3 Future Lake Development

279 Each of the 10 study glaciers was assigned a number, according to the stage of lake development described in  
280 established lake classification schemes (Komori, 2008; Robertson, 2012; Table 3). In 2015 (Fig. 10b), the study  
281 region was dominated by glaciers in Stage 2 of lake development, with 60% showing ponds that have coalesced,  
282 were ice-dammed and have large areas ( $>20,000\text{m}^2$ ). Only Ngozumpa Glacier was defined as Stage 3, due to the  
283 presence of the large terminal Spillway Lake (Fig. 10). The remaining three glaciers (Sumna, Lhotse Nup and Ama  
284 Dablam Glaciers) were all classified as Stage 1, with supraglacial ponds forming in their lower ablation zones (Fig.  
285 10b).

286 Over the four-year study period, (December 2015- April 2018) two glaciers (Ama Dablam and Lhotse Nup)  
287 transitioned to a new stage of lake development (Fig. 10). Both progressed from Stage 1, where a few  
288 supraglacial ponds were identified, to Stage 2, where ponds had begun to coalesce (Fig. 10). Two glaciers  
289 (Pangbung and Lhotse Shar) partially transitioned from Stage 1 to Stage 2, and from Stage 2 to Stage 3,  
290 respectively (Fig. 10). Features that fit more than one stage were identified on these two glaciers, meaning that

291 they could not be assigned a stage using the current classification. For example, on Pangbung Glacier, ponds  
292 were appearing on the lower ablation zone (characteristic of Stage 1), but some ponds were also beginning to  
293 coalesce (Stage 2; Fig. 3c). On Lhotse Shar Glacier, coalescing was observed (Stage 2), but there was also stable  
294 expansion of its proglacial lake (Stage 3). As a result, the current development cannot be captured by existing  
295 schemes.

## 296 **4. Discussion**

### 297 4.1. Changes in Supraglacial Pond Area 2015-2018

298 The area of supraglacial ponds in the study increased markedly between 2015 and 2018, ranging from a 13.6 %  
299 increase on Sumna Glacier to a 108.1 % increase on Lhotse Nup Glacier, despite showing large inter-annual  
300 variations (Fig. 2). These increases show a marked acceleration in pond growth, compared to 2000-2015 (Watson  
301 et al., 2016). For instance, ponded area increased at a rate of  $1.6 \% a^{-1}$  on Nuptse Glacier between 2000 and 2015  
302 (Watson et al., 2016), whereas our study measured a rate of  $22.6 \% a^{-1}$  between 2015 and 2018 (SI. Table 2).  
303 Similarly, the rate of expansion on Lhotse Glacier in our study ( $17.0 \% a^{-1}$ ) is four times greater than that found  
304 by Watson et al. (2016;  $4.1 \% a^{-1}$ ). This is a major concern in terms of risks to downstream communities, as these  
305 very high rates of pond growth will rapidly increase the water volumes available for outburst floods and will also  
306 encourage pond coalescence and lake formation.

307 One potential explanation for the observed acceleration in pond growth is climatic controls: warmer  
308 temperatures should increase melt rates and hence encourage pond expansion, whilst increased precipitation  
309 could add water directly to the ponds. Data on climate trends proximal to our study glaciers are very limited.  
310 However, available data (Salerno et al., 2015) suggest that minimum and mean air temperatures have risen in  
311 the Everest area between 1994 and 2013, at elevations above 5000 m. However, warming was most marked in  
312 spring and winter, and would thus have a more limited impact on ice melt, and it was concurrent with a reduction  
313 precipitation, which would decrease direct inputs to the ponds (Salerno et al., 2015). As such, we suggest that  
314 the observed increase in ponding may at least partly reflect changes in the dynamics of our study glaciers, which  
315 provide the conditions that promote pond formation. Between 2000 and 2017, glaciers in East Nepal decelerated  
316 by  $-1.8 \pm 0.1 ma^{-1}$  ( $17.0 \pm 1\% a^{-1}$ ) and thinned, which in turn reduced driving stresses (Dehecq et al., 2018). Down-  
317 wasting and deceleration creates an inverted mass balance gradient and an uneven glacier surface, which  
318 together facilitate pond formation (Reynolds, 2000; Miles et al., 2016 and 2017). Furthermore, slow flow is likely  
319 to reduce the number of crevasses forming and would thus reduce the chance of pond drainage (Immerzeel et  
320 al., 2014; Miles et al., 2017; Watson et al., 2017a and 2017b). As such, we suggest that recent changes in ice  
321 dynamics in the Everest region are likely to be contributing to observed pond growth and that this may lead to  
322 a positive feedback, whereby rapid pond growth accelerates down wasting, leading to further pond expansion.

### 323 4.2. Glacier-scale Ponded Area Patterns

324 The three larger glaciers (Ngozumpa, Khumbu and Pangbung) increased in ponded area between 2015 and 2018,  
325 but decreased in pond number (Fig. 2). This suggests that pond area is increasing via coalescence, as seen at the  
326 terminus of Ngozumpa Glacier (Fig. 3a) and the eastern margins of Khumbu Glacier (Fig. 3b). This has implications

327 for both glacier lake related hazards and ice loss rates in the future. As ponds continue to join, and the area of  
328 supraglacial ponds increases, the likelihood of proglacial lakes formation is increased (Komori, 2008; Robertson,  
329 2012), which in turn increases the risk of GLOFs (e.g. Richardson, 2000; Quincey et al., 2007; Benn et al., 2012;  
330 Rounce et al., 2017). In addition, the larger ponded surface area increases the wind fetch, which may lead to  
331 enhanced undercutting at pond margins and thus increase glacier melt rate (Benn et al., 2001; Röhl, 2006, 2008;  
332 Sakai et al., 2009). Our data demonstrate that the three largest glaciers had low velocities across their tongues:  
333 on Ngouzmpa and Pangbung glaciers, velocity remained below  $12 \text{ m a}^{-1}$  for all 10 bands, whilst on Khumbu Glacier  
334 velocities started high ( $> 30 \text{ m a}^{-1}$ ) in band 1, but rapidly reduced to below  $12 \text{ m a}^{-1}$  for the remaining 9 bands (Fig.  
335 7 and 8). We suggest that these slow velocities promoted pond coalescence and growth (Benn et al., 2012, 2017)  
336 and also reduced the frequency of pond drainage, by limiting the number of open crevasses (e.g. Miles et al.,  
337 2017), and thus increasing pond area.

338 Generally, the smaller glaciers in the region do not have large, extensive ponding at their termini, but  
339 the ponded area on all seven of the smaller glaciers increased during the study period (Fig. 2). For example, the  
340 ponded area and pond number found on Ama Dablam Glacier increased by  $38,958 \text{ m}^2$  (48 %) and 29 (43 %)   
341 respectively 2015 to 2018 (Fig. 2b, c). Given the increase in area of the ponds but variations in pond number, we  
342 suggest that this is primarily due to the coalescing of smaller area ponds (e.g. Ama Dablam Glacier SI. Fig. 3) but  
343 also through the formation of new ponds, which is consistent with earlier observations (Watson et al., 2016).  
344 This may enhance melt rates, by increasing the fetch across the pond and increasing the ponded area (Sakai et  
345 al., 2009), and potentially lead to proglacial lake development. Ice velocities are generally higher on the smaller  
346 glaciers, and more spatially variable, which may limit the opportunity for pond growth and/or encourage  
347 crevasses formation and thus promote drainage (e.g. Immerzeel et al., 2014; Miles et al., 2017; Watson et al.,  
348 2017a).

#### 349 4.3. Evaluating the use of Sentinel Data for Remote Sensing Studies

350 Previous studies of supraglacial and proglacial lakes changes in the Everest region have largely used Landsat  
351 imagery (30 m, 1 pixel =  $900 \text{ m}^2$ ) (Table 1; e.g. Gardelle et al., 2011; Nie et al., 2013; Zhang et al., 2015). Here we  
352 use Sentinel 2, which is 10 m resolution (1 pixel =  $100 \text{ m}^2$ ). Our results demonstrate that ponds  $< 100 \text{ m}^2$  (one  
353 pixel in Sentinel data) accounted for 3% - 8% and those  $< 400 \text{ m}^2$  (four pixels in Sentinel data) comprised 28% -  
354 59% of total ponds found in the region (SI. Fig. 4). Of the total ponds identified, between 55 % and 86 % were  
355 below  $900 \text{ m}^2$  (one pixel in Landsat data). As such, using Landsat imagery to map pond changes in the region  
356 would have missed the majority of the total number of ponds. Whilst these ponds are comparatively small in  
357 area, including them in assessments is vital, as they inform us about where ponds are nucleating, and hence  
358 controls on their formation. These data also indicate locations that may become ponded in the future, and  
359 therefore subject to enhanced melt rates, and/or areas that may eventually coalesce with other ponded  
360 sections. Furthermore, recent work has demonstrated that Sentinel-2 imagery has a better spectral contrast  
361 between debris-cover ice and supraglacial ponds than Landsat or RapidEye, which affirms its suitability (Watson  
362 et al., 2018).

#### 363 4.4 Controls on Pond Formation

364 4.4.1. *Glacier Elevation Profile*

365 Previous work suggests that supraglacial ponds typically begin to form on slopes  $< 10^\circ$  and larger ponds occur  
366 where surface gradients are less than  $2^\circ$ , typically found close to the glacier terminus (e.g. Quincey et al., 2007;  
367 Bolch et al., 2012). In some areas, large number of ponds coincided with slopes of  $2-4^\circ$  and pond frequency was  
368 highest at the termini of Imja, Lhotse and Sumna glaciers (Figs. 5 and 6). However, contrary to theory,  
369 supraglacial ponds were also found in areas with much greater slope gradients ( $> 10^\circ$ ), both at the termini and  
370 further up-glacier (Figs. 5 and 6). For instance, the highest number of ponds on Ama Dablam Glacier (76 %) were  
371 located in the high accumulation zone (band 1-2). We also observed changes in the number of ponds after breaks  
372 in slope (Fig. 5), which may reflect localised areas of extensional / compressional flow that would open or close  
373 crevasses, and thus facilitate or reduce pond drainage (Benn et al., 2001, 2012; Gulley and Benn, 2007; Röhl,  
374 2008; Thompson et al., 2012; Mertes et al., 2017).

375 Our data show that the greatest number and area of ponds occur in the mid-elevation bands (bands 5-  
376 6) and not at the termini (bands 8-10) of our study glaciers (Figs. 5-6). We suggest this is because of the inverted  
377 mass balance gradient observed on debris-covered glaciers in the Everest region: thick debris at the terminus  
378 suppresses melt, whereas thinner debris further up glacier enhances melt (Bolch et al., 2008; Quincey et al., 2009;  
379 King et al., 2018) This pattern of mass balance results in near-stagnant ice velocities over much of the tongue  
380 and rapid down-wasting at mid-elevations (Quincey et al., 2007, 2009; Nicholson and Benn, 2013; Juen et al.,  
381 2014; King et al., 2017), which produces uneven 'ablation topography' (Nicholson and Benn, 2013). Together,  
382 this creates ideal conditions for pond formation. Our data suggest that these conditions are now predominantly  
383 found at mid-elevations (bands 5 to 6) on glaciers in the Everest region, and that the area where ponds are able  
384 to form may be advancing up-glacier. This has important implications for total ice loss rates, as ponds strongly  
385 enhance glacier melting (Sakai et al., 2000; Benn et al., 2001; Miles et al., 2016; Watson et al., 2016; Mertes et  
386 al., 2017), and could thus expand the area of enhanced ice loss further up glacier.

387 4.4.2. *Velocity*

388 Ice velocities strongly influence pond size, pond drainage and pond persistence (Miles et al., 2017). Where  
389 velocities are low, little reorganisation of internal water pathways can occur, resulting in ineffective englacial  
390 drainage from the glacier surface, thus promoting surface ponding (Jordan and Stark, 2001; Benn et al., 2012,  
391 2017). As a result, low surface gradients at glacier tongues generally encourage pond formation and low surface  
392 velocities facilitate storage, as opposed to drainage (Reynolds, 2000; Quincey et al., 2007; Watson et al., 2017a).  
393 Our results generally support this, as there were more ponds and a greater ponded area closer to the terminus  
394 (bands 8-10), where velocities were lower (Figs. 7 and 8) For example, on Sumna, Imja and Lhoste Shar glaciers  
395 no ponds were found in bands 1 or 2 where velocity was  $> 20 \text{ m a}^{-1}$ , but as velocity decreased ( $< 20 \text{ m a}^{-1}$ ) pond  
396 numbers began to increase (Fig. 7). However, as with glacier slope, there are exceptions to this relationship. For  
397 example, the peak number of ponds corresponds with peak velocities on Lhotse and near-peak velocities on  
398 Nupste (Fig. 7). The cause of these velocity patterns is difficult to elucidate, but we suggest that it may relate to  
399 the pattern of debris inputs from the surrounding hillslopes (and hence the debris characteristics and  
400 distribution), the location of tributary glaciers and/or meltwater inputs from the surrounding slopes. Overall our

401 data indicate that ice velocities influence pond area and number, but that the relationship is far from simple and  
402 requires further detailed study.

#### 403 *4.4.3. Ice-Cliffs*

404 All ten of the glaciers included in this study had surface ice-cliffs, and the majority of ponds were associated with  
405 adjacent ice cliffs (54 %). This follows theory, as ice cliffs experience enhanced melt at a rate three to six times  
406 greater than that of debris-covered ice (Kirkbride, 1993; Benn et al., 2001; Röhl, 2008; Watson et al., 2016). Thus,  
407 ice cliffs can not only initiate pond formation through surface melt (Sakai et al., 2002; Buri et al., 2016a; Watson  
408 et al., 2017b) but also facilitate pond expansion in process of thermo-erosional notching (Josberger, 1978; Röhl,  
409 2006). Despite this, our results show that the proportion of ponds without an ice cliff increased substantially  
410 during the study period (2015-2018; Fig. 9). For example on Sumna Glacier, ponded area without an adjacent  
411 cliff almost doubled during the study, whereas ponded area with a cliff actually decreased, meaning that there  
412 were fewer ponds with an ice cliff than without in 2018 (Fig. 9). On Ngozumpa, in contrast, the percentage of  
413 ponds with cliffs were on average three times higher (4.3 %) than those without (1.25 %). Overall, our results  
414 show that ponds are expanding and that they are usually associated with ice cliffs (Figs. 2 and 9). However, as  
415 ponded area increases, the proportion of the ponds with ice cliffs reduces (Fig. 9). We speculate that this may  
416 be because pond growth is outstripping ice cliff formation, but this requires further investigation.

#### 417 *4.5. Future Lake Development*

##### 418 *4.5.1 Applicability of Lake Classification Models*

419 During this study, two of the ten study glaciers progressed to a new stage of the Komori/Robertson (2008; 2012)  
420 proglacial lake classification scheme; Ama Dablam and Lhotse Nup, both progressed from Stage 1 to Stage 2.  
421 However, Komori (2008) found glaciers in Bhutan took on average 40 years to pass through Stage 1 and 2, and  
422 enter into Stage 3, whilst those in the Aoraki/Mt Cook region took 8-30 years for the same transition (Robertson,  
423 2012). Given the short time-frame of this study (4 years), our results demonstrate that proglacial lakes in the  
424 Everest region are evolving rapidly and quicker than other regions, both in the Himalaya and globally. This has  
425 important implications for hazard assessments in the region, as these rapid changes require high temporal  
426 resolution monitoring to determine potential changes in GLOF risk and could result in hazardous lakes forming  
427 quickly in new locations. Our results therefore highlight the need for frequent monitoring of the hazards posed  
428 by glacier lake growth in the Everest region, particularly given that Nepal (along with Bhutan) is at the greatest  
429 economic risk from GLOFs (Carrivick and Tweed, 2016).

430 Two of our study glaciers exhibited characteristics of multiple classes of proglacial lake development  
431 (Fig. 10). Specifically, on Pangbung Glacier, ponds were appearing on the lower ablation zone, which is indicative  
432 of Stage 1 but some ponds were beginning to coalesce (Stage 2) (Table 3). On Lhotse Shar Glacier, we observed  
433 coalescing (Stage 2), but also the stable expansion of the proglacial lake (Stage 3). As such, the proglacial lake  
434 development observed in our study region does not fit within the four stage classification system of Komori  
435 (2008) and Robertson (2012; Table 3). Additionally, changes in ponded area, number of ponds and ice-cliffs were  
436 observed on all glaciers (Figs. 2 and 9), but these substantial differences were not accounted for in the current

437 model. For example, both Khumbu and Nuptse glaciers remained in Stage 2 throughout the study period, but  
438 experienced increases in ice-cliffs (45 % and 76 % respectively; Fig. 9). Ngozumpa Glacier remained in Stage 3,  
439 but showed a marked increase in both ponded area (33 %) and the number of ponds with ice-cliffs (24 %) (Fig.  
440 2c and Fig. 9). As such, the current classification model does not account for many of the observed changes in  
441 pond area and characteristics, which could contribute to proglacial lake growth, and does not account for  
442 glaciers that bridge different categories. Thus, on the basis of our observations, we propose a six-stage  
443 categorisation which accounts for the 'in-stage' changes observed, such as appearance of ice-cliffs and marginal  
444 pond expansion (Fig. 11). The inclusion of two additional stages (shown in red), now include 'in-stage' changes  
445 observed in this study, such as marginal expansion, ice-cliff formation and pond drainage. The inclusion of these  
446 stages would enable us to assign our study glaciers to just one stage, making the model more suitable for  
447 evaluating and communicating glacier lake hazard potential.

#### 448 *4.4.2 Lake Development Trajectories and Outburst Risk*

449 A key observation of this study was the coalescing of smaller area ponds on the eastern margins of Khumbu  
450 Glacier and the Ngozumpa terminus (e.g. Fig. 3a and b, i and ii). Despite this, both glaciers remained in the same  
451 stage of proglacial lake classification during the period (Stage 3 and Stage 2 respectively). Using our new  
452 classification scheme, Khumbu Glacier would be classified as Stage 3 (compared to currently assigned Stage 2),  
453 suggesting that Khumbu Glacier is further along the progression of proglacial lake formation than previous  
454 classifications have indicated. Our data show that Spillway Lake grew by 40,565 m<sup>2</sup> between 2015 and 2018.  
455 Previous work showed that Spillway Lake grew by ~ 10% per year (2001 to 2010) to reach 258,000 m<sup>2</sup> in  
456 December 2009 (Thompson et al., 2012) but more recently shrank by 8,345 m<sup>2</sup> between 2009 and 2015 (Watson  
457 et al., 2016). It was suggested that the reduction in area resulted from supraglacial drainage channel evolution  
458 and lowering of the hydrological base level, which paused the lake's expansion (Watson et al., 2016). Given  
459 Spillway lake grew by 40,565 m<sup>2</sup> to a size of 286,367 m<sup>2</sup> (December 2015- April 2018), our results indicate the  
460 channel has since stabilised, allowing Spillway Lake to expand up-glacier.

461 During the 4 years of our study, Imja Lake expanded by 456 m up-glacier and Spillway Lake increased in  
462 area by ~ 14% a<sup>-1</sup>. At the same time, we saw large ponds form via coalescing on the margins of Khumbu and  
463 Ngozumpa glaciers, and coalescence begin on Ama Dablam Glacier (Fig. 3 and SI. Fig. 2a). We also observed an  
464 increase ponded area increased on all of our study glaciers (Fig. 2). As such, our results show that increasing  
465 volumes of water are being stored on, and in front of glaciers in the Everest region, and comparison to previous  
466 work (Watson et al., 2016) suggests that trend is accelerating. The glaciers in our study region appear to be  
467 rapidly moving along the trajectory of proglacial lake formation, and may be doing so quicker than other regions  
468 (Komori, 2008; Robertson, 2012). These developments have major implications for downstream GLOF risk, as  
469 lake volume and speed of expansion are key factors in the hazard potential of proglacial lakes (Rounce et al.,  
470 2017b). As such, there is an urgent need for high resolution monitoring of ice-surface water volumes and  
471 proglacial lake development in the Everest Region.

472

## 473 5. Conclusions

474 This study represents the most up-to-date assessment of supraglacial ponds in the Everest region of Nepal. All  
475 10 glaciers demonstrated an overall ponded area increase over the period 2015 to 2018, ranging from 13.6 % to  
476 108 %. Given the short time span of this study, this is a marked acceleration in the rate of pond growth compared  
477 to that of 2000 – 2015 (Watson et al., 2016). This is a major concern for proglacial lake formation and thus future  
478 risk to downstream communities. A shift towards pond coalescing was observed, most notably on the three  
479 larger glaciers (Ngozumpa, Khumbu and Pangbung). This suggests a transition towards larger surface ponds and  
480 lakes. Despite this, smaller area ponds (< 900 m<sup>2</sup>) continue to form (accounting for between 55 % and 86 % of  
481 the ponds identified in this study) highlighting the need for higher resolution imagery for future remote sensing  
482 studies. Our data show the conditions for pond formations are now predominantly found at mid-elevations on  
483 glaciers in the Everest region, and the area where ponds can form may be advancing up-glacier. This has  
484 important implications for future ice loss in the region. Velocity has been shown to influence both ponded area  
485 and pond number, however this relationship requires further study. Ice cliffs were found on all 10 study glaciers,  
486 with the highest proportion found on the three larger glaciers, however the rate of formation appears lower  
487 than the rate of pond formation in this region. Two glaciers (Ama Dablam and Lhotse Nup) progressed to a new  
488 stage of proglacial lake development during the four year study period. This transition is markedly faster than  
489 observations elsewhere (e.g. Komori, 2008; Robertson, 2012), and suggests proglacial lakes are evolving much  
490 quicker here than other regions, and as such the situation requires continued high temporal resolution  
491 monitoring. Given this, our results show existing classification schemes are too simplistic to suitably evaluate  
492 and communicate glacier lake hazards, leading to the proposal of a new, six-stage model more suited to  
493 evaluating and communicating GLOF hazards.

## 494 6. Conflict of Interest

495 The authors declare that the research was conducted in the absence of any commercial or financial relationships  
496 that could be construed as a potential conflict of interest.

## 497 7. Acknowledgements

498 We acknowledge a number of freely available datasets used in this study. We are also grateful to Dehecq et al.  
499 (2015) for velocity data.

## 500 8. References

- 501 Bajracharya, S. R., Mool, P. K. and Shrestha, B. R. (2007) *Impact of climate change on Himalayan glaciers and*  
502 *glacial lakes: Case studies on GLOF and associated hazards in Nepal and Bhutan*, Kathmandu: International  
503 Centre for Integrated Mountain Development.
- 504 Benn, D. I. and Lehmkuhl, F. (2000) Mass balance and equilibrium-line altitudes of glaciers in high-mountain  
505 environments, *Quaternary International*, 65, 15-29.
- 506 Benn, D.I., Wiseman, S. and Hands, K.A. (2001) Growth and drainage of supraglacial lakes on debris-mantled  
507 Ngozumpa Glacier, Khumbu Himal, Nepal, *Journal of Glaciology*, 47(159), 626-638.

508 Benn, D.I., Bolch, T., Hands, K., Gulley, J., Luckman, A., Nicholson, L.I., Quincey, D., Thompson, S., Toumi, R. and  
509 Wiseman, S. (2012) Response of debris-covered glaciers in the Mount Everest region to recent warming, and  
510 implications for outburst flood hazards, *Earth-Science Reviews*, 114, 156–174.

511 Benn, D.I., Thompson, S., Gulley, J., Mertes, J., Luckman, A. and Nicholson, L. (2017) Structure and evolution of  
512 the drainage system of a Himalayan debris-covered glacier, and its relationship with patterns of mass loss,  
513 *Cryosphere*.

514 Bolch, T. and Kamp, U. (2006) Glacier mapping in high mountains using DEMs, Landsat and ASTER data. *Grazer  
515 Schriften der Geographie und Raumforschung*, 41, 37-48.

516 Bolch, T., Buchroithner, M. F., Peters, J., Baessler, M., and Bajracharya, S. (2008) Identification of glacier  
517 motion and potentially dangerous glacial lakes in the Mt. Everest region/Nepal using spaceborne imagery,  
518 *Natural Hazards and Earth System Sciences*, 8(6), 1329-1340.

519 Bolch, T., Pieczonka, T. and Benn, D. I. (2011) Multi-decadal mass loss of glaciers in the Everest area (Nepal  
520 Himalaya) derived from stereo imagery, *The Cryosphere*, 5(2), 349-358.

521 Bolch, T., Kulkarni, A., Kääh, A., Huggel, C., Paul, F., Cogley, J.G., Frey, H., Kargel, J.S., Fujita, K., Scheel, M.,  
522 Bajracharya, S. and Stoffel, M. (2012a) The State and Fate of Himalayan Glaciers, *Science*, 336(6079), 310.

523 Bolch, T., Peters, J., Yegorov, A., Pradhan, B., Buchroithner, M. and Blagoveshchensky, V. (2012b) Identification  
524 of potentially dangerous glacial lakes in the northern Tien Shan, *Natural Hazards*, 59(3), 1691-1714.

525 Brun, F., Berthier, E., Wagnon, P., Kääh, A. and Treichler, D. (2017) A spatially resolved estimate of High  
526 Mountain Asia glacier mass balances from 2000 to 2016, *Nature geoscience*, 10(9), 668-673.

527 Buri, P., Miles, E.S., Steiner, J.F., Immerzeel, W.W., Wagnon, P. and Pellicciotti, F. (2016a) A physically based 3-  
528 D model of ice cliff evolution over debris-covered glaciers, *Journal of Geophysical Research: Earth Surface*,  
529 121(12), 2471-2493.

530  
531 Buri, P., Pellicciotti, F., Steiner, J. F., Miles, E. S., & Immerzeel, W. W. (2016b) A grid-based model of  
532 backwasting of supraglacial ice cliffs on debris-covered glaciers, *Annals of Glaciology*, 57(71), 199-211.

533  
534 Carrivick, J. L., and Tweed, F. S. (2016) A global assessment of the societal impacts of glacier outburst floods,  
535 *Global and Planetary Change*, 144, 1-16.

536 Dehecq, A., Gourmelen, N. and Trouvé, E. (2015) Deriving large-scale glacier velocities from a complete  
537 satellite archive: Application to the Pamir–Karakoram–Himalaya, *Remote Sensing of Environment*, 162, 55-66.

538 Dehecq, A., Gourmelen, N., Gardner, A.S., Brun, F., Goldberg, D., Nienow, P.W., Berthier, E., Vincent, C.,  
539 Wagnon, P. and Trouvé, E. (2018) Twenty-first century glacier slowdown driven by mass loss in High Mountain  
540 Asia, *Nature Geoscience*, 1.

541 Fujita, K. (2008) Influence of precipitation seasonality on glacier mass balance and its sensitivity to climate  
542 change, *Annals of Glaciology*, 48, 88-92.

543 Fushimi, H., Yoshida, M., Watanabe, O. and Upadhyay, B. P. (1980) Distributions and grain sizes of supraglacial  
544 debris in the Khumbu glacier, Khumbu Region, East Nepal, *Journal of the Japanese Society of Snow and Ice*,  
545 41(Special), 18-25.

546 Gardelle, J., Arnaud, Y. and Berthier, E. (2011) Contrasted evolution of glacial lakes along the Hindu Kush  
547 Himalaya mountain range between 1990 and 2009, *Global and Planetary Change*, 75(1-2), 47-55.

548 Gardelle, J., Berthier, E., Arnaud, Y., and Kaab, A. (2013) Region-wide glacier mass balances over the Pamir-  
549 Karakoram-Himalaya during 1999-2011 (vol 7, pg 1263, 2013), *The Cryosphere*, 7(6), 1885-1886.

550 Gulley, J. and Benn, D. I. (2007) Structural control of englacial drainage systems in Himalayan debris-covered  
551 glaciers, *Journal of Glaciology*, 53(182), 399-412.

552 Immerzeel, W.W., van Beek, L.P.H. and Bierkens, M.F.P. (2010) Climate Change Will Affect the Asian Water  
553 Towers, *Science*, 328(5984), 1382.



554 Immerzeel, W. W., Kraaijenbrink, P. D. A., Shea, J. M., Shrestha, A. B., Pellicciotti, F., Bierkens, M. F. P., and De  
555 Jong, S. M. (2014) High-resolution monitoring of Himalayan glacier dynamics using unmanned aerial vehicles,  
556 *Remote Sensing of Environment*, 150, 93-103.

557 Iwata, S., Aoki, T., Kadota, T., Seko, K. and Yamaguchi, S. (2000) Morphological evolution of the debris cover on  
558 Khumbu Glacier, Nepal, between 1978 and 1995, *IAHS-AISH Publication*, 3-11.

559 Jordan, R.E. and Stark, J.A. (2001) *Capillary tension in rotting ice layers* (No. ERDC/CRREL-TR-01-13). ENGINEER  
560 RESEARCH AND DEVELOPMENT CENTER HANOVER NH COLD REGIONS RESEARCH AND ENGINEERING LAB.

561 Josberger, E. G. (1978) A laboratory and field study of iceberg deterioration, In *Iceberg Utilization* (pp. 245-  
562 264).

563 Juen, M., Mayer, C., Lambrecht, A., Han, H., and Liu, S. (2014). Impact of varying debris cover thickness on  
564 ablation: a case study for Koxkar Glacier in the Tien Shan, *The Cryosphere*, 8(2), 377-386.

565 Kääb, A. (2005) Combination of SRTM3 and repeat ASTER data for deriving alpine glacier flow velocities in the  
566 Bhutan Himalaya, *Remote Sensing of Environment*, 94(4), 463-474.

567 Kääb, A., Berthier, E., Nuth, C., Gardelle, J. and Arnaud, Y. (2012) Contrasting patterns of early twenty-first-  
568 century glacier mass change in the Himalayas, *Nature*, 488, 495.

569 Kattelman, R. (2003) Glacial lake outburst floods in the Nepal Himalaya: a manageable hazard?, *Natural*  
570 *Hazards*, 28(1), 145-154.

571 King, O., Quincey, D. J., Carrivick, J. L. and Rowan, A. V. (2017) Spatial variability in mass loss of glaciers in the  
572 Everest region, central Himalayas, between 2000 and 2015, *The Cryosphere*, 11(1), 407-426.

573 Kirkbride, M.P. (1993) The temporal significance of transitions from melting to calving termini at glaciers in the  
574 central Southern Alps of New Zealand, *The Holocene*, 3(3), 232-240.

575 Komori, J. (2008) Recent expansions of glacial lakes in the Bhutan Himalayas, *Quaternary International*, 184(1),  
576 177-186.

577 Mertes, J. R., Thompson, S. S., Booth, A. D., Gulley, J. D. and Benn, D. I. (2017) A conceptual model of supra-  
578 glacial lake formation on debris-covered glaciers based on GPR facies analysis, *Earth Surface Processes and*  
579 *Landforms*, 42(6), 903-914.

580 Miles, E.S., Willis, I.C., Arnold, N.S., Steiner, J. and Pellicciotti, F. (2016) Spatial, seasonal and interannual  
581 variability of supraglacial ponds in the Langtang Valley of Nepal, 1999–2013, *Journal of Glaciology*, 63(237), 88-  
582 105.

583 Miles, E.S., Steiner, J., Willis, I., Buri, P., Immerzeel, W.W., Chesnokova, A. and Pellicciotti, F. (2017a) Pond  
584 Dynamics and Supraglacial-Englacial Connectivity on Debris-Covered Lirung Glacier, Nepal, *Frontiers in Earth*  
585 *Science*, 5(69).

586 Nicholson, L., and Benn, D. I. (2013). Properties of natural supraglacial debris in relation to modelling sub-  
587 debris ice ablation, *Earth Surface Processes and Landforms*, 38(5), 490-501.

588 Nie, Y., Liu, Q. and Liu, S. (2013) Glacial lake expansion in the Central Himalayas by Landsat images, 1990–2010,  
589 *PLoS One*, 8(12), e83973.

590 Quincey, D.J., Richardson, S.D., Luckman, A., Lucas, R.M., Reynolds, J.M., Hambrey, M.J. and Glasser, N.F.  
591 (2007) Early recognition of glacial lake hazards in the Himalaya using remote sensing datasets, *Global and*  
592 *Planetary Change*, 56(1), 137-152.

593 Quincey, D.J., Luckman, A. and Benn, D. (2009) Quantification of Everest region glacier velocities between 1992  
594 and 2002, using satellite radar interferometry and feature tracking, *Journal of Glaciology*, 55(192), 596-606.

595 Reynolds, J.M. (2000) On the formation of supraglacial lakes on debris-covered glaciers, *IAHS publication*, 153-  
596 164.

597 Richardson, S.D. and Reynolds, J.M. (2000) An overview of glacial hazards in the Himalayas, *Quaternary*  
598 *International*, 65-66, 31-47.

599 Robertson, C. M. (2012) *Temporal evolution of the termini and subaqueous morphologies of lake-calving*  
600 *glaciers in Aoraki/Mount Cook National Park, New Zealand: a thesis presented in partial fulfilment of the*  
601 *requirements for the degree of Doctor of Philosophy in Geography at Massey University, Palmerston North,*  
602 *New Zealand, Doctoral dissertation, Massey University.*

603 Röhl, K. (2006) Thermo-erosional notch development at fresh-water-calving Tasman Glacier, New Zealand,  
604 *Journal of Glaciology*, 52(177), 203-213.

605 Röhl, K. (2008) Characteristics and evolution of supraglacial ponds on debris-covered Tasman Glacier, New  
606 Zealand, *Journal of Glaciology*, 54(188), 867-880.

607 Rounce, D. R., Byers, A. C., Byers, E. A. and McKinney, D. C. (2017a) Brief communication: Observations of a  
608 glacier outburst flood from Lhotse Glacier, Everest area, Nepal, *The Cryosphere*, 11(1), 443.

609 Rounce, D. R., Watson, C. S. and McKinney, D. C. (2017b) Identification of hazard and risk for glacial lakes in the  
610 Nepal Himalaya using satellite imagery from 2000–2015, *Remote Sensing*, 9(7), 654.

611 Sakai, A., Takeuchi, N., Fujita, K. and Nakawo, M. (2000) Role of supraglacial ponds in the ablation process of a  
612 debris-covered glacier in the Nepal Himalayas, *IAHS PUBLICATION*, 119-132.

613 Sakai, A., Nakawo, M., and Fujita, K. (2002). Distribution characteristics and energy balance of ice cliffs on  
614 debris-covered glaciers, Nepal Himalaya. *Arctic, Antarctic, and Alpine Research*, 34(1), 12-19.

615 Sakai, A., Nishimura, K., Kadota, T. and Takeuchi, N. (2009) Onset of calving at supraglacial lakes on debris-  
616 covered glaciers of the Nepal Himalaya, *Journal of Glaciology*, 55(193), 909-917.

617 Salerno, F., Guyennon, N., Thakuri, S., Viviano, G., Romano, E., Vuillermoz, E., Cristofanelli, P., Stocchi, P.,  
618 Agrillo, G., Ma, Y. and Tartari, G. (2015) Weak precipitation, warm winters and springs impact glaciers of south  
619 slopes of Mt. Everest (central Himalaya) in the last 2 decades (1994–2013), *The Cryosphere*, 9(3), 1229-1247.

620 Salerno, F., Thakuri, S., Tartari, G., Nuimura, T., Sunako, S., Sakai, A. and Fujita, K. (2017) Debris-covered glacier  
621 anomaly? Morphological factors controlling changes in the mass balance, surface area, terminus position, and  
622 snow line altitude of Himalayan glaciers, *Earth and Planetary Science Letters*, 471, 19-31.

623 Shrestha, A. B., Wake, C. P., Mayewski, P. A. and Dibb, J. E. (1999) Maximum temperature trends in the  
624 Himalaya and its vicinity: an analysis based on temperature records from Nepal for the period 1971–94,  
625 *Journal of climate*, 12(9), 2775-2786.

626 Shrestha, A.B. and Aryal, R. (2011) Climate change in Nepal and its impact on Himalayan glaciers, *Regional*  
627 *Environmental Change*, 11(1), 65-77.

628 Thompson, S., Benn, D., Luckman, A. and Kulesa, B. (2010) The evolution of supraglacial lakes, Ngozumpa  
629 Glacier, Khumbu Himal, Nepal, In *EGU General Assembly Conference Abstracts* (Vol. 12, p. 11396).

630 Thompson, S. S., Benn, D. I., Dennis, K., and Luckman, A. (2012) A rapidly growing moraine-dammed glacial  
631 lake on Ngozumpa Glacier, Nepal, *Geomorphology*, 145, 1-11.

632 Thompson, S., Benn, D.I., Mertes, J. and Luckman, A. (2016) Stagnation and mass loss on a Himalayan debris-  
633 covered glacier: processes, patterns and rates, *Journal of Glaciology*, 62(233), 467-485.


634 Tiwari, R. K., Garg, P. K., Saini, V. and Shukla, A. (2016) Comparisons of different methods for debris covered  
635 glacier classification, In Khanbilvardi, R., Ganju, A., Rajawat, A.S. and Chen, J.M, *Land Surface and Cryosphere*  
636 *Remote Sensing III*, International Society for Optics and Photonics, 9877-64.

637 Watson, C.S., Quincey, D.J., Carrivick, J.L. and Smith, M.W. (2016) The dynamics of supraglacial ponds in the  
638 Everest region, central Himalaya, *Global and Planetary Change*, 142(Supplement C), 14-27.

639 Watson, C. S., Quincey, D. J., Carrivick, J. L., and Smith, M. W. (2017) Ice cliff dynamics in the Everest region of  
640 the Central Himalaya, *Geomorphology*, 278, 238-251.

641

- 642 Watson, C.S., King, O., Miles, E.S. and Quincey, D.J. (2018) Optimising NDWI supraglacial pond classification on  
643 Himalayan debris-covered glaciers, *Remote sensing of environment*, 217, 414-425.
- 644 Wessels, R. L., Kargel, J. S. and Kieffer, H. H. (2002) ASTER measurement of supraglacial lakes in the Mount  
645 Everest region of the Himalaya, *Annals of Glaciology*, 34(1), 399-408.
- 646 Xu, J., Grumbine, R.E., Shrestha, A., Eriksson, M., Yang, X., Wang, Y.U.N. and Wilkes, A. (2009) The Melting  
647 Himalayas: Cascading Effects of Climate Change on Water, Biodiversity, and Livelihoods, *Conservation Biology*,  
648 23(3), 520-530.
- 649 Zhang, G., Yao, T., Xie, H., Wang, W. and Yang, W. (2015) An inventory of glacial lakes in the Third Pole region  
650 and their changes in response to global warming, *Global and Planetary Change*, 131, 148-157.




651 **FIGURE 1: LOCATION OF THE 10 STUDY GLACIERS (BLUE OUTLINE) WITHIN THE EVEREST REGION. ARROWS DICTATE**  
 652 **DIRECTION OF ICE FLOW. GLACIER NAMES ARE AS FOLLOWS: S- SUMNA GLACIER, P- PANGBUNG GLACIER, N- NGOZUMPA**  
 653 **GLACIER, K- KHUMBU GLACIER, NU- NUPTSE GLACIER, LN- LHOTSE NUP GLACIER, L- LHOTSE GLACIER, LS- LHOTSE**  
 654 **SHAR GLACIER, I- IMJA GLACIER, AD- AMA DABLAM GLACIER. INSET: LOCATION OF THE EVEREST REGION WITHIN NEPAL.**  
 655 **BACKGROUND IMAGE IS SURFACE ELEVATION, DERIVED FROM ASTER DEM (AVAILABLE AT**  
 656 **[HTTP://EARTHEXPLORER.USGS.GOV/](http://earthexplorer.usgs.gov/)).**

Reference	Date	Imagery	Resolution	Focus
Iwata et al. (2000)	1978-1995	SPOT	Not specified	-Sketch map from SPOT imagery compared to 1987 field survey.
Wessels et al. (2002)	2000	ASTER	15m	-Water delineated for a single time period.
Bolch et al. (2008)	1962-2005	ASTER	15m	-Normalized Difference Water Index (NDWI) and manual delineation used to classify water bodies.
		Landsat	30m	
Gardelle et al. (2011)	1990-2009	Landsat	30m	-Decision tree used to classify lakes using the NDWI.
Salerno et al. (2012)	2008	AVNIR-2	10m	-Manual digitalisation of water bodies.
Thompson et al. (2012)	1984-2010	Aerial Photos	<1m	-Multi-temporal analysis of Spillway Lake expansion. Glacier area change not reported.
		ASTER	15m	
		Landsat	30m	
Nie et al. (2013)	1990-2010	Landsat	30m	-NDWI based classification, but no area changes reported.
Zhang et al. (2015)	1990-2010	Landsat	30m	-Water bodies manually digitalised but ponded area changes not reported.
Mertes et al. (2016)	2014	GeoEye	<1m	-Conceptual model of supraglacial lake evolution based on Ground Penetrating Radar (GPR) facies analysis of Spillway Lake.
Watson et al. (2016)	2000-2015	GoogleEarth	2m	-Water bodies semi-automatically or manually digitised. Area changes are quantified.
		WorldView 1&2, GeoEye, QuickBird-2	0.5-0.6m	


657 **TABLE 1: PREVIOUS REMOTE SENSING STUDIES OF SUPRAGLACIAL WATER STORAGE IN THE EVEREST REGION (AFTER**  
658 **WATSON ET AL., 2016)**

<b>Glacier</b>	<b>Area (km<sup>2</sup>)</b>	<b>Length (km)</b>	<b>Min-Max Elevation (range) (m)</b>
Ama Dablam	2.13	4.40	4769 – 5084 (315)
Imja	1.08	2.46	5023 – 5187 (164)
Khumbu	6.64	10.82	4956 – 5246 (290)
Lhotse	5.74	6.69	4715 – 5245 (530)
Lhotse Nup	1.38	3.82	4954 – 5310 (356)
Lhotse Shar	3.00	4.03	5008 – 5429 (421)
Ngozumpa	15.10	15.76	4868 – 5541 (673)
Nuptse	3.06	6.20	4662 – 5354 (692)
Pangbung*	11.34	13.14	4742 – 5380 (638)
Sumna*	5.34	7.39	4888 – 5502 (614)

659 **TABLE 2: CHARACTERISTICS OF THE 10 STUDY GLACIERS, SHOWING AREA, LENGTH, ELEVATION AND**  
660 **DIRECTION OF FLOW. \*INDICATES GLACIERS NOT INCLUDED IN THE WATSON ET AL. (2016) STUDY.**




661 **FIGURE 2: CHANGES IN SUPRAGLACIAL PONDS OBSERVED ON ALL 10 GLACIERS MARCH 2015- APRIL**  
 662 **2018; (A) PONDED AREA AS A PERCENTAGE OF TOTAL GLACIER AREA, (B) TOTAL NUMBER OF PONDS**  
 663 **AND (C) TOTAL PONDED AREA. BARS ARE COLOUR CODED BY DATE. AD-AMA DABLAM GLACIER, I-IMJA**  
 664 **GLACIER, K-KHUMBU GLACIER, N-NGOZUMPA GLACIER, NU-NUPTSE GLACIER, L-LHOTSE GLACIER, LN-**  
 665 **LHOTSE NUP GLACIER, LS-LHOTSE SHAR GLACIER, P-PANGBUNG GLACIER, S-SUMNA GLACIER.**




666 **FIGURE 3: PONDED AREA CHANGE FOR THE NGOZUMPA (A) KHUMBU (B) PANGBUNG (C) AND SUMNA (D) GLACIERS, FOR**  
 667 **THE PERIOD 2015-2018. INSET (A) UP-GLACIER EXPANSION OF SPILLWAY LAKE, INSET (B) POND EXPANSION ALONG THE**  
 668 **LOWER EASTERN MARGINS OF KHUMBU GLACIER, INSET (C) POND EXPANSION ON THE GLACIER TONGUE, (D) POND**  
 669 **PROLIFERATION ON THE LOWER ABLATION ZONE 2015-2017 AND DRAINAGE ALONG THE WESTERN MARGINS 2017-2018.**  
 670 **ARROWS INDICATE DIRECTION OF ICE FLOW. BACKGROUND IMAGE IS 2018 SENTINEL-2 IMAGERY (FROM USGS AT**  
 671 **[HTTP://EARTHEXPLORER.USGS.GOV/](http://earthexplorer.usgs.gov/)).**






672  
 673 **FIGURE 4: AREA CHANGE OF IMJA LAKE, IN THE SOUTHEAST OF THE REGION FOR THE PERIOD 2015-2018.**  
 674 **ARROWS INDICATE DIRECTION OF ICE FLOW. BACKGROUND IMAGE IS 2018 SENTINEL-2 IMAGERY (FREELY**  
 675 **AVAILABLE FROM USGS AT [HTTP://EARTHEXPLORER.USGS.GOV/](http://earthexplorer.usgs.gov/)).**




676  
677  
678  
679

**FIGURE 5: NUMBER OF PONDS FOUND ON THE GLACIER SURFACE APRIL 2018 COMPARED TO GLACIER ELEVATION/SLOPE AT 10% GLACIER AREA DISTANCE BANDS (0-10 = BAND 1, 10-20= BAND 2 ETC.) DASHED RED LINE INDICATES BREAKS OF SLOPE.**




680  
681  
682

**FIGURE 6: PONDED AREA FOUND ON THE GLACIER SURFACE APRIL 2018 COMPARED TO GLACIER ELEVATION/SLOPE AT 10% GLACIER AREA DISTANCE BANDS (0-10 = BAND 1, 10-20= BAND 2 ETC.)**




683  
684  
685

**FIGURE 7: NUMBER OF PONDS FOUND ON THE GLACIER SURFACE APRIL 2018 COMPARED TO GLACIER VELOCITY AT 10% GLACIER AREA DISTANCE BANDS (0-10 = BAND 1, 10-20 = BAND 2 ETC.**



686  
687  
688


**FIGURE 8: PONDED AREA FOUND ON THE GLACIER SURFACE APRIL 2018 COMPARED TO GLACIER VELOCITY AT 10% GLACIER AREA DISTANCE BANDS (0-10 = BAND 1, 10-20 = BAND 2 ETC.)**



689  
 690 **FIGURE 9: PERCENTAGE OF PONDS WITH AND WITHOUT ICE-CLIFFS, DISPLAYED AS PERCENTAGE OF TOTAL GLACIER**  
 691 **SURFACE AREA. AD- AMA DABLAM GLACIER, I- IMJA GLACIER, K- KHUMBU GLACIER, N- NGOZUMPA GLACIER, NU-**  
 692 **NUPTSE GLACIER, L- LHOTSE GLACIER, LN- LHOTSE NUP GLACIER, LS- LHOTSE SHAR GLACIER, P- PANGBUNG GLACIER,**  
 693 **S- SUMNA GLACIER.**


Stage of Lake Development	Description
0	No supraglacial ponds.
1	Appearance and growth of supraglacial ponds in the lower ablation zones
2	Coalescing of supraglacial ponds to form larger, ice dammed ponds.
3	Stable expansion of ponds up-glacier and shift to moraine dammed proglacial lake.
4	Glacier retreats out of the proglacial lake.

694 **TABLE 3: STAGE OF LAKE DEVELOPMENT FOLLOWING KOMORI (2008) AND ROBERTSON (2012).**



695 **FIGURE 10: STAGE OF LAKE DEVELOPMENT CHANGE OBSERVED OVER THE STUDY PERIOD, FROM 2015 (BOTTOM) TO 2018**  
 696 **(TOP). AD- AMA DABLAM GLACIER, I- IMJA GLACIER, K- KHUMBU GLACIER, N- NGOZUMPA GLACIER, NU- NUPTSE**  
 697 **GLACIER, L- LHOTSE GLACIER, LN- LHOTSE NUP GLACIER, LS- LHOTSE SHAR GLACIER, P- PANGBUNG GLACIER, S-**  
 698 **SUMNA GLACIER. BACKGROUND IMAGE IS 2018 SENTINEL-2 IMAGE (AVAILABLE FROM USGS AT**  
 699 **[HTTP://EARTHEXPLORER.USGS.GOV/](http://earthexplorer.usgs.gov/)).**





700 **FIGURE 11: UPDATED KOMORI (2008) AND ROBERTSON (2012) CONCEPTUAL LAKE CLASSIFICATION MODEL, TO INCLUDE**  
 701 **2 ADDITIONAL STAGES ACCOUNTING FOR ‘IN-STAGE’ CHANGES OBSERVED IN THIS STUDY; STAGE 2: RECOGNISES THE ROLE**  
 702 **OF ICE-CLIFFS AND CHANGES IN SURFACE DEBRIS LEADING TO POND FORMATION AND DECAY; STAGE 3: GIVES MARGINAL**  
 703 **BASED EXPANSION OF PONDS AND POND DRAINAGE EVENTS THEIR OWN CATEGORY (AFTER KOMORI 2008; ROBERTSON**  
 704 **2012; MERTES ET AL., 2016).**

705