

Can Artificial Reoxygenation Revitalize Dying Coastal Seas?

C.P. Slomp¹, A. Oschlies², A.H. Altieri³, L.T. Bach⁴, L. Bopp⁵, D. Breitburg⁶, A. Canning^{7,8}, D. J. Conley⁹, M. Dai¹⁰, B. Dewitte¹¹⁻¹³, H. Enevoldsen¹⁴, E.M. Ferrer¹⁵, A. Galan^{16, 17}, V. Garcon¹⁸, M. Gregoire¹⁹, B. Gustafsson²⁰, D. Gutierrez²¹, P. Handmann⁷, A. Hylén²², K. Isensee²³, R. Lamond²⁴, M. Li²⁵, K. Limburg²⁶, I. Montes²⁷, J. Sterling²⁸, A. Tan Shau Hwai²⁹, J.M. Testa²⁵, D. Wallace³⁰, J.J. Waniek³¹, M. Yasuhara³²

1. Department of Microbiology, Radboud University, Nijmegen, the Netherlands. 2. GEOMAR Helmholtz Centre for Ocean Research Kiel, Germany. 3. Department of Environmental Engineering Sciences, University of Florida, USA. 4. Institute for Marine and Antarctic Studies, University of Tasmania, Australia. 5. CNRS - Laboratoire de Météorologie Dynamique, CNRS/ENS-PL, France. 6. Smithsonian Environmental Research Center, USA. 7. Lhyfe SA, France. 8. Atrium Environmental UG, Kiel, Germany. 9. Lund University, Sweden. 10. Xiamen University, China. 11. Centro de Estudios Avanzado en Zonas Áridas (CEAZA), La Serena, Chile. 12. Departamento de Biología, Facultad de Ciencias del Mar, Universidad Católica del Norte, Coquimbo, Chile. CEAZA, Chili. 13. CECI, Université de Toulouse, CERFACS/CNRS, Toulouse, France. 14. IOC-UNESCO, Copenhagen, Denmark. 15. National Center for Ecological Analysis and Synthesis, UC Santa Barbara USA. 16. Centro de Investigación de Estudios Avanzados del Maule (CIEAM), Vicerrectoría de Investigación y Postgrado, Universidad Católica del Maule, Talca, Chile. 17. Centro Regional de Estudios Ambientales (CREA), Universidad Católica de la Santísima Concepción, Concepción, Chile. 18. CNRS, Institut de Physique du Globe de Paris, France. 19. Liège University, Belgium. 20. Stockholm University, Sweden. 21. Instituto del Mar del Peru, Peru. 22. University of Antwerp, Belgium. 23. IOC-UNESCO, Paris, France. 24. Air Liquide, Belgium. 25. University of Maryland Center for Environmental Science, USA. 26. SUNY College of Environmental Science and Forestry, USA. 27. Instituto Geofísico del Perú, Peru. 28. IOC-UNESCO, The Netherlands. 29. Centre For Marine & Coastal Studies, Universiti Sains Malaysia, Malaysia. 30. Dalhousie University, Canada. 31. Leibniz Institute for Baltic Sea Research, Warnemünde (IOW), Germany. 32. School of Energy and Environment and State Key Laboratory of Marine Environmental Health, City University of Hong Kong, Hong Kong SAR.

Abstract

Eutrophication and global warming are key drivers of oxygen loss, also termed deoxygenation, in coastal ecosystems worldwide. Artificial reoxygenation has been put forward as a local or regional solution to increase oxygen concentrations and improve water quality. Three main approaches have been proposed: (1) bubbling with air with the aim to destratify and mix the water column (2) injection of pure oxygen and (3) pumping of oxygenated water to greater water depths (downwelling). In this review, we summarize the results of recent field trials and other implementations of artificial reoxygenation in coastal systems, which, to date, only involve small bays and estuaries. We also discuss potential benefits and risks. While the recent trials indicate that reoxygenation of the water column can be achieved, low oxygen conditions returned rapidly within days to months of discontinuing operations, illustrating that artificial reoxygenation typically only provides a temporary solution. Potential side effects of artificial reoxygenation could include enhanced emissions of the greenhouse gas carbon dioxide and, upon bubbling and destratification in shallow waters, also of methane. Additionally, downwelling could lead to warming and an associated increased oxygen demand near the seafloor. Reoxygenation will not necessarily affect nutrient availability for phytoplankton, implying that water quality may remain poor. We recommend a careful, case-by-case assessment of the suitability of artificial reoxygenation in coastal systems prior to implementation and monitoring before, during and after each intervention. Any field trials should involve all relevant parties, including scientists and local communities, and results should be reported with full transparency. While in the short term artificial reoxygenation may be useful to alleviate oxygen loss, long-term improvements in the oxygen levels and quality of coastal waters require reductions in nutrient inputs and greenhouse gas emissions.

1. Introduction

Coastal waters worldwide are rapidly losing oxygen. This is altering their productivity, biogeochemistry and biodiversity (Diaz and Rosenberg, 2008; Breitburg et al., 2018). Low oxygen zones are often termed “dead zones” because they are unable to support most forms of marine life, such as many species of fish and marine invertebrates (Diaz and Rosenberg, 2008). The causes of low oxygen are well-known: since the 1950’s, phosphorus and nitrogen from terrestrial sources (such as fertilizer, wastewater and, for nitrogen, also fossil fuels) are increasingly entering coastal seas where they are stimulating phytoplankton blooms that, upon their demise and the decay of the organic matter, remove oxygen. Moreover, global warming is reducing the solubility of oxygen in seawater, is increasing water column stratification and is slowing down circulation, which all limit ventilation of deeper waters (Breitburg et al. 2018; Dai et al., 2023). Oxygen loss in coastal waters can be sporadic, diel, seasonal or permanent, depending on the type of system, and the frequency, duration and intensity are key in determining the environmental impact (Diaz and Rosenberg, 2008; Tyler et al., 2009; Pezner et al., 2023).

The Baltic Sea is a well-investigated example of a eutrophic coastal sea where nutrient pollution has led to the development of a near-permanent low oxygen zone and major ecosystem change. Anoxic

and sulfidic bottom waters cover most of the seafloor at water depths >70 m, creating a large “dead zone” devoid of higher life. Cyanobacterial blooms have become frequent and widespread (Conley et al., 2009a; Reusch et al., 2018) and spawning grounds for cod have been lost (Bryhn et al., 2022). Trapping of fish in low oxygen areas has been observed to lead to fish kills, which are localized mass die-offs (Grégoire et al., 2023). Examples of regions where widespread low oxygen is seasonal include the Gulf of Mexico, Adriatic Sea and East China Sea (Pitcher et al., 2021; Dai et al., 2023). Some coastal systems, such as the Gulf of St. Lawrence in eastern Canada, may be under increased threat of low oxygen because of climate-change driven alterations in oceanic circulation that are lowering local oxygen supply (Jutras et al., 2020; Wallace et al., 2023).

Reductions in nutrient inputs from land and greenhouse gas emissions are essential to solve the problem of coastal oxygen loss, but this will take time to have an effect (Conley et al., 2009a). Nutrients have been accumulating in the water column and the sediments of coastal systems for decades and reducing nutrient inputs will not immediately lead to lower concentrations because of nutrient recycling (Kemp et al., 2009; Kulinski et al., 2022). Moreover, nutrient release from agricultural lands continues in many regions because of excess fertilizer use and release of nutrients that have already accumulated in soils and groundwater (Liu et al., 2024). Current efforts to reduce the global use of fossil fuels and related greenhouse gas emissions remain insufficient to stabilize global temperatures and hence fall short of limiting the impact on ocean oxygen associated with global warming (IPCC, 2023).

Artificial reoxygenation has been suggested as a potential solution to coastal oxygen loss (Stigebrandt and Gustafsson, 2007; Stigebrandt et al., 2015a; Stigebrandt and Andersson, 2022) but the possible benefits come with risks that need to be assessed carefully before implementation (Conley et al., 2009a). Such assessments are becoming urgent in the light of the emergence of potential new technologies for artificial reoxygenation of coastal seas linked to green hydrogen production, which, as a byproduct, generates oxygen (Beghoura et al., 2023; Wallace et al., 2023; Handmann and Wallace, 2024). This oxygen could be used to increase the oxygen supply to coastal waters, particularly when green hydrogen production facilities are located close to the sea.

In this article, we first briefly discuss the controls on oxygen concentrations in coastal waters and describe the most common approaches proposed for their artificial reoxygenation. We then discuss the results of field trials in coastal systems and their limitations and potential side effects. Finally, we discuss whether artificial reoxygenation is likely to be a long-term solution to coastal oxygen loss and provide recommendations for the formulation of best practices when assessing the suitability and application of artificial reoxygenation.

2. Artificial reoxygenation: increasing the oxygen supply

Coastal seas gain oxygen naturally through air-sea exchange, vertical mixing of the water column, lateral advection and photosynthetic production of organic matter (Figure 1). They lose oxygen through respiration in the water column and uptake by the underlying sediment. Surface waters typically remain

oxygenated because of rapid air-sea exchange and primary productivity but, in deeper waters, oxygen removal dominates, especially in systems with limited vertical mixing (Fennel and Testa, 2019).

Artificial reoxygenation aims to increase the supply of oxygen to deeper waters (Conley et al., 2009a). The primary goal is to increase the oxygen supply to deeper waters to such an extent that the water and surface sediment become or remain oxygenated. The oxygen supply needed to achieve this depends on the oxygen demand of the deeper water and the sediment. This, in turn, depends on the input of organic matter and the eutrophication history of the system: for instance, if the input of organic matter remains high and/or a lot of organic matter has accumulated on the seafloor, the sediment oxygen demand may remain high for a long time. This “legacy” effect can hinder the reoxygenation of coastal systems (Turner et al., 2008). Importantly, the reoxidation of products of anaerobic degradation of organic matter, such as ammonium and hydrogen sulfide, can become the dominant oxygen sink (e.g. Burdige, 2006; Testa et al., 2018).

A secondary goal of artificial reoxygenation can relate to phosphorus dynamics: reoxygenation may reduce phosphorus recycling through increased binding of phosphorus to iron oxides in sediments (Conley et al., 2009b; Jilbert et al., 2011; Testa et al., 2025). This, in turn, may reduce the availability of phosphorus for phytoplankton in surface waters. By lowering the production and sinking flux of organic matter, this may lead to a lower oxygen demand and, hence, higher oxygen concentrations in deeper waters, constituting a positive feedback loop (Conley et al., 2009a). A change in bottom water oxygen can thus lead to an improvement in water quality by promoting the retention of phosphorus in the sediment (Norkko et al., 2012; Stigebrandt et al., 2015a).

Because the same positive feedback loop would release phosphorus upon return to low-oxygen conditions, e.g. after termination of reoxygenation, a persistent improvement of water quality will only be achieved if the phosphorus is sequestered in the sediment permanently (Katsev et al., 2006). This requires the transformation of the iron-oxide bound phosphorus to sediment phosphorus forms that are not remobilized upon burial, such as the calcium-phosphate mineral apatite or, when there is excess reactive iron over reduced sulfur, the Fe(II) phosphate mineral vivianite (Egger et al., 2015; Slomp, 2011). This illustrates that sediment phosphorus burial is not a simple function of bottom water oxygen but instead depends on a range of factors, including the history of the iron supply and organic matter input to and burial in a system.

Reoxygenation affects many other biogeochemical processes. For example, nitrogen loss via coupled nitrification-denitrification in the sediment can be stimulated, improving water quality (De Brabandere et al., 2015; Harris et al., 2015; van Helmond et al., 2018). Reoxygenation can also change the balance between aerobic and anaerobic degradation of organic matter in coastal waters, impacting seawater pH, carbon dioxide concentrations and alkalinity. The effects are expected to be site-specific and will depend on factors such as the buffering capacity of the water and whether the water column remains stratified (Cai et al., 2017). Seasonally hypoxic, stratified systems can be used to illustrate some of the effects. During summer stratification, strong depth gradients in pH and alkalinity can develop,

with distinct pH minima below the redoxcline due to oxic respiration or sulfide oxidation. These gradients disappear upon water column mixing (Hagens et al., 2015; Cai et al., 2017).




Figure 1. Principal terms in the oxygen budget for coastal systems (modified from Fennel and Testa, 2019). Green shapes: living phytoplankton. Brown shapes: other organic matter. Purple: deeper waters prone to low oxygen

3. Approaches to artificially reoxygenate coastal systems

Inspired by methods used to reoxygenate lakes (Singleton and Little, 2006; Mobley et al., 2019), three main approaches have been proposed for artificial reoxygenation of coastal systems (Figure 2). These are (1) bubbling with air, (2) bubbling of pure oxygen (Kowek et al., 2020; Wallace et al., 2023) and (3) artificial downwelling, which refers to pumping of oxygenated surface water into deeper water (Stigebrandt et al., 2015b; Lehtoranta et al., 2022).

Bubble diffusers with air aim to destratify and aerate the water column by inducing vertical mixing. Such aerators have successfully been applied to reoxygenate shallow coastal systems (e.g. small estuaries with water depths of up to several meters) where tides contribute to lateral advection of the oxygen (Harris et al., 2015) and coastal impoundments (Lamping et al., 2005).

Bubbling with pure oxygen is required to oxygenate deeper coastal waters, based on methods for lakes (Mobley et al., 2019). To date, there are no known applications of injection with pure oxygen in coastal waters described in the literature.

Pumps used for coastal reoxygenation through downwelling take in near-surface water and eject it in deeper waters. Depending on the buoyancy of the injected water and the stratification of the ambient

water, downwelling may also affect stratification and mixing (Stigebrandt et al., 2015b; Lehtoranta et al., 2022). Downwelling experiments in Swedish coastal waters have demonstrated that oxygenation can be achieved with this method (Forth et al., 2015; Stigebrandt et al., 2015b; Lehtoranta et al., 2022). Notably, the oxygenation of the deeper water column in both cases was the combined result of increased oxygen supply through downwelling and deepwater renewal through lateral inflow from an adjacent basin. This highlights that oxygen levels in neighbouring water bodies can determine the outcome of artificial reoxygenation efforts and should be considered when planning such interventions (Stigebrandt et al., 2015b).

We note that there are also alternative approaches for reoxygenation for shallow waters, such as, for example, side stream supersaturation where water is pumped to a dissolution device and supersaturated with pure oxygen prior to discharging back into the waterbody. This approach has successfully been used to reoxygenate the water column and sediment-water interface in the upper reaches of the Swan River Estuary, with continued pumping essential to maintain the oxic state (Larsen et al., 2019).




Figure 2. Simplified representation of three main methods proposed for artificial reoxygenation of coastal waters: (A) (1) bubble diffusion with air with the aim to destratify the water column and (2) bubble diffusion to inject pure oxygen, with the aim to directly reoxygenate the deeper water and (B) (3) downwelling (modified from Kowek et al., 2020).

4. Limitations and potential side effects of artificial reoxygenation

Studies of the effects of artificial oxygenation show that, if no other measures are taken, low oxygen conditions return within days or several months when operations are halted, irrespective of the method

employed (Harris et al., 2015; Stigebrandt et al., 2015b; Larsen et al., 2019; Lehtoranta et al., 2022). The return of anoxia upon discontinuing operations is well-known for lakes (Tammgeorg et al., 2023) and illustrates that artificial reoxygenation alone does not provide a permanent solution to deoxygenation because it does not address its root causes.

Reoxygenation via artificial downwelling of coastal waters may be less successful during operation if it causes warming of deeper waters. This is especially a risk when surface water pumps are used to reoxygenate thermally-stratified coastal waters. The transfer of warm surface water to depths near the seafloor may weaken stratification and enhance vertical mixing (or induce lateral inflows of water as noted above). While this may increase the downward transfer of oxygen further, it can also increase the rate of upward mixing of nutrients. This may enhance productivity and, ultimately, could increase the oxygen demand in deeper waters to such an extent that a net decrease in oxygen results (Conley et al., 2009a; Lehtoranta et al., 2022). Another potential effect is increased respiration near the seafloor because of a higher metabolic activity of microbes at higher temperatures (Thamdrup et al., 1998). This may further decrease oxygen concentrations, instead of increasing them as intended (Fig. 3A).

Upon artificial reoxygenation of the water column, wider water quality problems will not necessarily disappear. Deoxygenation is accompanied by nuisance algal blooms in many coastal areas (Breitburg et al., 2018). Such blooms will remain upon reoxygenation if the availability of nutrients for phytoplankton remains high. In the Baltic Sea, for example, natural decadal-scale reoxygenation of deeper waters linked to lateral inflow of oxygenated North Sea water, does not lead to strong removal of phosphorus to the sediment (Sommer et al., 2017; Hermans et al., 2019; Hylén et al., 2021). This lack of a strong effect of the reoxygenation event, which after the most recent inflow lasted >1 year, results from the highly reducing nature of the sediment. This allows benthic release fluxes of ammonium and hydrogen sulfide to exceed the oxidative capacity provided by the supply of oxygen. The continued flux of reductants hinders the formation of phosphorus-containing minerals in the surface sediment upon bottom water oxygenation. Consequently, recycling of phosphorus is not decreased (Fig. 3B; Sommer et al., 2017; Hermans et al., 2019; Hylén et al., 2021), allowing the cyanobacterial blooms to be sustained in what has been denoted as a vicious cycle (Conley et al., 2009a).

In the pumping experiments in Swedish coastal waters described above (Stigebrandt et al., 2015b; Lehtoranta et al., 2022), phosphorus concentrations in the water column decreased during operation of the pumps but increased again within one year when operations were discontinued. The decrease was attributed to the combined effect of export of phosphorus to an adjacent basin through lateral flow and retention of phosphorus in the sediment. Unfortunately, the sediment phosphorus speciation and rate of phosphorus burial were not determined, hindering a detailed evaluation of the relative roles of both processes.

Studies of phosphorus dynamics in a range of coastal systems indicate that reoxygenation does not necessarily stimulate permanent burial of phosphorus (Slomp et al., 2011; Reed et al., 2011; Sulu-

Gambari et al., 2018). In some coastal systems, artificial reoxygenation with continued high nutrient loading could create a large, mobile Fe-bound P pool in the surface sediment. Upon termination of the artificial reoxygenation, this Fe-bound P pool could be released to the water column, exacerbating the water quality problems (Conley et al., 2009a; Reed et al., 2011). These observations are strikingly similar to those for eutrophic Swiss lakes where a lasting lake recovery is not achieved through artificial reoxygenation when phosphorus loads are not reduced (Gächter and Wehrli, 1998; Gächter and Müller, 2003).




Figure 3. Simplified schematic of potential caveats of reoxygenation initiatives. (A) An increased oxygen demand because of warming-induced increased respiration and increased productivity because of enhanced vertical mixing upon downwelling; (B) Limited impact of reoxygenation because of a sustained oxygen demand from the sediment, allowing the vicious cycle between P recycling and bottom water anoxia to be maintained in case of bubbling and downwelling; (C) Increased methane

emissions linked to bubble transport when using air and increased carbon dioxide emissions due to enhanced aerobic respiration; (D) Altered habitat: underwater noise upon bubbling, salinity change and remobilization of toxic substances such as organic pollutants and trace metals.

Artificial reoxygenation may alter the dynamics of greenhouse gases in coastal waters. Increased aerobic respiration will increase carbon dioxide production. This could potentially impact the role of coastal waters as a source of carbon dioxide to the atmosphere (Fig. 3C; Laruelle et al., 2010; Cai et al., 2021). Furthermore, bubbling with air (with destratification) has been shown to enhance the upward transport of methane in the water column and thereby its emission to the atmosphere upon application in a shallow estuary (Lapham et al., 2022). There is accumulating evidence that sediments in eutrophic coastal systems can leak methane because of less efficient removal via microbial oxidation of methane (Lapham et al., 2024; Żygadłowska et al., 2024a). While reoxygenation will typically reduce dissolved methane concentrations in the water column of aquatic systems (Hounshell et al., 2021; Żygadłowska et al., 2024b), this will not necessarily suppress the release of methane from the sediment (Żygadłowska et al., 2024a and b). This implies that, if eutrophication is not addressed in parallel, methane emissions from coastal waters could increase upon reoxygenation via bubbling with air. Further work is needed to quantify these effects (Lapham et al., 2022).

Habitats for marine organisms may also be altered. For example, pumps used for bubbling and downwelling generate underwater noise, which may interfere with natural auditory signal processing by marine animals, with potential impacts on feeding, mating and avoidance of predation (Fig. 3D; Duarte et al., 2021; Zhang et al., 2023). Reoxygenation through bubbling creates turbulence, gas-water interfaces and gradients in oxygen pressure in the water column that may differ from those that occur otherwise, which potentially could affect marine animals. There is also a risk of gas bubble disease in fish, especially when air rather than pure oxygen is used (e.g. Weitkamp and Katz, 1980; Wallace et al., 2023). Artificial downwelling not only changes deep-water temperatures but also alters vertical distributions of salinity. Marine fish and invertebrates are known to be sensitive to changes in salinity and could be affected (Farias et al., 2024). Increased sediment mixing upon return of bottom-dwelling animals may remobilize sediment contaminants, such as organic pollutants and trace metals (Conley et al., 2009a). These examples all illustrate that artificial reoxygenation can have unintended consequences for marine ecosystems that should be considered prior to implementation.

5. Assessing artificial reoxygenation as a solution to oxygen loss

According to the precautionary principle, one should, in the absence of full scientific certainty, act to avoid harm, which, in the case of reoxygenation could arise in various dimensions (Fig. 3). In some cases, this will imply that no measures should be taken. In other cases, this will imply that measures should not be postponed because delay could lead to even more harm. This creates a need for a very careful case-by-case assessment of the suitability of artificial reoxygenation at a given site, and, upon a

positive assessment, careful monitoring. Suitability assessments should look beyond impacts on oxygen only and should preferably be embedded in wider ecosystem restoration efforts (Oliveira et al., 2024). Since rapidly increasing quantities of oxygen might become available as a byproduct of green hydrogen production, there is an urgency to determine if, when, and where artificial reoxygenation can be applied to mitigate the consequences of coastal deoxygenation.

In heavily managed, small and shallow coastal systems, reoxygenation could be a temporary solution, as illustrated by the successful application of aeration (with destratification) in the sub-estuary of Chesapeake Bay (Harris et al., 2015; Lapham et al., 2022) and side stream supersaturation with pure oxygen in the upper reaches of the Swan River Estuary (Larsen et al., 2019) mentioned above. Success in the case of the Chesapeake Bay sub-estuary is defined as an increase in water column oxygen concentrations and a decrease in hydrogen sulphide and fish kills. Decades of aeration without a reduction in organic matter loading makes the water quality in this system directly dependent on the continuous operation of the aeration infrastructure supported by local communities. As noted by Harris et al. (2015), widespread applicability of aeration through destratification should not be inferred from this case and, also in this system, a permanent solution to the water quality problems relies on a reduction of the organic matter supply. In general, systems with such short residence times are expected to respond rapidly to both artificial oxygenation (e.g. Fennel and Testa, 2018) and nutrient input reductions. The Scheldt estuary is an example of a coastal system where the latter was achieved: in the 1980s, low oxygen conditions were frequent but nutrient reductions were successful in restoring an oxic state (Soetaert et al., 2006). Artificial reoxygenation could help bridge the time until recovery in such a relatively small system.

The long-term effects of reoxygenation on large eutrophic coastal systems with strong legacy effects, such as the Baltic Sea, where sediments act as a source of nutrients and sink for oxygen, are more difficult to predict (Reed et al., 2011; Conley et al., 2009a; Hermans et al., 2019). A key reason is that the potential for continued release of nutrients and reductants from the sediments cannot be projected from easily measurable parameters. Lateral exchange of water, solutes and particles will also affect the results of reoxygenation and, in many systems, are difficult to control and monitor. The large size of such systems also makes reoxygenation challenging (Conley et al., 2009a) while also delaying the response to nutrient load reductions. Notably, taking the Baltic Sea as an example, model results suggest that continued nutrient reductions will lead to a future improvement of water quality on time scales of several decades (Meier et al., 2022). This emphasizes the value of continued nutrient reduction strategies. The increasingly important role of climatic warming as a driver of low oxygen in the Baltic Sea (Carstensen et al., 2014) emphasizes the need to also reduce greenhouse gas emissions (IPCC, 2023).

In the Gulf of St. Lawrence, a coastal system without strong legacy effects because of its less eutrophic nature, reoxygenation might be harnessed to allow the current oxygen state of the system to be maintained (Jutras et al., 2020; Wallace et al. 2023). Subsurface waters (water depths > 150 m) in

the region are losing oxygen because of eutrophication and, since 2008, primarily via a change in the lateral input of oxygen from the Atlantic. Climatic-related decreases in the supply of oxygen from the southward flowing Labrador Current and a slowing of the Atlantic Meridional Overturning Circulation are key drivers of the deoxygenation, with the oxygen decline thought to be amplified by increased microbial oxygen uptake due to warming (Jutras et al., 2020; 2023). The annual loss of oxygen in the Gulf and adjacent estuary in recent decades and oxygen production of a proposed green hydrogen plant in the region are of the same order of magnitude (2×10^5 and 5×10^5 tons yr^{-1} , respectively), suggesting possibilities for mitigation of the oxygen decline by oxygenation with pure oxygen. The proposed pure oxygen injection would target the subsurface flows from the Atlantic upon entry into the Gulf. At present, no reoxygenation systems have been applied to the water depths and spatial scales relevant to this setting (Wallace et al., 2023).

Taken together, long-term, sustainable increases in oxygen levels and improved water quality of coastal systems require reductions in nutrient inputs and reductions of greenhouse gas emissions. Hence, artificial reoxygenation, if applied, should always be only one of various measures to improve the state of coastal waters.

6. Best practices for artificial reoxygenation needed

Analogous to best practices formulated for research on marine carbon dioxide removal (Oschlies et al., 2023), and associated governance considerations (Johnson et al., 2024), we point towards the need for the development of a suitability assessment framework (GESAMP, 2025) and best practices for scientific research on, and (possible) implementation, of artificial reoxygenation. When sites are found to be potentially suitable for artificial reoxygenation in the form of field trials or deployment, i.e. when the benefits of reoxygenation are perceived to be greater than the drawbacks and reoxygenation is both feasible and desirable, the following aspects should be considered in such best practices:

- (1) Performance criteria to define success should be formulated prior to an intervention. This implies that one should decide whether the intervention will only target oxygen concentrations or whether wider water quality measures are a goal such as a reduction in nutrient levels. When evaluating success, potential exchange with water masses in adjacent areas should be considered, since displacement of nutrients is not necessarily a solution. Where possible, historical baselines should be established. This can allow an assessment of whether return to such baselines is achievable.
- (2) Pilot studies of artificial reoxygenation should involve relevant government bodies, such as water management authorities, stakeholders, including representatives of local communities, industry partners, and scientists. Feasibility studies should be based on scientific principles and preferably include independent desk studies and lab experiments besides field implementation. Plans should be openly shared and discussed among all involved. Planned field trials or deployments should be announced via a public field-trial registry prior to the experiment or

deployment. This should foster knowledge exchange, accelerate learning and ensure maximum information gain of any interventions in the environment.

(3) Any field trial or implementation should respect environmental safety and consider perceived benefits and risks of the intended intervention in the environment including any relevant ethical issues, taking the intrinsic value of nature into account. Governance frameworks are needed to facilitate informed decision-making. In many countries, systematic decision support systems (e.g. Environmental Impact Assessments) are in place to evaluate proposed interventions in water bodies. Such assessments are critical to ensure the best possible approach in view of protecting the environment and human wellbeing. Risks and benefits should be evaluated against the counterfactual of no deployment and should take ongoing environmental change into account.

(4) Key parameters regarding the biology, chemistry and physics of a system should be monitored before, during and after each intervention to determine baseline conditions and assess the potential impacts on water quality and ecology, including termination effects after an intervention ceases and ecological effects. Continued measurements over years to decades are critical to determine long term effects, also considering ongoing global change and its impacts. A reference or control site, preferably in the vicinity, should be included in each implementation.

Besides oxygen profiling, potential changes in circulation and stratification should be assessed by determining salinity and temperature in parallel. Determining chlorophyll is critical to assess impacts on productivity, whereas hydrogen sulfide is an essential indicator of water quality given its toxic nature. Water column profiles of nutrients (ammonium, nitrate, phosphate, silicate) are key to understanding the level of eutrophication and potential changes in nutrient limitation. Data should have an appropriate spatial and temporal resolution, which will depend on the size of the system and daily or seasonal variability in system characteristics. If, for example, a coastal system is seasonally hypoxic, more frequent sampling is required when compared to a permanently anoxic system.

To assess potential legacy effects related to benthic oxygen demand and nutrient release, the chemical composition of the sediment and porewater and the history of deposition should be assessed. Key parameters that should be included are sediment contents of organic carbon, sulfur, iron oxides and phosphorus and porewater concentrations of ammonium and hydrogen sulfide. Ecological impacts should be assessed by monitoring organisms relevant for a given ecosystem. This could include various types of phytoplankton (e.g. diatoms, cyanobacteria), fish species and benthic macro- and meiofauna.

(5) The costs and greenhouse gas footprint of the intervention should be carefully assessed. This includes that of the required power for the reoxygenation, the infrastructure and oxygen

transport. The information on costs, energy demand and greenhouse gas footprint should be communicated to all parties involved prior to an intervention.

(6) Results of any field trials, including perceived failures, should be reported with full transparency and publicly in open data bases and, preferably, in the scientific literature.

(7) When deciding on a potential implementation, the impacts of successful reoxygenation of a given location on the willingness to continue nutrient load reductions should be considered, since the latter have wider benefits.

7. Conclusions

With science-based suitability assessments and ethical and environmentally safe practices, artificial reoxygenation could be useful as a temporary measure at carefully selected sites and might allow detrimental consequences of coastal deoxygenation to be alleviated or avoided. Artificial reoxygenation will rarely be a permanent solution to coastal deoxygenation, however, because it does not address its root causes. Hence, reoxygenation efforts cannot replace essential measures to reduce the input of nutrients to coastal waters nor can it replace reductions of greenhouse gas emissions.

Acknowledgements

This paper summarizes the discussion of a workshop on Marine Reoxygenation organized by the IOC-UNESCO Working Group Global Oxygen Network (GO2NE), September 10-11, 2024. CPS acknowledges funding from ERC MARIX 854088. A short summary of the findings of the workshop was published previously as a feature in EOS. BD acknowledges support from ANID (Concurso de Fortalecimiento al Desarrollo Científico de Centros Regionales 2020-R20F0008-CEAZA and Centro de Investigación Oceanográfica en el Pacífico Sur-Oriental COPAS COASTAL FB210021). EMF acknowledges support from the Chancellor's Postdoctoral Fellowship Program at the University of California Santa Cruz.

References

- Beghoua H, Gorgues T, Fransner F, Auger PA and Memery L 2023 Contrasting responses of the ocean's oxygen minimum zones to artificial re-oxygenation *Environ Res Lett* **18** 084012
- Breitburg D, Levin, LA, Oschlies A, Grégoire M, Chavez FP, Conley DJ, et al 2018 Declining oxygen in the global ocean and coastal waters *Science* **359** 1–13
- Bryhn AC, Bergek S, Bergström U, Casini M, Dahlgren E, Ek C et al 2022 Which factors can affect the productivity and dynamics of cod stocks in the Baltic Sea, Kattegat and Skagerrak? *Ocean Coast Manag* **223** 106154
- Burdige D 2006 *Geochemistry of marine sediments* Princeton University Press
- Cai, W-J, Feely RA, Testa JM, Li M, Evans W, Alin SR et al 2021 Natural and anthropogenic drivers of acidification in large estuaries *Annu Rev Mar Sci* **13** 23–55

- 430 Carstensen J, Andersen JH, Gustafsson BG and Conley DJ 2014 Deoxygenation of the Baltic Sea
431 during the last century *Proc Natl Acad Sci USA* **111** 5628–5633
- 432 Conley D J, Bonsdorff E, Carstensen J, Destouni G, Gustafsson BG, Hansson L-A, et al 2009a
433 Tackling hypoxia in the Baltic Sea: Is Engineering a Solution? *Environ Sci Technol* **43** 3407–
434 3411
- 435 Conley D, Bjork S, Bonsdorff E, Carstensen J, Destouni G, Gustafsson BG et al 2009b Hypoxia-
436 related processes in the Baltic Sea *Environ Sci Technol* **43** 3412-3420
- 437 Dai M, Zhao Y, Chai F, Chen M, Chen N, Chen Y et al 2023 Persistent eutrophication and hypoxia in
438 the coastal ocean *Cambridge Prisms: Coastal Futures* **1** e19 1–28
- 439 De Brabandere L, Bonaglia S, Kononets MY, Viktorsson L, Stigebrandt A, Thamdrup B and Hall POJ
440 2015 Oxygenation of an anoxic fjord basin strongly stimulates benthic denitrification and DNRA
441 *Biogeochemistry* **126** 131–152
- 442 Diaz RJ and Rosenberg R 2008 Spreading dead zones and consequences for marine ecosystems
443 *Science* **321** 926–929
- 444 Duarte CM, Chapuis L, Collin SP, Costa DP, Devassy RP, Eguiluz VM, et al 2021 The soundscape of
445 the anthropocene ocean *Science* **371** eaba4658
- 446 Egger M, Jilbert T, Behrends T, Rivard C and Slomp CP 2015 Vivianite is a major sink for
447 phosphorus in methanogenic coastal surface sediments *Geochim Cosmochim Acta* **169** 217-235
- 448 Farias L, Beszteri B, Burfeld Castellanos AM, Doliwa A, Enss J, Feld CK et al 2024 Influence of
449 salinity on the thermal tolerance of aquatic organisms *Sci Total Env* **953** 176120
- 450 Fennel K and Testa JM 2019 Biogeochemical controls on coastal hypoxia *Annu Rev Mar Sci* **11** 105–
451 130
- 452 Forth M, Liljebladh B, Stigebrandt A, Hall POJ and Treusch AH 2015 Effects of ecological
453 engineered oxygenation on the bacterial community structure in an anoxic fjord in western
454 Sweden *ISME J* **9** 656–669
- 455 Gächter R and Wehrli B 1998 Ten years of artificial mixing and oxygenation: no effect on the internal
456 phosphorus loading of two eutrophic lakes *Environ Sci Technol* **32** 3659-3665
- 457 Gächter R and Müller B 2003 Why the phosphorus retention of lakes does not necessarily depend on
458 the oxygen supply to their sediment surface *Limnology and Oceanography* **48** 929-933
- 459 GESAMP 2025 The state of the science for marine carbon dioxide removal (mCDR) – A scientific
460 summary for policy-makers Paris, UNESCO-IOC (IOC Technical Series, 209)
- 461 Grégoire M, Oeschlies A, Canfield D, Castro C, Ciglenc̆ki I, Croot P et al 2023 Ocean Oxygen: the
462 role of the Ocean in the oxygen we breathe and the threat of deoxygenation Rodriguez Perez A,
463 Kellett P, Alexander B, Muñiz Piniella Á, Van Elslander J, Heymans JJ [Eds] Future Science
464 Brief No 10 of the European Marine Board, Ostend, Belgium ISSN: 2593-5232 ISBN:
465 9789464206180

- Hagens, M, Slomp CP, Meysman F, Seitaj D, Harlay J, Borges A and Middelburg J 2015 Biogeochemical processes and buffering capacity concurrently affect acidification in a seasonally hypoxic coastal marine basin *Biogeosciences* **12** 1561-1583
- Handmann P, Wallace D 2024 The global energy transition offers new options for mitigation of coastal hypoxia: Do we know enough? *Global Change Biology* 2024 **30** 17228
- Harris, LA, Hodgkins CLS, Day MC, Austin D, Testa JM, Boynton W et al 2015 Optimizing recovery of eutrophic estuaries: Impact of destratification and re-aeration on nutrient and dissolved oxygen dynamics *Ecol Eng* **75** 470–483
- Hermans M, Lenstra WK, van Helmond NAGM, Behrends T, Egger M, Seguret MJM et al 2019 Impact of natural re-oxygenation on the sediment dynamics of manganese, iron and phosphorus in a euxinic Baltic Sea basin *Geochim Cosmochim Acta* **246** 174–196
- Hounshell AG, McClure RP, Lofton ME and Carey CC 2021. Whole-ecosystem oxygenation experiments reveal substantially greater hypolimnetic methane concentrations in reservoirs during anoxia *Limnol Oceanogr Lett* **6** 33-42
- Hylén A, van de Velde S J, Kononets M, Luo M, Almroth-Rosell E, and Hall POJ: Deep-water inflow event increases sedimentary phosphorus release on a multi-year scale, *Biogeosciences* **18** 2981–3004
- IPCC 2023 Summary for Policymakers In: Climate Change 2023: Synthesis Report Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H Lee and J Romero (eds)] IPCC, Geneva, Switzerland, pp1-34
- Jilbert T, Slomp CP, Gustafsson BG and Boer W 2011 Beyond the Fe-P-redox connection: preferential regeneration of phosphorus from organic matter as a key control on Baltic Sea nutrient cycles *Biogeosciences* **8** 1699-1720
- Johnson M, van Doorn E, Hilmi N, Marandino C, McDonald N, Thomas H et al 2024 Can coastal and marine carbon dioxide removal help to close the emissions gap? Scientific, legal, economic, and governance considerations *Elem Sci Anth* **12** 00071
- Jutras M, Dufour CO, Mucci A, Cyr F, Gilbert D 2020 Temporal changes in the causes of the observed oxygen decline in the St Lawrence Estuary *J Geophys Res: Oceans* **125** e2020JC016577
- Jutras M, Mucci A, Chaillou G, Nesbitt WA and Wallace DWR 2023 Temporal and spatial evolution of bottom-water hypoxia in the St Lawrence estuarine system *Biogeosciences* **20** 839–849
- Katsev S, Tsandev I, L’Heureux I and Rancourt DG 2006 Factors controlling long-term phosphorus efflux from lake sediments: Exploratory reactive-transport modeling *Chem Geol* **234** 127-147
- Kemp, WM, Testa, JM, Conley, DJ, Gilbert, D and Hagy, JD 2009 Temporal responses of coastal hypoxia to nutrient loading and physical controls *Biogeosciences* **6** 2985–3008

- 502 Kowalik DA, García-Sánchez C, Brodrick PG, Gassett P and Caldeira K 2020 Evaluating hypoxia
503 alleviation through induced downwelling *Sci Total Environ* **719** 137334–137412
- 504 Kuliński K, Rehder G, Asmala E, Bartosova A, Carstensen J, Gustafsson BG et al 2022
505 Biogeochemical functioning of the Baltic Sea *Earth Syst Dynam* **13** 633–685
- 506 Lamping J, Worrall F, Morgan HUW, Taylor S 2005 Effectiveness of aeration and mixing in the
507 remediation of a saline stratified river *Environ Sci Technol* **39** 7269–7278
- 508 Lapham LL, Hobbs EA, Testa JM, Heyes A, Forsyth MK, Hodgkins C et al 2022 The effects of
509 engineered aeration on atmospheric methane flux from a Chesapeake Bay tidal tributary *Front*
510 *Environ Sci* **10** 866152
- 511 Lapham L, Lloyd KG, Fossing H, Flury S, Jensen JB, Alperin MJ et al 2024 Methane leakage through
512 the sulfate methane transition zone *Nature Geoscience* **17** 1277–1283
- 513 Larsen SJ, Kilminster KL, Mantovanelli A, Goss Z, Evans GC, Bryant LD, McGinnis DF 2019
514 Artificially oxygenating the Swan River estuary increases dissolved oxygen concentrations in the
515 water and at the sediment interface *Ecol Eng* **128** 112–121
- 516 Laruelle GG, Dürr HH, Slomp CP and Borges AV 2010 Evaluation of sinks and sources of CO₂ in the
517 global coastal ocean using a spatially-explicit typology of estuaries and continental shelves
518 *Geophys Res Lett* **37** L15607
- 519 Lehtoranta, J, Bendtsen J, Lännergren C, Saarijärvi E, Lindström M and Pitkänen H 2022 Different
520 responses to artificial ventilation in two stratified coastal basins *Ecol Eng* **179** 106611 1–13
- 521 Liu, X, Beusen AHW, van Grinsven HJM, Wang J, van Hoek WJ, Ran X et al 2024 Impact of
522 groundwater nitrogen legacy on water quality *Nat Sustain* **7** 891–900
- 523 Meier HEM, Dieterich C, Gröger M, Dutheil C, Börgel F, Safonova K et al Oceanographic regional
524 climate projections for the Baltic Sea until 2100 *Earth Syst Dynam* **13** 159–199
- 525 Mobley M, Gantzer P, Benskin P, Hannoun I, McMahon S, Austin D and Scharf R 2019
526 Hypolimnetic oxygenation of water supply reservoirs using bubble plume diffusers *Lake and*
527 *Reservoir Management* **35** 247–265
- 528 Norkko J, Reed, DC, Timmermann K, Norkko A, Gustafsson BG, Bonsdorff E et al 2012 A welcome
529 can of worms? Hypoxia mitigation by an invasive species *Global Change Biology* **18** 422–434
- 530 Oliveira, VH, Díez, S, Dolbeth, M, Coelho, JP, 2024 Restoration of degraded estuarine and marine
531 ecosystems: a systematic review of rehabilitation methods in Europe *J Hazard Mater* 469,
532 133863
- 533 Oeschies A, Stevenson A, Bach LT, Fennel K, Rickaby REM, Satterfield T et al (Eds) 2023 *Guide to*
534 *Best Practices in Ocean Alkalinity Enhancement Research* (OAE Guide 23), Copernicus
535 Publications, State Planet, 2-oae2023
- 536 Pezner AK, Courtney TA, Barkley HC, Chou, W-C, Chu H-C, Clements SM et al 2023 Increasing
537 hypoxia on global coral reefs under ocean warming *Nat Clim Chang* **13** 403–409

- Pitcher, GC, Aguirre-Velarde, A, Breitburg, D, Cardich, J, Carstensen, J, Conley, DJ et al 2021
System controls of coastal and open ocean oxygen depletion *Progress in Oceanography* **197**
102613
- Reed, DC, Slomp CP and Gustafsson BG 2011 Sedimentary phosphorus dynamics and the evolution
of bottom water hypoxia: A coupled benthic-pelagic model of a coastal system *Limnol Oceanogr*
56 1075-1092
- Reusch, TBH, Dierking J, Andersson HC, Bonsdorff E, Carstensen J, Casini M et al 2018 The Baltic
Sea as a time machine for the future coastal ocean *Science Advances* **4** eaar8195
- Singleton VL, Little JC 2006 Designing hypolimnetic aeration and oxygenation systems– a review
Environ Sci Technol **40** 7512–7520
- Slomp CP 2011 Phosphorus cycling in the estuarine and coastal zones: Sources, sinks, and
transformations In *Treatise on estuarine and coastal science*, vol 5, ed E Wolanski and DS
McLusky p201–229 London Academic Press
- Soetaert K, Middelburg JJ, Heip C, Meire P, Van Damme S and Maris P 2006 Long-term change in
dissolved inorganic nutrients in the heterotrophic Scheldt estuary (Belgium, The Netherlands)
Limnol Oceanogr **51** 409–423
- Sommer S, Clemens D, Yucel M, Pfannkuche O, Hall POJ, Almroth-Rosell E et al 2017 Major
bottom water ventilation events do not significantly reduce basin-wide benthic N and P release in
the Eastern Gotland Basin (Baltic Sea) *Front Mar Sci* **4** 18
- Stigebrandt A and Gustafsson BG 2007 Improvement of Baltic proper water quality using large-scale
ecological engineering *Ambio* **36** 280–286
- Stigebrandt A, Rosenberg R, Råman Vinnå L and Ödalen M 2015a Consequences of artificial
deepwater ventilation in the Bornholm Basin for oxygen conditions, cod reproduction, and
benthic biomass—a model study *Ocean Sci* **11** 93–110
- Stigebrandt A, Liljebladh B, De Brabandere L, Forth M, Granmo Å, Hall P et al 2015b An experiment
with forced oxygenation of the deepwater of the anoxic By Fjord, Western Sweden *Ambio* **44**
42–54
- Stigebrandt A and Andersson A 2022 Improving oxygen conditions in periodically stagnant basins
using sea-based measures - Illustrated by hypothetical applications to the By Fjord, Sweden
Continental Shelf Res **244** 104806
- Sulu-Gambari F, Hagens M, Behrends T, Seitaj D, Meysman, FJR, Middelburg J and Slomp CP 2018
Phosphorus cycling and burial in sediments of a seasonally hypoxic marine basin *Estuaries and
Coasts* **41** 921–939
- Tammeorg O, Chorus I, Spears B, Noges P, Nürnberg GK, Tammeorg P and Lüring M 2024
Sustainable lake restoration: from challenges to solutions *Wiley Interdiscip Rev Water* **11** e1689
- Testa JM, Kemp WM and Boynton WR 2018 Season-specific trends and linkages of nitrogen and
oxygen cycles in Chesapeake Bay *Limnol Oceanogr* **63** 2045–2064

- Testa JM, Hodgkins C, Forsyth MK, Moore A, Bailey E, Day MC et al 2025 Time scales of hypoxia-induced nutrient recycling in sediments: an experimental approach using engineered aeration *Environ Sci Technol* **59** 15115-15127
- Thamdrup B, Hansen W and Jørgensen BB 1998 Temperature dependence of aerobic respiration in a coastal sediment *FEMS Microbiol Ecol* **25** 189–200
- Turner RE, Rabalais NN and Justic D 2008 Gulf of Mexico hypoxia: alternate states and a legacy *Environ Sci Technol* **42** 2323–2327
- Tyler RM, Brady DC and Targett T, 2009 Temporal and spatial dynamics of diel-cycling hypoxia in estuarine tributaries *Estuaries and Coasts* **32** 123-145
- van Helmond, NAGM, Robertson EK, Conley DJ, Hermans M, Humborg C, Kubeneck LJ et al 2018 Removal of phosphorus and nitrogen in sediments of the eutrophic Stockholm archipelago, Baltic Sea *Biogeosciences* **17** 2745–2766
- Wallace, DWR, Jutras M, Nesbitt WA, Donaldson A, and Tanhua T 2023 Can green hydrogen production be used to mitigate ocean deoxygenation? A scenario from the Gulf of St Lawrence *Mitig Adapt Strateg Glob Change* **28** 56
- Weitkamp DE and Katz M 1980 A review of dissolved gas supersaturation literature *Trans Am Fish Soc* **109** 659-702.
- Zhang Y, Shitu A, Hang S, Ye Z, Zhao H, Xu W, Zhao J and Zhu S 2023 The effects of aerator noise on the swimming, feeding, and growth of *Micropterus salmoides* *Front Mar Sci* **10** 1242793
- Żygadłowska OM, Roth F, van Helmond NAGM, Lenstra WK, Venetz J, Dotisios N et al 2024a Eutrophication and deoxygenation drive high methane emissions from a brackish coastal system *Environ Sci Technol* **58** 10582-10590
- Żygadłowska OM, Venetz J, Lenstra WK, van Helmond NAGM, Klomp R, Röckmann T et al 2024b Ebullition drives high methane emissions from a eutrophic coastal basin *Geochim Cosmochim Acta* **384** 1-13