Peer-review status: This is the author's accepted manuscript (postprint) of a peer-reviewed article, posted to EarthArXiv. The version of record is published as:

K. Klemmer, E. Rolf, M. Rußwurm, *et al.*, "Earth Embeddings: Toward artificial intelligence-centric representations of our planet," *IEEE Geoscience and Remote Sensing Magazine*, 2026. DOI: 10.1109/MGRS.2026.3710416.

Please cite the published version.

Earth Embeddings: Toward artificial intelligence-centric representations of our planet

Konstantin Klemmer[1,2,*], **Esther Rolf**[3,*], **Marc Rußwurm**[4,*], **Gustau Camps-Valls**[5], **Mikolaj Czerkawski**[6], **Stefano Ermon**[7], **Alistair Francis**[6], **Nathan Jacobs**[8], **Hannah Kerner**[9], **Lester Mackey**[10], **Gengchen Mai**[11], **Oisin Mac Aodha**[12], **Markus Reichstein**[13], **Caleb Robinson**[14], **David Rolnick**[15], **Evan Shelhamer**[16], **Vincent Sitzmann**[17], **Devis Tuia**[18], **and Xiaoxiang Zhu**[19]

[1]LGND AI, Inc.
[2]University College London
[3]University of Colorado Boulder
[4]University of Bonn
[5]Universitat de València
[6]Asterisk Labs
[7]Stanford University
[8]Washington University in St. Louis
[9]Arizona State University
[10]Microsoft Research
[11]University of Texas at Austin
[12]University of Edinburgh
[13]Max-Planck-Institute for Biogeochemistry
[14]Microsoft AI for Good
[15]Mila, McGill University
[16]University of British Columbia and Vector Institute
[17]Massachusetts Institute of Technology
[18]École Polytechnique Fédérale de Lausanne
[19]Technische Universität München
[*]These authors jointly led this work, other authors are in alphabetical order.

Abstract

We present a new perspective for the flexible and efficient representation of geospatial data, tailored to and empowered by AI: *Earth embeddings*. Earth embeddings provide a unified and accessible vector representation of local geographic characteristics. They fuse different geospatial data sources across time and space, compress raw geospatial inputs into one dense representation, can be used to guide interpolation between observations, and can serve as a bridge to other foundation models like LLMs. This provides a powerful alternative to existing geospatial workflows that rely on heterogeneous data, hard-to-acquire expertise, and significant computation by the user: embeddings instead provide convenient representations, easily adaptable for downstream tasks, with minimal computation. We posit that Earth embeddings redefine geospatial analytics, transforming it from fragmented, task-specific modeling into a coherent, generalizable framework for AI. We approach this from both the users' and developers' perspectives, outlining a path for how the rapidly developing technology of Earth embeddings can reshape the way we store, represent, and use geospatial data, evidenced by recent research charting initial directions. We call on Earth embedding users and developers to align methodological and applied development and deployment within an interdisciplinary, open-source community.




Figure 1. The Earth is observed by diverse geospatial modalities—such as satellite imagery, sensor networks, and geotagged social media posts captured at specific times and locations. These multimodal data streams serve as inputs to AI models that learn to represent the state of the Earth in compact, unified forms known as Earth embeddings. Earth embeddings encode information for all locations in a lightweight and flexible format, enabling downstream analyses across domains, from terrestrial ecosystems to ocean dynamics.

Keywords: Geospatial Data, Deep Learning, Foundation Models, Earth Observation, AI for Science.

1 Introduction

Geospatial data are discrete snapshots acquired to capture information about the conditions and continuous dynamics of our planet. These data range from Earth observation or meterological imagery captured by hundreds of in-orbit satellites to human-generated social media posts and photos, tagged with geospatial metadata. At first glance, these data sources may appear completely different, but they share a fundamental commonality: all geospatial data are *indexed in space and time*. As a satellite image provides information about a particular geographic area at a particular point in time, so does a geotagged image or text description uploaded to point-of-interest databases like Open Street Map or social media platforms. Geospatial data are widely used to support modeling and decision making: Notable examples include disaster response [1], weather forecasting [2], transportation management [3], forest and wildfire management [4], water quality management [5], and disease spread modeling [6].

To process the ever-growing amounts of geospatial data, machine learning and artificial intelligence (AI) techniques have risen in popularity among geospatial practitioners, such as environmental scientists studying ecological changes [7], data scientists building flood maps at insurance companies [8], or urban planners forecasting travel demand [9]. Enabled by the key strengths of AI methods to represent complex data in flexible and generalizable formats, ***Earth embeddings* are vector representations of geospatial data at specific locations in space and time** (see Fig. 1) that allow us to capture the similarities and differences between locations according to their local features. They are produced by ***Earth embedding models***, which process data inputs into the vector shape of embeddings. The distinction between ***Earth embedding models*** which produce the ***Earth embeddings*** themselves is important throughout the paper, and we explain it in detail (and formally) in the box below.

Formal definition of *Earth embeddings*

Earth embedding vectors *emb* are produced by a family of embedding functions E that map continuous location inputs (i.e., longitude, latitude with optionally elevation, and

time) into a d-dimensional vector space:

$$E : (location) \rightarrow emb \in \mathbb{R}^d \ . \tag{1}$$

E, also referred to as *embedding models*, come in two forms: First, as *explicit models*, extracting embeddings from raw data (e.g. satellite imagery) associated with a location ($emb \sim E_{explicit}(data_{location})$). Examples for this include vision models combined with a global satellite image database. Second, as *implicit models*, returning embeddings from only location inputs ($emb \sim E_{implicit}(location)$). Here, location-specific information is stored directly withing the weights of e.g. a location encoder network $E_{implicit}$. This distinction is explained in detail in Section 4.1.

AI embeddings represent specific concepts contained within the data they are based on; words or images and in the case of Earth embeddings: any geographic data. Over the past decades, fields such as natural language processing (NLP) and computer vision (CV) have developed specialized neural network architectures to efficiently embed text and image data, including methods specific to Earth observation images.

The origin of "embeddings" in Natural Language Processing (NLP)

The concept of embedding spaces has its modern origins in natural language processing, where word embeddings techniques like Word2Vec [10] and GloVe [11] revolutionized the field by capturing semantic relationships between words. For example, in a word embedding space, the vector difference between "kitten" and "cat" is similar to the vector difference between "puppy" and "dog", illustrating how embeddings can capture meaningful semantic relationships. This idea of "distance" between words is also used in large language models (LLMs), which leverage embeddings to map similarity of words closer in embedding space.

Similar to how text or image embeddings capture semantic relationships, Earth embeddings map places and times that share similar properties closer together in embedding space. For example, an embedding of urban New York City may resemble that of urban Delhi more than that of rural Arkansas, reflecting functional rather than geographic similarity. Likewise, embeddings across time capture temporal variability—summer and winter embeddings in equatorial Costa Rica may be nearly identical, while those in highly seasonal Alaska may differ strongly—enabling quantitative comparison of environmental change and similarity across space and time.

Early versions of Earth embeddings are already enabling efficient, real-time monitoring of our planet and reducing barriers to accessing powerful geospatial analytics. Recent years have, for example, seen the publication of Earth embeddings based on satellite imagery [12–14], social media images [15], and demographic data [16]. The applications of Earth embeddings include image geo-localization [17], poverty mapping [18], detecting small-scale and artisanal mining [19], and satellite image super-resolution [20]. These applications also drive an increased commercial interest in the space, highlighted by a number of startups with Earth embeddings at the center of their product securing funding and e.g., Google Earth Engine launching their first Earth embedding product [21]. This leads to a rapidly growing, yet heterogeneous, ecosystem around Earth embeddings, with an increasing number of models and initiatives that share common objectives like data compression, fusion, interpolation, or increasing interoperability between

models.

Current efforts to develop and apply Earth embeddings are often not coordinated and lack a clear, community-sourced research agenda. One sign of this is that existing Earth embedding models are implemented in fundamentally different ways: many recent efforts focus on developing geospatial foundation models (GeoFMs)—such as Clay [22], Prithvi [23], Galileo [24], Smarties [25], and DOFA [26]. GeoFMs have been scaled into the billion parameter scale, for example with the SkySense family of models [27, 28]. Embeddings are more of a byproduct of these models, which are targeted at fine-tuning for specific downstream applications, such as image segmentation or classification. Recent foundation models, including Alpha Earth Foundation [21] and Tessera [29] focus explicitly on embeddings as a dedicated output and provide them as pre-computed *embedding databases*, either directly downloadable on e.g. Google Earth Engine, or available via an API interface. Simultaneously, a new class of pre-trained *implicit neural representation* models is emerging, such as GeoCLIP [15], SatCLIP [30], SINR [31], and Climplicit [32]. These models store geospatial information directly in their neural network weights, take location information (e.g. longitude, latitude and time) as inputs, and can be seamlessly integrated in existing deep models for geographic conditioning, image retrieval, or geo-sematic search and retrieval.

Earth Embeddings and Geospatial Foundation Models (GeoFMs)

Geospatial foundation models (GeoFMs) is the term used for large, multimodal deep learning systems developed for general purpose use and trained on diverse geospatial datasets [33] that ingest raw data (e.g. remote sensing imagery, climate records, socio-economic layers). Earth embeddings can be produced by many of these models, but are more a byproduct of GeoFMs, which are often meant to be finetuned for specific downstream tasks such as satellite image segmentation. They are one way–but not the only way–to produce Earth embeddings. In essence, GeoFMs are large-scale modeling and learning frameworks, whereas Earth embeddings constitute the interoperable, location-indexed data outputs that can be stored, shared, or queried independently of the model that created them.

While the increased attention and the emerging potential of Earth embeddings clearly demonstrate a need for further development, this conceptual heterogeneity in approaches impedes joint community building efforts and the organization of shared research directions that capture both the developers' and users' needs. Hence, harmonizing and formalizing Earth embeddings is necessary to unlock the emerging potential of Earth embeddings for both developers and end-users of geospatial AI systems.

This article provides this harmonization with concrete calls-to-action mapping out a research agenda for Earth embeddings. We posit that Earth embeddings will emerge as the dominant format of geospatial data in the AI age, enabling efficient deployment in critical use cases and connecting the current heterogeneous landscape of geospatial data modalities and formats. In the remainder of this paper, we present a unifying framework on Earth embeddings, defining their central functions in Section 2. We then outline specific application use-cases and goals for methodological development, capturing both user and the developer perspectives in Sections 3 and 4. Finally, in Section 5, we describe existing considerations and future steps to facilitate effective community building to advance a coherent, AI-native foundation for the next generation of geospatial analysis.




Figure 2. Earth embeddings provide different functions: (1) They compress high-dimensional data into a lower-dimensional vector format. (2) They fuse together different geospatial data modalities, from different types of images to text and tabular data. (3) They can interpolate to unseen spatiotemporal locations, where raw data is missing. (4) They are interoperable with other AI foundation models, such as LLMs, through aligned embedding spaces.

2 The Central Functions of Earth Embeddings

Current Earth embedding approaches are designed to serve one or more of four central functions, as illustrated in Fig. 2:

1. **Compression:** Earth embeddings can be interpreted as data compression, as they distill high-dimensional, location-specific, multi-modal data such as satellite images or geotagged text data into a smaller vector format [34]. While this compression loses some information and varies with training method, geospatial data's redundancy and spatio-temporal patterns make it easier to compress effectively. Some work in this area exists already, for example on the compression of climate data within deep networks [35]; however, dedicated compression models are needed, as current embedding models produce correlated embedding vectors [36] with a comparatively low intrinsic dimension [37].

2. **Fusion:** Earth embeddings fuse different geospatial data modalities–e.g.,images and text data–into a joint embedding space, leveraging their shared location and time information. For example, Earth embeddings trained on multi-modal geospatial data may represent both an optical satellite image and social media posts associated with a given location. This allows them to provide a more holistic location representation for general-purpose use. For instance, current multi-modal remote sensing foundation models like MMEarth [38], DOFA [26], Galileo [24], Smarties [25], or the SkySense model family [27, 28], which focuses on cross-modal alignment, fuse multiple data sources into feature embeddings representations at one location.

3. **Interpolation:** Some Earth embeddings–those obtained via location encoders–are available in continuous space and time. They are based on *neural implicit representations* (see Section 4.1), which learn a smooth surface in embedding space by interpolating between seen training locations. As such, any location (in continuous space and time) may be queried from location encoder models like SatCLIP [13], SINR [31], or GeoCLIP [15], even if it was not present in the training data. This returns an Earth embedding representing an interpolation of nearby locations with available training data.




Figure 3. Earth embeddings can be used for diverse applications: For example, **a)** Earth embeddings may be passed through a learned prediction model to generate wall-to-wall dense predictions of forest cover, population density, or other outcomes across the globe. They **b)** may be used as auxiliary or conditioning context in other predictive tasks, such as animal species classification. Similarly, they **c)** could be used as contextual input to a generative model, enabling image synthesis reflecting certain ground conditions, such as flooding. Lastly, **d)** Earth embeddings can be used to query similar locations (i.e. find other locations with solar farms) around the world by matching similar Earth embeddings. As Earth embeddings are lightweight, integrating them in existing models in any of these examples can be done quickly, without any need for processing or analyzing raw data.

4. **Interoperability:** Earth embeddings can be used to inject geospatial contextual information into other AI foundation models. They can be seen as interoperable "location tokens" (Fig. 2), allowing users to interact with Earth embeddings using text prompts directly, asking queries such as "What are similar locations to the location (lon, lat)?" and to augment current text-based assistants and foundation models efficiently with location information. An analogous approach is to exchange pre-trained implicit embedding models that are becoming increasingly integrated in other deep frameworks, for instance for image synthesis [39] of geography-aware super-resolution [20], to facilitate interoperability between models through a joint location context.

3 User Perspectives: Earth Embeddings Unlock New Use-cases and Expedite Existing Ones

Earth embeddings provide users with a compact way to access the multifaceted contextual information encoded in geospatial data. From a user's perspective, Earth embeddings function as a versatile input that can be easily plugged into existing or new workflow, often reducing data-processing demands while improving model performance. Because general-purpose Earth embeddings are designed for broad downstream use, they allow practitioners to enhance familiar tasks and explore entirely new applications (Fig. 3). Below, we outline key ways in which users can employ Earth embeddings for prediction, conditioning, simulation, and search:

1. **Direct prediction:** Earth embeddings can be used as the input to predictive models that

learn spatially distributed outcomes, in much the same way that raw data would be input into AI models. Examples include temperature, species presence, or population density prediction across space [13, 16, 31, 40, 41]. Earth embeddings also have shown to be competitive or even outperform hand-crafted specialized features with random forests in applications like tree species identification [42]. Since Earth embeddings already represent (often major) data pre-processing, they often allow for learning with smaller predictive models that one would need if using the raw data (e.g. satellite data, text data) as input [12]. Additionally, in contrast to traditional geospatial analytics and GeoAI workflows, Earth embeddings remove the need to query databases of raw data, which can significantly ease storage and accessibility requirements.

2. **Geographic conditioning:** Earth embeddings can serve as "geographic priors" which provide geospatial context to downstream applications. Earth embedding models can be integrated as auxiliary module in an existing workflow, conditioning GeoAI models to the specific area of interests (e.g., continents), resulting in location-aware prediction models [15, 24, 43]. Strategies employing Earth embeddings for geographic conditioning have been used to improve classification of animal images [44, 45] and to generate synthetic remote sensing imagery [20, 39, 46]. Earth embeddings can also serve as auxiliary data in settings with limited labels and in the face of domain shift [47], helping to boost performance. These successes are particularly inspiring as research into novel GeoAI architectures and training procedures that incorporate Earth embeddings as auxiliary contextual information is still in its infancy.

3. **Data synthesis & simulation:** Earth embeddings can be integrated in generative models to synthesize and simulate Earth processes and can serve as a building block for improved simulators of dynamical processes on the Earth. This forward-looking use case for Earth embeddings aims to incorporate not only observational data, but to also integrate scientific knowledge about physical processes, such as atmospheric dynamics. While no such systems exists to date, Earth embedding conditional generative models such as [39] may be expanded into simulators, while spatio-temporal location encoding methods may be tested for forecasting and scenario generation [32, 35, 48].

4. **Geo-semantic search and retrieval:** Earth embeddings can also be used for fine-grained, semantic search. They enable a "queryable Earth", acting as a lookup table for efficiently identifying specific concepts of interest, such as solar farms, at scale: Knowing the Earth embedding of a satellite image containing a solar farm, we can conduct a similarity search among all other available Earth and return locations with high similarity scores [49]. Furthermore, Earth embeddings aligned with text embedding spaces allow for text-based queries and retrieval [15, 50–52]. Due to their small size, Earth embeddings can even unlock global search, which is computationally infeasible with e.g. raw satellite imagery.

While general-purpose Earth embeddings are not tailored to any specific application or use case, they are fundamentally constrained by the raw data they are based on. SatCLIP Earth embeddings [13], for example, which are based on multi-spectral satellite imagery, may only represent phenomena that are captured by large-scale overhead imagery, such as climate zones or forest coverage, but not, for example, ocean currents. Conversely, GeoCLIP [15] embeddings are based on ground Flickr images, and naturally capture patterns that are visible from a street-view. Futhermore, both SatCLIP and GeoCLIP–as examples of implicit neural representations (detailed in Section 4.1)–provide large-scale, low-resolution embedding fields, while explicit

feature extraction approaches like AlphaEarth Foundations [21] or Tessera [29] provide much higher resolution embeddings, as they process the underlying geolocated raw data.

With strong evidence that Earth embeddings can improve existing GeoAI workflows and unlock new use-cases, it is critical to align research directions with user needs. We synthesize several steps along this path, highlighting how the user and research community can work together to improve the responsible and effective application of available Earth embeddings models and data products in the call to action below:

Call to action: Advancing analyses and applications with Earth embeddings.

While Earth embeddings show clear potential to push the frontier of geospatial analysis, there are several bottlenecks hindering their broader adoption into unexplored areas of use. Here we highlight the most pressing application-centric research gaps for advancing the utility of Earth embeddings for public and private sector use:

- **Evaluating and benchmarking Earth embeddings:** In order to advance the capabilities of Earth embeddings, we must be able to comprehensively measure and benchmark their performance on a wide range of use-cases. Expanding on current benchmarks that measure the capacity of Earth embeddings to capture specific quantities (e.g. forest cover or population densities) [41, 53], more varied types of evaluations are needed. Future evaluation frameworks could test Earth's embeddings' ability to capture interactions of different planetary processes, such as the connection between climate change and human migration, or their utility for search and retrieval. In contrast to many existing GeoAI benchmarks that are primarily for image inputs, good benchmarks for Earth embeddings will focus on challenges of prediction when location and time references are used as inputs [54]. This could include evaluating spatial interpolation and extrapolation, performance under different degrees of spatial bias in the labels, and performance across different spatiotemporal scales.
- **Explainable and interpretable Earth embeddings:** Embeddings obtained by current deep networks often lack interpretability [55]. That is, we don't know which part of an Earth embedding represents a given ground condition, such as a tree or a building– or if the embedding captures these ground conditions at all. This impedes our ability to use Earth embeddings to understand interactions or explore drivers of variations, such as seasonal versus geographic distances. Black-box methods complicate decision making [56]–a general challenge with modern AI. As such, improving Earth embedding interpretability (e.g., understanding which input modality contributes to which part of the Earth embedding) can help to make them more attractive for integration into decision-support and real-world deployment settings. Data provenance and metadata reporting standards are further key factors in supporting interpretability efforts.
- **Areas, resolutions and periods of applicability:** Future Earth embeddings might be produced with a specific area, resolution or time period of applicability in mind. For example, while SatCLIP [13] provides implicit Earth embeddings for the whole world that can be generated with infinitesimal resolution, its effective resolution is quite coarse, constrained by the capacity of its location encoder neural network and the smoothness the location-image matching objective enforces. To obtain high-resolution Earth embeddings from location encoders, it might be beneficial to train them on a

specific area of interest, as e.g. [57], who train a SatCLIP-style location encoder for Spain specifically. Similarly, explicit embeddings might be constrained by the availability of input data. For example, a recent study uses explicit embeddings obtained via National Agriculture Imagery Program (NAIP) imagery and the Clay [22] foundation model to map log yards [58]–a use case that, because of the availability of NAIP, is constrained to the US and specific years. Overall, we expect to see the emergence of more use-case specialized Earth embedding products, aimed at covering specific geographic areas, periods and resolutions.

- **Learning planetary processes with Earth embeddings:** It has long been a dream of geospatial researchers to create a digital version of our planet, a "digital twin" [59]. Earth embeddings may offer a step towards this dream: as they learn to fuse and compress multi-modal geospatial data into an succinct representation of the planet within their vectors, Earth embeddings contain information necessary to understand the inter-connections and dependencies of different planetary processes–from seasonal weather to bird migration and land-use change. For example, the Climplicit Earth embeddings that implicitly encode climate data [32] are shown to be competitive models for ecological tasks like species distribution modeling and and plant trait regression, underscoring the relatedness of data from seemingly very different domains.

4 Developer Perspective: Learning to Embed our Planet with AI

From a developer's perspective, the central challenges of Earth embeddings are how to learn and serve them effectively. Developers must choose between fundamentally different strategies for generating embeddings–explicit feature extraction or implicit neural representations–as well as between delivery interfaces such as embedding models or embedding databases. This section provides a unified conceptual view of these design choices, outlining how Earth embeddings can be built, stored, and accessed, and outlines objectives for future methodological advances.

4.1 Earth Embedding Models: Explicit Feature Extraction versus Implicit Neural Representation

To capture the characteristics of a given input location, Earth embeddings capture information from geospatial data sources such as satellite imagery and social media data. The Earth embedding models producing them can be split into two categories, outlined in Fig. 4: encoding data sources explicitly (Fig. 4, left-top) or implicitly via neural network weights (Fig. 4, left-bottom).

4.1.1 Explicit feature extraction

Explicit feature extraction methods aim to directly embed geospatial data into a vector representation. There is a breadth of research on feature extraction of Earth observation and spatio-temporal data, most recently using neural network architectures and learning objectives from computer vision or time series analysis [12, 14, 21, 24, 29]. Since the raw data that these models encode is always geolocated, their embeddings are inherently linked to specific locations and times (e.g. the spatial extents and time stamp of images), as a function of the content of the inputs.




Figure 4. There are different methods for obtaining Earth embeddings (left side) and interfaces for accessing them (right side). Explicit encoding approaches embed data from geospatial databases (indexed in space and time) using e.g. a GeoFM. The resulting pre-computed embeddings can be stored in a new embedding database which can be accessed using location queries at inference time. Both the GeoFM used for producing embeddings and / or the resulting embedding database may be released. The implicit encoding approach directly encodes geospatial information within the weights of a location encoder network. The location encoder serves both at "model" and "database" and, once released, can also be queried simply using location queries.

An early explicit embedding approach, MOSAIKS [12], demonstrates that embeddings of satellite imagery extracted with random convolutional features can be used for direct prediction of a variety of economically and environmentally-relevant downstream tasks. Similarly, deep learning models trained with self-supervised contrastive learning trained on BigEarthNet [60] or SSL4EO [61] can be also seen as explicit embedding models that extract deep embedding features from Sentinel-2 satellite imagery. More recently, multi-modal foundation models like DOFA [26], Smarties [25], or MMEarth [38] can extract joint embeddings from multiple remote sensing and meteorological modalities. These larger models are still often open-source but are typically resource-intensive to deploy. As such, embeddings are increasingly pre-computed and made available via embedding databases like AlphaEarth [21], Tessera [29], or MajorTom [14], which we detail in Section 4.2.

Embedding vectors produced by explicit embedding models are fine-grained and typically produced at high native resolution of the underlying data. That is, the explicit Earth embedding reflects exactly the location and extent (e.g. bounding box), resolution (e.g. ground sample distance) and generation time of the input data (e.g. satellite image). This makes them well-suited for geo-sematic search and retrieval and detailed direct prediction of visual tasks like land cover and land use classification, but less-suited for geographic conditioning and data synthesis. Explicit feature extraction approaches can produce embeddings at either the patch level (i.e. representing a whole image or image patch), such as Clay [22] or MMEarth [38] or the pixel level, such as AlphaEarth [21]. This difference further informs their specific applicability for downstream use.

4.1.2 Implicit Neural Representations

Implicit Neural Representations are an alternative approach that store Earth embeddings within the weights of deep networks. They are based on location encoder networks [62]–models that take geographic coordinates (and other potential metadata such as timestamps) as inputs and are inspired by approaches in computer vision [63, 64]. Spatial and temporal coordinates are treated as continuous inputs and encoded using functions that allow users to define the resolution at which dynamics are modeled. For example, geographic coordinates can be encoded using Spherical Harmonics functions that reflect the spherical nature of Earth and allow for varying

scales (e.g. meter vs kilometer scale). Likewise, temporal trends of Earth's processes with periodic nature, such as seasons (annual scale) or day versus night time (daily scale), can be encoded with functions of varying frequencies, such as Fourier functions.

Location encoders can be trained in a supervised fashion to reproduce a given location-specific value (e.g. temperature) and effectively interpolate between training points. Typical applications include the estimation of sea ice thickness [48], housing prices [45, 65], or species presence probabilities [31, 44, 66]. SINR [31] has demonstrated that even in a supervised setting, features from intermediate layers encode general semantic information following topographic and biome-specific boundaries–and, as such, could serve as Earth embeddings.

When trained in a self-supervised fashion through geolocalization, location encoder models return embedding vectors that capture the patterns that make a particular location unique, which makes these embeddings potentially effective for a wide range of applications. Examples of recent studies include GeoCLIP trained on geolocating ground-based Flickr images [15], or SatCLIP [30] and CSPNet [67] trained on overhead satellite imagery. These models are usually light-weight, open-source and can be easily downloaded and integrated into existing workflows. They typically provide embeddings at a lower resolution suitable for general geographic conditioning, data synthesis and direct prediction of larger-scale direct prediction tasks like air quality or disease mapping, but are not optimal for retrieval or fine-grained tasks like land cover classification.

4.1.3 Hybrid approaches

Hybrid approaches combine explicit feature extraction with implicit neural representations. A straightforward but effective approach is concatenating high-resolution explicit feature embeddings with low-resolution implicit embeddings. Such an approach, for example, won the NeucoBench challenge [53] at the 2025 CVPR Earthvision workshop. A more complex strategy involves a retrieval augmented generation (RAG). For example, the RANGE model [40] dynamically queries suitable high-resolution explicit image features using implicit embeddings as query. These examples highlight that hybrid approaches can achieve good results in practice, though possibly at the cost of additional methodological complexity.

4.2 Earth Embedding Interfaces: Embedding Models versus Embedding Databases

For widespread adoption, the interface through which users access Earth embeddings determines how easily these embeddings can be used and integrated into downstream applications. This holds regardless of whether embeddings are produced by "explicit" or "implicit" approaches (Section 4.1). Two conceptually distinct interfaces have emerged: releasing embedding models (Fig. 4, right-top) directly and releasing embedding databases (Fig. 4, right-bottom). Each enables different forms of access, affords different degrees of flexibility and scalability, and introduces its own constraints.

4.2.1 Embedding models

Embedding models deliver the model as code and parameters. More specifically, the user must download the model implementation with its trained parameters and infer the embeddings for given coordinates on an appropriate device (optimally, using a GPU). For the implicit approach, only the coordinates are needed as input, while an explicit feature extraction model additionally requires the imagery (or other applicable data) at the corresponding the location to be collected, loaded, and pre-processed into model-compatible inputs. This complexity requires users to bring

some familiarity with the principles of deep learning and the underlying data and software implementations, as well as sufficient resources, in terms of the computation time, memory, and type of device. When these requirements are met, this interface offers flexibility: the explicit embeddings can be extracted from any raw self-compiled dataset, while implicit embeddings can be queried for any new point location. The underlying model can be analyzed through interpretable methods like Grad-CAM [68] and altered by fine-tuning it to a given use case, which often increases the final downstream accuracy. Sharing pre-trained models is a common practice across the machine learning community with open-source code and open-licensed parameters available to download on public platforms like GitHub or HuggingFace. Examples for this approach include implicit approaches like SatCLIP [13], Climplicit [32] and GeoCLIP [15], or explicit approaches like AnySat [69], MMEarth [38], Galileo [24], and MOSAIKS [12].

4.2.2 Embedding databases

Embedding databases deliver the pre-computed model outputs as a database of static embedding vectors corresponding to locations, which can be built on existing geospatial data infrastructure and served from storage or over a network protocol. The storage format could be as simple as tabular data on disk or as sophisticated as global-scale backend systems like Google Earth Engine, and in the same fashion the protocol could provide a simple exchange of coordinates for vectors or a sophisticated set of options over spatiotemporal regions of interest. Creating and hosting Embedding databases require storage capacity or network bandwidth, which can incur substantial costs on the developer's side, which we detail more in Section 5. However, for the users' side they come with the clear advantage in that they do not require deep learning expertise or computation of the underlying model by the user. This makes embedding databases applicable to a broad user base of scientists and practitioners from a diversity of disciplines. However, this accessibility comes with limitations: given the static nature of the pre-computed embeddings, it is not possible to run, alter, or inspect the underlying model or its parameters themselves. While Embedding databases are easy to adopt, extending them is often impractical or impossible. Embeddings-as-data is an emerging interface, popularized by AlphaEarth [21] built on prior pioneering works like MOSAIKS [12] and Major TOM [14]. In practice, embedding databases are mainly relevant for explicit feature extraction models that are often large and computationally expensive to run, while implicit embedding representations are comparatively small and served as embedding models.

Call to action: Challenges and opportunities for improving Earth embeddings.

There are several open methodological challenges that motivate future research to improve the utility of Earth embeddings:

- **Model capacity:** The embedding model capacity required for producing effective yet informative Earth embeddings remains an open research problem: Current implicit encoding models (i.e., location encoders) are few-layer multi-layer perceptrons with relatively few parameters. This leads to limitations in their capacity for storing geospatial patterns. Future advances could, for example, take inspiration from natural language processing, where transformers with several blocks of positional embeddings and feed forward layers have proven highly effective in storing large amounts of text data [70]. Similarly, it is unclear whether explicit feature extraction models (e.g. GeoFMs) have reached sufficient capacity to model the full breadth and expressivity of Earth data. An

alternative approach to solve this challenge could involve hybrid approaches between precomputed embedding databases and implicit neural embedding representations: for instance, Dhakal et al. [71] used location encoders with pre-combined embedding fields for more fine-grained embedding retrieval.

- **Spatio-temporal encodings & heterogeneity:** Current Earth embedding models have statically defined parameter spaces, with the same number of parameters for every point in space and time. Geospatial information, however, can be complex in one location, but sparse in another, and manifest differently at different spatio-temporal resolutions. For example, while a satellite image over New York City and over the Pacific Ocean might be of the same size, the former is much more information dense than the latter. Future Earth embeddings models should be able to reflect this by varying their capacity based on location, otherwise, a globally trained model might smooth over a specific heterogeneous region leading to poorer local performance. Similarly, different sensors and data modalities might be more or less important in different areas of the world. For example, radar data might be more important in areas with high cloud coverage or in arctic regions during winter, when sunlight is scarce. Adaptations in positional embeddings or distinct deep network architectures might be taken from advances such as adaptive coordinate networks in computer vision [72]. Encoding strategies for spatial and temporal coordinates should reflect the resolutions required by downstream users.
- **Data curation and scaling:** The training and downstream usage of Earth embedding models need to be explored with respect to their data–and how additional data improves performance or not. These analyses should be conducted along the different dimensions of geospatial data: space, time and modality. Beyond intuitive guidelines, such as that Earth embeddings should be trained on data covering all regions of the world and all months of the year, the marginal benefits of e.g. adding one more spatial location, one more timestep, or one more sensor or modality are emerging as a research question [73] but remain largely unanswered. The discussion on data importance and scaling laws is directly related to that of *spatio-temporal heterogeneity* above, as it might be important to "overrepresent" heterogeneous areas in the training data compared to homogeneous areas [74], or to re-sample or re-weight the different dimensions according to better align with the spatial regions, temporal intervals, and input modalities of interest.
- **Learning objective:** The learning objectives used to train models that produce Earth embeddings remain largely unexplored. To date, location encoder approaches have largely relied on supervised training and CLIP-style contrastive learning (e.g., geolocalization). Models used for explicit embedding generation, often large-scale convolutional or transformer models, often simply adapt learning objectives from other machine learning areas (e.g. computer vision, NLP). Future work should focus on exploiting the unique characteristics of geospatial data, such as spatial autocorrelation or temporal dynamics, for designing dedicated learning objectives. There exist several promising proof-of-concept studies in this direction [30, 75, 76].

5 Community Perspective: Fostering the Earth Embedding Ecosystem

As Earth embeddings mature, a broader ecosystem is emerging around them–one that depends not only on scientific progress, but also on how models, data, and infrastructure are shared and sustained. Two key aspects shape this ecosystem that includes users and developers: the openness of embedding resources and the practical costs of producing and maintaining them. A collective effort is needed to maintain open-accessibility and cultivate the impactful applications required to justify and sustain the costs associated with creating and maintaining embedding models and databases.

5.1 Open source code, embedding models, and embedding datasets

Open source approaches enable the broad adoption of Earth embeddings. Whatever the modeling approach and interface, the options for adoption and extension are gated by how closed or open a project is. The Marin foundation model project and community (`http://marin.community`) define a trichotomy of degrees: open-weight (with weights but no code or data), open-source (with weights, code, data), and open-process (with weights, code, data, process details, and participation). Existing Earth embedding releases, both models and databases, mostly choose open-weight or open-source release. Major TOM [14] is uniquely open-process at this time, with its public standard and path to contributing. We encourage more similar open-process efforts to accelerate progress on Earth embeddings. Other projects releasing both the embedding model and embedding dataset include MOSAIKS [12], Tessera [29], Clay [22], or Presto [77]. AlphaEarth [21] constitutes a more restrictive open-output policy, releasing the embedding dataset but opting not to share the underlying model used to generate the embeddings. Our perspective is that sufficiently open sharing is a prerequisite for community-driven advancement of Earth embeddings, affording more rapid progress for research and development of the embeddings and for their applications.

5.2 Earth embedding cards

To help formalize reporting standards for and intended use of Earth embeddings, we propose "Earth embedding cards". This draws heavily on the templates for "datasheets for datasets" [78] and "model cards for model reporting" [79]. Since embeddings share properties of both datasets and models, we encourage authors draw from union of relevant items from both datasheets and model cards, and we highlight specific questions from those frameworks that might be especially relevant or take special care in the case of Earth embeddings. We further suggest key questions that are unique to embeddings and Earth data. We emphasize that this is intended as a starting place from which community-driven modifications can further improve reporting and transparency. We sketch out a template showing an example workflow for an "Earth embedding card" in Appendix A.

5.3 The computational & environmental costs of Earth embeddings

The computational costs are another major factor to consider. While embedding databases lower barriers for adoption and use, they introduce substantial and often underappreciated costs up front. This makes precomputation infeasible for most research groups, with the dominant expense in large-scale inference being not GPU compute but the data pipeline and storage.

To illustrate the computational costs associated with Earth embeddings, and to allow for

comparisons with compute and storage needs of existing geospatial data paradigms, let us consider the following example: A single NVIDIA V100 GPU can process an *embedding model* like a U-Net with an EfficientNet-B7 backbone at roughly 900,000 km^2 per hour using 4.7 m PlanetScope imagery—enough to process all land on Earth in about seven days for an estimated compute cost of only $514 on Microsoft Azure. Yet the corresponding imagery totals approximately 108 TB uncompressed, requiring co-located compute and storage to maintain high utilization, and generating roughly 432 TB of 64-dimensional embeddings (assuming 1 byte per dimension). Storing these embeddings in commercial cloud storage would cost around $8,000 per month ($2,000 per 100 TB of Azure Blob "hot" storage as of September 2025), or nearly $100,000 per year—far exceeding compute costs within days of operation. In comparison, an *embedding database* like AlphaEarth [21] faces similar challenges: maintaining multi-year, global-scale precomputed embeddings at 100 m resolution implies storage in the hundreds of terabytes (about 475 TB for five time steps), that can cost approximately $9,500 per month even without redundancy.

These calculations are a rough estimate, but explain the underlying challenges behind why precomputed embeddings remain rare outside of major corporations or space agencies and point to a key research challenge: developing efficient compression, caching, and dynamic delivery schemes that reduce the dependency on massive static storage, or rethinking embedding delivery as on-demand computation where compute acts as a substitute for long-term storage.

We can also expand our example estimation to include estimates of energy usage and carbon emissions. We might estimate that the one-time compute for the above example draws on the order of 50 kWh at the GPU and roughly 150 kWh at the facility level, accounting for host-server overhead and a typical data-center power usage efficiency. Depending on grid carbon intensity, this corresponds to between 10 kg CO_2e on a low-carbon grid, 75 kg CO_2e at the global-average intensity (480 gCO_2/kWh) and over 100 kg CO_2e on a carbon-intensive grid – comparable to driving a typical passenger car a few hundred kilometers.

We note that this carbon estimate counts operational energy only and excludes the embodied carbon of the hardware. The recurring power draw of storing hundreds of terabytes of embeddings repeats indefinitely, whereas the compute emissions are incurred once. As with the dollar costs, this reinforces that storage and delivery, not raw inference, dominate the sustainability profile of precomputed embedding products.

Call to action: Building an accessible, interdisciplinary and impactful Earth embedding community.

Broad institutional support for Earth embeddings depends on community efforts to establish them as a distinct geospatial data modality that requires its own infrastructure scaling efforts. In particular, the interfaces used to access Earth embeddings and the computational requirements for storing them dictates who is able to develop, use, and analyze them. We envision an open ecosystem that includes researchers and model developers, users, and data providers. We identify the following community objectives to support building the Earth embedding ecosystem accessibly, efficiently and with a diverse user-developer community at its center:

- **Interdisciplinarity:** We need durable structures that put domain scientists, AI researchers, product engineers, and data stewards in communication with each other

from the start. Such structures include funding for cross-appointments and exchange residencies, co-designed benchmarks drawn from real decision contexts, and community sprints where method design, labeling, and evaluation are co-developed. Priority should go to use cases proposed by practitioners outside AI, with dedicated resources for inclusive participation from underrepresented communities and institutions.

- **Traceable computational infrastructure:** We need shared, sustainable infrastructure that makes access to Earth embeddings affordable and transparent. Rather than storing massive datasets permanently, we should promote on-demand generation where possible. Tracking the resources used is essential so that the costs of producing and distributing embeddings can be fairly shared among those who benefit from them. One key aspect to consider is implementing traceability of the computed embeddings to commercial downstream applications to fund the expensive but optimally open-source and open-access infrastructure.
- **Embedding standards and interfaces:** We need shared technical standards that link embedding models and embedding datasets, across both explicit feature embeddings and implicit neural representations. These standards should define how embeddings are described, versioned, and shared, with consistent metadata on location, time, provenance, and quality. Crucially, standardized protocols for how models and databases exchange geospatial context in a common format would allow different systems to exchange embeddings, making it easier to combine embeddings for specific downstream tasks and to move seamlessly between model-based generation and precomputed databases. Standardized conventions on embedding structures can help with efforts around versioning and cross-modal comparability of embedding products.
- **Reporting and documentation:** We envision "Embedding Cards" (see section 5.2)–a combination of model cards [79] and data sheets [78]–that report training data coverage, objectives, known failure modes, spatial/temporal biases, uncertainty calibration, and safety/privacy considerations. Releases should include exact recipes: code, configs, hashes, container images, and seeds—and make use of evaluation suites spanning scale transfer, domain shift, and interpolation/extrapolation (see the Call to action box for users in Section 3).

6 Conclusion

AI is rapidly transforming the geospatial analytics landscape and Earth embeddings are emerging as a central data format within this evolution. Earth embeddings allow us to compress and process the vast amounts of Earth data more efficiently, improving existing geospatial analytics and unlocking new applications.

This perspectives paper harmonizes a rapidly evolving and heterogeneous field and establishes Earth embeddings as an emergent research domain at the intersection of geospatial sciences and AI. We formalize Earth embeddings within a clear technical definition, harmonizing research over the past several years. We define functions and properties of Earth embeddings and outline current and emerging use cases. From these, we derive open research challenges centered around concrete actions items for the scientific community:

- We call on the community to tackle **application-centric** research challenges, ranging from

better evaluations for Earth embeddings to making them more interpretable. The goal of this line of work is to *improve accessibility and enable new use cases for practitioners.*

- We call on the community to tackle **methodological** challenges for improving and scaling future-generation Earth embeddings. Work on encoding our planet is in its infancy and methodological development is essential for *enabling more efficient, expressive and adaptable* Earth embeddings.
- We call on the **community of users and developers** to center considerations around *interdisciplinarity, infrastructure, standards, reporting and documentation* to support the development of Earth embeddings that are accessible and effective.

The research agenda we outline is fundamentally interdisciplinary: Earth embeddings will rely on feedback from domain scientists, e.g. in ecological, geological, oceanographic, and atmospheric sciences, that incorporate Earth embeddings into their analyses and from data practitioners applying Earth embeddings in their workflows and products. This research agenda is also far-reaching and ambitious: We believe that Earth embeddings are an important component in enabling real-time Earth monitoring and that they can help to unlock a deeper understanding of planetary processes. We hope that our perspectives and proposed research avenues can help inform the directions Earth embeddings research and development will take in the years to come, focusing on maximizing their utility for scientific discovery and deployment in critical use cases.

Acknowledgements

This publication was supported by the Open Access Publication Fund of the University of Bonn. Rußwurm acknowledges funding by the Deutsche Forschungsgemeinschaft, under Project 572735710. Esther Rolf acknowledges NSF Award 2153040, The Environmental Data Science Innovation and Impact Lab data synthesis center. Gengchen Mai is supported by NSF Award #2521631. The authors thank Taylor Geospatial for their support. Konstantin Klemmer, Esther Rolf, and Marc Rußwurm jointly led this work. The other authors are listed in alphabetical order in the byline and author biographies.

References

[1] Majda Ivić. Artificial intelligence and geospatial analysis in disaster management. *The International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, 42:161–166, 2019.

[2] Remi Lam, Alvaro Sanchez-Gonzalez, Matthew Willson, Peter Wirnsberger, Meire Fortunato, Ferran Alet, Suman Ravuri, Timo Ewalds, Zach Eaton-Rosen, Weihua Hu, et al. Learning skillful medium-range global weather forecasting. *Science*, 382(6677):1416–1421, 2023.

[3] Christoph Willing, Konstantin Klemmer, Tobias Brandt, and Dirk Neumann. Moving in time and space–location intelligence for carsharing decision support. *Decision Support Systems*, 99:75–85, 2017.

[4] Jamie Tolan, Hung-I Yang, Benjamin Nosarzewski, Guillaume Couairon, Huy V Vo, John Brandt, Justine Spore, Sayantan Majumdar, Daniel Haziza, Janaki Vamaraju, et al. Very high resolution canopy height maps from RGB imagery using self-supervised vision trans-

former and convolutional decoder trained on aerial lidar. *Remote Sensing of Environment*, 300:113888, 2024.

[5] Rachida El Morabet, Larbi Barhazi, Soufiane Bouhafa, Mohammed Abdullah Dahim, Roohul Abad Khan, and Nadeem A Khan. Geospatial distribution and machine learning algorithms for assessing water quality in surface water bodies of Morocco. *Scientific Reports*, 13(1):20599, 2023.

[6] Arman Hossain Chowdhury and Md Siddikur Rahman. Machine learning and spatio-temporal analysis of meteorological factors on waterborne diseases in Bangladesh. *PLoS Neglected Tropical Diseases*, 19(1):e0012800, 2025.

[7] Maximilian Pichler and Florian Hartig. Machine learning and deep learning—a review for ecologists. *Methods in Ecology and Evolution*, 14(4):994–1016, 2023.

[8] Mohammed Khalaf, Abir Jaafar Hussain, Dhiya Al-Jumeily, Thar Baker, Robert Keight, Paulo Lisboa, Paul Fergus, and Ala S Al Kafri. A data science methodology based on machine learning algorithms for flood severity prediction. In *2018 IEEE Congress on Evolutionary Computation (CEC)*, pages 1–8. IEEE, 2018.

[9] Sana Haery, Alireza Mahpour, and Alireza Vafaeinejad. Forecasting urban travel demand with geo-AI: a combination of GIS and machine learning techniques utilizing uber data in New York City. *Environmental Earth Sciences*, 83(20):594, 2024.

[10] Tomas Mikolov, Kai Chen, Greg Corrado, and Jeffrey Dean. Efficient estimation of word representations in vector space. *arXiv preprint arXiv:1301.3781*, 2013.

[11] Jeffrey Pennington, Richard Socher, and Christopher D Manning. Glove: Global vectors for word representation. In *Proceedings of the 2014 conference on empirical methods in natural language processing (EMNLP)*, pages 1532–1543, 2014.

[12] Esther Rolf, Jonathan Proctor, Tamma Carleton, Ian Bolliger, Vaishaal Shankar, Miyabi Ishihara, Benjamin Recht, and Solomon Hsiang. A generalizable and accessible approach to machine learning with global satellite imagery. *Nature communications*, 12(1):4392, 2021.

[13] Konstantin Klemmer, Esther Rolf, Caleb Robinson, Lester Mackey, and Marc Rußwurm. SatClip: Global, general-purpose location embeddings with satellite imagery. *AAAI*, 2025.

[14] Mikolaj Czerkawski, Marcin Kluczek, and Jędrzej S. Bojanowski. Global and dense embeddings of Earth: Major TOM floating in the latent space. *arXiv preprint arXiv:2412.05600*, 2024.

[15] Vicente Vivanco Cepeda, Gaurav Kumar Nayak, and Mubarak Shah. GeoCLIP: Clip-inspired alignment between locations and images for effective worldwide geo-localization. *Advances in Neural Information Processing Systems*, 36:8690–8701, 2023.

[16] Mohit Agarwal, Mimi Sun, Chaitanya Kamath, Arbaaz Muslim, Prithul Sarker, Joydeep Paul, Hector Yee, Marcin Sieniek, Kim Jablonski, Yael Mayer, et al. General geospatial inference with a population dynamics foundation model. *arXiv preprint arXiv:2411.07207*, 2024.

[17] Ling Li, Yu Ye, Bingchuan Jiang, and Wei Zeng. Georeasoner: Geo-localization with reasoning in street views using a large vision-language model. In *Forty-first International Conference on Machine Learning*, 2024.

[18] Makkunda Sharma, Fan Yang, Duy-Nhat Vo, Esra Suel, Swapnil Mishra, Samir Bhatt, Oliver Fiala, William Rudgard, and Seth Flaxman. KidSat: satellite imagery to map childhood poverty dataset and benchmark. *arXiv preprint arXiv:2407.05986*, 2024.

[19] Darin Christensen, Tamma Carleton, Esther Rolf, Cullen Molitor, Shopnavo Biswas, Karena Yan, and Graeme Blair. Estimating the footprint of artisanal mining in Africa. Working Paper 33646, National Bureau of Economic Research, April 2025. URL http://www.nber.org/papers/w33646.

[20] Daniel Panangian and Ksenia Bittner. Can location embeddings enhance super-resolution of satellite imagery? *arXiv preprint arXiv:2501.15847*, 2025.

[21] Christopher F Brown, Michal R Kazmierski, Valerie J Pasquarella, William J Rucklidge, Masha Samsikova, Chenhui Zhang, Evan Shelhamer, Estefania Lahera, Olivia Wiles, Simon Ilyushchenko, Noel Gorelick, Lihui Lydia Zhang, Sophia Alj, Emily Schechter, Sean Askay, Oliver Guinan, Rebecca More, Alexis Boukouvalas, and Pushmeet Kohli. AlphaEarth Foundations: An embedding field model for accurate and efficient global mapping from sparse label data. *arXiv preprint arXiv:2507.22291*, 2025.

[22] Clay: A geospatial foundation model for Earth observation. https://madewithclay.org/ / https://github.com/Clay-foundation/model, 2025. Open-source model, Apache-2.0 license. Accessed: 2025-10-17.

[23] Daniela Szwarcman, Sujit Roy, Paolo Fraccaro, Þorsteinn Elí Gíslason, Benedikt Blumenstiel, Rinki Ghosal, Pedro Henrique de Oliveira, Joao Lucas de Sousa Almeida, Rocco Sedona, Yanghui Kang, et al. Prithvi-EO-2.0: A versatile multi-temporal foundation model for Earth observation applications. *arXiv preprint arXiv:2412.02732*, 2024.

[24] Gabriel Tseng, Anthony Fuller, Marlena Reil, Henry Herzog, Patrick Beukema, Favyen Bastani, James R Green, Evan Shelhamer, Hannah Kerner, and David Rolnick. Galileo: Learning global & local features of many remote sensing modalities. In *Forty-second International Conference on Machine Learning*, 2025.

[25] Gencer Sumbul, Chang Xu, Emanuele Dalsasso, and Devis Tuia. Smarties: Spectrum-aware multi-sensor auto-encoder for remote sensing images. *arXiv preprint arXiv:2506.19585*, 2025.

[26] Zhitong Xiong, Yi Wang, Fahong Zhang, Adam J Stewart, Joëlle Hanna, Damian Borth, Ioannis Papoutsis, Bertrand Le Saux, Gustau Camps-Valls, and Xiao Xiang Zhu. Neural plasticity-inspired foundation model for observing the Earth crossing modalities. *CoRR*, 2024.

[27] Xin Guo, Jiangwei Lao, Bo Dang, Yingying Zhang, Lei Yu, Lixiang Ru, Liheng Zhong, Ziyuan Huang, Kang Wu, Dingxiang Hu, Huimei He, Jian Wang, Jingdong Chen, Ming Yang, Yongjun Zhang, and Yansheng Li. Skysense: A multi-modal remote sensing foundation model towards universal interpretation for earth observation imagery. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*, pages 27672–27683, June 2024.

[28] Kang Wu, Yingying Zhang, Lixiang Ru, Bo Dang, Jiangwei Lao, Lei Yu, Junwei Luo, Zifan Zhu, Yue Sun, Jiahao Zhang, Qi Zhu, Jian Wang, Ming Yang, Jingdong Chen, Yongjun Zhang, and Yansheng Li. A semantic-enhanced multi-modal remote sensing foundation model

for earth observation. *Nature Machine Intelligence*, 2025. doi: 10.1038/s42256-025-01078-8. URL https://doi.org/10.1038/s42256-025-01078-8.

[29] Zhengpeng Feng, Sadiq Jaffer, Jovana Knezevic, Silja Sormunen, Robin Young, Madeline Lisaius, Markus Immitzer, James Ball, Clement Atzberger, David A Coomes, et al. TESSERA: Temporal embeddings of surface spectra for earth representation and analysis. *arXiv preprint arXiv:2506.20380*, 2025.

[30] Konstantin Klemmer and Daniel B Neill. Auxiliary-task learning for geographic data with autoregressive embeddings. In *Proceedings of the 29th International Conference on Advances in Geographic Information Systems*, pages 141–144, 2021.

[31] Elijah Cole, Grant Van Horn, Christian Lange, Alexander Shepard, Patrick Leary, Pietro Perona, Scott Loarie, and Oisin Mac Aodha. Spatial implicit neural representations for global-scale species mapping. In *International conference on machine learning*, pages 6320–6342. PMLR, 2023.

[32] Johannes Dollinger, Damien Robert, Elena Plekhanova, Lukas Drees, and Jan Dirk Wegner. Climplicit: Climatic implicit embeddings for global ecological tasks. *arXiv preprint arXiv:2504.05089*, 2025.

[33] Xiao Xiang Zhu, Zhitong Xiong, Yi Wang, Adam J Stewart, Konrad Heidler, Yuanyuan Wang, Zhenghang Yuan, Thomas Dujardin, Qingsong Xu, and Yilei Shi. On the foundations of Earth and climate foundation models. *arXiv preprint arXiv:2405.04285*, 2024.

[34] Carlos Gomes, Isabelle Wittmann, Damien Robert, Johannes Jakubik, Tim Reichelt, Stefano Maurogiovanni, Rikard Vinge, Jonas Hurst, Erik Scheurer, Rocco Sedona, et al. Lossy neural compression for geospatial analytics: A review. *IEEE Geoscience and Remote Sensing Magazine*, 2025.

[35] Langwen Huang and Torsten Hoefler. Compressing multidimensional weather and climate data into neural networks. In *The Eleventh International Conference on Learning Representations*, 2023.

[36] Thijs L Van der Plas, Jacob JW Bakermans, and Ioannis N Athanasiadis. NeurEO: dissecting Earth observation embeddings with computational neuroscience. In *EurIPS REO workshop*, 2025.

[37] Arjun Rao, Marc Rußwurm, Konstantin Klemmer, and Esther Rolf. Measuring the intrinsic dimension of Earth representations. *arXiv preprint arXiv:2511.02101*, 2025.

[38] Vishal Nedungadi, Ankit Kariryaa, Stefan Oehmcke, Serge Belongie, Christian Igel, and Nico Lang. MMEarth: Exploring multi-modal pretext tasks for geospatial representation learning. In *European Conference on Computer Vision*, pages 164–182. Springer, 2024.

[39] Srikumar Sastry, Subash Khanal, Aayush Dhakal, and Nathan Jacobs. GeoSynth: Contextually-aware high-resolution satellite image synthesis. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pages 460–470, 2024.

[40] Aayush Dhakal, Subash Khanal, Srikumar Sastry, Adeel Ahmad, and Nathan Jacobs. GeoBind: Binding text, image, and audio through satellite images. In *IGARSS 2024-2024*

IEEE International Geoscience and Remote Sensing Symposium, pages 2729–2733. IEEE, 2024.

[41] Nemin Wu, Qian Cao, Zhangyu Wang, Zeping Liu, Yanlin Qi, Jielu Zhang, Joshua Ni, Xiaobai Yao, Hongxu Ma, Lan Mu, Stefano Ermon, Tanuja Ganu, Akshay Nambi, Ni Lao, and Gengchen Mai. TorchSpatial: A location encoding framework and benchmark for spatial representation learning. In *NeurIPS 2024 Data and Benchamrk Track*, 2024.

[42] Takayuki Ishikawa, Carmelo Bonannella, Bas JW Lerink, and Marc Rußwurm. Deep pre-trained time series features for tree species classification in the Dutch forest inventory. *arXiv preprint arXiv:2508.18829*, 2025.

[43] Zhongliang Zhou, Jielu Zhang, Zihan Guan, Mengxuan Hu, Ni Lao, Lan Mu, Sheng Li, and Gengchen Mai. Img2Loc: Revisiting image geolocalization using multi-modality foundation models and image-based retrieval-augmented generation. In *Proceedings of the 47th International ACM SIGIR Conference on Research and Development in Information Retrieval*, pages 2749–2754, 2024.

[44] Oisin Mac Aodha, Elijah Cole, and Pietro Perona. Presence-only geographical priors for fine-grained image classification. In *Proceedings of the IEEE/CVF International Conference on Computer Vision*, pages 9596–9606, 2019.

[45] Gengchen Mai, Yao Xuan, Wenyun Zuo, Yutong He, Jiaming Song, Stefano Ermon, Krzysztof Janowicz, and Ni Lao. Sphere2Vec: A general-purpose location representation learning over a spherical surface for large-scale geospatial predictions. *ISPRS Journal of Photogrammetry and Remote Sensing*, 202:439–462, 2023.

[46] Samar Khanna, Patrick Liu, Linqi Zhou, Chenlin Meng, Robin Rombach, Marshall Burke, David Lobell, and Stefano Ermon. DiffusionSat: A generative foundation model for satellite imagery. In *The Twelfth International Conference on Learning Representations*, 2024.

[47] Arjun Rao and Esther Rolf. Using multiple input modalities can improve data-efficiency and ood generalization for ml with satellite imagery. *arXiv preprint arXiv:2507.13385*, 2025.

[48] Weibin Chen, Azhir Mahmood, Michel Tsamados, and So Takao. Deep random features for scalable interpolation of spatiotemporal data. In *The Thirteenth International Conference on Learning Representations*, 2024.

[49] Gengchen Mai, Krzysztof Janowicz, Ling Cai, Rui Zhu, Blake Regalia, Bo Yan, Meilin Shi, and Ni Lao. Se-kge: A location-aware knowledge graph embedding model for geographic question answering and spatial semantic lifting. *Transactions in GIS*, 24(3):623–655, 2020.

[50] Valérie Zermatten, Javiera Castillo-Navarro, Diego Marcos, and Devis Tuia. Learning transferable land cover semantics for open vocabulary interactions with remote sensing images. *ISPRS Journal of Photogrammetry and Remote Sensing*, 220:621–636, 2025. ISSN 0924-2716. doi: https://doi.org/10.1016/j.isprsjprs.2025.01.006.

[51] Utkarsh Mall, Cheng Perng Phoo, Meilin Kelsey Liu, Carl Vondrick, Bharath Hariharan, and Kavita Bala. Remote sensing vision-language foundation models without annotations via ground remote alignment. In *The Twelfth International Conference on Learning Representations*, 2024. URL https://openreview.net/forum?id=w9tc699w3Z.

[52] Aayush Dhakal, Adeel Ahmad, Subash Khanal, Srikumar Sastry, Hannah Kerner, and Nathan Jacobs. Sat2Cap: Mapping fine-grained textual descriptions from satellite images. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pages 533–542, 2024.

[53] Rikard Vinge, Isabelle Wittmann, Jannik Schneider, Michael Marszalek, Luis Gilch, Thomas Brunschwiler, and Conrad M Albrecht. Neuco-bench: A novel benchmark framework for neural embeddings in earth observation. *arXiv preprint arXiv:2510.17914*, 2025.

[54] Esther Rolf. Evaluation challenges for geospatial ML. *arXiv preprint arXiv:2303.18087*, 2023.

[55] Ribana Roscher, Bastian Bohn, Marco F Duarte, and Jochen Garcke. Explainable machine learning for scientific insights and discoveries. *IEEE Access*, 8:42200–42216, 2020.

[56] Cynthia Rudin. Stop explaining black box machine learning models for high stakes decisions and use interpretable models instead. *Nature machine intelligence*, 1(5):206–215, 2019.

[57] Alexandros Dimitrios Timoleon. What pattern makes a place unique? training neural networks to play geoguesser with satellite images. MSc thesis, Wageningen University and Research, Wageningen, The Netherlands, February 2026. Laboratory of Geo-Information Science and Remote Sensing. Report number GIRS-2026-11, thesis course code GRS-80436. Registration number 1566245. Supervisor: M. C. Rußwurm.

[58] Alison L Ritz and Jesse D Henderson. Using naip embeddings for a conus log yard map. *Journal of Forestry*, pages 1–19, 2026.

[59] W Hazeleger, JPM Aerts, Peter Bauer, MFP Bierkens, Gustau Camps-Valls, MM Dekker, FJ Doblas-Reyes, Veronika Eyring, C Finkenauer, Arthur Grundner, et al. Digital twins of the Earth with and for humans. *Communications earth & environment*, 5(1):463, 2024.

[60] Gencer Sumbul, Marcela Charfuelan, Begüm Demir, and Volker Markl. BigEarthNet: A large-scale benchmark archive for remote sensing image understanding. In *IGARSS 2019-2019 IEEE international geoscience and remote sensing symposium*, pages 5901–5904. IEEE, 2019.

[61] Yi Wang, Nassim Ait Ali Braham, Zhitong Xiong, Chenying Liu, Conrad M Albrecht, and Xiao Xiang Zhu. SSL4EO-S12: A large-scale multimodal, multitemporal dataset for self-supervised learning in Earth observation [software and data sets]. *IEEE Geoscience and Remote Sensing Magazine*, 11(3):98–106, 2023.

[62] Gengchen Mai, Krzysztof Janowicz, Yingjie Hu, Song Gao, Bo Yan, Rui Zhu, Ling Cai, and Ni Lao. A review of location encoding for GeoAI: methods and applications. *International Journal of Geographical Information Science*, 36(4):639–673, 2022.

[63] Vincent Sitzmann, Julien Martel, Alexander Bergman, David Lindell, and Gordon Wetzstein. Implicit neural representations with periodic activation functions. *Advances in neural information processing systems*, 33:7462–7473, 2020.

[64] Ben Mildenhall, Pratul P Srinivasan, Matthew Tancik, Jonathan T Barron, Ravi Ramamoorthi, and Ren Ng. Nerf: Representing scenes as neural radiance fields for view synthesis. *Communications of the ACM*, 65(1):99–106, 2021.

[65] Marc Rußwurm, Konstantin Klemmer, Esther Rolf, Robin Zbinden, and Devis Tuia. Geographic location encoding with spherical harmonics and sinusoidal representation networks. In *The Twelfth International Conference on Learning Representations*, 2024.

[66] Max Hamilton, Christian Lange, Elijah Cole, Alexander Shepard, Samuel Heinrich, Oisin Mac Aodha, Grant Van Horn, and Subhransu Maji. Combining observational data and language for species range estimation. *Advances in Neural Information Processing Systems*, 37:17719–17742, 2024.

[67] Gengchen Mai, Ni Lao, Yutong He, Jiaming Song, and Stefano Ermon. Csp: Self-supervised contrastive spatial pre-training for geospatial-visual representations. In *International Conference on Machine Learning*, pages 23498–23515. PMLR, 2023.

[68] Ramprasaath R Selvaraju, Michael Cogswell, Abhishek Das, Ramakrishna Vedantam, Devi Parikh, and Dhruv Batra. Grad-CAM: Visual explanations from deep networks via gradient-based localization. In *Proceedings of the IEEE international conference on computer vision*, pages 618–626, 2017.

[69] Guillaume Astruc, Nicolas Gonthier, Clement Mallet, and Loic Landrieu. AnySat: One Earth observation model for many resolutions, scales, and modalities. In *Proceedings of the Computer Vision and Pattern Recognition Conference*, pages 19530–19540, 2025.

[70] Jared Kaplan, Sam McCandlish, Tom Henighan, Tom B Brown, Benjamin Chess, Rewon Child, Scott Gray, Alec Radford, Jeffrey Wu, and Dario Amodei. Scaling laws for neural language models. *arXiv preprint arXiv:2001.08361*, 2020.

[71] Aayush Dhakal, Srikumar Sastry, Subash Khanal, Adeel Ahmad, Eric Xing, and Nathan Jacobs. RANGE: Retrieval augmented neural fields for multi-resolution geo-embeddings. In *Proceedings of the Computer Vision and Pattern Recognition Conference*, pages 24680–24689, 2025.

[72] Julien NP Martel, David B Lindell, Connor Z Lin, Eric R Chan, Marco Monteiro, and Gordon Wetzstein. Acorn: adaptive coordinate networks for neural scene representation. *ACM Transactions on Graphics (TOG)*, 40(4):1–13, 2021.

[73] Mirali Purohit, Gedeon Muhawenayo, Esther Rolf, and Hannah Kerner. How does the spatial distribution of pre-training data affect geospatial foundation models? *arXiv preprint arXiv:2501.12535*, 2025.

[74] Livia Betti, Farooq Sanni, Gnouyaro Sogoyou, Togbe Agbagla, Cullen Molitor, Tamma Carleton, and Esther Rolf. Mapping on a budget: Optimizing spatial data collection for ml. In *Proceedings of the AAAI Conference on Artificial Intelligence*, 2025.

[75] Nina Wiedemann, Théo Uscidda, and Martin Raubal. Geot: a spatially explicit framework for evaluating spatio-temporal predictions. *International Journal of Geographical Information Science*, pages 1–31, 2025.

[76] Konstantin Klemmer, Tianlin Xu, Beatrice Acciaio, and Daniel B Neill. Spate-gan: Improved generative modeling of dynamic spatio-temporal patterns with an autoregressive embedding loss. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 36, pages 4523–4531, 2022.

[77] Gabriel Tseng, Ruben Cartuyvels, Ivan Zvonkov, Mirali Purohit, David Rolnick, and Hannah Kerner. Lightweight, pre-trained transformers for remote sensing timeseries, 2023.

[78] Timnit Gebru, Jamie Morgenstern, Briana Vecchione, Jennifer Wortman Vaughan, Hanna Wallach, Hal Daumé Iii, and Kate Crawford. Datasheets for datasets. *Communications of the ACM*, 64(12):86–92, 2021.

[79] Margaret Mitchell, Simone Wu, Andrew Zaldivar, Parker Barnes, Lucy Vasserman, Ben Hutchinson, Elena Spitzer, Inioluwa Deborah Raji, and Timnit Gebru. Model cards for model reporting. In *Proceedings of the conference on fairness, accountability, and transparency*, pages 220–229, 2019.

[80] Hugging Face. Displaying carbon emissions for your model. https://huggingface.co/docs/hub/model-cards-co2, 2026. Accessed: 2026-06-05.

[81] Alexandre Lacoste, Alexandra Luccioni, Victor Schmidt, and Thomas Dandres. Quantifying the carbon emissions of machine learning. *arXiv preprint arXiv:1910.09700*, 2019.

A Earth Embedding Card Template

Embedding Overview. Include at a minimum:

1. **Who made the embeddings?**
   - **a.** Authors
   - **b.** Maintainers (including where to send questions about the embeddings)
   - **c.** Funders
   - **d.** Citation details, if applicable
2. **How are the embeddings accessed?**
   - **a. Are the embedding model, embedding datasets, or both made publicly available?** Provide links when possible
   - **b. What license are the embedding model and/or dataset available under?**
3. **How are the embeddings generated?**
   - **a. Resolution of embeddings:** For Earth embedding datasets, at what spatiotemporal resolution are the embeddings released? For embedding models, at what spatiotemporal resolution can the embeddings be generated?
   - **b. Data used:** What data modalities (including spatiotemporal indices) are used to generate the Earth embeddings?
     - i. How were the data accessed, what were their licenses, and what preprocessing was done?
     - ii. Are the embeddings an explicit transformation of these data, an implicit representation, or something else? Differentiate between data used during pretraining and used at generation-time.
4. **How are the embeddings intended to be used?** Consider types of prediction tasks and/or spatiotemporal resolutions for which the embeddings are intended to be used or for which they might not be suited.
   - **a. What type of tasks or use cases are the embeddings designed for?**
   - **b. Are there any tasks or use cases the embeddings should not be used for?**
   - **c. Are there any tasks or use cases that require extra validation before use?**

+

Include all relevant items from "Datasheets for Datasets" [78] and "Model Cards for Model Reporting" [79] frameworks. For evaluation across groups in the model cards framework, consider evaluation across geographic groups (regions, biomes, urban/rural groupings, etc.) as relevant to the Earth embeddings being described. For items from either framework that are not applicable to the Earth embeddings, skip them or put "not applicable".

+

Additional Earth embeddings information For some Earth embeddings, these questions may already be answered in responses to the boxes above.

5. **How are space and time encoded in the model?**
    a. **How are space or time incorporated into the Earth embeddings?** E.g., as direct inputs, metadata used for sampling during pretraining, etc.
    b. **Are there any implicit or explicit spatial or temporal assumptions made in the generation or use of the Earth embeddings?** For example, this might include smoothness constraints.
6. **What are the computational and environmental costs of generating, hosting and using the Earth embeddings?**
    a. **What were the computational costs associated with generating the embeddings?** Are there costs associated with storing the embeddings centrally, and if so, what are they and who pays those costs.
    b. **What were the estimated energy use of generating the embeddings?** See for example, Huggingface instructions for "Displaying carbon emissions for your model" [80] and Machine Learning Impact calculator presented in [81], for tracking, computing, and reporting carbon use.
    c. **What are the computational costs users incur to use the Earth embeddings?**