

The Coupling Cloud: A community database of megathrust kinematic coupling models

This is a non-peer reviewed preprint submitted to EarthArXiv

Authors:

Bar Oryan¹, Alice-Agnes Gabriel¹⁺², Roland Bürgmann³, Eric Calais⁴, Guo Cheng⁵, Mohamed Chlieh⁶, Beatriz Cosenza-Murales⁷, Víctor M. Cruz-Atienza⁸, Luca Dal Zilio⁹⁺¹⁰, Charles DeMets¹¹, Julie Elliott¹², Andria Ellis¹³, Lujia Feng⁹⁺¹⁰, Jeffrey T. Freymueller¹², Endra Gunawan¹⁴, Nuraini R. Hanifa¹⁵, George E. Hilley¹⁶, Ya-Ju Hsu¹⁷, Takeshi Iinuma¹⁸, Yuji Itoh¹⁹, Jorge Jara²⁰, Kaj M. Johnson²¹, Romain Jolivet⁴⁺²², Masayuki Kano²³, Emilie Klein⁴, Shanshan Li²⁴, Shaoyang Li²⁵, Eric O. Lindsey²⁶, Zhen Liu²⁷, John P. Loveless²⁸, Bertrand Løvery⁶, Louise Maubant¹⁹, Sylvain Michel²⁹⁺³⁰, Cyril Muller³¹, Marianne Métois³², Takuya Nishimura²³, Akemi Noda³³, Dibyashakti Panda³⁴, Mason Perry⁹, Raymundo Plata-Martinez⁸, Mathilde Radiguet⁶, Baptiste Rousset³⁵, Elizabeth M. Sherrill³⁶, Anne Socquet⁶, Juan Carlos Villegas-Lanza³⁷, Laura M. Wallace³⁶⁺³⁸, Lian Xue³⁹, Yusuke Yokota⁴⁰, Shoichi Yoshioka⁴¹⁺⁴² and Shui-Beih Yu¹⁷

Affiliations

- (1) Scripps Institution of Oceanography, University of California at San Diego, La Jolla, CA 92093, USA
- (2) Department of Earth and Environmental Sciences, Ludwig-Maximilians-University, 80333 Munich, Germany
- (3) Department of Earth and Planetary Science, UC Berkeley
- (4) Laboratoire de Géologie, Ecole normale supérieure - PSL, CNRS UMR 8538, Paris France
- (5) Nevada Bureau of Mines and Geology, University of Nevada, Reno, NV, USA
- (6) Université Grenoble Alpes, Université Savoie Mont Blanc, CNRS, IRD, Université Gustave Eiffel, STerre, Grenoble, France
- (7) Instituto de Investigación en Ciencias Físicas y Matemáticas (IFIM), Escuela de Ciencias Físicas y Matemáticas, Universidad de San Carlos de Guatemala
- (8) Instituto de Geofísica, Universidad Nacional Autónoma de México
- (9) Earth Observatory of Singapore, Nanyang Technological University, Singapore, Singapore
- (10) Asian School of the Environment, Nanyang Technological University, Singapore, Singapore
- (11) Department of Geoscience, University of Wisconsin-Madison, Madison WI 53706 USA
- (12) Department of Earth and Environmental Sciences, Michigan State University, East Lansing, MI, USA
- (13) USGS Hawaiian Volcano Observatory
- (14) Institut Teknologi Bandung, Indonesia
- (15) Research Center for Geological Disaster, National Research and Innovation Agency, Bandung, Indonesia
- (16) Stanford University, Stanford, CA, USA
- (17) Institute of Earth Sciences, Academia Sinica

- 45 (18) Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan
- 46 (19) Earthquake Research Institute, The University of Tokyo, Tokyo, Japan
- 47 (20) GFZ Helmholtz Centre for Geosciences, Potsdam, Germany
- 48 (21) Department of Earth and Atmospheric Sciences, Indiana University, Bloomington, IN, USA
- 49 (22) Institut Universitaire de France, 1 rue Descartes, 75006 Paris, France
- 50 (23) Disaster Prevention Research Institute, Kyoto University, Japan
- 51 (24) School of Computing at University of Wyoming
- 52 (25) State Key Laboratory of Lithospheric and Environmental Coevolution, Institute of Geology
- 53 and Geophysics, Chinese Academy of Sciences, Beijing, China
- 54 (26) Department of Earth & Planetary Sciences, University of New Mexico, Albuquerque, NM,
- 55 USA
- 56 (27) Jet Propulsion Laboratory, California Institute of Technology
- 57 (28) Department of Geosciences, Smith College, Northampton, MA, USA
- 58 (29) Université Côte d'Azur, IRD, CNRS, Observatoire de la Côte d'Azur, Valbonne, France
- 59 (30) Sorbonne Université, CNRS-INSU, Institut des Sciences de la Terre Paris, ISTeP UMR 7193,
- 60 Paris, France
- 61 (31) Observatorio Vulcanológico y Sismológico de Costa Rica, Universidad Nacional de Costa
- 62 Rica Heredia, Costa Rica
- 63 (32) Université Claude Bernard Lyon 1, ENS de Lyon, Université Jean Monnet, CNRS, LGL-TPE,
- 64 UMR5276
- 65 (33) Japan Meteorological Agency, Tokyo, Japan
- 66 (34) Department of Geology, School of Earth Sciences, Central University of Tamil Nadu,
- 67 Thiruvavur, India, 610005
- 68 (35) Institut Terre et Environnement de Strasbourg UMR7063, Université de
- 69 Strasbourg/CNRS/ENGESS, 5 Rue René Descartes, Strasbourg, 67000, France
- 70 (36) GEOMAR Helmholtz Centre for Ocean Research Kiel, Germany
- 71 (37) Instituto Geofísico del Perú (IGP), Lima, Peru
- 72 (38) Institute for Geophysics, University of Texas, Austin, TX, USA
- 73 (39) School of Earth and Space Science, Peking University, Beijing, China
- 74 (40) Institute of Industrial Science, University of Tokyo, Japan
- 75 (41) Research Center for Urban Safety and Security, Kobe University, Rokkodai-cho 1-1, Nada
- 76 ward, Kobe 657-8501, Japan
- 77 (42) Department of Planetology, Graduate School of Science, Kobe University, Rokkodai-cho 1-
- 78 1, Nada ward, Kobe 657-8501, Japan
- 79

80 Authors email

81 Bar Oryan:bar.oryan@columbia.edu
 82 Alice-Agnes Gabriel:algabriel@ucsd.edu
 83 Roland Bürgmann:burgmann@berkeley.edu
 84 Eric Calais:eric.calais@ens.fr
 85 Guo Cheng:gcheng@unr.edu
 86 Mohamed Chlieh:mohamed.chlieh@univ-grenoble-alpes.fr; mohamed.chlieh@ird.fr
 87 Beatriz Cosenza-Murales:bcosenza@ecfm.usac.edu.gt
 88 Victor M. Cruz-Atienza:cruz@igeofisica.unam.mx

89 Luca Dal Zilio:dalzilio.luca@gmail.com
 90 Charles DeMets:dcdemets@wisc.edu
 91 Julie Elliott:ellio372@msu.edu
 92 Andria Ellis:apellis@usgs.gov
 93 Lujia Feng:lfeng@ntu.edu.sg
 94 Jeffrey T. Freymueller:freymuel@msu.edu
 95 Endra Gunawan:endra.gunawan@itb.ac.id
 96 Nuraini R. Hanifa:nura010@brin.go.id
 97 George E. Hilley:hilley@stanford.edu
 98 Ya-Ju Hsu:yar@earth.sinica.edu.tw
 99 Takeshi Iinuma:iinuma@jamstec.go.jp
 100 Yuji Itoh:yitoh@eri.u-tokyo.ac.jp
 101 Jorge Jara:jorge@gfz.de
 102 Kaj M. Johnson:kajjohns@iu.edu
 103 Romain Jolivet:romain.jolivet@ens.fr
 104 Masayuki Kano:kano.masayuki.4m@kyoto-u.ac.jp
 105 Emilie Klein:klein@geologie.ens.fr
 106 Shanshan Li:shli11@outlook.com
 107 Shaoyang Li:shaoyangli@mail.iggcas.ac.cn
 108 Eric O. Lindsey:eol@unm.edu
 109 Zhen Liu:zhen.liu@jpl.nasa.gov
 110 John P. Loveless:jloveless@smith.edu
 111 Bertrand Loversy:bertrand.lovery@univ-grenoble-alpes.fr
 112 Louise Maubant:maubantl@mit.edu
 113 Sylvain Michel:sylvain_michel@live.fr
 114 Cyril Muller:cyril.muller@una.ac.cr
 115 Marianne Métois:marianne.metois@univ-lyon1.fr
 116 Takuya Nishimura:nishimura.takuya.4s@kyoto-u.ac.jp
 117 Akemi Noda:a.noda@met.kishou.go.jp
 118 Dibyashakti Panda:dibyashakti1@gmail.com
 119 Mason Perry:masonkyle.perry@ntu.edu.sg
 120 Raymundo Plata-Martinez:raymundo.plata@igeofisica.unam.mx
 121 Mathilde Radiguet:mathilde.radiguet@univ-grenoble-alpes.fr
 122 Baptiste Rousset:baptiste.rousset@unistra.fr
 123 Elizabeth M. Sherrill:esherrill@geomar.de
 124 Anne Socquet:anne.socquet@univ-grenoble-alpes.fr
 125 Juan Carlos Villegas-Lanza:jvillegas@igp.gob.pe
 126 Laura M. Wallace:llwallace@utexas.edu;llwallace@geomar.de
 127 Lian Xue:lian.xue@pku.edu.cn
 128 Yusuke Yokota:yyokota@iis.u-tokyo.ac.jp
 129 Shoichi Yoshioka:yoshioka@port.kobe-u.ac.jp
 130 Shui-Beih Yu:yusb@earth.sinica.edu.tw

131 Author ORCIDs

132 Bar Oryan:0000-0001-5000-5668

133 Alice-Agnes Gabriel:0000-0003-0112-8412
134 Roland Bürgmann:0000-0002-3560-044X
135 Eric Calais:0000-0002-5935-8117
136 Guo Cheng:0000-0002-4144-8448
137 Mohamed Chlieh:0000-0003-2252-2187
138 Beatriz Cosenza-Murales:0000-0002-4626-2757
139 Victor M. Cruz-Atienza:0000-0001-7067-2636
140 Luca Dal Zilio:0000-0002-5642-0894
141 Charles DeMets:0000-0001-7460-1165
142 Julie Elliott:0000-0001-6996-5706
143 Andria Ellis:0000-0003-2543-0640
144 Lujia Feng:0000-0002-3736-5025
145 Jeffrey T. Freymueller:0000-0003-0614-0306
146 Endra Gunawan:0000-0002-7187-1466
147 Nuraini R. Hanifa:0000-0002-8012-5385
148 George E. Hilley:0000-0002-1761-7547
149 Ya-Ju Hsu:0000-0003-1389-9994
150 Takeshi Iinuma:0000-0003-0386-2055
151 Yuji Itoh:0000-0002-7848-1399
152 Jorge Jara:0000-0003-3176-0689
153 Kaj M. Johnson:0000-0003-1511-5241
154 Romain Jolivet:0000-0002-9896-3651
155 Masayuki Kano:0000-0002-7288-4760
156 Emilie Klein:0000-0003-3239-5118
157 Shanshan Li:0000-0002-8986-431X
158 Shaoyang Li:0000-0001-9832-4561
159 Eric O. Lindsey:0000-0003-2274-8215
160 Zhen Liu:0000-0002-6313-823X
161 John P. Loveless:0000-0003-0416-8727
162 Bertrand Lavery:0000-0002-3671-0608
163 Louise Maubant:0000-0003-2077-169X
164 Sylvain Michel:0000-0001-7878-6603
165 Cyril Muller:0000-0003-3744-5700
166 Marianne Métois:0000-0002-1489-0513
167 Takuya Nishimura:0000-0002-2469-8146
168 Akemi Noda:0000-0002-9825-568X
169 Dibyashakti Panda:0000-0002-3190-9693
170 Mason Perry:0000-0003-1719-9004
171 Raymundo Plata-Martinez:0000-0001-9272-0926
172 Mathilde Radiguet:0000-0002-3877-9393
173 Baptiste Rousset:0000-0001-9304-0498
174 Elizabeth M. Sherrill:0000-0002-5439-6464
175 Anne Socquet:0000-0002-9208-7136
176 Juan Carlos Villegas-Lanza:0000-0002-2772-1508
177 Laura M. Wallace:0000-0003-2070-0891
178 Lian Xue:0000-0002-0739-7616

179 Yusuke Yokota:0000-0003-2969-9110
180 Shoichi Yoshioka:0000-0002-0238-1033
181 Shui-Beih Yu:0000-0002-1754-6208

182 Author contributions

183 Conceptualization: B. Oryan, A. Gabriel

184

185 Data Curation: B. Oryan, A. Gabriel, R. Bürgmann, E. Calais, G. Cheng, M. Chlieh, B. Cosenza-
186 Muralles, V. Cruz-Atienza, L. Dal Zilio, C. DeMets, J. Elliott, A. Ellis, L. Feng, J. Freymueller,
187 E. Gunawan, N. Hanifa, G. Hilley, Y. Hsu, T. Iinuma, Y. Itoh, J. Jara, K. Johnson, R. Jolivet, M.
188 Kano, E. Klein, S. Li, S. Li, E. Lindsey, Z. Liu, J. Loveless, B. Lavery, L. Maubant, S. Michel, C.
189 Muller, M. Métois, T. Nishimura, A. Noda, D. Panda, M. Perry, R. Plata-Martinez, M. Radiguet,
190 B. Rousset, E. Sherrill, A. Socquet, J. Villegas-Lanza, L. Wallace, L. Xue, Y. Yokota, S. Yoshioka,
191 S. Yu

192

193 Validation: B. Oryan, A. Gabriel, R. Bürgmann, E. Calais, G. Cheng, M. Chlieh, B. Cosenza-
194 Muralles, V. Cruz-Atienza, L. Dal Zilio, C. DeMets, J. Elliott, A. Ellis, L. Feng, J. Freymueller,
195 E. Gunawan, N. Hanifa, G. Hilley, Y. Hsu, T. Iinuma, Y. Itoh, J. Jara, K. Johnson, R. Jolivet, M.
196 Kano, E. Klein, S. Li, S. Li, E. Lindsey, Z. Liu, J. Loveless, B. Lavery, L. Maubant, S. Michel, C.
197 Muller, M. Métois, T. Nishimura, A. Noda, D. Panda, M. Perry, R. Plata-Martinez, M. Radiguet,
198 B. Rousset, E. Sherrill, A. Socquet, J. Villegas-Lanza, L. Wallace, L. Xue, Y. Yokota, S. Yoshioka,
199 S. Yu

200

201 Funding Acquisition: B. Oryan, A. Gabriel

202

203 Methodology: B. Oryan, A. Gabriel

204

205 Project Administration: B. Oryan, A. Gabriel

206

207 Software: B. Oryan

208

209 Supervision: A. Gabriel

210

211 Visualization: B. Oryan

212

213 Writing – original draft: B. Oryan, A. Gabriel

214

215 Writing – review & editing: B. Lavery, Cyril Muller, Roland Bürgmann, Eric Calais, Jeffrey T.
216 Freymueller, Marianne Métois, Víctor M. Cruz-Atienza, Takuya Nishimura, Emilie Klein, Charles
217 DeMets, Dibyashakti Panda, Louise Maubant, Shoichi Yoshioka, Jorge Jara, Mathilde Radiguet,
218 Sylvain Michel, Ya-Ju Hsu, Yuji Itoh, Lujia Feng, Shaoyang Li, Romain Jolivet, Jack Loveless, J.
219 Villegas-Lanza, Yusuke Yokota, Mason Perry, Baptiste Rousset, Zhen Liu, Beatriz Cosenza-
220 Muralles, Raymundo Plata-Martinez, N.R. Hanifa, Eric O. Lindsey, Elizabeth M. Sherrill

221

222

Abstract

Kinematic coupling models inverted from geodetic data are widely used to evaluate how slip deficit is distributed along subduction megathrusts during the interseismic period, and are central to earthquake and tsunami hazard assessment. Yet, existing coupling models differ widely in methodology and inputs, lack common community standards, and are scattered across publications and repositories. Here, we introduce the “Coupling Cloud” (<https://couplingcloud.ucsd.edu>), an open, extensible, community-driven platform that curates, standardizes, documents, and disseminates more than 95 kinematic coupling models from 56 publications across 21 subduction margins. The platform provides interactive 2D and 3D plate-interface viewers to inspect coupling models together with associated information such as slab geometry, uncertainty estimates and metadata. All datasets can be downloaded directly in standardized formats: surface-projected coupling values as NetCDF, plate-interface dislocation geometries as VTU, and model metadata as YAML files. We demonstrate the advantages of centralized and standardized coupling data through a Cascadia subduction zone example, where synthesizing eight full-margin models reveals along-strike patterns that are not apparent when models are examined individually. Consolidating coupling models within a coherent, version-controlled framework enables systematic cross-margin comparison and FAIR-compliant data sharing, opening the door to more comprehensive assessment of megathrust mechanics.

Introduction

Community databases can accelerate Earth science research by standardizing how data products are archived and shared, enabling synthesis across models and systematic evaluation of methodological choices. Several initiatives illustrate the scientific value of open, standardized data infrastructures. For example, SubMachine compiles global seismic tomography models (Hosseini et al., 2018), the SubMap initiative assembles long-term subduction zone observations (<https://submap.gm.umontpellier.fr/>), and the Nevada Geodetic Laboratory provides processed GNSS time series and velocity fields (Blewitt et al., 2018). Similarly, the SRCMOD database unified finite-fault source inversion formats and metadata to aggregate hundreds of finite slip models (Mai & Thingbaijam, 2014), directly motivating community benchmarks such as the Source Inversion Validation exercises (SIV, Mai et al., 2016). The USGS finite-fault model archive provides another example of how openly accessible, consistently formatted datasets accelerate scientific discovery (Goldberg et al., 2022; Hayes, 2017; <https://earthquake.usgs.gov/data/finitefault/>). In contrast, despite the central role of coupling models in subduction zone science and hazard assessment, the geodetic megathrust coupling community has lacked a centralized, standardized infrastructure for archiving, comparing, and reusing published models in consistent, interoperable data formats.

Kinematic coupling models identify regions along subduction interfaces that accommodate little interseismic slip, commonly inferred as frictionally locked, amid regions that creep steadily or slip transiently during the interseismic period. These models are constructed from inversions of interseismic or inter-slow slip events (inter-SSE) surface displacements most commonly derived from GNSS (Dixon, 1991; Feigl et al., 1993) and InSAR (Bürgmann et al., 2000), but also incorporating leveling (Burgette et al., 2009; Jackson & Bilham, 1994), coral-based reconstructions (Chlieh et al., 2008; Tsang et al., 2015) and offshore geodetic measurements such

as GNSS-Acoustic techniques (DeSanto et al., 2025; Gagnon et al., 2005; Yokota et al., 2016). Such inversions commonly use Green's functions from forward models based on elastic dislocation theory in a half-space (Meade, 2007; Okada, 1985, 1992), or viscoelastic earthquake-cycle formulations (Itoh et al., 2021; Li et al., 2015, 2018; Pollitz, 1997) and the backslip formulation to infer slip deficit (Savage, 1983). The inverse problem is then solved with linear schemes (Tarantola, 2005; Tarantola & Valette, 1982) or Bayesian approaches that integrate priors with data to quantify posterior uncertainty (e.g., Fukuda & Johnson, 2008; Minson et al., 2013; Tomita et al., 2021; Yabuki & Matsu'ura, 1992). Expressing the inferred slip deficit rate relative to the long-term plate convergence rate yields a kinematic coupling model (Wang & Dixon, 2004) that describes a continuous spectrum of fault behavior, ranging from areas creeping at the long-term slip rate (coupling ≈ 0) to effectively fixed regions with no creep (coupling ≈ 1).

Because kinematic coupling models delineate regions where elastic strain accumulates and may be released in future megathrust earthquakes, they are an important tool for assessing seismic and tsunami hazard in subduction zones (e.g., Giardini et al., 1999; Glehman et al., 2025; Ramos et al., 2021; Small & Melgar, 2021; Wang et al., 2015; Wang, 2011; Widiyantoro et al., 2020). However, despite their scientific value, these models face methodological challenges. Differences in assumed slab geometry and viscoelastic earth structure (Laske et al., 2013; Loversy et al., 2025; Luo et al., 2025; Wang et al., 2021), the period studied, density and quality of geodetic observations, velocity plate model, and the imposed regularization applied during coupling inversion affect the resulting coupling distributions (Fukuda & Johnson, 2008; Ide, 2007; Minson et al., 2013). Moreover, kinematic coupling models have traditionally lacked a unified, FAIR-compliant (Findable, Accessible, Interoperable, Reusable; Wilkinson et al., 2016) data format and are often scattered across publications in forms that are difficult to extract, evaluate, or systematically compare, limiting reproducibility and efforts to assess how modeling assumptions affect coupling estimates.

Here, we present the Coupling Cloud (<https://couplingcloud.ucsd.edu>, hereafter abbreviated to CC), an open online platform that aggregates a growing collection of published kinematic coupling models from subduction zones worldwide. The CC provides an intuitive interface to explore, visualize, and download these version controlled models, featuring a 2D map viewer for surface-projected coupling values and a 3D viewer that displays coupling on the underlying slab geometry. By centralizing and unifying these datasets and providing both 2D and 3D visualization tools within a single platform, the CC facilitates comparative analysis, strengthens reproducibility, and broadens community access to essential megathrust earthquake science resources. We demonstrate the value of this unified framework with a Cascadia case study, where along-strike patterns emerging from averaged models are not apparent when models are examined individually.

Data selection

As of January 12, 2026 the CC database comprises 96 kinematic coupling datasets compiled from 55 publications covering 21 subduction zones worldwide, for a total of 219 spatial data products. These include (in alphabetical order) two models in Alaska and the Aleutians (Drooff & Freymueller, 2021; Shanshan Li & Freymueller, 2018), two in the Caribbean (van Rijnsingen et al., 2021), eleven in Cascadia (Li et al., 2018; Lindsey et al., 2021; Materna et al., 2023; Michel et al., 2019; Pollitz & Evans, 2017; Pollitz, 2025; Schmalzle et al., 2014; Sherrill et al., 2024), eight in

Chile (Jara et al., 2024; Jolivet et al., 2020; Klein et al., 2018; Li et al., 2015; Métois et al., 2012, 2013, 2016), four in Costa Rica (Feng et al., 2012; Perry et al., 2025; Xue et al., 2015), two in Ecuador-Colombia (Chlieh et al., 2021; Gombert et al., 2018), one in Guatemala (Ellis et al., 2019), six in Hikurangi (Maubant et al., 2023; Michel et al., 2025; Wallace et al., 2012, three in Himalayas (Dal Zilio et al., 2020; Panda & Lindsey, 2024; Stevens & Avouac, 2015), one in Indo-Burma (Lindsey et al., 2023), nineteen in the Japan Trench (Abe & Yoshioka, 2022; Itoh et al., 2021; Lindsey et al., 2021; Loveless & Meade, 2016), two in Java (Hanifa et al., 2014; Widiyantoro et al., 2020), four in Kamchatka (Bürgmann et al., 2005), two in Manila (Hsu et al., 2016), two in Makran (Cheng et al., 2024), six in Mexico (Cosenza-Muralles et al., 2022; Cruz-Atienza et al., 2025; Louise Maubant et al., 2022; Radiguet et al., 2016; Rousset et al., 2016), nine in Nankai (Liu et al., 2010; Loveless & Meade, 2016; Nishimura et al., 2018; Noda et al., 2018; Plata-Martinez et al., 2024; Sherrill et al., 2024; Yokota et al., 2016), ten in Peru (Jara et al., 2024; Lavery et al., 2024; Villegas-Lanza et al., 2016), one in Ryukyu (Kano et al., 2021), three in the Sagami trough (Loveless & Meade, 2016; Nishimura et al., 2018), and one in Sumatra (Chlieh et al., 2008). While the Coupling Cloud provides standardized access to these datasets, users are expected to cite the original publications associated with each model when using the data.

Data standardization and unified representation

Published coupling models vary widely in format and metadata, reflecting the diversity of data coverage, methods, and conventions used across studies and time. To standardize access and promote FAIR use (Wilkinson et al., 2016), we map each model onto a uniform latitude–longitude grid and store it as a NetCDF file. (Rew & Davis, 1990). When gridded outputs are not provided, interpolation is applied using a scheme chosen to match the representations in the original publications. Every file contains latitude, longitude, the coupling values, and additional optional parameters, such as standard deviation, interface depth, and slip deficit. The NetCDF format is machine-readable and supports direct analyses and visualization using open-source packages such as Matplotlib (Hunter, 2007), GMT (Wessel et al., 2019), and Xarray (Hoyer & Hamman, 2017). Because NetCDF files do not inherently preserve the original fault-dislocation geometry used to invert the geodetic observations, we reformat all available model dislocation geometries as VTU PolyData files (Schroeder et al., 1998). VTU is a widely adopted, self-describing format for unstructured meshes that natively supports complex 3D slab geometries and mixed element types, making it well suited for downstream use in numerical modeling workflows. This standardized format allows for direct visualization and processing using open-source tools such as ParaView (Ahrens et al., 2005), PyVista (Sullivan & Kaszynski, 2019), and Meshio (Schlömer, 2022) and preserves node coordinates, mesh connectivity, and associated attributes such as coupling values on both node and element levels. Together, these machine-readable formats remove the need to parse ASCII text files and allow users to work directly with the models using the tools listed above. They allow for quantitative operations, e.g., clipping by depth, computing isosurfaces, resampling to regular grids, decimation, and provide efficient binary input/output (I/O) with compression for large models. Both the regular-grid NetCDF products and the unstructured VTU meshes integrate with standard parallel analysis and visualization tools used in high-performance computing (HPC) workflows, enabling future scalable processing of large coupling model ensembles.

Beyond the spatial formats, we collect metadata on each study’s modeling choices and assumptions to provide context for every hosted coupling model. Each dataset includes metadata

detailing the source publication DOI, allowing users to readily identify and cite the original studies, as well as a short description, region, and model type. The latter distinguishes between interseismic coupling models that represent time-averaged strain accumulation between earthquakes, inter-SSE models estimated during periods between slow-slip events, and kinematic block-model formulations that infer coupling from relative motions between tectonic blocks. Additional optional metadata includes the quantity and type of geodetic observations, the forward model formulation, and the inversion framework, including parameters such as the regularization type and associated weights (see Table 1 for a list of recommended metadata). These metadata are stored both within the NetCDF file and in an accompanying YAML file (Ben-Kiki et al., 2009), whose hierarchical structure separates observational, forward-model, and inversion information into human- and machine-readable records of model parameters. Lastly, when available, we include the original velocity fields and associated uncertainties in ASCII format as part of the model dataset. All CC data products, including NetCDF coupling values, VTU geometries, YAML metadata, and geodetic observations freely accessible through the CC interface as well as an example Jupyter notebook that demonstrates data access, metadata parsing, and visualization.

Platform architecture

The CC platform operates on a virtual machine within an eight-node Proxmox virtualization cluster hosted at the Scripps Institution of Oceanography, University of California San Diego. The system is configured with two virtual CPUs and 8 GB of RAM, running AlmaLinux 9.6, with Apache serving the web application and a FastAPI backend that handles all dynamic data requests (Fig. 1). The CC features two viewers: a 2D map viewer and a 3D geometry viewer. The 2D viewer displays surface-projected coupling values in a map interface and uses real-time server-side tiling via TiTiler, which converts NetCDF datasets into “tiles”, small image segments, at multiple zoom levels that form the full map. These tiles are added as data layers in Leaflet, an open-source JavaScript library that renders the interactive basemap and enables users to explore and compare coupling fields in geographic context.

The 3D viewer provides an interactive environment for exploring coupling fields on the original plate-interface geometry. For models where the slab depth is available, we construct 3D representations of the interface by converting geographic coordinates (latitude, longitude, and depth) to Earth-Centered Earth-Fixed coordinates and exporting the resulting surfaces as VTK PolyData (.vtp) files (Schroeder et al., 1998). The Coupling Cloud 3D viewer loads these files directly in a browser using [vtk.js](#) (Fig. 1), an open-source JavaScript library for scientific visualization. For each subduction zone, the 3D viewer includes an Earth surface layer derived from GEBCO (Becker et al., 2009), providing geographic context as users rotate, zoom, and explore coupling models, the megathrust interface and the overlying topography.

The Coupling Cloud platform

The CC is a browser-based workspace for finding, visualizing in 2D and 3D, comparing, and downloading published kinematic coupling models across subduction zones. Using the 2D viewer, users can select a margin from the dataset panel, display one or several models for that region, and compare their spatial distributions by toggling through layers, and adjusting the colorbar, allowing

for examination of differences in the extent of coupling (Fig. 2A). In addition to the coupling fields, the CC documents supplementary fields such as uncertainty, which can become substantial where geodetic constraints are sparse, particularly in offshore regions (e.g., DeSanto et al., 2025; Lindsey et al., 2021; Nishimura et al., 2018). If provided, users can visualize these uncertainty data sets alongside the coupling maps (Fig. 2B) or download them, allowing assessment of the reliability of specific features and propagation of uncertainty into their own analyses.

Kinematic coupling models assume plate-interface geometries that range from planar surfaces to detailed 3D geometries, often based on the Slab2 model (Hayes et al., 2018), which may lead to variations in the inferred slip deficit distributions (Baba et al., 2002; Elston et al., 2025; Moreno et al., 2009). Within the CC 2D viewer, users can visualize and compare these geometric choices (Fig. 3C). In addition to the 2D viewer, the CC includes a 3D viewer for visualizing coupling on the 3D plate-interface geometry. Users can activate the 3D viewer by selecting a subduction zone and entering its 3D view, where all models with available geometry can be explored (Fig. 3). This tool provides an intuitive way to examine how geometric assumptions influence inferred coupling, complementing the 2D surface view and enhancing comparison across models that use different representations of the plate interface. Users can download dislocation geometries as VTU files for offline visualization and analyses.

Variability among kinematic coupling models arises from differences in inversion setup and observational constraints (Avouac, 2015; Kaneko et al., 2010). These differences include the geodetic measurement tools used, ranging from GNSS to InSAR, leveling, paleogeodetic, and offshore measurements, as well as the time periods over which interseismic velocities are estimated and how aseismic processes such as postseismic deformation and interseismic transients are treated (Bürgmann et al., 2000; Chlieh et al., 2008; Dixon, 1991; Gagnon et al., 2005; Jackson & Bilham, 1994). The inversion strategies themselves span regularized least-squares formulations to fully Bayesian frameworks that explicitly probe the posterior distribution. Differences in regularization schemes and associated weights, fault parameterization, a priori constraints (e.g., imposed creep at depth or fixed rake), the adopted kinematic block models and plate reference frame further influence the inferred coupling distribution (Lovery et al., 2024; Métois et al., 2016; Villegas-Lanza et al., 2016). To document these differences, we compiled detailed metadata for each model, which can be viewed directly on the platform (Fig. 4) or downloaded as YAML files.

Performance

To assess how well the server handles real-time tiling and how many users it can support simultaneously, we benchmark the number of concurrent tile requests the backend can sustain. We issue 3,000 requests for the same tile at progressively increasing concurrency levels and measure the time required for TiTiler to serve all requests. Under our current configuration, which includes two virtual CPUs and 8 GB of RAM, the total processing time decreases with increasing concurrency as CPU resources are utilized more fully, until it reaches a plateau where the server is operating at maximum capacity with sustaining a throughput of approximately 45 tile requests per second (Fig. 5). A typical 2D map viewer user interaction such as panning or zooming triggers approximately 15 new tile requests. In practice, tile caching means that retrieved tiles are stored locally in the user's browser. Thus, interactions within already viewed areas impose no additional load on the backend, and the CC 2D viewer can support many more users than implied by the baseline throughput for uncached tiles.

Even in cases where the dynamic server becomes slow to respond due to high user load, our benchmark demonstrates that the architecture is scalable at least for small numbers of CPUs. A single-CPU configuration delivers roughly half the throughput of the two-CPU setup (Fig. 5), indicating near-linear performance scaling in this test. This suggests that adding additional CPU resources can increase system capacity, with the caveat that allocating more virtual CPUs incur higher purchasing and maintenance costs. We highlight that the 3D viewer requires no backend computation, as all visualization is handled directly in the user's browser. The only server-side operation is transferring the VTK PolyData geometry files to the client, which are typically a few megabytes in size and load within seconds, depending on the user's internet connection.

A simple demonstration: along-strike segmentation of the Cascadia megathrust in kinematic coupling models

To illustrate the advantages of assembling many published coupling models within a single standardized framework, we present a case study for the Cascadia subduction zone. The CC currently aggregates eight full-margin Cascadia interseismic coupling models that include plate-interface depth information (Fig. 6A). These models span more than a decade of geodetic inversion studies and employ a range of elastic, viscoelastic, and boundary-based forward and inverse modeling approaches, including regularized, parameterized slip-rate deficit inversions (Schmalzle et al., 2014), regularized linear inversions (Michel et al., 2019), regularized inversions with physical constraints (Lindsey et al., 2021), Bayesian boundary inversions (Sherrill et al., 2024), and viscoelastic earthquake-cycle modeling (Pollitz & Evans, 2017; Pollitz, 2025). As a brief example of the utility of the CC platform, we view these models together (Fig. 6A1-8) to assess how their inferred coupling distributions relate to proposed along-strike segmentation of the Cascadia margin.

Whether a megathrust is segmented along strike controls rupture behavior and whether slip deficit is released in single margin-wide earthquakes or in a sequence of smaller events (e.g., Arnulf et al., 2022; Bassett et al., 2025; Melnick et al., 2009; Philibosian & Meltzner, 2020). Cascadia is a classic example where this question remains central (e.g., Melgar, 2021). Slow-slip and tremor catalogs reveal along-strike sections whose recurrence patterns and spatial distributions change abruptly along strike, implying structural or frictional contrasts (Bartlow, 2020; Brudzinski & Allen, 2007). High-resolution seismic imaging reveals sharp changes in megathrust morphology that align with these inferred segment boundaries (Carbotte et al., 2024). Together with along-strike patterns in turbidite records, these observations have been interpreted as evidence for Cascadia's along-strike segmentation, a pattern also suggested by early geodetic coupling estimates, with higher coupling in the northern and southern segments and lower coupling in the central portion (Goldfinger et al., 2012; Schmalzle et al., 2014).

We define three commonly used along-strike subsections of the Cascadia margin (e.g., Ramos et al., 2021): a northern segment (Vancouver Island to $\sim 46^\circ\text{N}$), a central segment (~ 46 – 43.5°N), and a southern segment ($\sim 43.5^\circ\text{N}$ to Mendocino Triple Junction). For each model, we extract Coupling Cloud-stored gridded coupling values shallower than 20 km, where along-strike variations in coupling are most commonly inferred (Fig. 6A1–2). These values correspond to spatially distinct fault patches within each along-strike segment. We describe how coupling values are distributed

across these spatial units by computing a probability density function (PDF) for each model and segment (e.g., Oryan & Gabriel, 2025) using a Gaussian kernel density estimator with bandwidth selected according to Scott's rule (Scott, 2015). We note that the PDF is used as a normalized representation of the density and relative weighting of coupling values across the coupling spectrum and not as a result of repeated measurements of a stochastic process). Finally, we express each segment's PDF as a deviation from the margin-wide PDF for the same model, so that positive deviations highlight coupling values that are overrepresented within a given segment relative to the margin as a whole.

Our analysis shows little sensitivity to kernel bandwidth and grid resolution, and reveals that the northern-segment PDF curve rises by up to ~ 1 in probability density above the margin-wide PDF at coupling values larger than 0.7. This indicates that high-coupling values are more concentrated in the northern segment than across the Cascadia subduction zone as a whole. Similarly, the central segment PDF remains below the margin-wide PDF for coupling values larger than ~ 0.8 , indicating a lower density of high-coupling values relative to the margin-wide distribution (Fig. 6B2). In contrast, the southern segment does not show a higher concentration of high coupling values, and its PDF remains below the margin-wide distribution, reaching negative deviations of up to ~ 1 in probability density, similar in magnitude to the positive deviations observed in the northern segment (Fig. 6B3).

This example averages distributions across multiple coupling models spanning a range of inversion strategies and forward-modeling approaches. These models are not strictly independent, however, as they share geodetic datasets and related regularization strategies, and the ensemble spread should not be interpreted as a quantitative measure of uncertainty. Rather, the ensemble is intended to illustrate first-order behavior rather than provide a formal statistical treatment of inter-model variability and uncertainty. Even so, it shows that the concentration of coupling values in the northern and central segments aligns with previous interpretations of Cascadia's along-strike segmentation (Brudzinski & Allen, 2007; Carbotte et al., 2024; Goldfinger et al., 2012), whereas the southern segment does not exhibit a similarly coherent high-coupling pattern. This example also demonstrates how synthesizing ensemble models within the Coupling Cloud framework could help distinguish features that persist across modeling approaches from those that may depend on individual methodological choices. We therefore urge users to use the associated metadata and uncertainty information when comparing models and, most importantly, to consult the original publications describing each model, where the underlying assumptions and methodological choices are fully documented.

Current and future roles of the Coupling Cloud platform

The CC is designed with long-term sustainability in mind, hosted at Scripps / UC San Diego's institutional infrastructure with redundant backups and grounded in stable, widely supported data formats and libraries, ensuring continued accessibility and minimal maintenance. Our goal is for the CC to serve as a long-term community-driven global repository for kinematic coupling models, an up-to-date archive that the community can rely on for synthesis, comparison, and reproducibility. We also envision the CC to form a valuable teaching resource, offering standardized datasets for training the next generation of geodesists and subduction zone scientists, as well as providing an easy entry point for researchers new to the field. In addition, the platform

is extensible and supports user-driven uploads, allowing researchers to easily contribute new coupling models, metadata, and geometries directly. We invite all authors preparing new or forthcoming coupling models to upload their datasets to the CC so that the community can benefit from timely, open, and standardized access to the latest coupling models.

In addition, we hope that the unified file formats adopted here, NetCDF for surface-projected coupling fields, VTU (VTK PolyData) for plate-interface dislocation geometries, and YAML for FAIR-compliant metadata, will help providing a working standard for the community. By encouraging authors to express new coupling models directly in these formats, we aim to streamline data exchange and improve reproducibility across tools and research groups. At the same time, if the community converges on alternative standards, the CC framework can readily adopt them through simple import workflows, ensuring continued support for whichever formats best serve the megathrust research community.

Looking ahead, we see the Coupling Cloud not only as a repository of coupling models but as an evolving platform designed to reduce methodological variability and to bridge our understanding of how slip is accumulated and released in subduction zones (e.g., Wang et al., 2012). One promising direction is to host community validated Green's functions for subduction margins (e.g., Hori et al., 2021), providing a consistent set of forward-modeling kernels that removes a major source of variability among existing coupling models (Li et al., 2015; Loversy et al., 2025; Wang et al., 2021). A near-term direction for future developments is to link coupling models more directly to the geodetic observations used in the underlying inversions. Although users can already download the coupling-model associated observational datasets, if available, future versions of the CC may place greater emphasis on graphical representation, e.g., allowing GNSS velocities and time series, InSAR displacement fields, and offshore GNSS-Acoustic measurements to be viewed alongside coupling fields.

Finally, seamless integration of the CC interseismic coupling archive with the SRCMOD coseismic slip repository (Mai & Thingbaijam, 2014) and the Slow Earthquake Database (Kano et al., 2018) would unify complementary perspectives on megathrust behavior. Realizing this vision would require transforming the text-based FSP (Mai, Shearer, et al., 2016) and CSV formats used by SRCMOD and Slow Earthquake Databases into the VTU and NetCDF formats adopted by the Coupling Cloud, harmonizing metadata structures across both archives, and establishing an interface that links ruptures and slow slip events to corresponding interseismic coupling models. With such integration, users could examine coseismic and slow earthquake slip distributions alongside interseismic coupling fields, offering a unified perspective on strain accumulation and release across the earthquake cycle.

Conclusions

We have developed the Coupling Cloud, a unified and FAIR-compliant platform for standardizing, distributing, visualizing, and comparing kinematic geodetic coupling models across global subduction zones. The CC currently hosts 97 coupling models from 56 publications covering 21 subduction margins, each standardized into common file formats with preserved plate-interface geometries and complete metadata. By providing interactive 2D and 3D visualization tools and unified formats, the platform aims to remove long-standing barriers to reproducibility and to

572 enable more transparent comparison among models constructed with differing data sets and
573 methodologies, while associated metadata enables users to consult the original publications for
574 methodological details. As the Coupling Cloud grows through community contributions and future
575 integration of observational datasets and models, it is well positioned to serve as a long-lived
576 resource for advancing understanding of megathrust deformation and seismic hazard.

577

Parameter	Description
Model type	Interseismic/Inter-SSE/Block model or a combination of these.
Inversion type	Method of inversion used (e.g. Tikhonov regularization, Bayesian, stochastic Bayesian, least-squares).
Forward model formulation	Earth model used to compute surface displacement, such as a uniform elastic half-space, layered elastic structure, or a viscoelastic rheology.
Type of dislocation	Geometry of the dislocation elements (triangular, rectangular, or piecewise-linear triangular).
Number of dislocations	Total number of dislocation elements used in the forward model.
Degrees of freedom	Total number of parameters solved for in the inversion.
Type of Observations	Type(s) of geodetic observations used (e.g., Campaign GNSS, permanent GNSS, InSAR, leveling, coral etc.).
Observation Components	East/North/Up (or E/N/U)
Number of sites	Number of sites (or pixels in case of InSAR) used in the inversion.
Number of observations	Total number of individual observation components (e.g., if one site uses North/East components and another uses Up/Down, the total is 3).
Reference frame	Global geodetic reference frame (e.g., ITRF2014, ITRF2008).
Plate-fixed frame	Tectonic or plate-relative reference frame used to interpret interseismic deformation (e.g., Australian plate) including when possible.
Earliest data used	Start date of the geodetic observations incorporated in the inversion (format: YYYY or YYYY-MM or YYYY-MM-DD).
Latest data used	End date of the geodetic observations incorporated in the inversion (format: YYYY or YYYY-MM or YYYY-MM-DD).
Regularization type	Form of regularization applied during the inversion (e.g., Laplacian smoothing, Tikhonov regularization. etc.).
Smoothing weight	Value of the regularization weights used in the inversion.
Couple values computed	Indicates whether coupling was computed at dislocation nodes or over full dislocation patches.


Poisson ratio	Value of the Poisson's ratio assumed if a single layer model is used.
Number of layers	Number of layers used in the model, if applicable.
Convergence model	Relative plate velocity model (required for normalization of slip deficit to obtain coupling)
Other kinematic constraints	A priori assumptions about slip deficit rate (e.g., rake direction etc).

579

580 Table 1. Recommended metadata fields for coupling models.

581 Figure captions

582 **Figure 1 - CC platform architecture describing the server and user sides.**



583
584




Figure 2. Coupling Cloud 2D viewer overview. The 2D viewer provides an interactive map-based interface for exploring kinematic coupling models and their associated metadata. Users can select models, adjust visualization settings, and switch seamlessly between the 2D and 3D viewers. A. The upper-left panel shows the dataset selector, which allows users to choose coupling models to display in the 2D map viewer. The 3D button switches to the 3D viewer, with the red polygon

591 indicating the region covered by the 3D viewer. The lower-left panel contains the brush controls
592 used to configure colormap settings. The upper-right panel displays the metadata associated with
593 the selected coupling model, and a link to download the dataset. The “About” panel contains the
594 full reference list and additional platform information. Black dots indicate the position of GNSS
595 stations associated with time series in the Nevada Geodetic Laboratory (NGL). Examples showing
596 the selection of supplementary fields, including standard deviation and interface depth, for the
597 same coupling model in Hikurangi (Michel et al., 2025).
598




Figure 3 - Coupling Cloud 3D viewer overview. The 3D viewer enables interactive exploration of coupling models on the actual plate-interface geometry used in each inversion. It displays the megathrust surface in Earth-centered coordinates and overlays Earth-surface layer that provides geographic context. Users can select which geometric or coupling layers to display and customize their colormaps and opacity settings through the *Layers* and *Active Layers* control panels. The example shown here is from the Hikurangi subduction zone, illustrating the coupling field for the same model displayed in Fig 2 with the horizontal extent of the surface layer matching the red polygon shown in Fig. 2A.




Figure 4. Example of the metadata interface in the Coupling Cloud platform. The metadata panel provides a detailed hierarchical summary of each coupling model, including DOIs, global attributes, per-variable attributes, observational constraints, forward-model settings, and inversion parameters. Shown here is the complete metadata for the Hikurangi coupling model of Michel et al. (2025), organized into its main components: the reference, observations, forward model, inversion, per-variable metadata, and download.




Figure 5. Coupling Cloud backend performance benchmark showing response times for increasing levels of concurrent tile requests. Points show the total time required to process 3,000 HTTP tile requests under different CPU configurations and concurrency levels.




Figure 6. Cascadia case study. A - Eight Cascadia wide interseismic coupling models with slab depth available on the CC platform (Lindsey et al., 2021; Michel et al., 2019; Pollitz & Evans, 2017; Pollitz, 2025; Schmalzle et al., 2014; Sherrill et al., 2024), shown for interface depths shallower than 60 km. Contours indicate the megathrust interface depth used in each model. Horizontal dashed lines mark segments. B - Thin curves show the deviation of each model's segment-specific coupling probability density from that model's margin-wide probability density (Δ PDF). Positive Δ PDF values indicate coupling values that are more frequent within a given segment than across the margin. The shaded band denotes the 25th–75th percentile range of Δ PDF across the eight models.

Acknowledgements

We thank Kathryn Materna, Fred Pollitz, Eileen Evans, Jean-Philippe Avouac and Adriano Gualandi for sharing their coupling models. The Coupling Cloud server was built using TiTiler (<https://github.com/developmentseed/titiler>), FastAPI (<https://github.com/fastapi/fastapi>), Leaflet (<https://leafletjs.com/>) and vtk.js (<https://kitware.github.io/vtk-js/docs/>). B.O. and A.G. acknowledges support from the IGPP Green Foundation, National Science Foundation (grant no. OAC-2311208) and CRESCENT Award 2017Y0SD. A.G. further acknowledges support from Horizon Europe (ChEESE-2P, grant number 101093038 and Geo-INQUIRE, grant number 101058518) and the NSF (grant nos. OAC-2139536, EAR-2225286, EAR-2121568). E.C thanks the IGS, UNAVCO, Institut Universitaire de France and the FEDER European Community program within the Interreg Caraïbes “PREST” project. M.C. and C.D. acknowledge the Agence Nationale de la Recherche (ANR-07-BLAN-0143-01) and the IRD. B.C. and C.D. acknowledge NSF grants EAR-9526419, EAR-9804905, EAR-9909321, EAR-0510553, EAR-1114174, the University of Wisconsin-Madison and the UW-Madison Department of Geoscience Weeks endowment funds. L.D. and L.F. acknowledge Singapore Ministry of Education Tier 3b project “Investigating Volcano and Earthquake Science and Technology (InVEST)” (MOE-MOET32021-0002). A.E. acknowledges UNAVCO Graduate COCONet Fellowship and NSF grant EAR-1144418. J.T.F. acknowledges NSF grant EAR-1457361, and USGS grant G15AP00051. E.G. acknowledges the Ministry of Higher Education, Science and Technology of Indonesia (Kemendikisaintek). N.R.H. acknowledges Geospatial Information Agency of Indonesia, MEXT scholarship, Nagoya University, and Research and Innovation for Advanced Indonesia (RIIM) scheme. Y.H. acknowledges grant AS-102-SS-A09 and MOST 102-2116-M-001-028-MY3. T.I. acknowledges support from the Japan Society of the Promotion of Science (JSPS) Grants-in Aid for Scientific Research (KAKENHI) grants JP21H05200 and JP16H06472. J.J. acknowledges Chilean National Science Cooperation (CONICYT) through “Becas Chile” Program and MSCA Postdoctoral Fellowship (MSCA-101066069, project ERASMUS). R.J. acknowledges support from the Institut Universitaire de France and the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (Grant Agreement 758210 and 805256, Geo4D and iQuake projects) as well as NASA (Grant NNX16AK58G). E.K. acknowledges the French-Chilean LiA ‘Montessus de Ballore’, the Agence Nationale de la Recherche (ANR-2012-BS06-004) and the French INSU Tellus program, all data were collected by the instruments of the French mobile GNSS park GPSMob of Epos-France managed by CNRS Terre & Univers. S.M. acknowledges the ERC under the European Union's Horizon 2020 research and innovation program (Grant 758210 for project Geo4D). S.M. acknowledges NSF award EAR-1821853. M.M. acknowledges the French-Chilean LiA “Montessus de Ballore”, CNRS, MAE, IPGP, UJF and ANR project MEGA-Chile (project number 12-BS06-004). R.P. acknowledges DGAPA–UNAM PAPIIT project IA104525. E.M.S. thanks NSF grants EAR-2121631 and EAR-2045291 and USGS grant G20AP00017, and in part by Lilly Endowment, Inc., through its support for the Indiana University Pervasive Technology Institute. J.V-L. acknowledges the Agence Nationale de la Recherche (ANR-07-BLAN-0143-01) and IRD and the Instituto Geofísico del Perú (IGP). E.O.L. acknowledges support from NSF grant OCE-2314273 and NASA (grant 80NSSC23K0654). B.L. acknowledges support from the French national space agency (CNES). A.S. acknowledges support from the ERC Consolidator grant 865963 DEEP-trigger. Instrumentation and data from OVSICORI-UNA used in this research was supported by Costa Rican Emergency Law 8933 and by Universidad Nacional de Costa Rica through the project 0097–2020 “Sistema de monitoreo geodésico (SiMoGeod) de los volcanes y de la tectónica de Costa

Rica.” D.P. thanks the United States-India Educational Foundation (USIEF) for the financial support through the Fulbright-Nehru Postdoctoral Research Programme (Award 2783 FNPDR/2022). L.M. acknowledges support from NASA under Grant No.~80NSSC21K0874. S.Y. acknowledges support by JSPS KAKENHI grants JP15H01140, JP16H06477 and JP21H05203, and The Project for Hazard Assessment of Large Earthquakes and Tsunamis in the Mexican Pacific Coast for Disaster Mitigation, SATREPS funded by JST-JICA (#1554361). M.R. acknowledges support by the Agence Nationale de la Recherche (ANR21-CE49-0023) SSDYN project. Y.I. acknowledges JSPS KAKENHI grant JP17J00173 and Kyoto University Foundation. T.N. acknowledges JSPS KAKENHI grants JP21H05200, JP16H06472, JP15K21755 and JP26109007. Z. L. acknowledges support from NASA’s Earth Surface and Interior Focus Area. Part of the work was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration (80NM0018F0591). We acknowledge GNSS data from Alaska and Cascadia provided by the NSF-supported GAGE Facility (awards 1724794, 1724509).

Data and code availability

All coupling datasets are freely available to download from couplingcloud.ucsd.edu. The code to run the coupling cloud can be found in Oryan, 2025 (<https://zenodo.org/records/17821569>).

Competing interests

All authors declare no competing interests.

Manuscript References

- Abe, D., & Yoshioka, S. (2022). Spatiotemporal distributions of interplate coupling in Tohoku, northeast Japan, for 14 years prior to the 2011 Tohoku-oki earthquake inverted from GNSS data. *Tectonophysics*, 838, 229479. <https://doi.org/10.1016/j.tecto.2022.229479>
- Ahrens, J., Geveci, B., & Law, C. (2005). ParaView: An End-User Tool for Large-Data Visualization. In *Visualization Handbook* (pp. 717–731). Elsevier. <https://doi.org/10.1016/B978-012387582-2/50038-1>
- Arnulf, A. F., Bassett, D., Harding, A. J., Kodaira, S., Nakanishi, A., & Moore, G. (2022). Upper-plate controls on subduction zone geometry, hydration and earthquake behaviour. *Nature Geoscience*, 15(2), 143–148. <https://doi.org/10.1038/s41561-021-00879-x>
- Avouac, J.-P. (2015). From Geodetic Imaging of Seismic and Aseismic Fault Slip to Dynamic Modeling of the Seismic Cycle. *Annual Review of Earth and Planetary Sciences*, 43(Volume 43, 2015), 233–271. <https://doi.org/10.1146/annurev-earth-060614-105302>
- Baba, T., Tanioka, Y., Cummins, P. R., & Uhira, K. (2002). The slip distribution of the 1946 Nankai earthquake estimated from tsunami inversion using a new plate model. *Physics of the Earth and Planetary Interiors*, 132(1–3), 59–73.
- Bartlow, N. M. (2020). A Long-Term View of Episodic Tremor and Slip in Cascadia. *Geophysical Research Letters*, 47(3), e2019GL085303. <https://doi.org/10.1029/2019GL085303>
- Bassett, D., Shillington, D. J., Wallace, L. M., & Elliott, J. L. (2025). Variation in slip behaviour along megathrusts controlled by multiple physical properties. *Nature Geoscience*, 18(1), 20–31. <https://doi.org/10.1038/s41561-024-01617-9>

Becker, J. J., Sandwell, D. T., Smith, W. H. F., Braud, J., Binder, B., Depner, J., et al. (2009). Global Bathymetry and Elevation Data at 30 Arc Seconds Resolution: SRTM30_PLUS. *Marine Geodesy*, 32(4), 355–371. <https://doi.org/10.1080/01490410903297766>

Ben-Kiki, O., Evans, C., & Ingerson, B. (2009). Yaml ain't markup language (yamlTM) version 1.1. *Working Draft 2008*, 5(11).

Blewitt, B., Hammond, W. C., & Kreemer. (2018, September 24). Harnessing the GPS Data Explosion for Interdisciplinary Science. Retrieved October 20, 2025, from <https://eos.org/science-updates/harnessing-the-gps-data-explosion-for-interdisciplinary-science>

Brudzinski, M. R., & Allen, R. M. (2007). Segmentation in episodic tremor and slip all along Cascadia. *Geology*, 35(10), 907. <https://doi.org/10.1130/G23740A.1>

Burgette, R. J., Weldon, R. J., & Schmidt, D. A. (2009). Interseismic uplift rates for western Oregon and along-strike variation in locking on the Cascadia subduction zone: CASCADIA INTERSEISMIC UPLIFT AND LOCKING. *Journal of Geophysical Research: Solid Earth*, 114(B1). <https://doi.org/10.1029/2008JB005679>

Bürgmann, R., Rosen, P. A., & Fielding, E. J. (2000). Synthetic Aperture Radar Interferometry to Measure Earth's Surface Topography and Its Deformation. *Annual Review of Earth and Planetary Sciences*, 28(1), 169–209. <https://doi.org/10.1146/annurev.earth.28.1.169>

Bürgmann, R., Kogan, M. G., Steblov, G. M., Hilley, G., Levin, V. E., & Apel, E. (2005). Interseismic coupling and asperity distribution along the Kamchatka subduction zone.

Journal of Geophysical Research: Solid Earth, 110(B7).
<https://doi.org/10.1029/2005JB003648>

Carbotte, S. M., Boston, B., Han, S., Shuck, B., Beeson, J., Canales, J. P., et al. (2024). Subducting plate structure and megathrust morphology from deep seismic imaging linked to earthquake rupture segmentation at Cascadia. *Science Advances*, 10(23), eadl3198. <https://doi.org/10.1126/sciadv.adl3198>

Cheng, G., Barnhart, W. D., & Small, D. (2024). Constraints from GPS measurements on plate coupling within the Makran subduction zone and tsunami scenarios in the western Indian Ocean. *Geophysical Journal International*, 237(1), 288–301. <https://doi.org/10.1093/gji/ggae046>

Chlieh, M., Avouac, J. P., Sieh, K., Natawidjaja, D. H., & Galetzka, J. (2008). Heterogeneous coupling of the Sumatran megathrust constrained by geodetic and paleogeodetic measurements. *Journal of Geophysical Research: Solid Earth*, 113(B5). <https://doi.org/10.1029/2007JB004981>

Chlieh, M., Beauval, C., Yepes, H., Marinière, J., Saillard, M., & Audin, L. (2021). Seismic and Aseismic Cycle of the Ecuador–Colombia Subduction Zone. *Frontiers in Earth Science*, 9. <https://doi.org/10.3389/feart.2021.701720>

Cosenza-Murales, B., DeMets, C., Márquez-Azúa, B., Sánchez, O., Stock, J., Cabral-Cano, E., & McCaffrey, R. (2022). GPS-derived interseismic fault locking along the Jalisco–Colima segment of the Mexico subduction zone. *Geophysical Journal International*, 228(3), 2174–2197. <https://doi.org/10.1093/gji/ggab436>

Cruz-Atienza, V. M., Tago, J., Domínguez, L. A., Kostoglodov, V., Ito, Y., Ovando-Shelley, E., et al. (2025). Seafloor geodesy unveils seismogenesis of large subduction

earthquakes in Mexico. *Science Advances*, 11(37), eadu8259.
<https://doi.org/10.1126/sciadv.adu8259>

Dal Zilio, L., Jolivet, R., & van Dinther, Y. (2020). Segmentation of the Main Himalayan Thrust Illuminated by Bayesian Inference of Interseismic Coupling. *Geophysical Research Letters*, 47(4), e2019GL086424. <https://doi.org/10.1029/2019GL086424>

DeSanto, J. B., Schmidt, D. A., Zumberge, M., Sasagawa, G., & Chadwell, C. D. (2025). Near full locking on the shallow megathrust of the central Cascadia subduction zone revealed by GNSS-Acoustic. *Earth and Planetary Science Letters*, 665, 119463. <https://doi.org/10.1016/j.epsl.2025.119463>

Dixon, T. H. (1991). An introduction to the global positioning system and some geological applications. *Reviews of Geophysics*, 29(2), 249–276. <https://doi.org/10.1029/91RG00152>

Drooff, C., & Freymueller, J. T. (2021). New Constraints on Slip Deficit on the Aleutian Megathrust and Inflation at Mt. Veniaminof, Alaska From Repeat GPS Measurements. *Geophysical Research Letters*, 48(4), e2020GL091787. <https://doi.org/10.1029/2020GL091787>

Ellis, A., DeMets, C., McCaffrey, R., Briole, P., Cosenza Muralles, B., Flores, O., et al. (2019). GPS constraints on deformation in northern Central America from 1999 to 2017, Part 2: Block rotations and fault slip rates, fault locking and distributed deformation. *Geophysical Journal International*, 218(2), 729–754. <https://doi.org/10.1093/gji/ggz173>

Elston, H. M., Loveless, J. P., & Delph, J. R. (2025). Influence of Subduction Interface Geometry on Surface Displacements and Slip Processes in Cascadia. *Earth and Space Science*, 12(10), e2025EA004623. <https://doi.org/10.1029/2025EA004623>

Feigl, K. L., Agnew, D. C., Bock, Y., Dong, D., Donnellan, A., Hager, B. H., et al. (1993). Space geodetic measurement of crustal deformation in central and southern California, 1984–1992. *Journal of Geophysical Research: Solid Earth*, 98(B12), 21677–21712. <https://doi.org/10.1029/93JB02405>

Feng, L., Newman, A. V., Protti, M., González, V., Jiang, Y., & Dixon, T. H. (2012). Active deformation near the Nicoya Peninsula, northwestern Costa Rica, between 1996 and 2010: Interseismic megathrust coupling. *Journal of Geophysical Research: Solid Earth*, 117(B6). <https://doi.org/10.1029/2012JB009230>

Fukuda, J., & Johnson, K. M. (2008). A Fully Bayesian Inversion for Spatial Distribution of Fault Slip with Objective Smoothing. *Bulletin of the Seismological Society of America*, 98(3), 1128–1146. <https://doi.org/10.1785/0120070194>

Gagnon, K., Chadwell, C. D., & Norabuena, E. (2005). Measuring the onset of locking in the Peru–Chile trench with GPS and acoustic measurements. *Nature*, 434(7030), 205–208. <https://doi.org/10.1038/nature03412>

Giardini, D., Grünthal, G., Shedlock, K. M., & Zhang, P. (1999). The GSHAP Global Seismic Hazard Map. *Annals of Geophysics*, 42(6). <https://doi.org/10.4401/ag-3784>

Glehman, J., Gabriel, A., Ulrich, T., Ramos, M., Huang, Y., & Lindsey, E. (2025). Partial ruptures governed by the complex interplay between geodetic slip deficit, rigidity, and pore fluid pressure in 3D Cascadia dynamic rupture simulations. *Seismica*, 2(4). <https://doi.org/10.26443/seismica.v2i4.1427>

Goldberg, D. E., Koch, P., Melgar, D., Riquelme, S., & Yeck, W. L. (2022). Beyond the Teleseism: Introducing Regional Seismic and Geodetic Data into Routine USGS Finite-

Fault Modeling. *Seismological Research Letters*, 93(6), 3308–3323.
<https://doi.org/10.1785/0220220047>

Goldfinger, C., Nelson, C. H., Morey, A. E., Johnson, J. E., Patton, J. R., Karabanov, E. B., et al. (2012). *Turbidite event history—Methods and implications for Holocene paleoseismicity of the Cascadia subduction zone* (No. 1661- F). *Professional Paper*. U.S. Geological Survey. <https://doi.org/10.3133/pp1661F>

Gombert, B., Duputel, Z., Jolivet, R., Simons, M., Jiang, J., Liang, C., et al. (2018). Strain budget of the Ecuador–Colombia subduction zone: A stochastic view. *Earth and Planetary Science Letters*, 498, 288–299. <https://doi.org/10.1016/j.epsl.2018.06.046>

Hanifa, N. R., Sagiya, T., Kimata, F., Efendi, J., Abidin, H. Z., & Meilano, I. (2014). Interplate coupling model off the southwestern coast of Java, Indonesia, based on continuous GPS data in 2008–2010. *Earth and Planetary Science Letters*, 401, 159–171. <https://doi.org/10.1016/j.epsl.2014.06.010>

Hayes, G. P. (2017). The finite, kinematic rupture properties of great-sized earthquakes since 1990. *Earth and Planetary Science Letters*, 468, 94–100. <https://doi.org/10.1016/j.epsl.2017.04.003>

Hayes, G. P., Moore, G. L., Portner, D. E., Hearne, M., Flamme, H., Furtney, M., & Smoczyk, G. M. (2018). Slab2, a comprehensive subduction zone geometry model. *Science*, 362(6410), 58–61. <https://doi.org/10.1126/science.aat4723>

Hori, T., Agata, R., Ichimura, T., Fujita, K., Yamaguchi, T., & Iinuma, T. (2021). High-fidelity elastic Green’s functions for subduction zone models consistent with the global standard geodetic reference system. *Earth, Planets and Space*, 73(1), 41. <https://doi.org/10.1186/s40623-021-01370-y>

Hosseini, K., Matthews, K. J., Sigloch, K., Shephard, G. E., Domeier, M., & Tsekhmistrenko, M. (2018). SubMachine: Web-Based Tools for Exploring Seismic Tomography and Other Models of Earth's Deep Interior. *Geochemistry, Geophysics, Geosystems*, 19(5), 1464–1483. <https://doi.org/10.1029/2018GC007431>

Hoyer, S., & Hamman, J. (2017). xarray: N-D labeled Arrays and Datasets in Python. *Journal of Open Research Software*, 5(1), 10–10. <https://doi.org/10.5334/jors.148>

Hsu, Y.-J., Yu, S.-B., Loveless, J. P., Bacolcol, T., Solidum, R., Luis Jr, A., et al. (2016). Interseismic deformation and moment deficit along the Manila subduction zone and the Philippine Fault system. *Journal of Geophysical Research: Solid Earth*, 121(10), 7639–7665. <https://doi.org/10.1002/2016JB013082>

Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science & Engineering*, 9(03), 90–95.

Ide, S. (2007). 4.07 - Slip Inversion. In G. Schubert (Ed.), *Treatise on Geophysics* (pp. 193–223). Amsterdam: Elsevier. <https://doi.org/10.1016/B978-044452748-6.00068-7>

Itoh, Y., Nishimura, T., Wang, K., & He, J. (2021). New Megathrust Locking Model for the Southern Kurile Subduction Zone Incorporating Viscoelastic Relaxation and Non-Uniform Compliance of Upper Plate. *Journal of Geophysical Research: Solid Earth*, 126(5), e2020JB019981. <https://doi.org/10.1029/2020JB019981>

Jackson, M., & Bilham, R. (1994). Constraints on Himalayan deformation inferred from vertical velocity fields in Nepal and Tibet. *Journal of Geophysical Research: Solid Earth*, 99(B7), 13897–13912. <https://doi.org/10.1029/94JB00714>

Jara, J., Jolivet, R., Socquet, A., Comte, D., & Norabuena, E. (2024). Detection of slow slip events along the southern Peru - northern Chile subduction zone. *Seismica*, 3(1). <https://doi.org/10.26443/seismica.v3i1.980>

Jolivet, R., Simons, M., Duputel, Z., Olive, J., Bhat, H. S., & Bletery, Q. (2020). Interseismic loading of subduction megathrust drives long term uplift in northern Chile, 1–21. <https://doi.org/10.1029/2019GL085377>

Kaneko, Y., Avouac, J.-P., & Lapusta, N. (2010). Towards inferring earthquake patterns from geodetic observations of interseismic coupling. *Nature Geoscience*, 3(5), 363–369. <https://doi.org/10.1038/ngeo843>

Kano, M., Aso, N., Matsuzawa, T., Ide, S., Annoura, S., Arai, R., et al. (2018). Development of a Slow Earthquake Database. *Seismological Research Letters*, 89(4), 1566–1575. <https://doi.org/10.1785/0220180021>

Kano, M., Ikeuchi, A., Nishimura, T., Miyazaki, S., & Matsushima, T. (2021). Potential of megathrust earthquakes along the southern Ryukyu Trench inferred from GNSS data. *Earth, Planets and Space*, 73(1), 199. <https://doi.org/10.1186/s40623-021-01531-z>

Klein, E., Métois, M., Meneses, G., Vigny, C., & Delorme, A. (2018). Bridging the gap between North and Central Chile: insight from new GPS data on coupling complexities and the Andean sliver motion. *Geophysical Journal International*, 213(3), 1924–1933. <https://doi.org/10.1093/gji/ggy094>

Laske, G., Masters, G., Ma, Z., & Pasyanos, M. (2013). Update on CRUST1.0 - A 1-degree Global Model of Earth's Crust. In *Geophys. Res. Abstract.*

Li, Shanshan, & Freymueller, J. T. (2018). Spatial Variation of Slip Behavior Beneath the Alaska Peninsula Along Alaska-Aleutian Subduction Zone. *Geophysical Research Letters*, 45(8), 3453–3460. <https://doi.org/10.1002/2017GL076761>

Li, Shaoyang, Moreno, M., Bedford, J., Rosenau, M., & Oncken, O. (2015). Revisiting viscoelastic effects on interseismic deformation and locking degree: A case study of the Peru-North Chile subduction zone. *Journal of Geophysical Research: Solid Earth*, 120(6), 4522–4538. <https://doi.org/10.1002/2015JB011903>

Li, Shaoyang, Wang, K., Wang, Y., Jiang, Y., & Dosso, S. E. (2018). Geodetically Inferred Locking State of the Cascadia Megathrust Based on a Viscoelastic Earth Model. *Journal of Geophysical Research: Solid Earth*, 123(9), 8056–8072. <https://doi.org/10.1029/2018JB015620>

Lindsey, E. O., Mallick, R., Hubbard, J. A., Bradley, K. E., Almeida, R. V., Moore, J. D. P., et al. (2021). Slip rate deficit and earthquake potential on shallow megathrusts. *Nature Geoscience*, 14(5), 321–326. <https://doi.org/10.1038/s41561-021-00736-x>

Lindsey, E. O., Wang, Y., Aung, L. T., Chong, J.-H., Qiu, Q., Mallick, R., et al. (2023). Active subduction and strain partitioning in western Myanmar revealed by a dense survey GNSS network. *Earth and Planetary Science Letters*, 622, 118384. <https://doi.org/10.1016/j.epsl.2023.118384>

Liu, Z., Owen, S., Dong, D., Lundgren, P., Webb, F., Hetland, E., & Simons, M. (2010). Estimation of interplate coupling in the Nankai trough, Japan using GPS data from 1996 to 2006. *Geophysical Journal International*. <https://doi.org/10.1111/j.1365-246X.2010.04600.x>

Loveless, J. P., & Meade, B. J. (2016). Two decades of spatiotemporal variations in subduction zone coupling offshore Japan. *Earth and Planetary Science Letters*, 436, 19–30. <https://doi.org/10.1016/j.epsl.2015.12.033>

Loverly, B., Chlieh, M., Norabuena, E., Villegas-Lanza, J. C., Radiguet, M., Cotte, N., et al. (2024). Heterogeneous Locking and Earthquake Potential on the South Peru Megathrust From Dense GNSS Network. *Journal of Geophysical Research: Solid Earth*, 129(2), e2023JB027114. <https://doi.org/10.1029/2023JB027114>

Loverly, B., Radiguet, M., Chlieh, M., Norabuena, E., Villegas-Lanza, J. C., Cresseaux, J., et al. (2025). Viscoelastic Relaxation Following the 2001 Mw 8.4 Arequipa Earthquake and Its Impact on the Interseismic Coupling of the South Peru Megathrust. *Geophysical Research Letters*, 52(12), e2024GL113879. <https://doi.org/10.1029/2024GL113879>

Luo, H., Wang, K., Feng, L., & Hill, E. M. (2025). Interseismic secondary zone of subsidence during earthquake cycles in subduction zones. *Nature Geoscience*, 18(10), 1027–1033. <https://doi.org/10.1038/s41561-025-01778-1>

Mai, Shearer, P., Ampuero, J., & Lay, T. (2016). Standards for Documenting Finite-Fault Earthquake Rupture Models. *Seismological Research Letters*, 87(3), 712–718. <https://doi.org/10.1785/0220150204>

Mai, P. M., & Thingbaijam, K. K. S. (2014). SRCMOD: An Online Database of Finite-Fault Rupture Models. *Seismological Research Letters*, 85(6), 1348–1357. <https://doi.org/10.1785/0220140077>

Mai, P. M., Schorlemmer, D., Page, M., Ampuero, J., Asano, K., Causse, M., et al. (2016). The Earthquake-Source Inversion Validation (SIV) Project. *Seismological Research Letters*, 87(3), 690–708. <https://doi.org/10.1785/0220150231>

Materna, K., Murray, J. R., Pollitz, F., & Patton, J. R. (2023). Slip Deficit Rates on Southern Cascadia Faults Resolved with Viscoelastic Earthquake Cycle Modeling of Geodetic Deformation. *Bulletin of the Seismological Society of America*, 113(6), 2505–2518. <https://doi.org/10.1785/0120230007>

Maubant, L., Frank, W. B., Wallace, L. M., Williams, C. A., & Hamling, I. (2023). Imaging the Spatiotemporal Evolution of Plate Coupling With Interferometric Radar (InSAR) in the Hikurangi Subduction Zone. *Geophysical Research Letters*, 50(19), e2023GL105388. <https://doi.org/10.1029/2023GL105388>

Maubant, Louise, Radiguet, M., Pathier, E., Doin, M.-P., Cotte, N., Kazachkina, E., & Kostoglodov, V. (2022). Interseismic coupling along the Mexican subduction zone seen by InSAR and GNSS. *Earth and Planetary Science Letters*, 586, 117534. <https://doi.org/10.1016/j.epsl.2022.117534>

Meade, B. J. (2007). Algorithms for the calculation of exact displacements, strains, and stresses for triangular dislocation elements in a uniform elastic half space. *Computers & Geosciences*, 33(8), 1064–1075. <https://doi.org/10.1016/j.cageo.2006.12.003>

Melgar, D. (2021). Was the January 26th, 1700 Cascadia Earthquake Part of a Rupture Sequence? *Journal of Geophysical Research: Solid Earth*, 126(10), e2021JB021822. <https://doi.org/10.1029/2021JB021822>

Melnick, D., Bookhagen, B., Strecker, M. R., & Echtler, H. P. (2009). Segmentation of megathrust rupture zones from fore-arc deformation patterns over hundreds to millions of years, Arauco peninsula, Chile. *Journal of Geophysical Research: Solid Earth*, 114(B1). <https://doi.org/10.1029/2008JB005788>

Métois, M., Socquet, A., & Vigny, C. (2012). Interseismic coupling, segmentation and mechanical behavior of the central Chile subduction zone. *Journal of Geophysical Research: Solid Earth*, 117(B3). <https://doi.org/10.1029/2011JB008736>

Métois, M., Socquet, A., Vigny, C., Carrizo, D., Peyrat, S., Delorme, A., et al. (2013). Revisiting the North Chile seismic gap segmentation using GPS-derived interseismic coupling. *Geophysical Journal International*, 194(3), 1283–1294. <https://doi.org/10.1093/gji/ggt183>

Métois, M., Vigny, C., & Socquet, A. (2016). Interseismic Coupling, Megathrust Earthquakes and Seismic Swarms Along the Chilean Subduction Zone (38°–18°S). *Pure and Applied Geophysics*, 173(5), 1431–1449. <https://doi.org/10.1007/s00024-016-1280-5>

Michel, S., Gualandi, A., & Avouac, J.-P. (2019). Interseismic Coupling and Slow Slip Events on the Cascadia Megathrust. *Pure and Applied Geophysics*, 176(9), 3867–3891. <https://doi.org/10.1007/s00024-018-1991-x>

Michel, S., Jolivet, R., Klein, E., & Maubant, L. (2025). 14 Years of Slip on the Hikurangi Subduction Zone. *Journal of Geophysical Research: Solid Earth*, 130(7), e2024JB030865. <https://doi.org/10.1029/2024JB030865>

Minson, S. E., Simons, M., & Beck, J. L. (2013). Bayesian inversion for finite fault earthquake source models I—theory and algorithm. *Geophysical Journal International*, 194(3), 1701–1726. <https://doi.org/10.1093/gji/ggt180>

Moreno, M. S., Bolte, J., Klotz, J., & Melnick, D. (2009). Impact of megathrust geometry on inversion of coseismic slip from geodetic data: Application to the 1960 Chile earthquake. *Geophysical Research Letters*, 36(16). <https://doi.org/10.1029/2009GL039276>

Nishimura, T., Yokota, Y., Tadokoro, K., & Ochi, T. (2018). Strain partitioning and interplate coupling along the northern margin of the Philippine Sea plate, estimated from Global Navigation Satellite System and Global Positioning System-Acoustic data. *Geosphere*, 14(2), 535–551. <https://doi.org/10.1130/GES01529.1>

Noda, A., Saito, T., & Fukuyama, E. (2018). Slip-Deficit Rate Distribution Along the Nankai Trough, Southwest Japan, With Elastic Lithosphere and Viscoelastic Asthenosphere. *Journal of Geophysical Research: Solid Earth*, 123(9), 8125–8142. <https://doi.org/10.1029/2018JB015515>

Okada, Y. (1985). Surface deformation due to shear and tensile faults in a half-space. *International Journal of Rock Mechanics and Mining Sciences & Geomechanics Abstracts*, 23(4), 128. [https://doi.org/10.1016/0148-9062\(86\)90674-1](https://doi.org/10.1016/0148-9062(86)90674-1)

Okada, Y. (1992). Internal deformation due to shear and tensile faults in a half-space. *Bulletin - Seismological Society of America*.

Oryan, B. (2025, December 4). Code for the coupling cloud. Zenodo. Retrieved from <https://zenodo.org/records/17821569>

Panda, D., & Lindsey, E. O. (2024). Overriding Plate Deformation Controls Inferences of Interseismic Coupling Along the Himalayan Megathrust. *Journal of Geophysical Research: Solid Earth*, 129(9), e2024JB029819. <https://doi.org/10.1029/2024JB029819>

Perry, M., Muller, C., Protti, M., Feng, L., & Hill, E. M. (2025). Interseismic Megathrust Coupling at the Osa Peninsula, Costa Rica. *Journal of Geophysical Research: Solid Earth*, 130(7), e2024JB030641. <https://doi.org/10.1029/2024JB030641>

Philibosian, B., & Meltzner, A. J. (2020). Segmentation and supercycles: A catalog of earthquake rupture patterns from the Sumatran Sunda Megathrust and other well-studied

faults worldwide. *Quaternary Science Reviews*, 241, 106390.
<https://doi.org/10.1016/j.quascirev.2020.106390>
 Plata-Martinez, R., Iinuma, T., Tomita, F., Nakamura, Y., Nishimura, T., & Hori, T.
 (2024). Revisiting Slip Deficit Rates and Its Insights Into Large and Slow Earthquakes at
 the Nankai Subduction Zone. *Journal of Geophysical Research: Solid Earth*, 129(12),
 e2023JB027942. <https://doi.org/10.1029/2023JB027942>
 Pollitz, F.F., & Evans, E. L. (2017). Implications of the earthquake cycle for inferring fault
 locking on the Cascadia megathrust. *Geophysical Journal International*, 209(1), 167–185.
<https://doi.org/10.1093/gji/ggx009>
 Pollitz, Fred F. (1997). Gravitational viscoelastic postseismic relaxation on a layered
 spherical Earth. *Journal of Geophysical Research: Solid Earth*, 102(B8), 17921–17941.
<https://doi.org/10.1029/97JB01277>
 Pollitz, Fred F. (2025). 3D Viscoelastic Models of Slip-Deficit Rate Along the Cascadia
 Subduction Zone. *Journal of Geophysical Research: Solid Earth*, 130(1), e2024JB029847.
<https://doi.org/10.1029/2024JB029847>
 Radiguet, M., Perfettini, H., Cotte, N., Gualandi, A., Valette, B., Kostoglodov, V., et al.
 (2016). Triggering of the 2014 Mw7.3 Papanoa earthquake by a slow slip event in
 Guerrero, Mexico. *Nature Geoscience*, 9(11), 829–833. <https://doi.org/10.1038/ngeo2817>
 Ramos, M. D., Huang, Y., Ulrich, T., Li, D., Gabriel, A.-A., & Thomas, A. M. (2021).
 Assessing Margin-Wide Rupture Behaviors Along the Cascadia Megathrust With 3-D
 Dynamic Rupture Simulations. *Journal of Geophysical Research: Solid Earth*, 126(7),
 e2021JB022005. <https://doi.org/10.1029/2021JB022005>

Rew, R., & Davis, G. (1990). NetCDF: an interface for scientific data access. *IEEE Computer Graphics and Applications*, 10(4), 76–82. <https://doi.org/10.1109/38.56302>

van Rijnsingen, E. M., Calais, E., Jolivet, R., Robertson, R., & Ryan, G. A. (2021). Inferring Interseismic Coupling Along the Lesser Antilles Arc: A Bayesian Approach. *Journal of Geophysical Research*, 21.

Rousset, B., Lasserre, C., Cubas, N., Graham, S., Radiguet, M., DeMets, C., et al. (2016). Lateral Variations of Interplate Coupling along the Mexican Subduction Interface: Relationships with Long-Term Morphology and Fault Zone Mechanical Properties. *Pure and Applied Geophysics*, 173(10), 3467–3486. <https://doi.org/10.1007/s00024-015-1215-6>

Savage, J. C. (1983). A dislocation model of strain accumulation and release at a subduction zone. *Journal of Geophysical Research*. <https://doi.org/10.1029/JB088iB06p04984>

Schlömer, N. (2022). meshio: Tools for mesh files. *GitHub Repository*.

Schmalzle, G. M., McCaffrey, R., & Creager, K. C. (2014). Central Cascadia subduction zone creep. *Geochemistry, Geophysics, Geosystems*, 15(4), 1515–1532. <https://doi.org/10.1002/2013GC005172>

Schroeder, W., Martin, K. M., & Lorensen, W. E. (1998). *The visualization toolkit (2nd ed.): an object-oriented approach to 3D graphics*. USA: Prentice-Hall, Inc.

Sherrill, E. M., Johnson, K. M., & Jackson, N. M. (2024). Locating Boundaries Between Locked and Creeping Regions at Nankai and Cascadia Subduction Zones. *Journal of Geophysical Research: Solid Earth*, 129(10), e2024JB029346. <https://doi.org/10.1029/2024JB029346>

Small, D. T., & Melgar, D. (2021). Geodetic Coupling Models as Constraints on Stochastic Earthquake Ruptures: An Example Application to PTHA in Cascadia. *Journal of Geophysical Research: Solid Earth*, 126(7), e2020JB021149. <https://doi.org/10.1029/2020JB021149>

Stevens, V. L., & Avouac, J. P. (2015). Interseismic coupling on the main Himalayan thrust. *Geophysical Research Letters*, 42(14), 5828–5837. <https://doi.org/10.1002/2015GL064845>

Sullivan, C., & Kaszynski, A. (2019). PyVista: 3D plotting and mesh analysis through a streamlined interface for the Visualization Toolkit (VTK). *Journal of Open Source Software*, 4(37), 1450. <https://doi.org/10.21105/joss.01450>

Tarantola, A. (2005). *Inverse Problem Theory and Methods for Model Parameter Estimation*. Society for Industrial and Applied Mathematics. <https://doi.org/10.1137/1.9780898717921>

Tarantola, A., & Valette, B. (1982). Generalized nonlinear inverse problems solved using the least squares criterion. *Reviews of Geophysics*, 20(2), 219–232. <https://doi.org/10.1029/RG020i002p00219>

Tomita, F., Iinuma, T., Agata, R., & Hori, T. (2021). Development of a Trans-Dimensional Fault Slip Inversion for Geodetic Data. *Journal of Geophysical Research: Solid Earth*, 126(5), e2020JB020991. <https://doi.org/10.1029/2020JB020991>

Tsang, L. L. H., Meltzner, A. J., Hill, E. M., Freymueller, J. T., & Sieh, K. (2015). A paleogeodetic record of variable interseismic rates and megathrust coupling at Simeulue Island, Sumatra. *Geophysical Research Letters*, 42(24), 10,585–10,594. <https://doi.org/10.1002/2015GL066366>

1061 Villegas-Lanza, J. C., Chlieh, M., Cavalié, O., Tavera, H., Baby, P., Chire-Chira, J., &
 1062 Nocquet, J.-M. (2016). Active tectonics of Peru: Heterogeneous interseismic coupling
 1063 along the Nazca megathrust, rigid motion of the Peruvian Sliver, and Subandean shortening
 1064 accommodation. *Journal of Geophysical Research: Solid Earth*, 121(10), 7371–7394.
 1065 <https://doi.org/10.1002/2016JB013080>
 1066 Wallace, L. M., Barnes, P., Beavan, J., Van Dissen, R., Litchfield, N., Mountjoy, J., et al.
 1067 (2012). The kinematics of a transition from subduction to strike-slip: An example from the
 1068 central New Zealand plate boundary. *Journal of Geophysical Research: Solid Earth*,
 1069 117(B2). <https://doi.org/10.1029/2011JB008640>
 1070 Wang, K., & Dixon, T. (2004). “Coupling” Semantics and science in earthquake research.
 1071 *Eos, Transactions American Geophysical Union*, 85(18), 180–180.
 1072 <https://doi.org/10.1029/2004EO180005>
 1073 Wang, K., Hu, Y., & He, J. (2012). Deformation cycles of subduction earthquakes in a
 1074 viscoelastic Earth. *Nature*, 484(7394), 327–332. <https://doi.org/10.1038/nature11032>
 1075 Wang, K., Zhu, Y., Nissen, E., & Shen, Z.-K. (2021). On the Relevance of Geodetic
 1076 Deformation Rates to Earthquake Potential. *Geophysical Research Letters*, 48(11),
 1077 e2021GL093231. <https://doi.org/10.1029/2021GL093231>
 1078 Wang, L., Hainzl, S., & Mai, P. M. (2015). Quantifying slip balance in the earthquake
 1079 cycle: Coseismic slip model constrained by interseismic coupling. *Journal of Geophysical*
 1080 *Research: Solid Earth*, 120(12), 8383–8403. <https://doi.org/10.1002/2015JB011987>
 1081 Wang, Z. (2011). Seismic Hazard Assessment: Issues and Alternatives. *Pure and Applied*
 1082 *Geophysics*, 168(1), 11–25. <https://doi.org/10.1007/s00024-010-0148-3>

Wessel, P., Luis, J. F., Uieda, L., Scharroo, R., Wobbe, F., Smith, W. H. F., & Tian, D. (2019). The Generic Mapping Tools Version 6. *Geochemistry, Geophysics, Geosystems*, 20(11), 5556–5564. <https://doi.org/10.1029/2019GC008515>

Widiyantoro, S., Gunawan, E., Muhari, A., Rawlinson, N., Mori, J., Hanifa, N. R., et al. (2020). Implications for megathrust earthquakes and tsunamis from seismic gaps south of Java Indonesia. *Scientific Reports*, 10(1), 15274. <https://doi.org/10.1038/s41598-020-72142-z>

Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A., et al. (2016). The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data*, 3(1), 160018. <https://doi.org/10.1038/sdata.2016.18>

Xue, L., Schwartz, S., Liu, Z., & Feng, L. (2015). Interseismic megathrust coupling beneath the Nicoya Peninsula, Costa Rica, from the joint inversion of InSAR and GPS data. *Journal of Geophysical Research: Solid Earth*, 120(5), 3707–3722. <https://doi.org/10.1002/2014JB011844>

Yabuki, T., & Matsu'ura, M. (1992). Geodetic data inversion using a Bayesian information criterion for spatial distribution of fault slip. *Geophysical Journal International*, 109(2), 363–375. <https://doi.org/10.1111/j.1365-246X.1992.tb00102.x>

Yokota, Y., Ishikawa, T., Watanabe, S., Tashiro, T., & Asada, A. (2016). Seafloor geodetic constraints on interplate coupling of the Nankai Trough megathrust zone. *Nature*, 534(7607), 374–377. <https://doi.org/10.1038/nature17632>