

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Title: Sea floor bedforms and their influence on slope accommodation.

Authors: Maselli, V.^{1,2}, Kneller, B.¹, Taiwo, O.L.¹, Iacopini, D.¹

¹School of Geosciences, University of Aberdeen, Meston Bld., King’s College, Aberdeen, AB24 3UF, United Kingdom.

²Department of Earth Sciences, Life Sciences Centre, Dalhousie University, 1355 Oxford Street, Halifax, Nova Scotia, B3H 4R2, Canada.

Keywords: Stoss accommodation space, intra-slope basin, turbidity currents, bedforms, offshore Brazil.

Highlights:

1- 3D seismic data are used to investigate sea floor morphology and underneath stratigraphy in the Potiguar Basin, offshore Brazil.

2- Large- and short-wavelength bedforms are recognized.

3- Poned lobes accumulate on the stoss side of the large-wavelength bedforms, and seismic attributes are used to characterize the associated lithology.

4- The concept of stoss accommodation space is introduced.

Abstract

In deep-water settings, the accommodation space for sediment transported by turbidity flows relates to the difference between the elevation of the depositional surface and its equilibrium profile. As a consequence, accommodation space creation, or disruption, may depend from

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

24 changes in the physiography of the receiving basin, or changes in the flow properties. In
25 topographically complex slopes, such where salt-withdrawal intra-slope basins occur, three
26 different types of accommodation space have been recognized. Among other parameters, the
27 ratio between flow thickness and depth of the intra-slope basin controls the partial, or full,
28 ponding of the sediment in suspension, and consequently, the lithology distribution within the
29 deposit. On a smaller spatial scale, the behavior of turbidity flows is affected by the
30 topography of the sea floor. Indeed, the presence of large-wavelength bedforms may generate
31 local topographic low compared to the adjacent sea floor that may trap part of the sediment
32 carried by sediment-laden flows, such as turbidity flows or bottom currents.

33 With a beautiful example from the offshore Brazil, we show how ponded lobes accumulate
34 on the convex-up stoss side of pre-existing sea floor bedforms and how the three-dimensional
35 topography of the sea floor controls the flow behavior and the deposition farther downslope.
36 In detail, using 3D seismic data and attributes we demonstrate that the stoss side of the
37 bedform traps the coarse-grained fraction of turbidity currents flowing downslope, while the
38 fine-grained cloud spills over its crest. Further studies are necessary to fully understand the
39 behavior of sediment-laden flows on a complex sea floor topography, the preservation
40 potential of such deposits and the role of *stoss accommodation* in the evolution of deep-water
41 depositional systems.

42 43 **1. Introduction**

44 Accommodation space was defined by Vail (1987) and Jervey (1988) as the space available
45 for sediment accumulation, with global sea level change and tectonic processes (driving
46 subsidence or uplift of the sea floor) considered as first order controls. In shallow water
47 shelfal systems, the accommodation space available is also dependent from the energy of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48 marine processes, such as waves, longshore drift or tides, and by the presence of topographic
49 lows, such as incised valleys (Dalrymple et al., 1992).

50 In deep-water settings, the concept of accommodation space was expanded considering the
51 analogy between subaerial (rivers) and submarine channels, both characterized by
52 downstream concave-up equilibrium profiles and a base level (Carter, 1988; Prather et al.,
53 1998; Pirmez et al., 2000; Kneller, 2003). A topographic profile is considered in equilibrium,
54 or at grade, when the kinetic energy distribution along the system is such that no net sediment
55 aggradation or erosion occurs. In fluvio-deltaic systems the base level coincides with sea or
56 lake levels, i.e. the channel mouth, while for submarine channels the base level was defined
57 as the deepest point reached by a gravity-driven flow (Carter, 1988), or the point where the
58 transition from confined to unconfined flow occurs (Kneller, 2003). Turbidity currents exert a
59 paramount control on the shape of the equilibrium profile with the gradient of submarine
60 channels directly related to the flow conditions (flow density, thickness, grain size, mud
61 content; Mutti et al., 1999; Kneller, 2003). Considering the above, the accommodation space
62 was defined by the difference between the topography of the depositional surface (i.e., the
63 thalweg of a slope turbidite channel) and its equilibrium profile (Prather et al., 1998; Pirmez
64 et al., 2000). When a submarine channel is at grade, the accommodation space is limited, a
65 meandering planform morphology develops, with no aggradation or incision (Kneller, 2003),
66 producing fluvial-like meander belts (Abreu et al., 2003; Kolla et al., 2012).

67 A disequilibrium between the channel thalweg and the graded profile will lead to
68 accommodation space creation or destruction that the system will exploit through deposition
69 within the channel or erosion of its thalweg and rejuvenation of the system (Pirmez et al.,
70 2000; Heiniö and Davies, 2007). Several mechanisms, mainly driven by tectonic processes
71 (Prather et al., 1998; Pirmez et al., 2000; Ferry et al., 2005) or emplacement of mass-transport
72 deposits (Armitage et al., 2009; Kneller et al., 2016 and references therein), may lead to the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

73 formation of accommodation space for sediment deposition. The topography of the slope may
74 change in response to shale or salt diapirism under loading by thick sediment accumulations,
75 or in response to gravitational tectonics driven by rapid sedimentation along passive margins
76 (Prather, 2003), or by crustal extension or compression, leading to the formation of ponded
77 and healed-slope accommodation space (*sensu* Prather et al., 1998). Ponded slope basins have
78 been recognized in different settings, both modern and ancient, and extensively investigated
79 in the Gulf of Mexico and in the Eastern Equatorial Atlantic margin (Prather et al., 1998;
80 Beaubouef and Friedmann, 2000; Pirmez et al., 2000; Sinclair and Tomasso, 2002; Booth et
81 al., 2003; Smith, 2004; Barton, 2012; Deptuck et al., 2012; Prather et al., 2012; Jobe et al.,
82 2015; Jobe et al., 2017; Hawie et al., 2018). Through integration of seismic and well data, the
83 motif of the sedimentary infill has been interpreted in terms of a process of fill-and-spill, i.e
84 filling of the mini-basin by ponded turbidites and associated deposits, and subsequent bypass
85 from the shallower mini-basin to the one downslope (Winker, 1996; Prather et al., 1998;
86 Badalini et al., 2000; Prather et al., 2012). Mass-transport deposits (MTDs), ubiquitously
87 recognized in all margin settings, have the potential to generate different styles of
88 accommodation space and to control deep-water sediment routing systems (Kneller et al.,
89 2016; Soutter et al., 2018). Sediment may accumulate along the evacuation zone of
90 submarine landslides or along the relative topographic lows generated atop the MTDs by the
91 presence of blocks, faults, folds and compaction (Kneller et al., 2016).

92 Local topographic changes of the sea floor (i.e., bedforms) have been observed on the slope
93 in association with the passage of gravity-driven flows such as turbidity currents, or of
94 bottom currents (Wynn and Stow, 2002; Smith et al., 2007; Piper and Normark, 2009;
95 Rebesco et al., 2014; Talling et al., 2015; Symons et al., 2016 and references therein).

96 Turbidity and bottom currents interacting with the sea floor may generate bedforms both
97 depositional (sediment waves), erosional (scours), or mixed, generated where both erosion

1 98 and deposition occurs (terminology *sensu* Symons et al., 2016). Bedforms of different shapes,
2
3 99 aspect ratio, direction of migration and grain size, from mud to gravel, have been recognized
4
5 100 in both confined and unconfined settings, such as shelfal systems (Berndt et al., 2006), pro-
6
7 101 delta slopes (Casalbore et al., 2017), channel axis (Paull et al., 2010; 2011), channel levees
8
9 102 (Normark et al., 2002), and channel-lobe transitions (Carvajal et al., 2017). After the seminal
10
11 103 work of Fildani et al. (2006) on the Monterey East Channel, increasing attention has been
12
13 104 dedicated to the study of bedforms, including those described as cyclic steps and their
14
15 105 associated supercritical flow regimes (i.e., densimetric Froude numbers >1). Cyclic steps and
16
17 106 antidunes have been increasingly recognized along delta fronts (Normandeau et al., 2016;
18
19 107 Hughes Clarke, 2016) and slope channel systems (Covault et al., 2017), and a growing body
20
21 108 of evidence has suggested that channels may evolve from a series of erosional bedforms
22
23 109 arranged in a cyclic manner (i.e., cyclic steps; Fildani et al., 2013; Covault et al., 2014). On
24
25 110 the sea floor, erosional bedforms, or those with an erosional component, may reach up to 10^3
26
27 111 m in length and width, and up to 10^2 m in height (Cartigny et al., 2011; Symons et al., 2016),
28
29 112 often showing circular to elliptical morphology, such as in the case of the Monterey East
30
31 113 channel (Fildani et al., 2006). With respect to the adjacent sea floor, the stoss side of such
32
33 114 bedforms constitutes an area of lower bathymetry, and consequently represent
34
35 115 accommodation space for sediment accumulation.
36
37
38
39
40
41
42
43

44 116 This study aims to understand the role of topographic lows generated by turbidity-current-
45
46 117 related bedforms (both erosional and depositional) in promoting slope accommodation space
47
48 118 and affecting sediment dispersal patterns and pathways. Using an example from the Brazilian
49
50 119 slope in the offshore Potiguar basin, we will demonstrate how the stoss side of large-
51
52 120 wavelength bedforms may act as a mini-basin where coarse-grained sediments transported by
53
54 121 turbidity currents may accumulate generating ponded lobes. Moreover, we will discuss how
55
56 122 the flow transformation of turbidity currents through flow stripping across the bedform-
57
58
59
60
61
62
63
64
65

123 related ponded mini-basin may affect the sedimentation pathways and related sea floor
124 topography of the slope farther downslope.

125

126 **2. Study Area and Geological Setting**

127 The present study focuses on a portion of the Brazilian slope just south of the Equator, in the
128 offshore Potiguar basin, in water depths between ca. 700 m and 1800 meters below mean sea
129 level (m bmsl; Fig. 1). The area is characterized by a ca. 60 km wide, low angle (0.04°), shelf,
130 and a steep slope, dipping towards NE at ca. 3.8° . Towards the basin, in deeper water, a series
131 of volcanic islands and structural highs develop, creating troughs that interrupt the continuity
132 of the slope (Fig. 1).

133 The Potiguar basin is a NE-trending aborted rift with ~6,000 m thick sedimentary infill,
134 structurally characterized by SW-NE-trending asymmetric grabens separated by internal
135 basement horsts (Matos, 2000; Jovane et al., 2016). The rifting process began in response to
136 continental breakup between the Borborema and Benin-Nigeria provinces during the South
137 Atlantic opening in the Early Cretaceous (Matos, 2000). Rift phase deposition during the
138 Aptian to Campanian, consisted of fluvial to shallow-marine transgressive sediments (Araripe
139 and Feijó, 1994). The drift phase, starting in the Campanian, is characterized by thermal
140 subsidence and deposition of fluvio-deltaic to deep-water clastic sediments, with the Neogene
141 mainly recording the onset and evolution of the submarine canyon systems still active today.

142

143 **3. Data and Methods**

144 The dataset from the Potiguar Basin used in the present study consists of a high-quality 3D
145 full stack, Kirchhoff time-migrated reflection seismic volume, covering about ~2000 km²,
146 and acquired by PGS in 2009 (Fig. 1). The line spacing is 12.5 m in both in-line and cross-

147 line directions. The sample interval is 2 milliseconds (ms). The data are zero-phase migrated
148 and displayed with Society for Exploration Geologists (SEG) normal polarity, so that an
149 increase in acoustic impedance is represented by a blue-red-blue reflection loop. The
150 dominant frequency (F) of the section of interest (upper 250 ms below the seabed) ranges
151 between 40 and 75 Hz. Sound velocities of 1,500 ms⁻¹ and 1,800-2,500 ms⁻¹ have been
152 respectively assigned for sea water and for the investigated interval below the sea floor, with
153 the latter velocity obtained from the sonic log of well CES-112, located 2 km to the SE (see
154 Fig. 1; Conde et al., 2007). Using those end-member velocities and frequencies, we estimate
155 a vertical resolution (defined as tuning thickness, or limit of separability, as $\lambda/4$, λ being the
156 wavelength of the P wave) as 5 m at the sea floor and 6 to 15.5 m for the units below. Taking
157 into account the focusing effect of Kirchhoff migration (Brown, 2004), a radius Fresnel zone
158 (with a radius equal to velocity $V_{\text{average}}/4F$) of 5 m to 15.5 m can be reasonably expected, and
159 therefore a minimum diameter 10-31 m represents the limit of our interpretation analysis on a
160 horizon (*sensu* Brown, 2004). However our ability to recognize sea floor features in plan
161 view, defined as detectability or limit of visibility (Brown, 2004), can go below the tuning
162 thickness limit (Brown, 2004); while in the 3D migrated dataset, reflectors (e.g., peak to peak)
163 will be at least 5 m thick, and two reflecting point in horizontal space need to be around 5-15
164 m a part, when we are mapping the seafloor as a surface horizon, geological features on
165 considerably smaller scales can be detected and visualized with greater detail (Reijenstein et
166 al., 2011). Thus we can describe geological and sedimentary features or patterns smaller than
167 the tuning thickness, although our capacity to define volumes is limited by the tuning
168 thickness.

169 The bathymetry of the sea floor, presented at a 12.50 × 12.5 m of horizontal resolution (Fig.
170 2), was generated picking the first reflection from the 3D seismic data. Two other seismic
171 horizons, H1 and H2, were identified on 2D arbitrary lines extracted from the 3D seismic

172 volume, based on the seismic facies and reflector terminations. The structural map of each
173 seismic horizon is presented as surface gridded at a 12.5×12.5 m of horizontal resolution.
174 Seismic attributes have been calculated and extracted from the sea floor horizon and include
175 both amplitude-derived (root-mean-square, RMS) and time-derived (variance) values. While
176 the variance, which measures the similarity of consecutive waveforms over a given sampling
177 window (3×3 traces in the present study), is useful for imaging lateral discontinuities (Chen
178 and Sidney, 1997; Brown, 2004), the RMS amplitude, which represents the square root of the
179 arithmetic mean of the squares of the amplitudes within a defined window interval (3
180 instantaneous traces in the present study), is helpful for revealing coarse-grained facies (Rijks
181 and Jauffred, 1991; Chen and Sidney, 1997; Brown, 2004).

182

183 **4. Results**

184 **4.1. Sea floor morphology**

185 The sea floor shows two main canyon-channel systems, named C-1 and C-3 (Fig. 2) that are
186 located towards NW and SE corners of the dataset, respectively. The depth of both channels
187 changes from ca. 400 m to less than 200 downslope, while the thalweg presents an average
188 gradient of 2.7° and a sinuosity index of 1.158, for C-1, and a gradient of 3.8° and a sinuosity
189 index of 1.028, for C-3 (Fig. 2). A smaller channel, C-2, ca. 90 m deep, crosses the slope with
190 an average thalweg gradient of 4.15° and a sinuosity index of 1.031. The present study
191 focuses in the slope area comprised between C-2 and C-3 (confined by the red line in Fig. 2).

192 Two narrow channel incisions (C-A and C-B, Fig. 2), up to 60 m deep and oriented SW-NE,
193 form upstream of a topographic step (slope break) oriented approximately NS (dashed red
194 line in Fig. 2). Farther downslope, the sea floor presents a series of large-wavelength
195 bedforms (Fig. 3), named LB1 to LB4, which are clearly highlighted by the variance attribute

196 extracted from the sea floor horizon (Fig. 3 right, and Fig. 4). The bedform wavelength
197 changes from ca. 4 km (LB1, Fig. 5) to less than 1 km (LB4, Fig. 5), while the bedform
198 height from ca. 150 m (LB1, Fig. 5) to less than 50 m (LB4, Fig. 5). The crests of the
199 bedforms show a sinuous shape, with dominant downslope convexity (Fig. 3), and are
200 progressively shifted towards East moving downdip, following the maximum gradient of the
201 sea floor.

202 In a cross section on the sea floor, the bedforms are downslope asymmetric, with seaward
203 dipping (LB1 and LB2, Fig. 3 bottom) or sub-horizontal (LB3 and LB4) stoss sides, and up to
204 8° dipping lee sides. A series of small channels (named gutter-like channels), up to 15 m deep,
205 incise the lee sides of LB1 to LB4, and the area farther downslope.

206 On the sea floor, the stoss side of LB1, just downdip of the slope break at the mouth of C-A
207 and C-B, is characterized by two fields of short-wavelength bedforms (SB1a and SB1b, Fig.
208 3), both symmetric (section a-b in Fig. 6) and asymmetric (section c-d in Fig. 6), and with
209 sinuous crests (Fig. 6, right). Wavelengths and heights are, on average, 120 m and 8 m,
210 respectively (Fig. 5). A third train of bedforms (named SB2) with linear crests and ca. 5 m
211 wave heights is present on the lee side of LB3 (SB2 in Fig. 3; section e-f in Fig. 6). We are
212 confident that the spatial (vertical and horizontal) resolution of the sea floor generated by
213 picking the sea floor horizon on the 3D seismic dataset is high enough to visualize such
214 small-scale sea floor features. Seismic artefacts are present in the data, as indicated the
215 contour-parallel undulations highlighted by the slope map of Fig. 6, but they are
216 characterized by a totally different seismic footprint (see section g-h in Fig. 6), unrelated to
217 the bathymetry, with wave height and length extremely short, which will not have any effect
218 on the interpreted structures.

219 The RMS amplitude extraction map of the sea floor (Fig. 7) shows lobe-shaped areas
220 characterized by high RMS values on the stoss side of each large-wavelength bedform

221 (highlighted in orange in the B/W version of the RMS map in Fig. 7C). In detail, high RMS
222 values can be found where SB1a, SB1b and SB2 fields develop (named Lobe A, Lobe B and
223 Lobe D, respectively, Figs. 6, 7), on the stoss side of LB2 (Lobe C, Figs. 6, 7), of LB4 (Lobe
224 E, Figs. 6, 7), and of LB3 towards the northern flank of channel C-3 (Lobe F in Figs. 6, 7).
225 High RMS amplitude values also characterize the southern flank of C-2, while low values can
226 be detected along the lee side of all the large-wavelength bedforms (Fig. 7)

227

228 **4.2. Seismic stratigraphy**

229 The stratigraphy of the study area, and in particular of the lobe-shaped areas identified in the
230 RMS attribute map (Fig. 7), has been revealed using a combination of 2D arbitrary lines
231 extracted from the 3D seismic cube, surface maps of key stratigraphic horizons and thickness
232 maps (Figs. 8, 9).

233 Horizon H1, identified by a continuous positive reflection, is the first continuous horizon
234 below the sea floor, which can be traced in much of the study area (Figs. 8, 9B). The horizon
235 forms at the base of a series of lobe-to-lens-shaped deposits (named Lobe A to Lobe F, in
236 Figs. 7, 8, 9), whose tops correspond to the sea floor and show high RMS amplitude values.
237 In the same position, corresponding to the stoss side of the large-wavelength bedforms (Figs.
238 3, 7), the surface map of H1 shows a series of topographic depressions, triangular to circular
239 in shape, with progressively reducing size downslope (Fig. 9B). Thickness map of the unit
240 between the sea floor and horizon H1 (Fig. 9D) shows a series of sediment depocenters up to
241 65 m thick (Lobe A, Fig. 9D), whose internal seismic character is highlighted in Figure 8
242 (Lobes A, B, D and F, as examples). Each lobe shows a positive relief respect to the adjacent
243 sea floor, and is confined basinward by the topography generated by the large-wavelength
244 bedforms (Fig. 8). Lobe A, in detail, is the largest sediment depocenter, covering a surface

245 area of ca. 3.5 km² (Fig. 9D), and is composed by thick, high-amplitude, and wavy reflections
246 (seismic lines 1, 2 and 5 in Fig. 8). The sea floor reflection on top of Lobe A is also wavy
247 (Fig. 8), and corresponds to the short-wavelength bedform field SB1a visible on the sea floor
248 maps of Figures 3 and 6. Thickness map (Fig. 9D) highlights that Lobe A is made up of two
249 bodies, with the shallower backstepping with respect to the deeper (see seismic line 5 in Fig.
250 8). Lobe A accumulates on the stoss side of the large-wavelength bedform LB1, which is
251 confined by horizon H1 at its top and horizon H2 at its base (Figs. 8, 9E). Horizon H2 shows
252 an erosional character, as highlighted by several truncated reflections (see the black arrows in
253 Fig. 8), and can be traced over part of the study area (Fig. 9C). The topographic depression
254 generated by H2 is exploited by the accumulation of LB1, which is a 90 m thick, L shaped,
255 sediment body (Fig. 9E), made by continuous and low amplitude reflections, showing lateral
256 (see seismic lines 3 and 4 in Fig. 8) and upslope (see seismic line 5 in Fig. 8) migration of
257 progradation, and internal erosional surfaces (highlighted by black dashed lines in the seismic
258 profiles of Fig. 8). The deposition of the large-wavelength bedform LB1 visible on the sea
259 floor (Figs. 3, 4) creates the accommodation space for the accumulation of Lobe A and its
260 downslope confinement, as shown by Figure 9F. Similar geometric relations are observed for
261 each Lobe B to F, where the deposition of a lower unit bounded by an erosional surface (see
262 the red dashed lines in seismic lines 6 and 7 in Fig. 8, and line 8 in Fig. 9) causes the
263 generation of the large-wavelength bedforms and for the formation of accommodation space
264 along the slope. RMS amplitude extraction of the sea floor integrated with the seismic facies
265 in cross section highlight that each lobe has high RMS values and is made by high-amplitude
266 reflections (Fig. 7). By contrast, the units beneath, which crop out on the seafloor along the
267 lee side of the large-wavelength bedforms, present low RMS amplitude and mainly low
268 amplitude seismic reflections (Figs. 7, 8, 9). Consequently, we can infer that the lobes are
269 made by coarser-grained (probably sandy) sediment compared to the deposit beneath that are

1 270 responsible for the formation of the large-wavelength bedforms, which are probably muddier.

2 271 Correlation of horizon H2 with the other erosional surfaces occurring farther downslope is

3 272 not straightforward, which poses problem is the development of a conceptual model for

4 273 explaining their evolution. In the supplementary material we present two models which take

5 274 into account the effect of different processes and the possibility that the erosional surfaces

6 275 occurred synchronously to H2, or not.

7 276

8 277 **5. Discussion**

9 278 Topographically complex slopes (*sensu* Smith, 2004) occur when tectonic processes or

10 279 deformations of the sea floor driven by sediment loading on a mobile substrate create

11 280 topographic lows or highs that can affect the path and behavior of gravity-driven flows

12 281 traveling downslope. In such contexts, different types of accommodation space may exist

13 282 (namely ponded, healed and slope accommodation; Prather et al., 1998), whose infill reflects

14 283 the effect of changing accommodation space through time (due to deposition), on the

15 284 behavior of gravity flows, and on the instability of the slope (Prather, 2003). Accommodation

16 285 space can be generated *a-priori*, and then filled by sediments, or can be increased by

17 286 sediment loading during basin infill, as in the case of salt withdrawal intra-slope basins

18 287 (Winker, 1996; Prather et al., 1998). It has been demonstrated that also sediment compaction

19 288 may significantly increase slope accommodation (Reynolds et al., 1991).

20 289 As sediment suspension in turbulent flows depends on bed shear stress, which is directly

21 290 related to flow velocity, 3D sea floor topography may control sediment deposition, erosion or

22 291 bypass through flow non-uniformity (Kneller and McCaffrey, 1995). Sea floor bedforms in

23 292 unconfined settings, normally generated by both erosional or depositional turbidity flows and

24 293 bottom currents (Rebesco et al., 2014; Symons et al., 2016 and references therein), may

294 create relative topographic lows (i.e., the convex-up stoss side of the bedform) where the
295 sediment transported by newly generated gravity flows may accumulate. Such lows can be up
296 to 10^2 m height and 10^3 m long, ca. an order of magnitude smaller, in both dimensions, than
297 the intra-slope basins in the Gulf of Mexico, and may act as an intra-slope mini-basin
298 generating what here we call *stoss accommodation*. In the study area, deposition from
299 unconfined flows or bottom currents was probably responsible for the creation of stoss
300 accommodation through the deposition of the large-wavelength bedforms (LB1 to LB4 in Fig.
301 3; see supplementary material). The ability of a turbidity current to flow across a
302 topographically complex slope, such as a salt withdrawal mini-basin or a large-wavelength
303 bedform, depends on the flow type (surging vs continuous; Lamb et al., 2004), the flow
304 thickness (Lane-Serff et al., 1995), the internal Froude number and the flow stratification
305 (Kneller and McCaffrey, 1999). Complete ponding of the flow occurs if the entire flow is
306 trapped within the topographic depression (Patacci et al., 2015, and references therein), and
307 sedimentation farther downslope is expected after its filling, partial or total, through a process
308 called fill-and-spill (Prather et al., 1998). If the depression is small enough compared to the
309 flow, the turbidity current may be able to surmount its downstream lip: the coarse-grained
310 part of the flow will accumulate within the topographic low while the fine-grained cloud will
311 be able to escape through a process called flow stripping (Piper and Normark, 1983).
312 Experimental results of Lane-Serff et al. (1995) demonstrated that a volume-limited flow (i.e.,
313 a surge-like turbidity current) may rise a topographic relief up to 5 times the flow thickness.
314 If we consider the case of one of the largest bedforms discovered so far on the modern sea
315 floor (i.e., the Monterey East channel which shows a maximum wave height of 220 m;
316 Fildani et al., 2006), all the incoming flows thicker than 44 m (with a supercritical regime, as
317 from Lane-Serff et al., 1995) will likely be able to generate overspill from the downstream lip
318 of the bedform. Of course this is an approximation based on Lane-Serff et al. (1995)

1 319 numerical results, as flow stratification (density and velocity) is also key in controlling the
2 320 maximum run-up height of a turbidity current (Kneller and McCaffrey, 1999; Kneller and
3
4 321 Buckee, 2000). Considering the example of the Monterey East channel as a conceptual end
5
6
7 322 member, we can argue that turbidity currents thicker than ca. 40 meters will be likely able to
8
9 323 overspill from any topographic depressions generated by pre-existing sea floor bedforms. In
10
11
12 324 such scenario, stoss accommodation will be mainly exploited by deposition of the coarser
13
14 325 part of the flow, while the fine-grained cloud of the turbidity current will be likely to
15
16 326 overspill, potentially creating sandy ponded lobes. This mechanism may affect sediment
17
18 327 deposition farther downslope, and the potential development of new sediment corridors.
19
20 328 Although with some limitations due to the lack of vertical resolution of the seismic data, this
21
22 329 conclusion is supported by the results of this study, which show high RMS amplitude values,
23
24 330 considered a proxy for coarse-grained sediment, on the stoss side of each large-wavelength
25
26 331 bedform (Fig. 7), and the presence of small channel incisions (named gutter-like channels)
27
28 332 along their lee sides, which will control deposition towards the basin. In addition, the short-
29
30 333 wavelength bedforms on Lobe A (SB1a, Figs. 3, 5) show crest directions perpendicular to the
31
32 334 local slope, probably reflecting deposition from turbidity currents radially spreading at the
33
34 335 mouth of channel C-A, on the flat surface generated after the infill of the stoss side of LB1.
35
36 336 Similar features have been observed in other context and linked to deposition from turbidity
37
38 337 currents (Normandeau et al., 2015). Turbidity currents are highly sensitive to changes in sea
39
40 338 floor topography, as divergence or convergence of the streamlines produces sediment
41
42 339 deposition or erosion/bypass, respectively (Kneller and McCaffrey, 1995). As in the case of
43
44 340 supra-MTD topography (Kneller et al., 2016), pre-existing bedforms may create a complex
45
46 341 sea floor topography that will undoubtedly have an effect on the turbidity currents (i.e.,
47
48 342 deposition, erosion, bypass) depending on the flow properties and direction (see Kneller et al.,
49
50 343 2016 for further discussion) with respect to the stoss accommodation space available. Further
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

344 work is needed to evaluate the flow behavior across pre-existing large-wavelength bedforms
345 and for turbidity currents unrelated with the deposition of the bedforms themselves, to
346 quantify the facies association of the ponded lobes through direct sediment sampling, and
347 their preservation potential. This may shed lights on defining the role of stoss accommodation
348 in hydrocarbon exploration and in the whole evolution of deep-water depositional systems.

349

350 **6. Conclusion**

351 Sea floor topography is a first order control on the flow behavior of turbidity currents flowing
352 down the slope of continental margins. Intra-slope basins are normally associated to large-
353 scale deformation of the sea floor, mainly promoted by salt or gravitational tectonics. With an
354 example from the offshore Brazil, here we show that smaller-scale topographic variations of
355 the sea floor associated with bedforms, may have a large effect on subsequent turbidity
356 currents and may promote the formation of coarse-grained ponded lobes. In detail, bedforms
357 characterized by convex-up stoss sides form topographic lows compared to the sea floor near-
358 by, generating stoss accommodation. Depending on the flow characteristics of newly sourced
359 turbidity currents, flow stripping or fill-and-spill may occur, in the first case promoting the
360 formation of sandy ponded lobes. The existence of a 3D topography may capture the coarse-
361 grained fractions of the flows in the relative lows while promoting the delivery of only the
362 fine-grained part downstream. Further studies are needed to understand the role of stoss
363 accommodation in the evolution of turbidity flows and associated deposits, and their
364 preservation potential in the stratigraphy of continental margins.

365

366 **Acknowledgments**

367 We sincerely thank Petroleum Geo-Services (PGS) and specifically David Hajovsky and
368 Scott Opdyke who kindly provided the dataset and allowed us to show these results. We
369 would also like to thank Schlumberger for providing academic licenses of their software
(Petrel), and for their support.

372 **Figure captions**

374 **Figure 1**

375 Top: Digital elevation model of the Equatorial Atlantic margin (data from GEBCO). Centre:
376 close-up on the Potiguar Basin in the offshore Brazil; white rectangle represents the full 3D
377 seismic data coverage, while the study area is highlighted in red; black and orange lines mark
378 the position of the bathymetric profiles presented below. Bottom: bathymetric profiles cross
379 the Ceará high and across an open slope setting.

381 **Figure 2**

382 Top: Bathymetric map with 75 m spaced contour lines; white lines oriented NW-SE are the
383 bathymetric profiles presented below, while the thick and continuous red line marks the study
384 area. Note the two large canyon-channel systems bordering the study area (named C-1 and C-
385 3) and the narrower incisional channel C-2. The thin dashed red line marks the slope break at
386 the mouth of incisions C-A and C-B. Bottom: Bathymetric profiles across section AB, CD,
387 EF and GH showing the change of channel depth of C-1, C-2 and C-3 with bathymetric depth;
388 note the position of the slope break in sections AB and CD.

390 Figure 3

1
2
3 391 A: Bathymetric map of the study area with 75 m spaced contour lines; the white continuous
4
5 392 line marks the position of the bathymetric profile IJ presented in C, while the white dashed
6
7 393 lines mark the crest of the large-wavelength bedforms, named LB1 to LB4; the red dashed
8
9 394 line marks the slope break at the mouth of C-A and C-B; note the gutter-like channels. B:
10
11
12 395 variance attribute map extracted from the sea floor horizon; note the short-wavelength
13
14 396 bedforms (SB1a, SB1b, SB2). C: the bathymetric profile IJ showing the large- and short-
15
16
17 397 wavelength bedforms (gray rectangle) shown in black; the sea floor gradient along the section
18
19
20 398 IJ shown in red, with highlighted the different bedform fields.
21
22

23 399

24
25 400 Figure 4

26
27
28 401 Variance attribute extracted from the sea floor and presented in a 3D view. The white dashed
29
30 402 lines mark the crest of the large-wavelength bedforms (LB1 to LB4), while the red dashed
31
32 403 line marks the slope break at the mouth of C-A and C-B. Note the gutter-like channels (Gc)
33
34 404 and the short-wavelength bedform SB2.
35
36
37
38

39 405

40
41
42 406 Figure 5

43
44
45 407 Wave length (in km) and wave height (m) of the different bedform fields recognized in this
46
47 408 study. The inset is a zoom of the left corner of the diagram to highlight bedforms SB1a, SB1b
48
49 409 and SB2.
50
51

52 410

53
54
55 411 Figure 6

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

412 Plan view of the sea floor slope map (left) and a close-up 3D of the stoss side of LB1 (right);
413 the red dashed line marks the slope break at the mouth of C-A and C-B; blue squares 1 and 2
414 highlight short-wavelength bedform fields SB1a and SB2 (zoom visible below), while the red
415 square is an example of seismic artefact. Bathymetric profiles across section ab and cd show
416 bedform styles on SB1, while profile ef shows the bedforms on SB2 (bedform's crests
417 pointed by the arrows). Profile gf highlights the seismic artefacts, almost invisible on a
418 bathymetric profile. All the bathymetric profiles are presented at the same scale.

419
420 Figure 7

421 A: Top view of the RMS attribute map extracted from the sea floor horizon; the white dashed
422 lines mark the crest of the large-wavelength bedforms (LB1 to LB4), while the red dashed
423 line marks the slope break. B: RMS amplitude map presented on a 3D view. Note that the
424 stoss side of the bedforms is repeatedly characterized by high RMS amplitude values (named
425 Lobe 1 to Lobe F), while the lee side by lower values. Note the gutter-like channels (Gc). C:
426 Grey-scale version of the RMS attribute map presented in A. This graphic solution is used to
427 better highlight the different features: lobes in orange, crests of the large-wavelength
428 bedforms in dashed white line, slope break in dashed red line.

429
430 Figure 8

431 2D arbitrary lines (see inset map for location) extracted from the 3D seismic cube, all
432 presented at the same scale. Lines 1 to 5 show Lobes A and B, and the internal stratigraphy of
433 the large-wavelength bedform LB1, while Lines 6 and 7 show Lobe D and F, respectively.
434 Horizon H2, highlighted in red (note the truncated reflections in sections 4 and 5, for
435 example), marks the base of bedform LB1. Internal reflections of LB1 are continuous and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

436 low-amplitude (internal erosional surfaces marked by black dashed lines), and present an
437 oblique to upslope direction of migration as seen on the 3D data. Horizon H1, highlighted in
438 black, can be traced at the base of the all the Lobes, which present a positive relief on the sea
439 floor and, when visible, the internal reflections are wavy and high-amplitude. Note the short-
440 wavelength bedform fields on the sea floor reflection. Red dashed lines on seismic Lines 6
441 and 7 highlight the erosional surface at the base of the LB3 and LB4, in analogy with horizon
442 H2.

444 Figure 9

445 Top: Seismic line across the large-wavelength bedforms LB1 to LB4, with highlighted
446 horizon H2 (continuous red line), horizon H1 (black line), and other erosional surfaces (red
447 dashed lines); see trackline in A. Horizon H1 can be traced in much of the study area and
448 forms the base of the ponded Lobes A to F. A: Sea floor bathymetry. B: Structural map of
449 horizon H1. C: Structural map of horizon H2, which highlights the base of LB1. D: Thickness
450 map generated by the difference between the sea floor and horizon H1, which highlights the
451 ponded Lobes A to F. E: Thickness map generated by the difference between H1 and H2
452 horizons, which highlights the large-wavelength bedform LB1. F: Combined thickness maps
453 showing how Lobe A is confined basinward by LB1, filling the accommodation space
454 generated by the stoss side of LB1.

456 References

457 Abreu, V., Sullivan, M., Pirmez, C., Mohrig, D., 2003. Lateral accretion packages (LAPs): an
458 important reservoir element in deep-water sinuous channels. *Marine and Petroleum*
459 *Geology* 20, 631-648.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- 460 Adeogba, A.A, McHargue, T.R., Graham, S.A., 2005. Transient fan architecture and
461 depositional controls from near-surface 3-D seismic data, Niger delta continental
462 slope. AAPG Bulletin 89, 627-643.
- 463 Araripe, P.T., Feijó, F.J., 1994. Carta estratigráfica da Bacia Potiguar. Boletim De
464 Geociências Da Petrobras 8, 127-141.
- 465 Armitage, D.A., Romans, B.W., Covault, J.A., Graham, S.A., 2009. The influence of mass-
466 transport-deposit surface topography on the evolution of turbidite architecture: The
467 Sierra Contreras, Tres Pasos Formation (Cretaceous), southern Chile. Journal of
468 Sedimentary Research 79, 287-301.
- 469 Badalini, G., Kneller, B., Winker, C.D., 2000. Architecture and processes in the Late
470 Pleistocene Brazos-Trinity turbidite system, Gulf of Mexico Continental Slope.
471 GCSSEPM Foundation 20th Annual Research Conference, Deep-Water Reservoirs of
472 the World, 40-103.
- 473 Barton, M.D., 2012. Evolution of an Intra-Slope Apron, Offshore Niger Delta Slope: Impact
474 of Step Geometry on Apron Architecture. In: Prather, B.E., Deptuck, M.E., Mohrig,
475 D., van Hoorn, B., Wynn, R.B., (Eds.), Application of the Principles of Seismic
476 Geomorphology to Continental -Slope and Base-of-Slope Systems: Case Studies from
477 Seafloor and Near-Seafloor Analogues. SEPM Special Publication 99, pp. 181-197.
- 478 Beaubouef, R.T., Friedmann, S.J., 2000. High Resolution Seismic/Sequence Stratigraphic
479 Framework for the Evolution of Pleistocene Intra Slope Basins, Western Gulf of
480 Mexico: Depositional Models and Reservoir Analogs. Deep-Water Reservoirs of the
481 World, Gulf Coast SEPM, 40-60.

- 1
2
3
4
5 482 Berndt, C., Cattaneo, A., Szuman, M., Trincardi, F., Masson, D., 2006. Sedimentary
6
7
8 483 structures offshore Ortona, Adriatic Sea - Deformation or sediment waves? Marine
9
10 484 Geology 234, 261-270.
11
12
13 485 Booth, J.R., Dean, M.C., Duvernay, III, A.E., Styzen, M.J., 2003. Paleo-bathymetric controls
14
15 486 on the stratigraphic architecture and reservoir development of confined fans in the
16
17 487 Auger Basin: central Gulf of Mexico slope. Marine and Petroleum Geology 20, 563-
18
19 488 586.
20
21
22 489 Brown, A.R., 2004: Interpretation of three-dimensional seismic data, 5th edition. AAPG
23
24 490 Memoir 42, Tulsa, Oklahoma, pp. 514.
25
26
27 491 Casalbore, D., Ridente, D., Bosman, A., Chiocci, F.L., 2017. Depositional and erosional
28
29 492 bedforms in Late Pleistocene-Holocene pro-delta deposits of the Gulf of Patti
30
31 493 (southern Tyrrhenian margin, Italy). Marine Geology 385, 216-227.
32
33
34 494 Carter, R.M., 1988. The nature and evolution of deep-sea channel systems. Basin Research 1,
35
36 495 41-54.
37
38
39 496 Cartigny, M.J.B., Postma, G., van den Berg, J. H., Mastbergen, D.R., 2011. A comparative
40
41 497 study of sediment waves and cyclic steps based on geometries, internal structures and
42
43 498 numerical modeling. Marine Geology 280, 40-56.
44
45
46 499 Carvajal, C., Paull, C.K., Caress, D.W., Fildani, A., Lundsten, E., Anderson, K., Maier, K.L.,
47
48 500 Mcgann, M., Gwiazda, R., Herguera, J.C., 2017. Unraveling the channel-lobe
49
50 501 transition zone with high-resolution AUV bathymetry: Navy Fan, Offshore Baja
51
52 502 California, Mexico. Journal of Sedimentary Research 87, 1049-1059.
53
54
55 503 Chen, Q., Sidney, S., 1997. Seismic attribute technology for reservoir forecasting and
56
57 504 monitoring. The Leading Edge 16, 445-448.
58
59
60
61
62
63
64
65

- 505 Condé, V.C., Lana, C.C., Pessoa Neto, O.C., Roesner, E.H., Morais Neto, J.M., Dutra, D.C.,
1
2 506 2007. Baciado Ceará. *Bol. Geoc. Petrobras* 15, 347-355.
3
4
5 507 Covault, J.A., Kostic, S., Paull, C.K., Ryan, H.F., Fildani, A., 2014. Submarine channel
6
7 508 initiation, filling and maintenance from sea-floor geomorphology and
8
9 morphodynamic modelling of cyclic steps. *Sedimentology* 61, 1031-1054.
10
11
12
13 510 Covault, J.A., Kostic, S., Paull, C.K., Sylvester, Z., Fildani, A., 2017. Cyclic steps and related
14
15 511 supercritical bedforms: Building blocks of deep-water depositional systems, western
16
17 North America. *Marine Geology* 393, 4-20.
18
19
20
21 513 Dalrymple, R.W., Zaitlin, B.A., Boyd, R., 1992. Estuarine facies models: conceptual basis
22
23 514 and stratigraphic implications. *Journal of Sedimentary Petrology* 62, 1130-1146.
24
25
26 515 Deptuck, M.E., Sylvester, Z., O'Byrne, C.J. Pleistocene seascape evolution above a “simple”
27
28 516 stepped slope-western Niger Delta. In: Prather, B.E., Deptuck, M.E., Mohrig, D., van
29
30 Hoorn, B., Wynn, R.B., (Eds.), *Application of the Principles of Seismic
31
32 Geomorphology to Continental -Slope and Base-of-Slope Systems: Case Studies from
33
34 518 Seafloor and Near-Seafloor Analogues. SEPM Special Publication 99, pp. 199-222.*
35
36 519
37
38
39 520 Ferry, J.N., Mulder, T., Parize, O., Raillard, S., 2005. Concept of equilibrium profile in deep-
40
41 521 water turbidite system: effects of local physiographic changes on the nature of
42
43 sedimentary process and the geometries of deposits. In: Hodgson, D.M., Flint, S.S.,
44 522 (Eds.), *Submarine Slope Systems: Processes and Products*, Geological Society of
45
46 523 London, Special Publications 244, pp. 181-193.
47
48
49 524
50
51
52 525 Fildani, A., Normark, W.R., Kostic, S., Parker, G., 2006. Channel formation by flow
53
54 526 stripping: large-scale scour features along the Monterey East Channel and their
55
56 527 relation to sediment waves. *Sedimentology* 53, 1265-1287.
57
58
59
60
61
62
63
64
65

- 528 Fildani, A., Hubbard, S.M., Covault, J.A., Maier, K.L., Romans, B.W., Traer, M., Rowland,
1
2 529 J.C., 2013. Erosion at inception of deep-sea channels. *Marine and Petroleum Geology*
3
4
5 530 41, 48-61.
6
7
8 531 Hawie, N., Covault, J.A., Dunlap, D., Sylvester, Z., 2018. Slope-fan depositional architecture
9
10 532 from high-resolution forward stratigraphic models. *Marine and Petroleum Geology* 91,
11
12 533 576-585.
13
14
15 534 Heiniö, P., Davies, R.J., 2007. Knickpoint migration in submarine channels in response to
16
17
18 535 fold growth, western Niger Delta. *Marine and Petroleum Geology* 24, 434-449.
19
20
21 536 Hughes Clarke, J.E., 2016. First wide-angle view of channelized turbidity currents links
22
23 537 migrating cyclic steps to flow characteristics. *Nature Communications* 7, 1-13.
24
25
26 538 Jervey, MT., 1988. Quantitative geological modeling of siliciclastic rock sequences and their
27
28
29 539 seismic expression. In: Wilgus, C.K., Hasting, B.S., Kendall, C.G.S.C., Posamentier,
30
31 540 H.W., Ross, C.A., Van Wagoner, J., (Eds.), *Sea-level Changes-an Integrated*
32
33 541 *Approach*, 42, SEPM Special Publication, pp. 47-69.
34
35
36 542 Jobe, Z.R., Sylvester, Z., Parker, A.O., Howes, N., Slowey, N., Pirmez, C., 2015. Rapid
37
38
39 543 adjustment of submarine channel architecture to changes in sediment supply. *Journal*
40
41 544 *of Sedimentary Research* 85, 729-753.
42
43
44 545 Jobe, Z.R., Sylvester, Z., Howes, N., Pirmez, C., Parker, A.O., Cantelli, A., 2017. High-
45
46
47 546 resolution, millennial-scale patterns of bed compensation on a sand-rich intraslope
48
49 547 submarine fan, western Niger Delta slope. *Geol. Soc. Am. Bull.* 129, 23-37.
50
51
52 548 Jovane, L., Figueiredo, J.J.P., Alves, D.V.P., Iacopini, D., Giorgioni, M., Vannucchi, P.,
53
54
55 549 Moura, D.S., Bezerra, F.H.R., Vital, H., Rios, I.L.A., Molina, E.C., 2016.
56
57 550 Seismostratigraphy of the Ceará Plateau: Clues to Decipher the Cenozoic Evolution of
58
59 551 Brazilian Equatorial Margin. *Frontiers in Earth Science* doi:10.3389/feart.2016.00090.
60
61
62
63
64
65

- 552 Kneller, B., McCaffrey, W.D., 1995. Modelling the effects of salt-induced topography on
1
2 553 deposition from turbidity currents. GCSSEPM Foundation 16th Annual Research
3
4 554 Conference Salt, Sediment and Hydrocarbons, 137-145.
5
6
7 555 Kneller, B., McCaffrey, W.D., 1999. Depositional effects of flow nonuniformity and
8
9 stratification within turbidity currents approaching a bounding slope: deflection,
10 556 reflection, and facies variation. *Journal of Sedimentary Research* 69, 980-991.
11
12 557
13
14 558 Kneller, B., Buckee, C., 2000. The structure and fluid mechanics of turbidity currents: a
15
16 559 review of some recent studies and their geological implications. *Sedimentology*, 47,
17
18 560 62-94.
19
20
21 561 Kneller, B., 2003. The influence of flow parameters on turbidite slope channel architecture.
22
23 562 *Marine and Petroleum Geology* 20, 901-910.
24
25
26 563 Kneller, B., Dykstra, M., Fairweather, L., Milana, J.P., 2016. Mass-transport and slope
27
28 564 accommodation: Implications for turbidite sandstone reservoirs. *AAPG Bulletin* 100,
29
30 565 213-235.
31
32
33 566 Kolla, V., Bandyopadhyay, A., Gupta, P., Mukherjee, B., Ramana, D.V., 2012. Morphology
34
35 567 and internal structure of a recent upper Bengal fan-valley complex. In: Prather, B.E.,
36
37 568 Deptuck, M.E., Mohrig, D., Van Horn, B., Wynn, R.B., (Eds.), *Application of the*
38
39 569 *Principles of Seismic Geomorphology to Continental-Slope and Base-of-Slope*
40
41 570 *Systems: Case Studies from Seafloor and Near-Seafloor Analogues*, SEPM, Special
42
43 571 *Publication* 99, pp. 347-369.
44
45
46 572 Lamb, M.P., Hickson, J.G., Marr, B., Sheets, B., Paola, C., Parker, G., 2004. Surging and
47
48 573 continuous turbidity currents: Flow dynamics and deposits in an experimental
49
50 574 intraslope minibasin. *Journal of Sedimentary Research* 74, 148-155.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5 575 Lane-Serff, G.F., Beal, L.M., Hadfield, T.D., 1995. Gravity current flow over obstacles.
6
7 576 Journal of Fluid Mechanics 292, 39-53.
8
9
10 577 Matos, R.M.D., 2000. Tectonic evolution of the Equatorial South Atlantic. In: Mohriak, W.U.,
11
12 578 Talwani, M. (Eds.), Atlantic Rifts and continental margins, Geophysical Monograph,
13
14 579 115, American Geophysical Union, pp. 332-354.
15
16 580 Mutti, E., Tinterri, R., Remacha, E., Mavilla, N., Angella, S., Fava, L., 1999. An introduction
17
18 581 to the analysis of ancient turbidite basins from an outcrop perspective AAPG Course
19
20 582 Note 39, pp. 93.
21
22 583 Normandeau, A., Lajeunesse, P., St-Onge, G., 2015. Submarine canyons and channels in the
23
24 584 Lower St. Lawrence Estuary (Eastern Canada): Morphology, classification and recent
25
26 585 sediment dynamics. Geomorphology 241, 1-18.
27
28
29 586 Normandeau, A., Lajeunesse, P., Poiré, A.G., Francus, P., 2016. Morphological expression of
30
31 587 bedforms formed by supercritical sediment density flows on four fjord-lake deltas of
32
33 588 the south-eastern Canadian Shield (Eastern Canada). Sedimentology 63, 2106-2129.
34
35
36 589 Normark, W.R., Piper, D.J.W., Posamentier, H., Pirmez, C., Migeon, S., 2002. Variability in
37
38 590 form and growth of sediment waves on turbidite channel levees. Marine Geology 192,
39
40 591 23-58.
41
42
43
44 592 Paull, C.K., Ussler III, W., Caress, D.W., Lundsten, E., Covault, J.A., Maier, K.L., Xu, J. and
45
46 593 Augenstein, S., 2010. Origins of large crescent-shaped bedforms within the axial
47
48 594 channel of Monterey Canyon, offshore California. Geosphere 6, 755-774.
49
50
51
52 595 Paull, C.K., Caress, D.W., Ussler III, W., Lundsten, E., Meiner-Johnson, M., 2011. High-
53
54 596 resolution bathymetry of the axial channels within Monterey and Soquel submarine
55
56 597 canyons, offshore central California. Geosphere 7, 1077-1101.
57
58
59
60
61
62
63
64
65

- 598 Patacci, M., Haughton, P.D.W., McCaffrey, W.D., 2015. Flow behavior of ponded turbidity
1
2
3 599 currents. *Journal of Sedimentary Research* 85, 885-902.
4
- 5 600 Piper, D.J.W., Normark, W.R., 1983. Turbidite depositional patterns and flow characteristics,
6
7 601 Navy submarine fan, California Borderland. *Sedimentology* 30, 681-694.
9
- 10 602 Piper, D.J.W., Normark, W.R., 2009. Processes that initiate turbidity currents and their
11
12 603 influence on turbidites: a marine geology perspective. *Journal of Sedimentary*
13
14 604 *Research* 79, 347-362.
15
16
17
- 18 605 Pirmez, C., Beauboeuf, R.T., Friedmann, S.J., Mohrig, D.C., 2000. Equilibrium profile and
19
20 606 baselevel in submarine channels: examples from Late Pleistocene systems and
21
22 607 implications for the architecture of deepwater reservoirs. In: Weimer, P., Slatt, R.M.,
23
24 608 Coleman, J., Rosen, N.C., Nelson, H., Bouma, A.H., Styzen, M.J., Lawrence, D.T.,
25
26 609 (Eds.), *Deep water reservoirs of the world*, GCSSEPM Foundation, Houston (2000),
27
28 610 pp. 782-805.
29
30
31
32
- 33 611 Prather, B.E., Booth, J.E., Steffens, G.S., Craig, P.A., 1998. Classification, lithologic
34
35 612 calibration, and stratigraphic succession of seismic facies of intraslope basins, deep-
36
37 613 water Gulf of Mexico. *American Association of Petroleum Geologists Bulletin* 82,
38
39 614 701-728.
40
41
42
- 43 615 Prather, B.E., 2003. Controls on reservoir distribution, architecture and stratigraphic trapping
44
45 616 in slope settings. *Marine and Petroleum Geology* 20, 529-545.
46
47
48
- 49 617 Prather, B.E., Pirmez, C., Winker, C.D. 2012. Stratigraphy of linked intraslope basins:
50
51 618 Brazos-Trinity system Western Gulf of Mexico. In: Prather, B.E., Deptuck, M.E.,
52
53 619 Mohrig, D., van Hoorn, B., Wynn, R.B. (Eds.), *Application of the Principles of*
54
55 620 *Seismic Geomorphology to Continental -Slope and Base-of-Slope Systems: Case*
56
57
58
59
60
61
62
63
64
65

- 621 Studies from Seafloor and Near-Seafloor Analogues. SEPM Special Publication 99,
1
2 622 pp. 83-109.
3
4
5 623 Rebesco, M., Hernández-Molina, F.J., van Rooij, D., Wåhlin, A., 2014. Contourites and
6
7 624 associates sediments controlled by deep-water circulation processes: State of the art
8
9 625 and future considerations. *Marine Geology* 352, 111-154.
10
11
12
13 626 Reijnenstein, H.M., Posamentier, H.W., Bhattacharya, J.P., 2011. Seismic geomorphology and
14
15 627 high-resolution seismic stratigraphy of inner-shelf fluvial, estuarine, deltaic, and
16
17 628 marine sequences, Gulf of Thailand. *AAPG Bulletin* 95, 1959-1990.
18
19
20
21 629 Rijks, E.J.H., Jauffred, J.C.E.M., 1991. Seismic interpretation 29; attribute extraction; an
22
23 630 important application in any detailed 3-D interpretation study. *The Leading Edge* 10,
24
25 631 11-19.
26
27
28
29 632 Reynolds, D.J., Steckler, M.S., Coakley, B.J., 1991. The role of the sediment load in
30
31 633 sequence stratigraphy: The influence of flexural isostasy and compaction. *Journal of*
32
33 634 *Geophysical Research* B01914.
34
35
36
37 635 Sinclair, H.D., Tomasso, M., 2002. Depositional Evolution of Confined Turbidite Basins.
38
39 636 *Journal of Sedimentary Research* 72, 451-456.
40
41
42 637 Smith, R., 2004. Silled sub-basins to connected tortuous corridors: sediment distribution
43
44 638 systems on topographically complex sub-aqueous slopes. *Geological Society of*
45
46 639 *London Special Publications* 222, 23-43.
47
48
49
50 640 Smith, D.P., Kvitek, R., Iampietro, P.J., Wong, K., 2007. Twenty-nine months of geomorphic
51
52 641 change in upper Monterey Canyon (2002–2005). *Marine Geology* 236, 79-94.
53
54
55 642 Soutter, E.L., Kane, I.A., Huuse, M., 2018. Giant submarine landslide triggered by Paleocene
56
57 643 mantle plume activity in the North Atlantic. *Geology* 46, 511-514.
58
59
60
61
62
63
64
65

- 644 Symons, W.O., Sumner, E.J., Talling, P.J., Cartigny, M.J.B., Clare, M.A., 2016. Large-scale
1 sediment waves and scours on the modern seafloor and their implications for the
2
3 645 prevalence of supercritical flows. *Marine Geology* 371, 130-148.
4
5 646
6
7 647 Talling, P.J., Allin, J., Armitage, D.A., Arnott, R.W.C., Cartigny, M.J.B., Clare, M.A.,
8
9
10 648 Felletti, F., Covault, J.A., Girardclos, S., Hansen, E., Hill, P.R., Hiscott, R.N., Hogg,
11
12 649 A.J., Hughes Clarke, J.E., Jobe, Z.R., Malgesini, G., Mozzato, A., Naruse, H.,
13
14 650 Parkinson, S., Peel, F.J., Piper, D.J.W., Pop, E., Postma, G., Rowley, P., Sguazzini, A.,
15
16 651 Stevenson, C.J., Sumner, E.J., Sylvester, Z., Watts, C., Xu, J., 2015. Key future
17
18 652 directions for research on turbidity currents and their deposits. *Journal Sedimentary*
19
20 653 *Research* 85, 153-169.
21
22
23
24
25 654 Vail, PR., 1987. Seismic stratigraphy interpretation using sequence stratigraphy. Part I:
26
27 655 Seismic stratigraphy interpretation procedure. In *Atlas of Seismic Stratigraphy*, ed.
28
29 656 AW Bally, pp. 1-10. *Am. Assoc. Petrol. Geol. Stud. Geol. No. 27, Vol. 1.* 125 pp.
30
31
32
33 657 Winker, C.D., 1996. High-resolution seismic stratigraphy of a late Pleistocene submarine fan
34
35 658 ponded by salt-withdrawal mini-basins on the Gulf of Mexico continental slope. 28th
36
37 659 Annual Offshore Technology Conference, Proceedings, pp. 619-628.
38
39
40
41 660 Wynn, R.B., Stow, D.A.V., 2002. Classification and characterisation of deep-water sediment
42
43 661 waves. *Marine Geology* 192, 7-22.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)





Figure 3
[Click here to download high resolution image](#)




Figure 4
[Click here to download high resolution image](#)




Figure 5

[Click here to download high resolution image](#)




Figure 6
[Click here to download high resolution image](#)




Figure 7
[Click here to download high resolution image](#)




Figure 8
[Click here to download high resolution image](#)




Figure 9

[Click here to download high resolution image](#)

