

1 **A Benchmark Dataset of Agricultural Weather Stations over the Contiguous United** 2 **States for Evapotranspiration Applications**

3 **Authors**

4 Christian Dunkerly¹, John M. Volk¹, Sayantan Majumdar¹, Justin L. Huntington¹, Richard G. Allen²,
5 Christopher Pearson¹, Yeonuk Kim¹, Charles G. Morton¹, Blake A. Minor¹, Peter ReVelle¹, Ayse Kilic³,
6 Forrest Melton⁴, Adam J. Purdy^{4,5}, Todd G. Caldwell¹

7 **Affiliations**

- 8 1. Desert Research Institute, Division of Hydrologic Sciences, Reno, Nevada, USA
- 9 2. (retired) University of Idaho, Kimberly, Idaho, USA
- 10 3. University of Nebraska-Lincoln, School of Natural Resources, Lincoln, Nebraska, USA
- 11 4. NASA Ames Research Center, Moffett Field, California, USA
- 12 5. California State University, Monterey Bay, Seaside, California, USA

13 **Corresponding author:** John M. Volk (john.volk@dri.edu)

14 **Abstract**

15 Agricultural weather data are fundamental for the accurate estimate of evapotranspiration (ET), irrigation
16 scheduling, and water-use accounting. In particular, reference ET provides a standardized atmospheric
17 demand for water loss from a hypothetical well-watered grass (ET_o) or alfalfa (ET_r); however, weather
18 stations may not adequately represent such climatic conditions. Weather data commonly contain errors
19 from poor siting, sensor drift, and network management deficiencies. No standardized dataset exists over
20 the contiguous United States (CONUS). Systematic errors affect ET_o/ET_r calculations and derived
21 products. Notably, satellite-based platforms like OpenET require agricultural weather data to bias correct
22 gridded reference ET to interpolate between satellite overpasses. CONUS-AgWeather is a benchmark
23 dataset of daily agricultural weather data (precipitation, solar radiation, air temperature, humidity, wind
24 speed, ET_o, ET_r) from 793 stations. This dataset contains 4,191,808 days (11,484 station-years, 1981-
25 2020) and was produced through standardized and systematic quality control procedures and open-source
26 software packages for time series inspection, outlier detection, corrections, and ET_o/ET_r calculations.
27 CONUS-AgWeather is intended primarily to support OpenET in the Western U.S. but has broader
28 applications.

29 **Background & Summary**

30 The sustainable management of water resources through measurement and optimization of agricultural
31 water use and crop productivity increasingly requires accurate and timely weather information. Weather
32 data, such as solar radiation, air temperature, humidity, wind speed, and precipitation, are fundamental
33 inputs for a multitude of agricultural applications. These data are necessary for calculating reference
34 evapotranspiration (ET), a key determinant of crop water requirements, which directly informs irrigation
35 system design, scheduling, and water allocation decisions^{1,2}. Beyond reference ET, weather data can be
36 directly used to assess land surface-atmospheric boundary layer feedbacks and regional actual ET^{3,4}.
37 Furthermore, high-quality weather data serve as essential direct input or ground-truth for calibration and
38 validation of hydrological models⁵, gridded weather datasets⁶⁻⁸, and satellite-based ET products used for

39 crop water use monitoring and reporting⁹⁻¹⁸.


40 Weather conditions, such as solar radiation, temperature, humidity, and wind speed, are closely linked to
41 land surface conditions. For instance, the air above irrigated croplands is typically cooler and more humid
42 than surrounding areas because a larger portion of net radiation is consumed by ET rather than surface
43 heating¹⁹. Consequently, weather stations located within croplands provide critical information for
44 accurately characterizing agricultural weather conditions and total evaporative demand. Despite their
45 importance, no unified national-scale agricultural weather station network or dataset exists in the United
46 States (U.S.), stations are sparse, and public data access is limited when compared to the assortment of
47 National Weather Service (NWS) stations located within towns, cities, roadways, and airport locations.
48 NWS stations typically collect select weather data of air temperature and precipitation, with airport
49 locations also collecting humidity and wind speed often at heights of 10 m or greater. Solar radiation
50 measurements are often limited to state mesonets only. Agricultural weather stations typically measure
51 the full suite of meteorological variables required to compute physically based evaporative demand or
52 standardized reference ET for short grass (ET_o) or tall alfalfa (ET_r). However, fragmented funding and
53 limited operations and maintenance often limit rigorous quality assurance, resulting in datasets that
54 contain random and systemic errors. Sensor degradation over time, calibration drift, physical obstructions,
55 inadequate maintenance, communication errors, and non-ideal station siting (e.g., insufficient fetch or
56 deviation from well-watered conditions) can all introduce errors and biases into the observational record
57²⁰⁻²². If unaddressed, these data quality issues can lead to erroneously high calculations of crop water
58 requirements, ET, flawed irrigation project design, irrigation scheduling, and ultimately, suboptimal crop
59 and water resource management^{20,21}.

60 To address these challenges, observational weather datasets should be subject to robust quality control
61 (QC) procedures before use. Such procedures aim to systematically identify and correct or remove
62 erroneous data, thereby enhancing the overall integrity and reliability of the dataset. While various QC
63 methodologies exist, the development and application of a consistent, transparent, and accessible
64 framework for QC of daily agricultural weather data has been a persistent need. The open-source
65 *agweather-qaqc* Python package²³ was developed to meet this need, offering a command-line interface
66 (CLI) tool that facilitates reading, visualization, and comprehensive QC of daily weather observations
67 from diverse sources, followed by the calculation of reference ET using standardized methods and
68 guidelines by the American Society of Civil Engineers and Environmental and Water resources Institute
69 (ASCE-EWRI)²⁴.

70 **Data Overview**


71 This paper presents CONUS-AgWeather, a high-quality benchmark dataset of daily weather and
72 reference ET variables, compiled from 793 agricultural weather stations from 19 networks across the
73 contiguous U.S. (CONUS) (Fig. 1). CONUS-AgWeather is the result of applying rigorous QC procedures
74 and standardized data processing within the *agweather-qaqc* package, along with visual inspections of all
75 weather station site locations to ensure stations are free from nearby obstructions (e.g., trees, buildings,
76 etc.), located within agricultural areas, and generally suitable for calculating reference ET^{22,24}. The
77 primary purpose of CONUS-AgWeather is to support and advance ET and agricultural water resources
78 research and applications using high-quality, consistent, and transparent weather and reference ET data.
79 Although CONUS-AgWeather was initially developed to support the OpenET^{15,16} project with a

80 particular focus on the Western U.S., this benchmark dataset is also valuable for validation and bias-
81 correction of gridded or modeled weather datasets, and potential integration within weather forecast,
82 reanalysis, land data assimilation systems, hydrologic models, and satellite remote sensing ET models to
83 improve and support agricultural and water management.



84
85 **Fig 1.** Map showing the distribution of the initial 1,078 agricultural weather stations, of which 793 were
86 included and 285 were removed after site evaluation and data QC, overlaid on major Köppen-Geiger
87 (KG) climate zones²⁵: cold and hot semi-arid steppe (Bsk + Bsh); hot and cold desert (Bwh + Bwk);
88 humid subtropical (Cfa); hot- and warm-summer Mediterranean (Csa + Csb); and hot- and warm-summer
89 humid continental (Dfa + Dfb). Here, stations over white no data areas are in other KG climate zones.
90 Note that CONUS-AgWeather was initially developed to support the OpenET^{15,16} project, with a
91 particular focus on the Western U.S, resulting in a higher weather station density compared to the Eastern
92 U.S.

93 The CONUS-AgWeather dataset includes a total of 4,191,808 days (11,484 years) of data spanning May
94 21, 1981, to December 31, 2020, with record lengths varying by each station. Fig. 1 illustrates stations
95 that were included and removed after the QC process and visual site inspections, along with the major
96 Köppen-Geiger (KG) climate zones²⁵. Fig. 2a shows included stations, colored by the number of valid
97 daily weather observations for each station after data QC. In addition, the number of years and the
98 average annual station record completeness of the available weather data are highlighted in Fig. 2b and
99 Fig. 2c, respectively. Here, the average annual record completeness (expressed in percentage) for an
100 individual station is defined as the ratio of the number of available weather observations to the total
101 number of days the station was active within that calendar year. This method normalizes for partial years
102 at the beginning or end of the record by using the actual period of record as the denominator rather than a
103 fixed 365/366-day calendar year. Note, there are two California Irrigation Management Information
104 System (CIMIS)²⁶ stations (currently inactive) located in Mexicali, Mexico and San Luis Río Colorado,
105 Mexico that were included given close proximity to the U.S. boarder and quality long-term records.



106
107
108

Fig. 2. The complete, quality-controlled CONUS-AgWeather station locations showing (a) the number of weather observations, (b) years of weather observations, and (c) the average annual record completeness

109 for each station. This dataset includes 793 stations with a total of 11,484 years, i.e., 4,191,808 days of
110 valid weather data across the CONUS, spanning from May 21, 1981, to December 31, 2020 (record
111 lengths vary by station).

112 **Methods**

113 Development of the CONUS-AgWeather dataset began with the acquisition of raw weather data from
114 1,078 weather stations illustrated in Fig. 1. Raw weather data underwent extensive QC data processing
115 using the *agweather-qaqc* Python package²³, with visual inspection of data and site conditions as
116 described below, resulting in 793 station datasets illustrated in Fig 1. While this is not an exhaustive list,
117 we present these stations as the first round of available agricultural weather sources for the Western U.S.
118 needed, foremost to bias correct gridded data sets used in the current Version 2 of OpenET^{15,16}. Future
119 efforts will expand this analysis to the Eastern U.S.

120 **Data Sources**

121 Agricultural weather station data used in the development of the CONUS-AgWeather dataset were
122 sourced from 22 networks operating throughout the CONUS. These include the U.S. Bureau of
123 Reclamation's AgriMet Network, Pacific Northwest and Great Plains regions²⁷, Arizona Meteorological
124 Network (AZMET)²⁸, CIMIS²⁶, Colorado Agricultural and Meteorological Network (CoAgMET)²⁹,
125 Enviroweather Network³⁰, Florida Automated Weather Network (FAWN)³¹, Georgia Automated
126 Environmental Monitoring Network (GAEMN)³², High Plains Regional Climate Center (HPRCC)³³,
127 Missouri Mesonet³⁴, North Dakota Agricultural Weather Network (NDAWN)³⁵, Nebraska Mesonet³⁶,
128 Nevada Integrated Climate and Evapotranspiration Network (NICE Net)³⁷, Oklahoma Mesonet³⁸, U.S.
129 Department of Agriculture (USDA) Soil Climate Analysis Network (SCAN)³⁹, South Dakota Mesonet⁴⁰,
130 National Oceanic and Atmospheric Administration's United States Climate Reference Network (USCRN)
131⁴¹, West Texas Mesonet⁴², Western Regional Climate Center (WRCC)⁴³, New Mexico's ZiaMet⁴⁴, Utah
132 Climate Center (UCC)⁴⁵, USDA Agricultural Research Service (USDA-ARS)⁴⁶, and Wyoming
133 Agricultural and Climate Network (WACNet)⁴⁷. Data were primarily acquired at a daily temporal
134 resolution. If data were acquired at higher temporal resolution (e.g., hourly), data were resampled to daily
135 timesteps following standardized methods prior to ingestion into the QC process²⁴. Table 1 shows the
136 number of stations present in the initial dataset (1,078) and the number of stations that were included
137 (793) in the final CONUS-AgWeather dataset after the QC process.

138 Weather data is collected for a variety of reasons for a variety of stakeholders and not all weather stations
139 are suitable for reference ET^{19,48}. A mesonet is a network of automated, fixed, surface weather observing
140 stations with a spatial density of ~one station per 1000 km² that monitors environmental variables in the
141 vertical domain between 10 m above and 1 m below ground surface and provides high data quality and
142 reliable near-real time weather data⁴⁹. Such stations have the specific objective of collecting observations
143 that are representative of the mesoscale environment on the scale of 3–100 km, whereas an agricultural
144 meteorological station provides detailed information on the very lowest layer of the atmosphere that may
145 include soil temperature, soil moisture, and ET_o⁵⁰. As such, many stations are not sited specifically
146 within the microclimate of agricultural footprint for the accurate calculation of ET^{19,48}.

147

148 **Table 1.** Network information for agricultural weather stations present in the initial dataset (1,078
149 stations) and the final CONUS-AgWeather dataset (793 stations).

Network	Initial Number of Stations	Number of Stations Removed	Number of Stations Included	Access Date (yyyy-mm-dd)
AgriMet, Columbia-Pacific Northwest Region ²⁷	132	45	87	2021-01-10
AgriMet, Missouri Basin ²⁷ Region	25	0	25	2020-02-14
AZMET ²⁸	29	4	25	2021-01-15
CIMIS ²⁶	161	23	138	2019-06-10
CoAgMET	101	33	68	2021-01-09
GAEMN ³²	19	0	19	2020-04-17
HPRCC ³³ †	249	33	216	2020-07-10
Missouri Mesonet ³⁴	37	6	31	2020-02-27
NICE Net ³⁷	19	6	13	2019-05-10
Oklahoma Mesonet ³⁸	120	64	56	2019-10-17
SCAN ³⁹	56	9	47	2021-04-17
USCRN ⁵¹	24	2	22	2020-03-20
WACNet ⁴⁷	17	3	14	2021-01-10
Other ‡	90	58	32	2021-01-20
Grand Total	1,078	285	793	

150 † Includes stations from HPRCC ³³, NDAWN ³⁵, Nebraska Mesonet ³⁶, and South Dakota Mesonet ⁴⁰.

151 ‡ Includes stations from Enviroweather ³⁰, FAWN ³¹, West Texas Mesonet ⁴², WRCC ⁴³, New Mexico's
152 ZiaMet ⁴⁴, UCC ⁴⁵, and USDA-ARS ⁴⁶.

153 **Data Processing and Quality Control**

154 The *agweather-qaqc* package ²³ was used for data visualization, screening, automated and manual QC,
155 and calculation of reference ET. It includes modules to ingest data from tabular files (e.g., CSV) and a
156 corresponding configuration file, which specifies station metadata (e.g., latitude, longitude, elevation,
157 anemometer height), weather variables, and respective units within the input data file. *agweather-qaqc*
158 was developed to handle most common input weather variables, units, and data formats, standardizing
159 variables, QC, and calculations for consistency and reproducibility as described below.

160 Data Pre-processing

161 Several pre-processing steps were performed using *agweather-qaqc*: 1) raw data were read and variable
162 names were standardized across all station files, 2) meteorological variables were converted into units
163 compliant with ASCE-EWRI standardized reference ET calculations (e.g., air temperature to °C, solar
164 radiation to MJ m⁻², vapor pressure to kPa, wind speed to m s⁻¹), and 3) data were systematically screened

165 and removed due to physical limits and obvious erroneous values. Examples include negative
166 precipitation, negative wind speed, daily total solar radiation values at or near zero during expected
167 daylight hours, or air temperature readings that fall outside extreme, historically plausible ranges for the
168 station's location (see the Technical Validation section).

169 Data Quality Control

170 The QC procedures applied to solar radiation, air temperature, humidity, and wind speed were applied
171 based on the established guidelines described below^{20,24,52}.

172 *Solar Radiation (R_s)*

173 For each day, theoretical clear sky solar radiation, R_{s0} , was calculated following ASCE-EWRI guidelines
174²⁴. R_{s0} represents the theoretical maximum daily incoming solar radiation a station received under clear
175 sky conditions, based on station latitude, elevation, day of the year, and atmospheric water vapor content
176 (derived from humidity data). R_s data were compared to R_{s0} and values that significantly exceed R_{s0}
177 (commonly due to data logger or sensor electrical issues) were removed²³. Along with progressive
178 calibration drift, solar radiation sensors are subject to maintenance issues related to the sensor being out-
179 of-level and dust or debris on the optics⁴⁹. To identify and correct R_s drift or anomalously low values due
180 to temporary obstructions, R_s was systematically compared to R_{s0} and adjusted based on the ratio of R_s
181 to R_{s0} ²³. The R_s record was divided into 60-day periods, and a percentile correction factor (CF_P) was
182 calculated and applied to all R_s data within each period based on the assumption that observed R_s should
183 approach R_{s0} on the clearest days. For each 60-day period, CF_P was calculated as the ratio of the average
184 R_{s0} to the average R_s as:

$$185 \quad CF_P = \frac{\overline{R_{s0P}}}{\overline{R_sP}}$$

186
187 where $\overline{R_{s0P}}$ and $\overline{R_sP}$ are the average R_{s0} and R_s values, respectively, for the selected clearest days in
188 period P. For the CONUS-AgWeather dataset, respective R_s and R_{s0} data within top 10th percentile of a
189 60-day period (i.e., 6 days) were selected and used to compute CF_P . This 60-day period was used to
190 account for seasonal variations at the stations while remaining responsive to potential sensor drift. CF_P
191 values were then multiplied by all R_s values within each respective 60-day period. If CF_P ranged from
192 0.97 to 1.03, no adjustment was applied, and if CF_P was <0.5 or >1.5 , R_s data for that period were
193 deemed erroneous and were removed.
194

195 *Air Temperature (T_{max} , T_{min})*

196 Temperature data were quality controlled using a modified Z-score approach, as detailed by Iglewicz and
197 Hoaglin (1993)⁵³. This method is minimally influenced by outliers, particularly in smaller samples, as it
198 utilizes the median and median absolute deviation (MAD)⁵⁴, i.e., $\text{median}_i(x_i - \tilde{x})$, where \tilde{x} is the
199 median of the sample observations x_i . The modified Z-score (M_i) for each observation x_i was then
200 computed as:

$$201 \quad M_i = \frac{0.6745(x_i - \tilde{x})}{\text{MAD}}$$

202

203 Here, observations with $|M_i| > 3.5$ were flagged as potential outliers and removed, as recommended by
204 Iglewicz and Hoaglin (1993)⁵³.

205 *Relative Humidity (RH_{max} , RH_{min})*

206 Daily RH_{max} should approach or reach 100% on at least a few days a year, usually coinciding with
207 precipitation events, early morning condensation or dew, or periods of high atmospheric moisture due to
208 irrigation-driven increases in ET^{1,21}. This is particularly true for an agricultural weather station to meet
209 the requirements of reference ET; however, less expensive capacitive hygrometers common to most
210 networks lose accuracy above 95%⁵⁵. To correct for sensor drift and inaccuracies, a yearly correction
211 factor ($CF_Y = 100/\overline{RH_{maxY}}$) was calculated, where $\overline{RH_{maxY}}$ is the average of the top 1%, or the 3 highest
212 values if sufficient data exist, of the highest RH_{max} values within year Y. This factor was derived by
213 comparing the topmost percentile of observed RH_{max} values within a calendar year to the expected 100%.
214 CF_Y was then applied to all RH_{max} and RH_{min} observations for that year. It should be noted that the CF_Y
215 also adjusts the observed values down to be in line with the expected 100% when the sensor has a high
216 bias over 100%. The number of points used for this calculation was adjusted for years with incomplete
217 data records.

218 *Wind Speed (u_Z)*

219 If wind speed (u_Z) was measured at an anemometer height (Z , in meters) other than the standard 2-meter
220 reference height, it was adjusted to u_{2m} using the logarithmic wind profile equation as specified by
221 ASCE-EWRI (2005)²⁴:

222

223

$$u_{2m} = u_Z \frac{4.87}{\ln(67.8Z - 5.42)}$$

224 The primary QC for wind speed relied on manual inspection of the interactive time series plots generated
225 by *agweather-qaqc*. We investigated wind speed patterns, such as trends or rapid changes in wind speed
226 (e.g., due to failing anemometer bearings, nearby obstructions, tree growth, etc.), prolonged periods of
227 zero or constant wind speed, or values that appear implausibly high or low relative to the typical wind
228 regime of the station and or nearby stations. Such identified periods were flagged and removed.

229 *Precipitation*

230 Precipitation data were screened for erroneous values (e.g., negative values) and extremely large events
231 (e.g., daily precipitation greater than 610 mm). These were removed during the initial data processing
232 phase. Additional visual inspection was performed to identify implausible values based on the
233 environment and historical record of the station, using approaches similar to those detailed in Durre et al.
234 (2010)⁵⁶.


235 **Gap-Filling**

236 Gap-filling was not performed in the CONUS-AgWeather dataset beyond creating a complete record of
237 R_{s0} to support QC observed R_s . The *agweather-qaqc* package incorporates optional routines for gap-

238 filling missing data in the daily records²³, but these were not applied to the CONUS-AgWeather dataset.

239 Calculation of Reference Evapotranspiration

240 Following the comprehensive QC, the daily grass (i.e., short) reference ET (ET_o) and alfalfa (tall)
241 reference ET (ET_r) were calculated according to the ASCE Standardized Penman-Monteith (ASCE-PM)
242 equation²⁴ using *RefET* Python library⁵⁷. For stations where vapor pressure was not available directly, it
243 was derived from dew point temperature or relative humidity data present in the record following
244 recommendations and guidelines of ASCE-EWRI (2005)²⁴. A histogram of daily ET_o data for all 793
245 stations, and violin plots of ET_o by KG climate zones²⁵, is illustrated in Fig. 3.



246 **Fig. 3** Histogram of daily ET_o in the CONUS-AgWeather dataset with corresponding violin plot
247 distributions across major KG climate zones²⁵: cold and hot semi-arid steppe (Bsk + Bsh); hot and cold
248 desert (Bwh + Bwk); humid subtropical (Cfa); hot- and warm-summer Mediterranean (Csa + Csb); and
249 hot- and warm-summer humid continental (Dfa + Dfb). ‘Other’ includes all other climate zones outside
250 the five major zones listed.
251

252 Output Generation and Data Archiving

253 The CONUS-AgWeather dataset includes comprehensive output that include processing log files,
254 interactive time series plots, and tabular data files, as described below:

255 a) Log Files: Detailed, human-readable log files for each station. These files record all automated QC
256 checks, user-initiated corrections, and parameters used for adjustments of data during the processing.

257 b) Interactive Plots: Interactive time series plots (rendered using Bokeh⁵⁸) that display both the pre-QC
258 (original) and post-QC (corrected) data for solar radiation, air temperature, humidity, wind speed, and
259 precipitation. These plots, saved as standalone HTML files for easy sharing and archiving, are invaluable
260 for visual assessment of data quality, the impact of applied corrections, and for facilitating manual review.

261 c) Data Files: The primary output for each station consists of structured data files (Microsoft Excel

262 spreadsheets) containing the fully quality-controlled observations, the numerical difference (delta values)
263 between original and corrected data where adjustments were made.

264 In summary, CONUS-AgWeather output is a collection of individual station log files, interactive plots,
265 and pre- and post-QA data files for all 793 weather stations, in the format text files, Bokeh HTML, and
266 Microsoft Excel spreadsheets, respectively.

267 **Data Records**

268 Each station record within the CONUS-AgWeather dataset includes the following 23 standardized
269 variables, with the ones in bold font used in the ASCE-PM reference ET²⁴ calculation:

- 270 1) Station ID: Unique identifier for the weather station (<Number>_<State abbreviation>, e.g.,
271 001_AR).
- 272 2) Latitude: Latitude of the weather station in decimal degrees (0.0001°). Some networks report at a
273 reduced precision to protect the privacy of the landowner.
- 274 3) Longitude: Longitude of the weather station in decimal degrees (0.0001°). Some networks report at a
275 reduced precision to protect the privacy of the landowner.
- 276 4) Elevation: Elevation of the weather station in meters.
- 277 5) Date: Observation date (YYYY-MM-DD).
- 278 6) **TMax**: Daily maximum air temperature (°C).
- 279 7) **TAvg**: Daily average air temperature (°C), computed as the average of the daily maximum and
280 minimum temperatures.
- 281 8) **TMin**: Daily minimum air temperature (°C).
- 282 9) **Ea**: Daily mean vapor pressure (kPa), either directly from the QC'd record or calculated using
283 *agweather-qaqc*. If Ea was calculated, *agweather-qaqc* used only the most preferred form of humidity
284 data available as specified in the ASCE-EWRI standard²⁴. Note that preferred humidity data was
285 QC'd prior to calculation of Ea if Ea was not provided directly (e.g., when only RHMax and RHMin
286 are provided in the observational record).
- 287 10) TDew: Daily average dew point temperature (°C), either from the QC'd record or calculated.
- 288 11) RHMax: Daily maximum relative humidity (%). This variable may not be present if it is not provided
289 by the weather station network. RHMax is used in the ASCE-PM reference ET²⁴ calculation only if
290 Ea or TDew are not provided in the observational record.
- 291 12) RHAvg: Daily average relative humidity (%). This variable may not be present if it is not provided by
292 the weather station network. RHAvg is only used in the ASCE-PM reference ET²⁴ calculation if no
293 other sources of humidity observations are provided in the record.
- 294 13) RHMin: Daily minimum relative humidity (%). This variable may not be present if it is not provided
295 by the weather station network. RHMin is used in the ASCE-PM reference ET²⁴ calculation only if
296 measured Ea or TDew are not provided in the observational record.
- 297 14) Compiled Ea: Daily mean water vapor (kPa). This is an aggregate record that uses all forms of
298 humidity data present, not just the most preferred, to calculate as complete a record of Rso as possible
299 for the QC of Rs.
- 300 15) **Rs**: Average daily incoming shortwave solar radiation (W m⁻²).
- 301 16) Optimized TR Rs: Optimized Thornton-Running solar radiation (W m⁻²), which computes a modeled
302 version of incoming shortwave radiation according to Thornton and Running's equation⁵⁹. This
303 variable is calculated to provide the option of a full record of solar radiation but is separate from

- 304 QC'd observations of solar radiation.
305 17) **Rso**: Computed clear-sky solar radiation (W m^{-2}) for QC of observed Rs. The values are computed
306 using station latitude and equations that model the effects of precipitable water in the atmosphere on
307 incoming solar radiation, as described in the ASCE-EWRI (2005)²⁴.
308 18) Measured Uz: Daily average wind speed at the actual height of the anemometer (m s^{-1}).
309 19) Anemometer Height: Height of the anemometer at the station (m).
310 20) **Uz at 2m**: Daily average wind speed adjusted to 2m height (m s^{-1}).
311 21) Precipitation: Daily total precipitation (mm).
312 22) ETo: Daily grass reference ET (mm).
313 23) ETr: Daily alfalfa reference ET (mm).

314 **Data Format and Availability**

315 The CONUS-AgWeather dataset⁶⁰ is available as a zip file, containing Microsoft Excel spreadsheets for
316 individual stations (<https://doi.org/10.5281/zenodo.18122157>). Detailed metadata with station-specific
317 QC notes, human readable plaintext log files, and interactive HTML Bokeh⁵⁸ plots of the pre-and post-
318 QC datasets. The period of record for each station varies, and this information is included in the metadata.

319 **Technical Validation**


320 The technical quality and reliability of the CONUS-AgWeather dataset was established through the
321 systematic, standardized, and reproducible QC procedures within the *agweather-qaqc* Python package²³.
322 Visual inspection of individual site locations using satellite imagery to confirm station-siting in
323 agricultural land was the first hurdle, then weather data time series further enhanced the validation of the
324 final dataset.

325 After the initial ingestion of weather station data, the records were carefully assessed for key variables
326 required to compute the ASCE-PM reference ET²⁴. These variables included air temperature, solar
327 radiation, wind speed, and humidity. Each weather station was evaluated based on its availability of data
328 for all four variables. Stations that did not have at least two years of continuous, quality data during the
329 growing season for all four variables were removed, resulting in a reduction of 21 stations from the initial
330 dataset of 1,078 stations.

331 We visually inspected the environment surrounding each weather using current and historical imagery
332 from Google Earth and Google Street View, to ensure the station was located within well-watered
333 agricultural areas, per recommendations and guidelines of ASCE-EWRI (2005)²⁴ (Fig. 4a-d). Poor station
334 siting was carefully considered, specifically stations located in urban or non-agricultural areas, and those
335 possibly affected by obstructions (e.g., trees, buildings) and microclimates (e.g., water bodies, barren
336 areas, urban heat), were flagged and removed. Location and visual screening resulted in a further
337 reduction of 264 stations, i.e., a total of 285 stations were removed from the initial dataset of 1,078
338 stations. Fig. 1 and Fig. 2 illustrate the spatial distribution of initial (1,078 stations) and final (793
339 stations) CONUS-AgWeather data, respectively.

340 Technical validation of automated statistical QC using *agweather-qaqc*²³ (detailed in the Methods
341 section) corrected or flagged suspicious time series data. Manual visual inspection of flagged data
342 identified any trends or abrupt shifts related to sensor malfunction or data logging errors. Stations with


343 frequent, suspect, or unresolved data quality issues were removed from the dataset.



344


345 **Fig. 4 (a)** show satellite and **(b)** street views for a ZiaMet⁴⁴ station in Los Lunas, New Mexico that was
346 removed from the CONUS-AgWeather dataset because of poor station siting with many obstructions
347 (e.g., trees) affecting measurements. In contrast, **(c)-(d)** show U.S. Bureau of Reclamation AgriMet²⁷
348 station (465_MT) in Glasgow, Montana, that is well-sited in an area surrounded by agriculture, with the
349 nearest structures being greater than 76 m (250 ft) away.

350 For example, common errors in measured R_s from pyranometers include calibration drift, improper
351 leveling, sensor degradation, or temporary obstructions from dust and debris^{22,52}. Measured R_s should
352 approach or slightly exceed the theoretical clear-sky solar radiation (R_{so}) at least a few days a season,
353 particularly in the western U.S. For example, the measured maximum daily R_s frequently approaches R_{so}
354 across all seasons at a NICE Net station in Reno, NV, as expected for this arid site location, but begins to
355 drift consistently lower than R_{so} in summer of 2014 (Fig. 4a). From 2014 onward, maximum daily R_s
356 was consistently lower than R_{so} , indicating sensor drift that was corrected to measured R_s during the QC
357 process (Fig. 4b) using 60-day correction factors (Fig. 4c). As part of the technical validation process,
358 time series plots of pre- and post-corrected R_s as well as percent change values, as illustrated in Fig. 5,
359 were visually inspected for every station to ensure corrections were justified and sensible. Automated
360 corrections to R_s made with *agweather-qaqc* are based on manual R_s corrections^{20,24,52}; however,
361 *agweather-qaqc* is less subjective, more reproducible, and fully documented in the CONUS-AgWeather
362 dataset.



363
 364 **Fig. 5 (a)** Original daily shortwave radiation (Rs) measurements and computed clear-sky solar radiation
 365 (Rso) at a NICE Net³⁷ station (635_NV) located in Reno, Nevada, illustrating consistent sensor drift
 366 beginning in summer of 2014. **(b)** Corrected daily Rs after QC adjustment using Rso as a limit. **(c)**
 367 Percent change between the original (pre-QC) and corrected (post-QC) Rs values for each sixty-day
 368 period (i.e., $((\text{corrected} - \text{original}) / \text{original}) * 100$)²³. Note that Rso is not corrected.

369 Capacitive humidity-sensing elements, common in combined air temperature and humidity sensors, drift
 370 over time and should be replaced every two to three years; however, that is often not practical in network
 371 operations. As shown in Fig. 6a, measured daily maximum relative humidity (RHMax) shows a clear
 372 trend starting near 100% and decreasing to ~80% over 7 years. RHMin is also decreasing through time at
 373 a similar rate. Fig. 6c shows the percent change in pre- and post-corrected RH as a result of yearly
 374 correction factors computed and applied by *agweather-qaqc* to both RHMax and RHMin, resulting in RH
 375 data that are within expected limits and are free from drift artifacts (Fig. 6b).



376
377 **Fig. 6** (a) Daily relative humidity maximum (RHMax) and minimum (RHMin) from a SCAN ³⁹ station
378 (1069_MT) in Sidney, Montana with pronounced sensor drift, (b) corrected data after a year-based
379 percentile adjustments, and (c) the percent change for both.

380 As described and illustrated above, manual site location and data time series inspections, along with the
381 automated QC procedures using *agweather-qaqc* are foundational to the CONUS-AgWeather dataset and
382 achieving benchmark data quality. Our technical validation approach focused on adherence to scientific
383 best practices based on standardized methods and guidelines. The following elements summarize our
384 technical validation:

385 a) Standardized and Documented Procedures: The application of consistent and standardized QC rules
386 and algorithms across all stations and variables, based on widely accepted meteorological and agricultural
387 engineering principles ^{20,24,61}. All QC procedures are documented and reproducible within the *agweather-*
388 *qaqc* package ²³.

389 b) Physically-Constrained Corrections: Adjustments are constrained by physical limits, expected values,
390 and well-established practices in the meteorological domain ^{20,52}. For example, Rs corrections are
391 calculated and limited by R_{so} , and RH adjustments consider the likelihood of relative water vapor
392 saturation in well-watered reference crop conditions ^{20,52}.

393 c) Robust Statistical Outlier Detection: The use of physical limits for Rs and RHMax/RHMin, and non-
394 parametric outlier detection methods, such as the modified Z-score for temperature data based on median

395 absolute deviation about the sample median (i.e., MAD), ensuring minimal influence of outliers when
396 statistically identifying and removal of outliers^{53,54}.


397 d) Visual Inspection and Expert Review: Remote visual inspection of site locations based on street view,
398 aerial and satellite imagery (Fig. 4), and generation of interactive time series plots for visualization and
399 expert review are a cornerstone of the QC process and *agweather-qaqc* output. These images and plots
400 allow for expert review, which is crucial for identifying subtle data quality issues that automated
401 algorithms often miss (e.g., anomalously low humidity for agricultural conditions, local obstructions
402 affecting wind speed and precipitation) and for verifying automated corrections (Fig. 5 and Fig. 6).

403 e) Quantifiable Impact of QC: The necessity and impact of QC steps and corrections can be substantial.
404 For instance, uncorrected R_s (Fig. 5a) directly impacts reference ET calculations. Comparing ETo data
405 values between the original and QC'd observations can result in substantial differences, potentially by a
406 factor (QC / original) of only ~ 0.03 (station: 814_SD, 1992-01-07) to ~ 581.84 (station: 956_WY, 2017-
407 04-29) and ~ 0.76 (station: 824_SD, 1984) to ~ 1.28 (station: 014_AZ, 2014) for daily and annual time
408 steps, respectively, with the QC'd values being higher or lower than the original data depending on the
409 underlying cause^{20,52}.

410 While the above factors represent the extreme ends of the daily and annual QC / original ETo
411 distributions, Fig. 7 illustrates (1st–99th percentile) the ETo change across 504 (out of 793) CONUS-
412 AgWeather stations with complete annual records, where QC was applied (i.e., post-QC / pre-QC $\neq 1$).
413 QC corrections resulted in a median increase of $\sim 0.8\%$ in daily ETo values, with a mean increase of \sim
414 1% , indicating that the original station data tended to slightly underestimate ETo prior to correction. The
415 distribution is slightly right skewed, with most corrections falling within $\pm 5\%$ of the original values.
416 Notably, $\sim 36\%$ of daily observations (345,495) required no correction (i.e., post-QC / pre-QC = 1), while
417 the remaining $\sim 64\%$ of site-days (606,270) showed measurable QC adjustments as highlighted in Fig. 7a.
418 At the annual scale (Fig. 7b), the aggregated effect of daily corrections is dampened, with a median
419 percent change of $\sim 0.5\%$ and mean of $\sim 0.8\%$, as positive and negative daily corrections offset each other
420 over the year.

421 In addition, Figure 7 reveals modest variations in corrected ETo across KG climate zones²⁵, though all
422 climate classes show positive and negative corrections centered near zero. Stations located in humid
423 subtropical (Cfa) regions exhibited the highest variability and largest positive QC corrections, with a
424 mean daily percent change in ETo due to QC of $+2.02 \pm 3.61\%$ and annual change of $+2.00 \pm 2.15\%$. In
425 contrast, stations in arid desert climates (Bwh + Bwk) showed minimal QC effects with near-zero mean
426 corrections and the lowest variability (daily: $+0.08 \pm 3.00\%$, annual: $-0.09 \pm 1.13\%$), indicating more
427 stable measurement conditions in these environments.


428 Results demonstrate that while individual daily QC corrections can be substantial, their cumulative annual
429 impact on ETo estimates are relatively modest, typically within $\pm 2\text{--}3\%$ for most stations. This indicates
430 that supporting the robustness of the QC methodology generally preserves the majority of the original
431 observations while correcting for sensor drift, data gaps, and measurement anomalies^{20,52}.



432 **Fig. 7** Distributions of (a) daily and (b) annual ETo change due to QC (in % change) across major KG
 433 climate zones²⁵ based on 504 (out of 793) CONUS-AgWeather stations with complete annual records.
 434 Left panels show histograms and kernel density estimates; right panels show violin plots stratified by KG
 435 climate zones: cold and hot semi-arid steppe (Bsk + Bsh); hot and cold desert (Bwh + Bwk); humid
 436 subtropical (Cfa); hot- and warm-summer Mediterranean (Csa + Csb); and hot- and warm-summer humid
 437 continental (Dfa + Dfb). ‘Other’ includes all other climate zones outside the five major zones listed.
 438 Reference lines indicate no change (0%, red dashed), median (blue solid), mean (green dotted), and ± 1
 439 standard deviation (green shaded region). Data are trimmed to the 1st–99th percentile range for
 440 visualization, while sample sizes reflect all QC-corrected records prior to trimming.

441 f) Achieving Benchmark Quality: The overarching goal of the comprehensive QC process was to produce
 442 a dataset of benchmark quality. This benchmark dataset can serve as a reliable reference for developing,

443 validating, and bias correcting ET and meteorological products (e.g., satellite-derived ET datasets,
 444 gridded weather data, and hydrologic models), for calculating accurate crop and irrigation water
 445 requirements, and for integration within water resource decision making processes. For example, the



446 **Fig. 8.** The CONUS-AgWeather dataset was validated using independent measurements from 79 flux
447 stations¹⁴ and used to bias-correct gridMET⁶². Scatter plots comparing monthly uncorrected and bias-
448 corrected gridMET ETo reveal substantial improvements in error metrics across all major land covers. Error
449 metrics include coefficient of determination (r^2), mean bias error (MBE), and mean absolute error (MAE).
450 Here, the subscripts ‘u’ and ‘c’ (e.g., MBE_u and MBE_c) denote uncorrected and bias-corrected ETo,
451 respectively. This evaluation demonstrates the application of the CONUS-AgWeather dataset for bias-
452 correcting gridded meteorological products used in operational workflows, such as OpenET^{15,16,62}.

453 OpenET Consortium relies on the CONUS-AgWeather dataset for bias correcting gridMET-based ETo⁷,
454 which is a key input to the majority of OpenET models^{15,16}. Based on the work of Volk et al. (2026)⁶²
455 that use 79 independent flux stations¹⁴ throughout the CONUS, gridMET ETo bias correction using the
456 CONUS-AgWeather dataset improves the accuracy of monthly ETo for cropland sites, and also forests,
457 shrublands, grasslands, and wetlands (Fig. 8).

458 Usage Notes

459 The CONUS-AgWeather dataset is intended to be a valuable resource for a diverse community of users,
460 including agricultural engineers, hydrologists, meteorologists, water resource managers, farmers, and
461 other users (e.g., educators). CONUS-AgWeather is a static data from 1981 to 2020. We plan to expand
462 eastward, incorporate more weather stations, improve the site screening algorithm to assess site aridity,
463 and update time series through more current years.

464 Potential Applications in Crop Management

465 a) Irrigation Scheduling: Providing high quality, bias-corrected weather and reference ET information as a
466 input for irrigation scheduling tools, leading to more precise water application and improved water use
467 efficiency⁶³, b) Crop Water Use and Stress Assessment: Using high quality, bias-corrected weather and
468 reference ET information in conjunction with crop models or remote sensing data to assess and monitor
469 crop water requirements, water use, and water stress^{9,10,13,64}, c) Yield Forecasting: Supplying quality-
470 controlled weather inputs for crop growth and yield forecasting models⁶⁵, d) Development and Validation
471 of Decision Support Systems: Serving as a benchmark dataset for developing, calibrating, and validating
472 new agricultural decision support tools⁶⁶, e) Climate Impact Studies on Agriculture: Providing quality
473 controlled historical weather data for analyzing the impacts of climate variability and change on crop
474 production and water demand^{67,68}, and f) Pest and Disease Modeling: Use of accurate weather data for
475 predicting the development and spread of crop pests and diseases⁶⁹.

476 General Meteorological and Hydrological Applications

477 These include a) validation and bias correction of numerical weather forecasts⁷⁰⁻⁷³, reanalysis products,
478 ^{8,74}, b) drought monitoring and assessment⁷⁵⁻⁷⁸, c) surface energy balance and ET studies¹⁶, and d)
479 surface water and groundwater modeling, water resource investigations, and national scale water use
480 reporting^{5,17,79-81}.

481 Limitations and Considerations for Users

482 a) Weather Station Networks: Weather station networks present in the CONUS-AgWeather dataset reflect
483 the strategic priorities of the OpenET Phase I project¹⁶, which focused primarily on developing an

484 operational actual ET product in the Western U.S. Consequently, the dataset exhibits a substantially
485 higher station density in the West compared to the East (Fig. 1-2). While we sought broader coverage,
486 many networks, particularly in the Eastern U.S. (e.g., North Carolina ECONet⁸², Alabama Mesonet⁸³,
487 New York State Mesonet⁸⁴, and others) were excluded to align with the Phase I timeline and specific
488 modeling objectives. We omitted data sources that were paywalled, unavailable prior to 2020, or
489 inaccessible during the project's operational window. Furthermore, because the dataset was specifically
490 created to bias-correct gridMET ETo for the OpenET actual ET models⁶², we enforced spatial filtering to
491 exclude stations located outside of well-watered agricultural areas, ensuring that only data representative
492 of agricultural microclimates was included in the dataset. Therefore, users focusing on the Eastern U.S.
493 may need to supplement CONUS-AgWeather with additional local or regional meteorological networks
494 that were either inaccessible at the time of creation or did not strictly meet the bias-correction criteria
495 adopted in the OpenET project.

496 b) Metadata Reliance: Some QC steps, calculation assumptions (e.g., standardized wind speed adjustment
497 to 2 m height based on reported anemometer height), depend on the accuracy of the station metadata
498 provided by the networks (i.e., latitude, longitude, elevation, anemometer height). Users should consult
499 the accompanying metadata for detailed site-specific information. It should be noted that the metadata
500 lists 19 weather station networks instead of the initial 22 as the HPRCC network includes stations from
501 HPRCC³³, NDAWN³⁵, Nebraska Mesonet³⁶, and South Dakota Mesonet⁴⁰ (see Table 1).

502 c) Station Siting: The ASCE Penman-Monteith reference ET equation²⁴ is used to calculate reference ET
503 for a hypothetical standardized reference crop surface (i.e., well-watered grass or alfalfa) and assumes
504 that weather data is representative of reference conditions. Out-of-specification siting and deviations from
505 reference conditions (e.g., excessive dryland within the station's fetch, tall crops, trees, or other
506 obstructions) can influence calculated reference ET. Extensive QC of station siting was conducted, but
507 ground-truthing was limited to a few stations.

508 d) Period of Record: The length of available data records varies by station but all end in 2020. Users
509 should verify that the period of record for selected stations is adequate for their intended applications.

510 **Code Availability**

511 The CONUS-AgWeather dataset⁶⁰ was generated using publicly available, open-source Python packages
512 to process and QC publicly available weather data. The primary package for quality assurance and quality
513 control is:

514 a) *agweather-qaqc*: The source code, documentation, and example usage are available on GitHub at
515 <https://github.com/WSWUP/agweather-qaqc>. Version 1.0.4²³ was instrumental in developing the
516 methodologies.

517 The calculation of standardized reference evapotranspiration (ETo and ETr) was performed using:

518 b) *RefET*: This package (Version 0.4.2), which implements the ASCE Standardized Reference
519 Evapotranspiration Equation²⁴, is available on GitHub (<https://github.com/WSWUP/RefET>) and PyPI
520 (<https://pypi.org/project/refet>)⁵⁷.

521 In addition, the Python scripts used for generating the maps (Fig. 2) and plots (Fig. 3 and Fig. 5-8) are
522 available on GitHub at <https://github.com/Open-ET/gridMET-bias-correction>. The open availability of

523 these software tools ensures transparency and reproducibility of the CONUS-AgWeather dataset
524 generation process and allows other researchers to apply identical or adapted methodologies to their own
525 agricultural weather datasets.

526 **Acknowledgements**

527 This dataset includes contributions from the following weather station networks:
528 The U.S. Bureau of Reclamation's AgriMet Network, Pacific Northwest and Great Plains regions,
529 Arizona Meteorological Network, California Irrigation Management Information System, Colorado
530 Agricultural and Meteorological Network, Enviroweather Network, Florida Automated Weather Network,
531 Georgia Automated Environmental Monitoring Network, High Plains Regional Climate Center, Missouri
532 Mesonet, North Dakota Agricultural Weather Network, Nebraska Mesonet, Nevada Integrated Climate
533 and Evapotranspiration Network, Oklahoma Mesonet, U.S. Department of Agriculture (USDA) Soil
534 Climate Analysis Network, South Dakota Mesonet, The National Oceanic and Atmospheric
535 Administration's United States Climate Reference Network, West Texas Mesonet, Western Regional
536 Climate Center, New Mexico's ZiaMet, Utah Climate Center, USDA Agricultural Research Service, and
537 Wyoming Agricultural and Climate Network.

538 Development of this dataset was supported by the S. D. Bechtel, Jr. Foundation; Walton Family
539 Foundation; Lyda Hill Philanthropies; U.S. Bureau of Reclamation; United States Geological Survey
540 (USGS) Water Resources Research Institute (grant G22AC00584-00); National Aeronautics and Space
541 Administration (NASA) Applied Science Program (grants NNX17AF53G and NNX12AD05A); USGS-
542 NASA Landsat Science Team (grant number 140G0118C0007); USGS Cooperative Ecosystem Studies
543 Units (CESU) (grant G23AC00568); NASA Western Water Applications Office (grant 1669431,
544 80NSSC23K0836); California State University Agricultural Research Institute (grant number 21-01-106);
545 Desert Research Institute Maki Endowment.

546 **Author contributions**

547 C.D. contributed to methodology, software, investigation, validation, data curation, visualization, writing
548 of the original draft, and writing – review & editing. J.M.V. contributed to conceptualization,
549 methodology, software, validation, formal analysis, investigation, data curation, and writing – review &
550 editing. S.M. contributed to writing of the original draft, software, visualization, formal analysis,
551 investigation, data curation, and writing – review & editing. J.L.H. was involved in funding acquisition,
552 supervision, project administration, conceptualization, methodology, data curation, formal analysis,
553 investigation, and writing – review & editing. R.G.A. contributed to conceptualization, methodology, data
554 curation, investigation, and project administration. C.P. contributed to formal analysis, investigation,
555 methodology, and data curation. Y.K. contributed to writing – review & editing and visualization.
556 C.G.M., B.A.M., and P.R. contributed to formal analysis and investigation. B.A.M. also contributed to
557 visualization and validation. A.K., F.M., and A.J.P. contributed to writing – review & editing, and project
558 administration. T.G.C. contributed to writing – review & editing, investigation, and visualization.

559 **Competing interests**

560 The authors declare no competing interests.

561 **References**

- 562 1. Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). *Crop Evapotranspiration - Guidelines for*
563 *Computing Crop Water Requirements - FAO Irrigation and drainage paper 56*. FAO - Food and
564 Agriculture Organization of the United Nations. <http://www.fao.org/3/X0490E/x0490e00.htm>
- 565 2. Bos, M. G., Kselik, R. A. L., Allen, R. G., & Molden, D. (2009). *Water Requirements for Irrigation*
566 *and the Environment*. Springer Netherlands. <https://doi.org/10.1007/978-1-4020-8948-0>
- 567 3. McColl, K. A., Salvucci, G. D., & Gentine, P. (2019). Surface Flux Equilibrium Theory Explains an
568 Empirical Estimate of Water-Limited Daily Evapotranspiration. *Journal of Advances in Modeling*
569 *Earth Systems*, 11(7), 2036–2049. <https://doi.org/10.1029/2019MS001685>
- 570 4. Salvucci, G. D., & Gentine, P. (2013). Emergent relation between surface vapor conductance and
571 relative humidity profiles yields evaporation rates from weather data. *Proceedings of the National*
572 *Academy of Sciences*, 110(16), 6287–6291. <https://doi.org/10.1073/pnas.1215844110>
- 573 5. Faunt, C. C. (2009). Groundwater availability of the Central Valley Aquifer, California. In C. C. Faunt
574 (Ed.), *U.S. Geological Survey Professional Paper 1766*. <https://doi.org/10.3133/pp1766>
- 575 6. Daly, C., Halbleib, M., Smith, J. I., Gibson, W. P., Doggett, M. K., Taylor, G. H., Curtis, J., & Pasteris,
576 P. P. (2008). Physiographically sensitive mapping of climatological temperature and precipitation
577 across the conterminous United States. *International Journal of Climatology*, 28(15), 2031–2064.
578 <https://doi.org/10.1002/joc.1688>
- 579 7. Abatzoglou, J. T. (2013). Development of gridded surface meteorological data for ecological
580 applications and modelling. *International Journal of Climatology*, 33(1), 121–131.
581 <https://doi.org/10.1002/joc.3413>
- 582 8. Blankenau, P. A., Kilic, A., & Allen, R. (2020). An evaluation of gridded weather data sets for the
583 purpose of estimating reference evapotranspiration in the United States. *Agricultural Water*
584 *Management*, 242, 106376. <https://doi.org/10.1016/j.agwat.2020.106376>
- 585 9. Ott, T. J., Majumdar, S., Huntington, J. L., Pearson, C., Bromley, M., Minor, B. A., ReVelle, P.,
586 Morton, C. G., Sueki, S., Beamer, J. P., & Jasoni, R. L. (2024). Toward field-scale groundwater
587 pumping and improved groundwater management using remote sensing and climate data.
588 *Agricultural Water Management*, 302, 109000. <https://doi.org/10.1016/j.agwat.2024.109000>
- 589 10. Huntington, J., Minor, B., Bromley, M., Pearson, C., Beamer, J., Ingwersen, K., Carrara, K., Atkin, J.,
590 Brito, J., Morton, C., Dunkerly, C., Volk, J., Ott, T., ReVelle, P., Fellows, A., & Hoskinson, M.
591 (2025). *Crop evapotranspiration, consumptive use, and open water evaporation for Oregon*. Desert
592 *Research Institute report 41306*. <https://doi.org/10.82269/DRI-DHS-41306>
- 593 11. Ji, L., Senay, G. B., Friedrichs, M., & Kagone, S. (2025). Estimating agricultural irrigation water
594 consumption for the High Plains aquifer region with integrated energy- and water-balance
595 evapotranspiration modeling approaches. *Agricultural Water Management*, 309, 109308.
596 <https://doi.org/10.1016/j.agwat.2025.109308>
- 597 12. Boser, A., Caylor, K., Larsen, A., Pascolini-Campbell, M., Reager, J. T., & Carleton, T. (2024). Field-
598 scale crop water consumption estimates reveal potential water savings in California agriculture.
599 *Nature Communications*, 15(1), 2366. <https://doi.org/10.1038/s41467-024-46031-2>
- 600 13. Majumdar, S., Smith, R. G., Hasan, M. F., Wilson, J. L., White, V. E., Bristow, E. L., Rigby, J. R.,
601 Kress, W. H., & Painter, J. A. (2024). Improving crop-specific groundwater use estimation in the
602 Mississippi Alluvial Plain: Implications for integrated remote sensing and machine learning
603 approaches in data-scarce regions. *Journal of Hydrology: Regional Studies*, 52, 101674.
604 <https://doi.org/10.1016/j.ejrh.2024.101674>

- 605 14. Volk, J. M., Huntington, J., Melton, F. S., Allen, R., Anderson, M. C., Fisher, J. B., Kilic, A., Senay,
606 G., Halverson, G., Knipper, K., Minor, B., Pearson, C., Wang, T., Yang, Y., Evett, S., French, A. N.,
607 Jasoni, R., & Kustas, W. (2023). Development of a Benchmark Eddy Flux Evapotranspiration
608 Dataset for Evaluation of Satellite-Driven Evapotranspiration Models Over the CONUS.
609 *Agricultural and Forest Meteorology*, 331, 109307. <https://doi.org/10.1016/j.agrformet.2023.109307>
- 610 15. Volk, J. M., Huntington, J. L., Melton, F. S., Allen, R., Anderson, M., Fisher, J. B., Kilic, A., Ruhoff,
611 A., Senay, G. B., Minor, B., Morton, C., Ott, T., Johnson, L., Comini de Andrade, B., Carrara, W.,
612 Doherty, C. T., Dunkerly, C., Friedrichs, M., Guzman, A., ... Yang, Y. (2024). Assessing the
613 accuracy of OpenET satellite-based evapotranspiration data to support water resource and land
614 management applications. *Nature Water*. <https://doi.org/10.1038/s44221-023-00181-7>
- 615 16. Melton, F., Huntington, J., Grimm, R., Herring, J., Hall, M., Rollison, D., Erickson, T., Allen, R.,
616 Anderson, M., Fisher, J. B., Kilic, A., Senay, G. B., Volk, J., Hain, C., Johnson, L., Ruhoff, A.,
617 Blankenau, P., Bromley, M., Carrara, W., ... Anderson, R. G. (2022). OpenET: Filling a Critical
618 Data Gap in Water Management for the Western United States. *JAWRA Journal of the American*
619 *Water Resources Association*. <https://doi.org/10.1111/1752-1688.12956>
- 620 17. Martin, D. J., Niswonger, R. G., Regan, R. S., Huntington, J. L., Ott, T., Morton, C., Senay, G. B.,
621 Friedrichs, M., Melton, F. S., Haynes, J., Henson, W., Read, A., Xie, Y., Lark, T., & Rush, M.
622 (2025). Estimating irrigation consumptive use for the conterminous United States: coupling satellite-
623 sourced estimates of actual evapotranspiration with a national hydrologic model. *Journal of*
624 *Hydrology*, 662, 133909. <https://doi.org/10.1016/j.jhydrol.2025.133909>
- 625 18. Ruehr, S., Bassiouni, M., Kang, Y., Socolar, Y., Magney, T., & Keenan, T. F. (2025). Crop
626 diversification improves water-use efficiency and regional water sustainability. *Environmental*
627 *Research Letters*, 20(11), 114062. <https://doi.org/10.1088/1748-9326/ae15a9>
- 628 19. Allen, R. G., Dhungel, R., Dhungana, B., Huntington, J., Kilic, A., & Morton, C. (2021).
629 Conditioning point and gridded weather data under aridity conditions for calculation of reference
630 evapotranspiration. *Agricultural Water Management*, 245, 106531.
631 <https://doi.org/10.1016/j.agwat.2020.106531>
- 632 20. Allen, R. G. (1996). Assessing Integrity of Weather Data for Reference Evapotranspiration
633 Estimation. *Journal of Irrigation and Drainage Engineering*, 122(2), 97–106.
634 [https://doi.org/10.1061/\(ASCE\)0733-9437\(1996\)122:2\(97\)](https://doi.org/10.1061/(ASCE)0733-9437(1996)122:2(97))
- 635 21. Allen, R. G., Pereira, L. S., Howell, T. A., & Jensen, M. E. (2011). Evapotranspiration information
636 reporting: I. Factors governing measurement accuracy. *Agricultural Water Management*, 98(6),
637 899–920. <https://doi.org/10.1016/j.agwat.2010.12.015>
- 638 22. Allen, R. G., Brockway, C. E., & Wright, J. L. (1983). Weather Station Siting and Consumptive Use
639 Estimates. *Journal of Water Resources Planning and Management*, 109(2), 134–136.
640 [https://doi.org/10.1061/\(ASCE\)0733-9496\(1983\)109:2\(134\)](https://doi.org/10.1061/(ASCE)0733-9496(1983)109:2(134))
- 641 23. Dunkerly, C., Huntington, J. L., McEvoy, D., Morway, A., & Allen, R. G. (2024). agweather-qaqc:
642 An Interactive Python Package for Quality Assurance and Quality Control of Daily Agricultural
643 Weather Data and Calculation of Reference Evapotranspiration. *Journal of Open Source Software*,
644 9(97), 6368. <https://doi.org/10.21105/joss.06368>
- 645 24. ASCE-EWRI. (2005). *The ASCE Standardized Reference Evapotranspiration Equation* (R. G. Allen,
646 I. A. Walter, R. L. Elliott, T. A. Howell, D. Itenfisu, M. E. Jensen, & R. L. Snyder, Eds.). American
647 Society of Civil Engineers. <https://doi.org/10.1061/9780784408056>
- 648 25. Kotttek, M., Grieser, J., Beck, C., Rudolf, B., & Rubel, F. (2006). World Map of the Köppen-Geiger

- 649 climate classification updated. *Meteorologische Zeitschrift*, 15(3), 259–263.
650 <https://doi.org/10.1127/0941-2948/2006/0130>
- 651 26. State of California. (2025). *CIMIS*. <https://cimis.water.ca.gov/Default.aspx>
- 652 27. Palmer, P. L. (2011). AgriMet: A Reclamation Tool for Irrigation Water Management. *World*
653 *Environmental and Water Resources Congress 2011*, 2682–2691.
654 [https://doi.org/10.1061/41173\(414\)279](https://doi.org/10.1061/41173(414)279)
- 655 28. University of Arizona. (2008). *Arizona Meteorological Network (AZMET) Data*. NSF NCAR Earth
656 Observing Laboratory. <https://doi.org/10.26023/Z7HE-7QP7-S40C>
- 657 29. Colorado State University (CSU). (2012). *DC3: Colorado Agricultural Meteorological Network*
658 *(CoAgMet) Data*. NSF NCAR Earth Observing Laboratory. [https://doi.org/10.26023/E18Y-7J8X-](https://doi.org/10.26023/E18Y-7J8X-AY02)
659 [AY02](https://doi.org/10.26023/E18Y-7J8X-AY02)
- 660 30. Michigan State University Enviroweather. (2025). *Enviroweather: Weather-based pest, natural*
661 *resource and production management tools*. <https://enviroweather.msu.edu/>
- 662 31. Lusher, W., Jackson, J., & Morgan, K. (2009). Florida Automated Weather Network (FAWN): Ten
663 Years of Providing Weather Information to Florida Growers. *EDIS*, 2009(7).
664 <https://doi.org/10.32473/edis-ss511-2009>
- 665 32. Hoogenboom, G. (1993, April). The Georgia Automated Environmental Monitoring Network.
666 *Proceedings of the 1993 Georgia Water Resources Conference*. <http://hdl.handle.net/10724/33288>
- 667 33. High Plains Regional Climate Center. (2019). *Automated Weather Data Network*.
668 <https://hprcc.unl.edu/awdn/index.php>
- 669 34. AgEBB - University of Missouri. (2025). *Missouri Mesonet*.
670 <http://agebb.missouri.edu/weather/realTime/maps/index.php>
- 671 35. NDAWN. (2025). *NDAWN - North Dakota Agricultural Weather Network*.
672 <https://ndawn.ndsu.nodak.edu/>
- 673 36. Shulski, M., Cooper, S., Roebke, G., & Dutcher, A. (2018). The Nebraska Mesonet: Technical
674 Overview of an Automated State Weather Network. *Journal of Atmospheric and Oceanic*
675 *Technology*, 35(11), 2189–2200. <https://doi.org/10.1175/JTECH-D-17-0181.1>
- 676 37. DRI. (2025). *NICE Net | Nevada Integrated Climate and Evapotranspiration Network*.
677 <https://nicenet.dri.edu/>
- 678 38. Brock, F. V., Crawford, K. C., Elliott, R. L., Cuperus, G. W., Stadler, S. J., Johnson, H. L., & Eilts,
679 M. D. (1995). The Oklahoma Mesonet: A Technical Overview. *Journal of Atmospheric and Oceanic*
680 *Technology*, 12(1), 5–19. [https://doi.org/10.1175/1520-0426\(1995\)012<0005:TOMATO>2.0.CO;2](https://doi.org/10.1175/1520-0426(1995)012<0005:TOMATO>2.0.CO;2)
- 681 39. Schaefer, G. L., Cosh, M. H., & Jackson, T. J. (2007). The USDA Natural Resources Conservation
682 Service Soil Climate Analysis Network (SCAN). *Journal of Atmospheric and Oceanic Technology*,
683 24(12), 2073–2077. <https://doi.org/10.1175/2007JTECHA930.1>
- 684 40. South Dakota Mesonet South Dakota State University. (2025). *South Dakota Mesonet Database*.
685 <https://climate.sdstate.edu/>
- 686 41. Diamond, H. J., Karl, T. R., Palecki, M. A., Baker, C. B., Bell, J. E., Leeper, R. D., Easterling, D. R.,
687 Lawrimore, J. H., Meyers, T. P., Helfert, M. R., Goodge, G., & Thorne, P. W. (2013). U.S. Climate
688 Reference Network after One Decade of Operations: Status and Assessment. *Bulletin of the*
689 *American Meteorological Society*, 94(4), 485–498. <https://doi.org/10.1175/BAMS-D-12-00170.1>

- 690 42. Schroeder, J. L., Burgett, W. S., Haynie, K. B., Sonmez, I., Skwira, G. D., Doggett, A. L., & Lipe, J.
691 W. (2005). The West Texas Mesonet: A Technical Overview. *Journal of Atmospheric and Oceanic*
692 *Technology*, 22(2), 211–222. <https://doi.org/10.1175/JTECH-1690.1>
- 693 43. WRCC. (2025). *Western Regional Climate Center*. <https://wrcc.dri.edu/>
- 694 44. DuBois, D. W., & Engle, S. (2022). Expansion of ZiaMet: New Mexico’s Mesonet. *American*
695 *Meteorological Society Meeting Abstracts*, 102, 293.
696 <https://ui.adsabs.harvard.edu/abs/2022AMS...10299559D/abstract>
- 697 45. Utah State University. (2026). *Utah Climate Center Mesonet*. <https://climate.usu.edu/mchd/>
- 698 46. Evett, S. R., Marek, G. W., Copeland, K. S., & Colaizzi, P. D. (2018). Quality Management for
699 Research Weather Data: USDA-ARS, Bushland, TX. *Agrosystems, Geosciences & Environment*,
700 1(1), 1–18. <https://doi.org/10.2134/age2018.09.0036>
- 701 47. University of Wyoming. (2025). *Wyoming Agricultural Climate Network (WACNet)*.
702 <https://www.wrds.uwyo.edu/WACNet/WACNet.html>
- 703 48. Singh, A., Taghvaeian, S., Mirchi, A., & Moriasi, D. N. (2023). Station Aridity in Weather
704 Monitoring Networks: Evidence from the Oklahoma Mesonet. *Applied Engineering in Agriculture*,
705 39(2), 167–177. <https://doi.org/10.13031/aea.15325>
- 706 49. Fiebrich, C. A., Brinson, K. R., Mahmood, R., Foster, S. A., Schargorodski, M., Edwards, N. L.,
707 Redmond, C. A., Atkins, J. R., Andresen, J. A., & Lin, X. (2020). Toward the Standardization of
708 Mesoscale Meteorological Networks. *Journal of Atmospheric and Oceanic Technology*, 37(11),
709 2033–2049. <https://doi.org/10.1175/JTECH-D-20-0078.1>
- 710 50. World Meteorological Organization (WMO). (2017). *Manual on the Global Observing System*
711 *(WMO-No. 544), Volume I*. <https://library.wmo.int/idurl/4/58672>
- 712 51. NOAA. (2025). *U.S. Climate Reference Network*. <https://www.ncei.noaa.gov/access/crn/>
- 713 52. Allen, R. (2008). Quality Assessment of Weather Data and Micrometeorological Flux-Impacts on
714 Evapotranspiration Calculation. *Journal of Agricultural Meteorology*, 64(4), 191–204.
715 <https://doi.org/10.2480/agrmet.64.4.5>
- 716 53. Iglewicz, B., & Hoaglin, D. (1993). How to Detect and Handle Outliers. In E. F. Mykytka (Ed.), *The*
717 *ASQC Basic References in Quality Control: Statistical Techniques* (1st ed., Vol. 16). ASQC Quality
718 Press.
- 719 54. Leys, C., Ley, C., Klein, O., Bernard, P., & Licata, L. (2013). Detecting outliers: Do not use standard
720 deviation around the mean, use absolute deviation around the median. *Journal of Experimental*
721 *Social Psychology*, 49(4), 764–766. <https://doi.org/10.1016/j.jesp.2013.03.013>
- 722 55. World Meteorological Organization. (2018). *Guide to Instruments and Methods of Observation*
723 *(WMO-No. 8)*. [https://community.wmo.int/site/knowledge-hub/programmes-and-](https://community.wmo.int/site/knowledge-hub/programmes-and-initiatives/instruments-and-methods-of-observation-programme-imop/guide-instruments-and-methods-of-observation-wmo-no-8)
724 [initiations/instruments-and-methods-of-observation-programme-imop/guide-instruments-and-](https://community.wmo.int/site/knowledge-hub/programmes-and-initiatives/instruments-and-methods-of-observation-programme-imop/guide-instruments-and-methods-of-observation-wmo-no-8)
725 [methods-of-observation-wmo-no-8](https://community.wmo.int/site/knowledge-hub/programmes-and-initiatives/instruments-and-methods-of-observation-programme-imop/guide-instruments-and-methods-of-observation-wmo-no-8)
- 726 56. Durre, I., Menne, M. J., Gleason, B. E., Houston, T. G., & Vose, R. S. (2010). Comprehensive
727 Automated Quality Assurance of Daily Surface Observations. *Journal of Applied Meteorology and*
728 *Climatology*, 49(8), 1615–1633. <https://doi.org/10.1175/2010JAMC2375.1>
- 729 57. Morton, C. G. (2023). *ASCE Standardized Reference Evapotranspiration Functions*.
730 <https://github.com/WSWUP/RefET>
- 731 58. Bokeh Development Team. (2025). *Bokeh documentation*. <https://docs.bokeh.org/en/latest/>

- 732 59. Thornton, P. E., & Running, S. W. (1999). An improved algorithm for estimating incident daily solar
733 radiation from measurements of temperature, humidity, and precipitation. *Agricultural and Forest*
734 *Meteorology*, 93(4), 211–228. [https://doi.org/10.1016/S0168-1923\(98\)00126-9](https://doi.org/10.1016/S0168-1923(98)00126-9)
- 735 60. Dunkerly, C., Volk, J. M., Majumdar, S., Huntington, J. L., Allen, R. G., Pearson, C., Kim, Y.,
736 Morton, C. G., Minor, B. A., ReVelle, P., Kilic, A., Melton, F., Purdy, A. J., & Caldwell, T. G.
737 (2026). CONUS-AgWeather, a high-quality benchmark daily agricultural weather station dataset for
738 evapotranspiration applications in the Contiguous United States (1.0.0) [Data set]. *Zenodo*.
739 <https://doi.org/10.5281/zenodo.18122157>
- 740 61. Allen, R. G., Irmak, A., Trezza, R., Hendrickx, J. M. H., Bastiaanssen, W., & Kjaersgaard, J. (2011).
741 Satellite-based ET estimation in agriculture using SEBAL and METRIC. *Hydrological Processes*,
742 25(26), 4011–4027. <https://doi.org/10.1002/hyp.8408>
- 743 62. Volk, J. M., Dunkerly, C., Majumdar, S., Huntington, J. L., Minor, B. A., Kim, Y., Morton, C. G.,
744 ReVelle, P., Kilic, A., Melton, F., Allen, R. G., Pearson, C., Purdy, A. J., & Caldwell, T. G. (2026).
745 CONUS Gridded Reference Evapotranspiration Bias Correction: Inputs, Station Validation, and
746 Outputs (gridMET/OpenET) [Data set]. *Zenodo*. <https://doi.org/10.5281/zenodo.18673484>
- 747 63. Stöckle, C. O., Liu, M., Kadam, S. A., Evett, S. R., Marek, G. W., & Colaizzi, P. D. (2025).
748 Comparing evapotranspiration estimations using crop model-data fusion and satellite data-based
749 models with lysimetric observations: Implications for irrigation scheduling. *Agricultural Water*
750 *Management*, 311, 109372. <https://doi.org/10.1016/j.agwat.2025.109372>
- 751 64. Smith, R., Oyler, L., Campbell, C., Woolley, E. A., Hopkins, B. G., Kerry, R., & Hansen, N. C.
752 (2021). A new approach for estimating and delineating within-field crop water stress zones with
753 satellite imagery. *International Journal of Remote Sensing*, 42(16), 6003–6022.
754 <https://doi.org/10.1080/01431161.2021.1931536>
- 755 65. Yang, Y., Anderson, M. C., Gao, F., Johnson, D. M., Yang, Y., Sun, L., Dulaney, W., Hain, C. R.,
756 Otkin, J. A., Prueger, J., Meyers, T. P., Bernacchi, C. J., & Moore, C. E. (2021). Phenological
757 corrections to a field-scale, ET-based crop stress indicator: An application to yield forecasting
758 across the U.S. Corn Belt. *Remote Sensing of Environment*, 257, 112337.
759 <https://doi.org/10.1016/j.rse.2021.112337>
- 760 66. Ara, I., Turner, L., Harrison, M. T., Monjardino, M., deVoil, P., & Rodriguez, D. (2021). Application,
761 adoption and opportunities for improving decision support systems in irrigated agriculture: A
762 review. *Agricultural Water Management*, 257, 107161. <https://doi.org/10.1016/j.agwat.2021.107161>
- 763 67. Meza, I., Siebert, S., Döll, P., Kusche, J., Herbert, C., Eyshi Rezaei, E., Nouri, H., Gerdener, H.,
764 Popat, E., Frischen, J., Naumann, G., Vogt, J. V., Walz, Y., Sebesvari, Z., & Hagenlocher, M.
765 (2020). Global-scale drought risk assessment for agricultural systems. *Natural Hazards and Earth*
766 *System Sciences*, 20(2), 695–712. <https://doi.org/10.5194/nhess-20-695-2020>
- 767 68. Huntington, J., Gangopadhyay, S., Spears, M., Allen, R. G., King, D., Morton, C., Harrison, A.,
768 McEvoy, D., Joros, A., & Pruitt, T. (2015). *West-Wide Climate Risk Assessments: Irrigation*
769 *Demand and Reservoir Evaporation Projections (Technical Memorandum No. 68-68210-2014-01)*
770 (U.S. Bureau of Reclamation, Ed.). U.S. Bureau of Reclamation.
771 <https://www.usbr.gov/watersmart/baseline/docs/irrigationdemand/irrigationdemands.pdf>
- 772 69. Olatinwo, R., & Hoogenboom, G. (2014). Weather-based Pest Forecasting for Efficient Crop
773 Protection. In *Integrated Pest Management* (pp. 59–78). Elsevier. [https://doi.org/10.1016/B978-0-](https://doi.org/10.1016/B978-0-12-398529-3.00005-1)
774 [12-398529-3.00005-1](https://doi.org/10.1016/B978-0-12-398529-3.00005-1)
- 775 70. Saminathan, S., & Mitra, S. (2025). Enhancing NWP-Based Reference Evapotranspiration Forecasts:

- 776 Role of ETo Approaches and Temperature Postprocessing. *Journal of Hydrologic Engineering*,
777 30(2). <https://doi.org/10.1061/JHYEFF.HEENG-6315>
- 778 71. Luo, Y., Chang, X., Peng, S., Khan, S., Wang, W., Zheng, Q., & Cai, X. (2014). Short-term
779 forecasting of daily reference evapotranspiration using the Hargreaves–Samani model and
780 temperature forecasts. *Agricultural Water Management*, 136, 42–51.
781 <https://doi.org/10.1016/j.agwat.2014.01.006>
- 782 72. McEvoy, D. J., Huntington, J. L., Mejia, J. F., & Hobbins, M. T. (2016). Improved seasonal drought
783 forecasts using reference evapotranspiration anomalies. *Geophysical Research Letters*, 43(1), 377–
784 385. <https://doi.org/10.1002/2015GL067009>
- 785 73. McEvoy, D. J., Roj, S., Dunkerly, C., McGraw, D., Huntington, J. L., Hobbins, M. T., & Ott, T.
786 (2022). Validation and Bias Correction of Forecast Reference Evapotranspiration for Agricultural
787 Applications in Nevada. *Journal of Water Resources Planning and Management*, 148(11).
788 [https://doi.org/10.1061/\(ASCE\)WR.1943-5452.0001595](https://doi.org/10.1061/(ASCE)WR.1943-5452.0001595)
- 789 74. Mankin, K. R., Mehan, S., Green, T. R., & Barnard, D. M. (2025). Review of gridded climate
790 products and their use in hydrological analyses reveals overlaps, gaps, and the need for a more
791 objective approach to selecting model forcing datasets. *Hydrology and Earth System Sciences*,
792 29(1), 85–108. <https://doi.org/10.5194/hess-29-85-2025>
- 793 75. Kukal, M. S., & Hobbins, M. (2025). Thirstwaves: Prolonged Periods of Agricultural Exposure to
794 Extreme Atmospheric Evaporative Demand for Water. *Earth's Future*, 13(3).
795 <https://doi.org/10.1029/2024EF004870>
- 796 76. Senay, G. B., Verdin, J. P., Lietzow, R., & Melesse, A. M. (2008). Global Daily Reference
797 Evapotranspiration Modeling and Evaluation 1. *JAWRA Journal of the American Water Resources*
798 *Association*, 44(4), 969–979. <https://doi.org/10.1111/j.1752-1688.2008.00195.x>
- 799 77. Hobbins, M., Jansma, T., Sarmiento, D. P., McNally, A., Magadzire, T., Jayanthi, H., Turner, W.,
800 Hoell, A., Husak, G., Senay, G., Boiko, O., Budde, M., Mogane, P., & Dewes, C. F. (2023). A
801 global long-term daily reanalysis of reference evapotranspiration for drought and food-security
802 monitoring. *Scientific Data*, 10(1), 746. <https://doi.org/10.1038/s41597-023-02648-4>
- 803 78. Hobbins, M. T., Wood, A., McEvoy, D. J., Huntington, J. L., Morton, C., Anderson, M., & Hain, C.
804 (2016). The Evaporative Demand Drought Index. Part I: Linking Drought Evolution to Variations in
805 Evaporative Demand. *Journal of Hydrometeorology*, 17(6), 1745–1761.
806 <https://doi.org/10.1175/JHM-D-15-0121.1>
- 807 79. Larsen, J. D., Langevin, C. D., Hughes, J. D., & Niswonger, R. G. (2024). An Agricultural Package
808 for MODFLOW 6 Using the Application Programming Interface. *Groundwater*, 62(1), 157–166.
809 <https://doi.org/10.1111/gwat.13367>
- 810 80. Hoekema, D. J., Ryu, J., & Abatzoglou, J. T. (2025). Validation of the Impacts of Recent Aquifer
811 Management on the Eastern Snake Plain Aquifer in Idaho, USA. *Groundwater*, 63(3), 387–398.
812 <https://doi.org/10.1111/gwat.13482>
- 813 81. Ebrahimi, E., & Shourian, M. (2025). Modeling farmer responses to reservoir operation policies using
814 agent based analysis of risk behavior and irrigation adoption. *Scientific Reports*, 15(1), 25591.
815 <https://doi.org/10.1038/s41598-025-11908-9>
- 816 82. Saia, S. M., Heuser, S. P., Neill, M. D., LaForce IV, W. A., McGuire, J. A., & Dello, K. D. (2023). A
817 Technical Overview of the North Carolina ECONet. *Journal of Atmospheric and Oceanic*
818 *Technology*, 40(6), 701–717. <https://doi.org/10.1175/JTECH-D-22-0079.1>

- 819 83. Kimball, S. K., Mulekar, M. S., Cummings, S., & Stamatatos, J. (2010). The University of South
820 Alabama Mesonet and Coastal Observing System: A Technical and Statistical Overview. *Journal of*
821 *Atmospheric and Oceanic Technology*, 27(9), 1417–1439.
822 <https://doi.org/10.1175/2010JTECHA1376.1>
- 823 84. Brotzge, J. A., Wang, J., Thorncroft, C. D., Joseph, E., Bain, N., Bassill, N., Farruggio, N., Freedman,
824 J. M., Hemker, K., Johnston, D., Kane, E., McKim, S., Miller, S. D., Minder, J. R., Naple, P., Perez,
825 S., Schwab, J. J., Schwab, M. J., & Sicker, J. (2020). A Technical Overview of the New York State
826 Mesonet Standard Network. *Journal of Atmospheric and Oceanic Technology*, 37(10), 1827–1845.
827 <https://doi.org/10.1175/JTECH-D-19-0220.1>

828