

# How do sea-level curves influence modeled marine terrace sequences?

Gino de Gelder<sup>1,2</sup>, Julius Jara-Muñoz<sup>3</sup>, Daniel Melnick<sup>4</sup>, David Fernández-Blanco<sup>1</sup>, Hélène Rouby<sup>1</sup>, Kevin Pedoja<sup>5</sup>, Laurent Husson<sup>2</sup>, Rolando Armijo<sup>1</sup> and Robin Lacassin<sup>1</sup>

<sup>1</sup>Université de Paris, Institut de physique du globe de Paris, CNRS, F-75005 Paris, France.

<sup>2</sup>ISTerre, CNRS, Université Grenoble Alpes, 1381 rue de la Piscine, 38400 Saint Martin d'Hères, France

<sup>3</sup>Institut für Geowissenschaften, Universität Potsdam, Karl-Liebknecht-Strasse 24, 14476 Potsdam, Germany.

<sup>4</sup>Instituto de Ciencias de la Tierra, Universidad Austral de Chile, 5111430 Valdivia, Chile

<sup>5</sup>Laboratoire de Morphodynamique Continentale et Côtière, CNRS, Université de Caen, 14000 Caen, France.

Corresponding author: Gino de Gelder ([Gino.De-Gelder@univ-grenoble-alpes.fr](mailto:Gino.De-Gelder@univ-grenoble-alpes.fr))

## Abstract

Sequences of uplifted marine terraces are widespread and reflect the interaction between climatic and tectonic processes at multiple scales, yet their analysis is typically biased by the chosen sea-level (SL) curve. Here we explore the influence of Quaternary SL curves on the geometry of marine terrace sequences using landscape evolution models (LEMs). First, we modeled the young, rapidly uplifting sequence at Xylokastro (Corinth Rift; <240 ka; ~1.5 mm/yr), which allowed us to constrain terrace ages, model parameters, and best-fitting SL curves. Models that better reproduced the terraced topography used a glacio-isostatically adjusted SL curve based on coral data (for ~125 ka), and a global mean SL curve based on ice-sheet models (for ~240 ka). Second, we explored the opposite end-member of older, slower uplifting sequences (2.6 Ma; 0.1-0.2 mm/yr). We find that cliff diffusion is important to model terrace sequence morphology, and that a hydraulic-model based SL curve reproduced observed terrace morphologies best. Third, we modeled the effect of SL noise with various amplitudes and wavelengths on our interpretations, finding that younger, faster uplifting sequences are less noise-sensitive and thus generally more promising for LEM studies. Our results emphasize the importance of testing a variety of SL-curves within marine terrace studies, and highlight that accurate modeling through LEMs may provide valuable insight on climatic and tectonic forcing to Quaternary coastal evolution.

**Keywords:** Quaternary; Sea level changes; Global; Coastal geomorphology; marine terraces; landscape evolution models; Corinth Rift


## 1 Introduction

Quantifying coastal uplift rates is essential for assessing tectonic dynamics and estimating seismic hazard (e.g. Merritts and Bull, 1989; Shaw et al., 2008), and quantifying glacio-eustatic sea-level (SL) variations is fundamental to estimate global ice-sheet volumes and their spatio-temporal response to climate change (e.g. Chappell and Shackleton, 1986; Lambeck et al., 2002, 2014). Sequences of fossil strandlines, resulting from the interplay between tectonic uplift and SL variations, cover much of the world's coastline (Pedoja et al., 2011; 2014) and thus provide a key archive to quantify both processes (e.g. Bradley, 1958; Lajoie, 1986, Anderson et al., 1999). Strandlines are markers of past SL position, reflecting global-to-regional tectonic and climatic processes since the Paleogene (Yamato et al., 2013; Henry et al., 2014; Pedoja et al., 2014), and are often expressed as marine terrace sequences. Marine terraces are fossil rocky shore platforms formed by coastal erosion, occasionally covered by a thin layer of coastal sediments and bounded inland by a fossil sea-cliff (e.g. Anderson et al., 1999). Sustained land uplift over several SL cycles leads to staircase morphologies comprising a series of marine terraces separated by fossil sea-cliffs. Shoreline angles, at the intersection of terraces and paleo-cliffs, are approximately time-equivalent to interglacial/interstadial SL highstands.

Marine terrace studies generally rely on terrace ages to constrain land uplift rates and/or relative SL history. Since typically only few ages within a terrace sequence are known (Pedoja et al., 2014), studies typically match undated terraces to Quaternary SL highstands using modeling strategies based on either statistical metrics (e.g. Zeuner, 1952; Bowles and Cowgill, 2012; Roberts et al., 2013) or landscape evolution models (LEMs; e.g. Quartau et al., 2010; Melnick, 2016; Jara-Muñoz et al., 2017, Pastier et al., 2019). These studies typically use a single SL curve, and do not always justify if the particular choice is based on resolution, timescale or geographic setting. However, the choice of SL curve can introduce significant uncertainties that might lead to biased results (e.g. Caputo, 2007, Sarr et al., 2019), as highstand estimates range in age by ~20 ka and SL elevation by ~30 meters (Fig. 1). Recent studies have explored the role of SL-curve choice for correlations of shoreline angles to SL highstands (e.g. Caputo et al., 2010; Pedoja et al., 2018a,b; Robertson et al., 2019), but the role of SL curves in LEMs and their impact on the full staircase morphology of a terrace sequence has not been investigated yet.


Here we apply a novel approach to model the development and age of marine terraces by investigating the influence of SL curves in LEMs. We test a spectrum of marine terrace sequences ranging from young and rapidly uplifting to old and slowly uplifting sequences (Fig. 2). To test our approach we initially focus on two end members. For the first end-member, we model the well-studied terrace sequence of Xylokaastro (Corinth Rift, Greece; Fig. 2), which is rapidly uplifting (~1.5 mm/yr) and relatively young (~240 ka). This allows us to constrain terrace ages, uplift rates, and other model parameters using 14 SL-curves, of which one is corrected for Glacial Isostatic Adjustments (GIA). Simultaneously, we evaluate which SL-curves can best reproduce the observed morphology. For the other end-member, we model older (Quaternary-Pliocene), slower uplifting (0.1-0.2 mm/yr) terrace sequences. We focus on the SL curve signature of the Mid-Pleistocene Transition (MPT), instead of modeling a specific site, as there are no precisely dated sequences covering the

time-spans of interest. The MPT relates to the change from dominantly 40 ky to 100 ky climate cycles at ~1250-700 ka (Clark et al., 2006), associated with contrasting pre- and post-MPT morphologies in nature (Fig. 2; Pedoja et al., 2014). Lastly we investigate for both end-member cases the influence of SL noise, i.e. the complex arrangement of variable wavelengths and amplitudes defining SL oscillations. Known analogues for such SL noise are meltwater pulses (e.g. Bard et al., 1996), Heinrich events (e.g. Andrews, 1998) and the Younger Dryas (e.g. Fairbanks, 1989) identified over the last glacial cycle and probably affecting SL on all timescales, but not clearly reflected in many SL curves (Fig. 1c). Testing SL noise allows us to verify to what extent interpretations from our modeling approach are robust, and to what extent they may be jeopardized by poorly constrained SL fluctuations. The combined case studies allow us to explore the intrinsic relation between terrace sequence morphology and Quaternary SL cycles on a range of timescales, and provide general recommendations on the use of SL-curves in marine terrace studies.






**Figure 1. Compilation of selected SL curves. (a)** The equatorial Pacific curve of Bates et al. (2014; curve 10, blue line) with the envelope of all other curves of our compilation (Table 1) in grey (see other curves in Supplementary Material). Boxes indicate age ranges for b and c **(b)** Elevation of SL highstands, with error bars as given in the original studies. Numbers indicate MIS and letters are substages as defined by Railsback et al. (2015). Black boxes encompass range of highstand estimates by the different SL curves **(c)** Zoom-in of SL curves since the LGM, comparing the curves of our compilation (numbered 1-14) to the detailed study of Lambeck et al. (2014). MWP1A, MWP1B and 8.2 are meltwater pulses, YD = Younger Dryas, H1 = Heinrich Event 1


**a** Sequences worldwide



**b** Rapidly uplifting sequences



**c** Slowly uplifting sequences



**Figure 2. Marine terrace/rasa sequences. (a)** Worldwide occurrence of marine terrace sequences (red), with white dots showing locations where rasas of Early to Mid-Pleistocene age or older have been described in the literature. **(b)** Examples of rapidly uplifting marine terrace sequences, as shown in the literature. **(c)** Examples of slower uplifting and older marine terrace/rasa sequences. UR = uplift rate

**2 Background**

**2.1 Sea-level curves**

Sea-level (SL) changes are driven by variations in volume and shape of oceans, as well as vertical land movement of coastal areas. During the Quaternary, glacio-eustasy has been a key long-term mechanism driving relative SL changes (Bloom, 1971). Long-term climatic cycles drive the periodic decay and growth of large ice-sheets, associated with global (eustatic) SL rise and fall, respectively. To quantify past SL, the earlier SL curves relied on the elevation of radiometrically dated geologic/geomorphic SL markers, typically corals (e.g. Chappell, 1974; Bard et al., 1990). A dense set of coral data from several locations worldwide have resulted in well-constrained global SL estimates since the Last Glacial Maximum (LGM) at ~21 ka (e.g. Lambeck and Chappell, 2001; Lambeck et al., 2014; Fig. 1c), but beyond this

age, coral data becomes progressively sparser and discontinuous, and has larger age uncertainties (e.g. Braithwaite, 2016). As an illustrative example, the coral database of Hibbert et al. (2016) contains 606 data points aged 0-21 ka (~30/ka), 1197 data points aged 21-130 ka (~10/ka), and 597 data points aged 130-850 ka (~1/ka).

Several methods have been developed to obtain longer and more continuous records of past SL, by constructing SL curves from marine sedimentary cores. Glacio-eustasy results in important changes within the hydrological cycle, as evaporation favors the removal of ocean water containing the lighter oxygen isotope ( $^{16}\text{O}$ ) over the heavier oxygen isotope ( $^{18}\text{O}$ ). As a consequence, the oxygen isotope ratio in seawater,  $\delta^{18}\text{OSW}$  ( $^{16}\text{O}/^{18}\text{O}$ ), can increase by ~1.5‰ when large amounts of water are stored within continental ice sheets during glacial periods. Most SL curves utilize this relationship between global ice-sheet volumes and  $\delta^{18}\text{OSW}$  to estimate past SL. Different techniques have been developed to this end, each relying on a different set of assumptions and sedimentary cores from various locations (Table 1), which results in a large scatter observed when comparing the different SL curves (Fig. 1).

**Table 1:** The different SL curves used in this study

Publication	Duration (ka)	Location	Average Resolution original data (ka)	Method
1. Shackleton 2000	400	Equatorial Pacific	1.3	$\delta^{18}\text{O}$ – temperature correction other proxy
2. Lea et al 2002	360	Cocos ridge	3.0	$\delta^{18}\text{O}$ – temperature correction other proxy
3. Waelbroeck et al 2002	430	Equatorial Pacific & N-Atlantic	1.5 (best 0.3)	$\delta^{18}\text{O}$ – coral regression
4. Bintanja et al 2005	1070	Global stack	1.4 (best 1)	Inverse ice volume model
5. Bintanja & V.d. Wal 2008	3000	Global stack	2.0 (best 1)	Inverse ice volume model
6. Rohling et al 2009	520	Red Sea	0.8 (best 0.3)	Hydraulic control models of semi-isolated basins
7. De Boer et al 2010	35000	Global stack	2.0 (best 1)	Inverse ice volume model
8. Elderfield et al 2012	1575	South Pacific	1.1	$\delta^{18}\text{O}$ – temperature correction other proxy
9. Bates et al 2014	5000	Equatorial Pacific*	2.8 (best 1.275)	$\delta^{18}\text{O}$ – coral regression
10. Grant et al 2014	500	Red Sea	0.2	Hydraulic control models of semi-isolated basins
11. Rohling et al 2014	5300	Mediterranean	1.0	Hydraulic control models of semi-isolated basins
12. Shakun et al 2015	800	Global stack	3.25 (best 1.5)	$\delta^{18}\text{O}$ – temperature correction other proxy
13. Spratt & Lisiecki 2016	800	Global stack	1.0	PCA on 7 existing records
14. This study	130	Local GIA-corrected		GIA-corrected, observation-calibrated ice volume models

\* Out of the 10 SL curves in Bates et al. (2014) this was used as their reference curve for comparisons

**Table 1. The different SL curves used in this study.**

### 2.1.1 Converting $\delta^{18}\text{OC}$ to $\delta^{18}\text{OSW}$

The  $\delta^{18}\text{O}$  of calcite ( $\delta^{18}\text{OC}$ ) is affected by both the  $\delta^{18}\text{OSW}$  and by temperature. Assuming no post-depositional alteration has occurred, the use of independent proxies to quantify the temperature component within  $\delta^{18}\text{OC}$ , should result in reliable estimates for the  $\delta^{18}\text{OSW}$  conditions in which the calcite formed. SL curves based on these assumptions typically use Mg/Ca ratios in ostracods (e.g. Dwyer et al., 1995) or foraminifera (e.g. Lea et al., 2002) as temperature proxies. The  $\delta^{18}\text{OC}$  is

derived from benthic (e.g. Elderfield et al., 2012; Sosdian and Rosenthal 2008) or planktonic (e.g. Lea et al., 2002; Shakun et al., 2015) foraminifera. The former mostly reflect the  $\delta^{18}\text{OSW}$  and temperature of deep water, whereas the latter reflects the sea surface temperature and the  $\delta^{18}\text{OSW}$  of shallow waters. As the shallow seas are subjected to stronger temperature variability, the conversion of benthic  $\delta^{18}\text{OC}$  to  $\delta^{18}\text{OSW}$  is less influenced by local conditions than planktonic  $\delta^{18}\text{OC}$ , and should be generally more reliable. An alternative way to convert  $\delta^{18}\text{OC}$  to  $\delta^{18}\text{OSW}$  was used by Shackleton (2000), pairing a Pacific  $\delta^{18}\text{OC}$  with an Arctic record of atmospheric  $\delta^{18}\text{O}$  trapped in ice-cores. For the final step of converting  $\delta^{18}\text{OSW}$  to SL, studies typically use a linear conversion of 0.08-0.1‰ m<sup>-1</sup> (e.g. Shackleton, 2000; Elderfield et al., 2012; Shakun et al., 2015), following calibration with a supposedly known SL like the LGM at -130 m (Lambeck et al., 2014). This introduces additional uncertainties in the final SL estimate, also as  $\delta^{18}\text{OSW}$  varies spatially, depending on local precipitation, evaporation, and deep water forming processes.

### *2.1.2 Coral regression*

Some studies have used coral datasets as benchmarks for  $\delta^{18}\text{O}$  records. Waelbroeck et al. (2002) constrained the direct relationship between coral-based paleo-SL estimates for the last glacial cycle (0-140 ka) and Pacific/N-Atlantic  $\delta^{18}\text{OC}$  records by fitting polynomial regressions. They separated the main glaciation from deglaciation interval, which were then applied to estimate SL for the  $\delta^{18}\text{OC}$  sections older than ~140 ka. The resulting SL curve is a composite curve constructed from the most reliable sections of the different sedimentary cores. A similar approach has been used on longer time-scales using transfer functions for 10 cores spanning 0-5 Ma (Siddall et al., 2010; Bates et al., 2014), although at lower resolution (Table 1). Bates et al. (2014) apply their methodology to different oceanic basins, and given major differences between Atlantic and Pacific curves prior to the MPT, they conclude that their approach is not appropriate for this interval.

### *2.1.3 Inverse ice-sheet modeling*

Sub-polar surface air temperature of the Northern Hemisphere plays a key role to two dominant factors affecting benthic  $\delta^{18}\text{OC}$ : the ice sheet size and the deep-water temperature. Bintanja et al. (2005) built on this notion using a global stack of 57 benthic  $\delta^{18}\text{OC}$  records (Lisiecki and Raymo, 2005) to drive a 3D ice-sheet-ocean-temperature model. The model results in estimates for ice sheet volume, temperature and average global SL from 1070 ka to present. A similar approach was subsequently used in studies covering 2.7 Ma (Bintanja and Van der Wal, 2008) and even 35 Ma (De Boer et al., 2010). The models over these longer timescales suffer from a lack of independent paleoclimate data to constrain their results, and the authors point out that the simplicity of their ocean-atmosphere temperature coupling module and other modeling assumptions lead to considerable uncertainties (Bintanja and Van der Wal, 2008).

### *2.1.4 Hydraulic models*

In semi-isolated basins like the Red Sea and the Mediterranean,  $\delta^{18}\text{O}_{\text{C}}$  is strongly influenced by both evaporation and exchange with the open ocean, the latter closely related to sill depth and thus relative SL. Siddall et al. (2003) used high-resolution  $\delta^{18}\text{O}_{\text{C}}$  records from the Red Sea to reconstruct the history of water residence times. Through a hydraulic model of the water exchange between the Red Sea and the open ocean, they calculated relative SL at the sill. Rohling et al. (2009) extended and improved this record using additional data, whereas Grant et al. (2014) improved the chronology through synchronization with an Asian monsoon record. Whereas the Red Sea records extend to  $\sim 500\text{ka}$  (Table 1), Rohling et al. (2014) applied the same methodology to a Mediterranean  $\delta^{18}\text{O}_{\text{C}}$  stack to obtain a relative SL record for the last 5.3 Ma. However, the stability of Mediterranean Sea surface temperatures and relative humidity are difficult to verify for this timescale, and so are the tectonic stability and isostatic response at the Gibraltar sill over several millions of years.

Contrarily to the other SL curves used in this study (Table 1), the hydraulic model curves represent relative sea-level (RSL) at the Hanish (Rohling et al., 2009; Grant et al., 2014) and Gibraltar (Rohling et al., 2014) sills, rather than global mean sea-level (GMSL). Using glacial isostatic adjustment models at those two locations, the RSL is expected to scale approximately linearly with GMSL, with factors of  $\sim 1.18$  (Hanish; Grant et al., 2014) and  $\sim 1.23$  (Gibraltar; Rohling et al., 2014).

### *2.1.5 Principal Component Analysis*

Principal Component Analysis (PCA) is a statistical technique to reduce the number of variables of a dataset, while retaining the most important trends within those variables. Spratt and Lisiecki (2016) selected 7 SL records constructed with the methods outlined above, and used PCA to identify the common SL signal. The resulting stack was then scaled based on Holocene and LGM estimates. Although the signal-to-noise ratio in this PCA-curve should be much better than that of the individual curves, the approach may have resulted in smoothed interglacials, thus underestimating the SL elevation of brief highstands within those interglacials.

### *2.2 Modeling marine terrace sequences*

Several studies explored the generation of marine terrace sequences through LEMs. Anderson et al. (1999) showed that wave dissipation is essential to the formation and preservation of marine terraces, and the formation of a terrace sequence is sensitive to a complex interaction of parameters, including SL history. They highlighted that the subsequent sub-aerial erosion of terraces is closely linked to stream incision rates, reflecting climatic conditions and limited by the rate of base-level fall. Similar terrace formation models by Trenhaile (2002, 2014) showed that terraces formed during interglacial and glacial periods are more likely to be recorded in uplifting and subsiding landmasses, respectively, in agreement with natural examples (e.g. Chappell, 1974; Ludwig et al., 1991). The author also modeled the positive correlation between uplift rates and the number of terraces preserved within a sequence, which had been previously recognized in nature (Merritts and Bull, 1989; Armijo et al., 1996). Along the same lines, LEMs of Melnick (2016) and Pastier et al. (2019) emphasized that for low uplift rates, the continuous reoccupation of abrasion

platforms, coupled to paleo-cliff diffusion, hinders the direct conversion of terrace elevations to uplift rates.

The amplitude and period of SL cycles has a distinct influence on the geometry of a modeled marine terrace sequence. Using synthetic SL curves, both Trenhaile (2002) and Husson et al. (2018) show that short-period, low amplitude SL oscillations before the MPT result in narrower terraces with smaller paleocliffs, compared to longer-period, higher amplitude SL swings after the MPT. Using the 3-Ma SL curve of Bintanja and Van der Wal (2008), Trenhaile (2014) drew similar conclusions, highlighting that the ~100-ky, ~120m amplitude SL cycles since ~1 Ma accomplish relatively more cliff erosion. Husson et al. (2018) performed similar tests for coral reef terraces, and found that SL noise is an influential factor in reef building, and infrequent, relatively long SL transgressions are important to the geometry of a sequence.

### *2.3 The coastal sequence at Xylokaastro*

The sequence of marine terraces at Xylokaastro (Fig. 4) is located in the SE Corinth Rift (Greece). High uplift rates of ~1.2-1.5 mm/yr (Armijo et al., 1996; De Gelder et al., 2019), low sub-aerial erosion rates and thin cover of cemented coastal deposits have resulted in a well-preserved sequence of 13 marine terraces (e.g. Dufaure and Zamanis, 1979; Keraudren and Sorel 1987; Armijo et al., 1996; De Gelder et al., 2019). The terrace sequence spans the last ~240 ka, and extends over an area of ~3 x 3 km, culminating at an elevation of ~400 m (Fig. 4a). The terraces were previously named after local towns (see Armijo et al., 1996), but herein we assign a simpler name designation of TH (Holocene terrace) and T1-T12 (Fig. 4a). The T7 (New Corinth; ~175 m elevation) and T11 (Old Corinth; ~320 m elevation) terraces have been dated using both U/Th on solitary corals (Collier et al., 1992; Dia et al., 1997; Leeder et al., 2005), and <sup>14</sup>C dating of Pecten (Pierini et al., 2016). These studies correlate T7 and T11 to the Marine Isotope Stage (MIS) 5e (~125 ka) and MIS 7e (~240 ka) highstands, respectively. An inactive alluvial fan at ~200-230 m elevation caps T8 and T9, hindering our map of terraces in this ~ 0.5 km<sup>2</sup> area (Fig. 4a).

## **3 Methods**

### *3.1 Landscape Evolution Model*

We use a LEM based on the wave erosion and wave energy dissipation model formulated by Sunamura (1992), and further developed by Anderson et al. (1999). The model simulates the evolution of rocky coasts by the retreat of sea-cliffs, driven by wave erosion and resulting in the genesis of rocky shore platforms. The model assumes that the vertical seabed erosion rate is a linear function of the rate of wave energy dissipation against the seabed (Sunamura, 1992). The energy available at the sea-cliff to drive horizontal erosion is defined by the far-field wave energy remaining after wave energy dissipation (Anderson et al., 1999). The water depth profile dictates the spatial pattern of dissipation rate, exponentially increasing landwards as the water depth decreases. We use a 2D model setup formed by a planar shelf of given slope and assume that the rate of rock uplift is uniform. Cliff retreat starts at an



initial rate and evolves as the platform is carved during SL oscillations, which depends on the SL curve. Detailed description and equations can be found in Anderson et al. (1999).

### 3.2 Sea-Level Curves

#### 3.2.1 Selected Sea-Level Curves

We systematically selected from literature GMSL curves that cover at least the last 3 major glacial cycles (~350 ka), and are based on data with a temporal resolution of <3 ka (Table 1). We also included three RSL curves based on hydraulic models. The 13 resulting SL curves (Table 1, Fig. 1) are subdivided into curves that; (i) convert  $\delta^{18}O_C$  to  $\delta^{18}O_{SW}$  (light blue, Fig. 1); (ii) use a regression analysis to fix  $\delta^{18}O$  curves to RSL estimates from corals (dark blue, Fig. 1); (iii) are based on global ice-sheet modeling (green, Fig. 1); (iv) are based on hydraulic models of water exchange between an evaporative sea and the ocean (pink, Fig. 1); and (v) result from principal component analysis (PCA) of 7 other curves (brown, Fig. 1). More details are given in Table 1 and section 2.1.

#### 3.2.2 Relative Sea-Level Curve Xylokaastro

GMSL fluctuations occur in response to the buildup and retreat of ice sheets, with strong local departures because of Glacial Isostatic Adjustment (GIA), i.e. solid earth and sea surface deformation under water and ice mass redistribution (e.g. Bloom, 1967; Walcott, 1972, Lambeck, 1995). Using the GIA models developed at the Australian National University (ANU) (e.g. Nakada and Lambeck, 1989; Johnston, 1993; Lambeck et al., 2003, 2014), we calculated a GIA-corrected relative SL (RSL) curve for Xylokaastro. The GIA model contains parameters describing both the deformation sources, i.e. the ice-volume distribution history, and the behavior of the deformed object, i.e. the rheology of the Earth mantle. The ANU CALSEA code allows computation of the RSL for a given location and time back to the last interglacial. It includes all deformational, gravitational and rotational changes induced by the global ice history during the last glacial cycle, and accounts for the laterally variable rheology of the Earth mantle.

The ANU model is constructed using an iterative procedure. The ice volume history of each ice-sheet is solved regionally together with mantle rheology under the formerly glaciated areas by inversion of direct RSL observations (near-field data from or within the former ice margins). Far-field RSL data are separately inverted for total melted-ice volume and mantle parameters under oceanic and continental margins. The sum of individual ice-sheets volumes and the global volume are then compared and corrections are made to the different ice-volumes. The operation is reiterated until reasonable convergence is obtained (Lambeck et al., 2001; 2010; 2014; 2017). Because there is sparse direct observational data before the last interglacial, the model is limited to 130 ka.

The impact of GIA on a region situated in intermediate field during the last glaciation has been demonstrated in Simms et al. (2016) for the US-Mexico Pacific coast. This is at similar intermediate distances from major northern ice-sheets as the

Mediterranean region, and therefore we adopted the same model parameters as Simms et al. (2016) and expect uncertainties of the same order of magnitude (~5-10 m). The effective parameters used for the Earth mantle reflect an intermediate region where the response to water and ice loads are comparable: a lithospheric thickness  $h = 80$  km, an upper-mantle viscosity  $\mu_{um} = 2 \times 10^{20}$  Pa·s and a lower-mantle viscosity  $\mu_{lm} = 10^{22}$  Pa·s. The uncertainty is estimated considering a range of parameters that reflect the lateral variations of the Earth's mantle. The uncertainty that can be imputed to ice sheet configuration is estimated by scaling the North American ice sheet up to +/- 20% during the last glaciation period. More details can be found in Simms et al. (2016). The purpose here is mostly demonstrative, and we note that uncertainties in the GIA-effects are relatively small in comparison to the differences amongst SL curves (Fig. 1). The calculated SL elevations for the RSL curve can be found in Dataset 2.

### 3.3 Modeling the coastal sequence at Xylokaastro

We constructed a representative cross-section of the Xylokaastro terraces to compare with the LEM simulated topography (Fig. 4b), by calculating average (i) shoreline angle elevations, (ii) terrace widths, (iii) terrace slopes and (iv) the modern rocky shore platform width. To determine shoreline angles, the intersection between the marine terrace and its associated fossil sea-cliff, we used a 2-m resolution Digital Surface Model (DSM) developed from Pleiades satellite imagery (De Gelder et al., 2015, 2019). From the DSM we calculated 100-m-wide swath profiles perpendicular to the fossil sea-cliffs and determined the shoreline angle positions and elevations using the fixed-slope method of TerraceM (Jara-Muñoz et al., 2016). We used the maximum swath profile topography and the modern sea-cliff slope angle of  $39^\circ \pm 10^\circ$  (De Gelder et al., 2017) as a proxy for the slope of the paleo sea-cliff (Dataset S1). To approximate the terrace average width, we used the distance between two successive shoreline angles (Fig. S1), since sub-aerial erosion of the paleo-cliffs has reduced the original terrace-width since they emerged. To estimate terrace slopes, we used the average slope of the modern terrace as less-eroded proxies for their older counterparts. To estimate the outer limit of the modern rocky shore platform, we assume that it has largely been carved during Holocene SL rise. Before ~12 ka (Moretti et al., 2003) the Corinth Gulf was a lake, its water exchange with the open sea limited by the 62 m deep Rion sill (Perissoratis et al., 2000). Assuming Holocene terrace carving started 12 ka at 62 m depth, and given an uplift rate of ~1.3 mm/yr (Armijo et al., 1996), the present depth contour of -46 m should approximately represent the outer limit of the modern rocky shore platform (Fig. S1).

We tied the cross-section shoreline angles of dated terraces T7 and T11 to the shoreline angles formed during the MIS 5e (~125 ka) and MIS 7e (~240 ka) highstands in the LEM. We fixed LEM uplift rates to reproduce the observed average shoreline angle elevations, and varied initial erosion rate and initial shelf slope with steps of 0.1 m/yr and  $0.25^\circ$ , respectively. This allowed us to select the best-fitting pair of values that resulted in the lowest Root-Mean-Squared (RMS) misfit on both MIS 5e and MIS 7e timescales. The RMS misfit takes into account the full 2D-geometries of the modeled and observed profiles, by comparing the vertical difference between both for every horizontal step (dx). To match modeled with observed terraces, we

pick the observed terrace of which the shoreline angle elevation is closest to that of the modeled terrace. Like the RMS-misfit, we consider the number of matched terraces as an indication for how well a SL curve can reproduce a terrace sequence. Our models used a spatial resolution of 2 m to match our DSM. We used time-steps of 50 years, following sensitivity tests on the two highest resolution SL curves (Fig. S2): smaller time-steps give similar results. In the modeling, we assumed that the SL in Corinth did not get lower than the Rion sill (62 m depth) during the past 240 ka. We used a wave height of 3 m, based on the highest waves recorded between 2010-2015 at the Gulf of Corinth with AVISO satellite altimetry measurements (Fig. S3; <https://www.aviso.altimetry.fr/>). Sensitivity tests for wave height and sill depth show that those parameters do not strongly affect our results (Fig. S3).

### *3.4 Modeling the Mid-Pleistocene Transition*

To constrain marine terrace geometries for older, slower uplifting sequences, we model the formation of marine terraces sequences over the whole Quaternary (2.6-0 Ma) using the four longest GMSL curves of our compilation (Bintanja and Van de Wal, 2008; De Boer et al., 2010, Bates et al., 2014, Rohling et al., 2014). In our modeling strategy, we use a relatively low initial slope of 4° and an uplift rate of 0.1 mm/yr, consistent with an approximate average of the examples shown in Fig. 2c. We used an initial sea-cliff erosion rate of 0.6 m/yr, consistent with our average estimate for Corinth (see results), and included a 0.1 m<sup>2</sup>/yr sub-aerial cliff diffusion rate to obtain a more realistic sequence morphology over this timescale.

### *3.5 Influence of noise in sea-level curves*

To test the influence of noise in SL curves on both end-member type marine terrace sequences, we start with a curve composed of 40 ky-period, 65 m-amplitude sine function to describe SL for the 2.6-1 Ma interval, and a 100 ky-period, 130 m-amplitude sine function for the 1-0 Ma interval. We then add noise to this SL curve with amplitudes of 4, 10 or 25 m spaced every 1, 5 or 20 ky, using the rand function in MATLAB®. We use the same erosion rate and initial slope as in section 3.4, test cases with and without a sub-aerial cliff diffusion rate of 0.1 m<sup>2</sup>/yr, and test both 0.1 and 1.5 mm/yr uplift rates over 2.6 and 0.4 Ma, to compare with the MPT and Xylokaastro case studies, respectively.

## **4 Results**


### *4.1 Modeling the coastal sequence at Xylokaastro*

The systematic comparison of observed topography in the Xylokaastro marine terraces sequence and their modeled topography allows us to 1) assess possible age ranges for undated terraces, 2) quantify the governing parameters of terrace formation, and 3) evaluate the best-fitting SL curves by means of the number of matched terraces and the RMS misfit. In Fig. 3 we show four examples of our analysis, displaying the curves resulting in the most terraces (Figs. 3a, 3c) and lowest RMS misfits (Figs. 3b, 3d) over ~125 ka (Figs. 3a, 3b) and ~240 ka (Figs. 3c, 3d). Fig. S4 contains the results for the other tested SL curves. Correlation of


modeled shoreline angle elevations to the nearest shoreline angles results in different ages estimates as a function of SL curve (Fig. 4b). Most curves suggest a chrono-stratigraphy in which T2 was formed during MIS 5a (~70-85 ka), T3 during MIS 5a or MIS 5c (~92-107 ka), T4 and T5 during MIS 5c, T6 and T7 during MIS 5e (~115-128 ka), T9 during MIS 7a (~190-205 ka), T10 during MIS 7a or 7c (~210-225 ka) and T11 during MIS 7e (~235-242 ka). T1 and T8 were only reproduced by SL curves 2 and 3, respectively, but relative to the other terraces would logically have an age of MIS 5a or younger (T1), and MIS 8 or MIS 9a (T8).

The uplift rate (Fig. 4c), initial erosion rate (Fig. 4d) and initial slope (Fig. 4e), vary strongly between different SL curves, and for the two different timescales. Despite this variation, the overall average values do not increase or decrease relevantly over time (Figs. 4c-4e). The number of matched terraces for the different curves has a broad range for both ~125 ka (2-6 terraces) and ~240 ka (4-7 terraces) timescales, but none of the selected curves could recreate the observed number of 11 successive terraces (Fig. 4f). Plotting the number of matched terraces against the temporal resolution of each SL curve shows a strong correlation between higher resolution curves and a larger number of matched terraces (Fig. S5), although higher curve resolution does not correlate with lower RMS misfits (Fig. S5). No SL curve gives consistently good results in terms of RMS misfits, but we note that for ~125 ka the two lowest misfit curves are based on corals (dark blue; 3 and 14 in Fig. 4g). Over the same timescale, the curves based on hydraulic models (pink; 6, 10 and 11 in Fig. 4g) give consistently high misfits.


We compared the GMSL and the RSL curves calculated by the ANU model (see section 3.2.2) to highlight how GIA corrections influence our approach (Fig. 5). The difference between these two curves is typically on the order of several meters (Fig. 5b), and the GIA-correction of the GMSL curve improves the RMS misfit by >3m (Fig. 5a), while reproducing an extra terrace. The GIA-corrected curve is also the overall best-fitting curve over the last ~125 ka.



**Figure 3. Examples of modeling the Xylokaastro terrace sequence. (a)** for the most matched terraces over ~125 ka; **(b)** for the lowest RMS-misfit over ~125 ka; **(c)** for the most matched terraces over ~240 ka, and; **(d)** for the lowest RMS-misfit over ~240 ka. In profile plots (above), red lines show average-derived terrace topography (see methods) with  $1\sigma$  uncertainty (grey envelope), black lines the modeled topography, and arrows indicate which SL highstand corresponds to which shoreline angle. Dashed lines connecting shoreline angle of modeled and observed shoreline angles indicate the correlation used for Fig. 4b. In the SL plots (below), numbers indicate MIS and letters are substages as defined by Railsback et al. (2015).



**Figure 4. Modeling the marine terrace sequence at Xylokastro. (a)** Map of Xylokastro marine terrace sequence. SA = shoreline angle. **(b)** Profile of the terrace sequence. The black line shows average-derived terrace topography (see methods) with  $1\sigma$  uncertainty (grey envelope) and range of possible terrace ages indicated by the different SL curves, including median age (m) and number of curves (n) reproducing a given marine terrace. **(c-g)** Different parameters and outcomes resulting from finding the lowest RMS-misfit over the two time-scales, shown for all 14 RSL curves. Models over 125 ka and 240 ka are shown in red and green, respectively. Numbers below plot correspond to SL curve numbering and colors of Fig. 1, and are sorted by type of SL curve for easy comparison. Solid and dashed lines indicate average values and  $1\sigma$  uncertainty. **(c)** Uplift rates required by the different SL curves to match the correct elevations of the dated T7 (red) and T11 (green) terraces. **(d)** Erosion rate for lowest RMS-misfit result. **(e)** Initial slope for lowest RMS-misfit result. **(f)** Total number of terraces for lowest RMS-misfit result. **(g)** Lowest RMS-misfits.




**Figure 5. Xylokaastro modeling results for the CALSEA GMSL curve and GIA-corrected relative SL curve. (a)** Profile comparison for lowest RMS-misfit modeling results. The red line shows average-derived terrace topography (see methods) with  $1\sigma$  uncertainty (grey envelope), the black dashed line shows the modeled topography with the CALSEA GMSL curve, and solid black line shows the modeled topography with the GIA-corrected relative SL curve. Arrows indicate which shoreline angle corresponds to which SL highstand in b. **(b)** The two SL curves, in which numbers indicate MIS and letters are substages as defined by Railsback et al. (2015).

#### 4.2 Modeling the Mid-Pleistocene Transition (MPT)

Lower uplift rates (0.1 mm/yr) in the 2.6-Ma models (Fig. 6) generally result in fewer terraces being formed than at Xylokaastro (Fig. 4). At such uplift rates, only terraces formed during the maxima of interglacial highstands (MIS 5e, 7e etc.) are preserved. The effect of the MPT is most pronounced when modeling the curve of Rohling et al. (2014), with 2 rasas (wide polygenic fossil strandlines) formed before the MPT and 5 marine terraces after the MPT (Figs. 6a, e). Sequences produced by other SL curves do not show a similarly clear contrast before and after the MPT. The curve with the most (Bates et al., 2014) and least (Rohling et al., 2014) pronounced change in cyclicity around the MPT (spectrograms in Fig. S6), correspond to the relative least and most pronounced change in sequence morphologies, respectively (Fig. 6). Additional tests with other values for uplift rates, erosion rates and initial slope vary in

shape of staircase sequences (Fig. S7), but consistently show the lowest ratio between terraces/rasas preserved before and after the MPT for the hydraulic curve (Rohling et al., 2014).




**Figure 6. Model results on Quaternary (2.6 Ma) timescale. (a)** Modeled geometry of the long-term sequences, dashed lines indicating the MPT. Inset shows typical morphology for a Quaternary staircase sequence, modified from Pedoja et al. (2014). **(b-e)** Modeled SL curves, with lettered arrows indicating SL highstands that result in preserved terraces in a.

#### 4.3 Influence of noise in sea-level curves

We estimated the effect of various degrees of noise added to synthetic SL curves for slowly uplifting coastlines ( $\sim 0.1$  mm/yr) (Fig. 7). The reference profile has no noise (black lines, Fig. 7a) and portrays a prominent cliff that separates an upper, pre-MPT section of narrow terraces with small cliffs, from a lower, post-MPT section of wider terraces with higher cliffs. Cliff-diffusion completely smoothens all pre-MPT cliffs, and partly smoothens post-MPT cliffs (thick black line, Fig. 7a). Adding 4 m of noise spread out every 20 ky already has a significant influence on the sequence morphology with respect to the reference profile, and leads to variations in terrace width and cliff heights both pre- and post-MPT. With higher noise amplitudes and lower noise wavelengths, the contrast between the two intervals (pre- and post-MPT) is less clear. The cliff separating the two intervals becomes less prominent, and the overall width of the pre-MPT section increases relative to the width of the post-MPT




section. In general the addition of cliff diffusion (thick lines in Fig. 7a) result in fewer and often composite terraces (rasas), especially for the pre-MPT section. Using the same curves over 400 ka with an uplift rate of 1.5 mm/yr (Fig. 8) results in very minor differences in comparison to the reference profile, and are only clearly visible for the SL curve with 25 m, 1 ky wavelength noise. For noise amplitudes of 10 and 25 m with 1 and 5 ky wavelengths, respectively, the overall shape of the terrace sequence is similar but minor sub-levels develop. Overall the noise tests indicate that terrace morphology is more affected by noise with higher noise amplitudes and shorter noise wavelengths. Sequences with lower uplift rates are more sensitive to the influence of noise.



**Figure 7. Modeling noise on Quaternary (2.6 Ma) timescale. (a)** Modeled geometry of the long-term sequences, thick lines representing model runs with 0.1 m<sup>2</sup>/yr cliff diffusion and thin lines without cliff diffusion **(b)** Modeled synthetic SL

curves, characterized by a 40 ky, 65 m amplitude sine function before 1 Ma, and a 100 ky, and a 130 m amplitude sine function after 1 Ma. Noise amplitudes are 4, 10 and 25 m, increasing to the right, and noise spacing 1, 5 and 20 ky increasing downwards.



**Figure 8. Modeling noise over 400 ka timescale, with a 1.5 mm/yr uplift rate. (a)**

Modeled geometry of the long-term sequences, thick lines representing model runs with 0.1 m<sup>2</sup>/yr cliff diffusion **(b)** Modeled synthetic SL curves, characterized by a 100 ky, 130 m amplitude sine function. Noise amplitudes are 4, 10 and 25 m, increasing to the right, and noise spacing 1, 5 and 20 ky increasing downwards.

## 5 Discussion

Here we re-evaluate different aspects of our analysis towards a unified objective: understanding how SL curves affect modeled marine terrace sequences. We first discuss results from the Xylokaastro sequence, which is a short yet well-defined sequence, and therefore appropriate as a test site at high resolution. We then discuss the MPT results and the possible pitfalls associated with longer time-scale modeling, including the influence of SL noise, and compare both end-member type marine terrace sequences. We then conclude with some general recommendations and perspectives on using SL-curves in marine terrace analyses based on our findings.

### 5.1 Modeling coastal sequences: the Xylokaastro lessons

#### 5.1.1 Approach limitations

Modeling tests on the Xylokaastro terrace sequence reveal the complexity of reproducing as many terraces as observed in nature, either as a consequence of model assumptions and/or SL curves used. The relatively simple model used (Anderson et al., 1999) does not take into account the abrasive effect of sediments, long-shore drift, or coastal sediments on the terraces. Additionally, the far field wave energy is considered constant in our models, even though it probably varies within glacial-interglacial climate cycles and thus affects erosion rates. Although these might all influence terrace sequence geometry, implementing such processes within our model is beyond the scope of this study, and in any model the influence of the chosen SL curve would remain of primary importance. Models by Trenhaile (2002, 2014) included a more sophisticated parameterization of wave regime and coastal cliff retreat, but the author concluded that the results were generally consistent with those of Anderson et al. (1999). Therefore, we expect a more detailed set of cliff erosion parameters would not have changed our results significantly. Our simple model assumptions of using a constant and uniform uplift rate, erosion rate and initial shelf slope for Xylokaastro appear justified, as none of those parameters have significantly changed on both timescales (Fig. 4c-e). Even if variations in the initial slope existed within the Xylokaastro sequence and affected the resulting terrace geometry, by averaging the terrace width and cliff height over a 1-5 km wide area we expect to have accounted for such possible lateral variations. Furthermore, the widest terraces in the Xylokaastro sequence are also the widest terraces up to ~25 km further east (Armijo et al., 1996; De Gelder et al., 2019), suggesting that terrace width is influenced more strongly by SL curves than by variable initial slopes.

The temporal resolution may restrict the number of marine terraces produced, given the limitations of the SL curves used. The amount of SL peaks within a curve logically

limit the number of terraces that can be formed. As such, relative lower-resolution curves with few SL peaks like the smooth curve of De Boer et al. (2010), lead to few distinct terraces. The curves with the highest resolution (Rohling et al., 2009; Grant et al., 2014, Fig. S4) show that sharp and short duration peaks within a SL curve can result in extra terraces, as confirmed by the positive correlation between resolution and number of matched terraces (Fig. S5). Detailed studies of MIS 5e show that multiple peaks may have occurred even within one interglacial (e.g. Hearty et al., 2007; Pedroja et al., 2011; Murray-Wallace and Woodroffe, 2014), which is not reflected in most of our selected GMSL and RSL curves. In terms of SL noise, this suggests that short-wavelength peaks are underrepresented in most SL curves and limit the number of matched terraces for the Xylokastro sequence.


### *5.1.2 Approach advantages*

Our analysis of the Xylokastro sequence provides clear advantages over more classic analyses that do not include modeling (e.g. Merritts and Bull, 1989; Armijo et al., 1996; Strobl et al., 2014). Using a range of curves is essential to check the robustness of uplift rate estimates and possible correlations between SL highstands and undated marine terraces (e.g. Caputo, 2007; Yildirim et al., 2013; Pedroja et al., 2018a,b). Our approach expands on these studies by not only using shoreline angles and SL highstands, but the full terrace sequence geometry and complete SL curves. In this manner, we take advantage of the model prediction that higher highstands and longer periods of preceding SL rise lead to wider terraces. Such geometrical trends, with some highstands leading to wider terraces, are also observed in nature (Regard et al., 2017). Additionally, our modeling approach allows for an evaluation of parameters like erosion rates and initial slopes, and their evolution through time, with possible climatic and paleogeographic implications.


Another advantage is that we can analyze which SL curves better reproduce the geometry of a studied marine terrace sequence. For example, it is reasonable that for the Xylokastro sequence, curves based on coral data (dark blue in Fig. 4g) have lower misfits on a 125 ka than 240 ka timescale, since the data on which they are based become sparser with increasing age. Possibly the lower resolution of Bates et al. (2014; curve 9 in Fig. 4g) with respect to Waelbroeck et al. (2002; curve 3 in Fig. 4g) results in higher RMS misfits over 125 ka. Considering curves based on the conversion of  $\delta^{18}O_C$  to  $\delta^{18}O_{SW}$  over 125 ka (light blue in Fig. 4g), the benthic-based curves (1 and 8 in Fig. 4g) have lower RMS-misfits than the planktonic-based curves (2 and 12 in Fig. 4g). This possibly relates to the stronger temperature variability in shallow seas (see Background section), although over 240 ka the planktonic-based curve of Shakun et al. (2015, curve 12 in Fig. 4g) has the lowest RMS misfit. For the inverse ice-sheet modeling curves (green in Fig. 4g), the lowest-resolution and longest timescale curve (35 Ma; curve 7 in Fig. 4g) has a much higher RMS-misfit than the other two, suggesting it is less appropriate for short timescales of 125-240 ka. The hydraulic model RSL curves generally have high RMS misfits (pink in Fig. 4g) on both timescales, whereas the PCA-based curve has relatively low RMS misfits on both timescales (curve 13 in Fig. 4g). Overall, the four SL curves with lowest RMS misfits over 240 ka (curves 4, 5, 12 and 13) are all based on globally distributed datasets, which argues in favor of using such curves.

We infer that the curves with the lowest RMS misfits are the most appropriate to describe SL variations at Xylokaastro to first order, but their lack of temporal resolution produces a lower number of terraces than observed in nature. Contrarily, the two highest-resolution SL curves (6, 10 in Fig. 4f) produce more matched terrace levels, but their high RMS misfits suggest that they are less appropriate to describe first order SL variations (Fig. 9a). The noise tests with high uplift rates (Fig. 8) support this inference: at high uplift rates, the overall sequence geometry is mostly a function of first-order SL variations. The type of short wavelength noise represented within high-resolution SL curves can lead to an increase in terrace levels, but does not significantly change the overall staircase morphology unless for unrealistically extreme values (Fig. 8; amplitude 25 m, wavelength 1 ka).

a Rapidly uplifting, young sequences



b Slowly uplifting, old sequences



**Figure 9. Schematic representation of main results (a)** In the case of rapidly uplifting, young sequences ( $\sim 1.5$  mm/yr and  $\sim 240$  ka in the Xylokaastro case study), a higher resolution SL-curve leads to more matching terraces, whereas a SL-curve with better first-order shape of RSL leads to a lower RMS-misfit with the observed sequence morphology. We list some of the SL-curves that roughly fall into the four categories illustrated for the Xylokaastro case study. **(b)** In the case of slowly uplifting, old sequences ( $\sim 0.1$  mm/yr and  $\sim 2.6$  Ma in the MPT case study), the observed contrast in sequence morphology pre- and post-MPT can be the result of (left) a

contrasting sensitivity to cliff diffusion for narrow terraces with small cliffs formed pre-MPT and wider terraces with larger cliffs formed post-MPT, or (right) the specific arrangement of SL-noise pre-and post-MPT, possibly better constrained by the Rohling et al. (2014) RSL-curve.

### 5.1.3 Glacio-isostatic adjustments

The use of GIA-corrections in the analysis of paleoshorelines has become increasingly popular (e.g. Raymo and Mitrovica, 2012; Creveling et al., 2015; Simms et al., 2016), although they are highly uncertain before the LGM given our limited knowledge of past ice-sheet configuration, discontinuous RSL data availability and uncertainties in ice model parameters. Within our study of the Xylokaastro sequence, the GIA-corrected RSL curve fits the terrace sequence morphology better than any of the other curves (Figs. 3, 4g), encouraging the use of such GIA corrections within coastal terrace studies. However, we note that the improvement that is achieved through the GIA correction is relatively minor in our study (3 m in RMS-misfit; Fig. 5) in comparison to the variations between different GMSL curves (21 m in RMS-misfit; Fig. 4g). Therefore, we do not think that applying GIA-corrections to all GMSL curves would have affected the first-order results of this study. Along the same lines, we conclude that GIA-corrections may be useful, but the use of a wide variety of SL curves is more important to terrace analyses at latitudes comparable to, or lower than Xylokaastro.

In the case of the hydraulic model Gibraltar/Hanish RSL curves, ideally they should be converted to GMSL curves, and then GIA-corrected to create Xylokaastro RSL curves. Following the RSL to GMSL conversions of Rohling et al. (2014) and Grant et al. (2014), the first step implies a decrease in glacial/interstadial SL of several meters, and following our GMSL to RSL conversion, the second step implies an increase in glacial/interstadial SL of several meters (Fig. 5). Therefore, we expect these corrections to roughly balance out, and make a relatively small impact on the results.

### 5.2 The MPT, SL noise and Quaternary Evolution of Staircase Coastal Landscapes

Continuous re-occupation and/or removal of older terraces at low uplift rates leading to relatively few terraces (Fig. 6) is confirmed by both natural observations (e.g. Merritts and Bull, 1989; Armijo et al., 1996) and previous LEM studies (Trenhaile 2002, 2014; Melnick 2016, Pastier et al., 2019). Based on global observations of Neogene-Quaternary strandline sequences, Pedoja et al. (2014) suggested that the change in cyclicity frequency from 40 to 100 ka during the MPT is probably causing a contrast between wide rasas before the MPT, and narrower and better individualized marine terraces after the MPT (inset in Fig. 6a). Within this context, modeling with low uplift rates over 2.6 Ma (Fig. 6) suggests that SL curves based on hydraulic models (Rohling et al., 2014) and corals (Bates et al., 2014) are the most and least successful, respectively, in recreating globally observed sequences. This is opposite to the best fitting ~125 ka results for Xylokaastro, and suggests that different types of SL curves may be suitable for different timescales and/or uplift rates.

Additional processes may become relevant at sites that exhibit a terrace sequence over the entire Quaternary or longer timescales. In cases where uplift is driven by magmatic processes, some studies have noticed that land uplift is episodic and spatially variable rather than continuous (Ramalho et al., 2010; Walker et al., 2016). We note that such episodic uplift would have similar effects as the SL noise we test (Figs. 7, 8), both influencing relative SL in an unpredictable manner. Other studies have focused on the influence of dynamic topography to relative SL (e.g. Conrad and Husson, 2009, and references therein). Over ~125 ka this may contribute to a few meters of uplift/subsidence (Austermann et al., 2017) whereas over Pliocene timescales this may increase to a few tens of meters (Rowley et al., 2013). Such corrections for dynamic topography can be relevant to studies that aim at reconstructing past GMSL from individual sites, or estimating the tectonic contribution to uplift/subsidence rates, but should not be particularly important to our MPT case study. The difference in terrace morphology pre- and post-MPT appears to be a global instead of a local feature (Fig. 2; Pedoja et al., 2014), and should be relatively insensitive to gradual and constant RSL changes associated with dynamic topography (Conrad and Husson, 2009).

The differences amongst SL curves after the MPT (Fig. 1) warrant caution on their reliability, and even more so for the SL curves covering longer timespans (Fig. 7). SL noise like meltwater pulses and Heinrich events are not well reflected in most SL curves on post-LGM timescales (Fig. 1c), and is probably reflected worse on longer timescales. We note that GIA-corrections like those applied for Xylokastro, on the order of a few meters, can similarly be considered as 'noise' within local studies. Different methods for SL-reconstruction rely on assumptions for which its reliability is difficult to verify over long timescales, given the limited independent data to compare against. For all four SL curves used on Quaternary timescales there are methodological concerns (see Background section), and as a consequence, differences amongst the curves are more striking pre- than post-MPT. Post-MPT oscillations are all on the order of ~100 m amplitude (Fig. 6b-e), but pre-MPT oscillations in SL occur with amplitudes of ~25 m (De Boer et al., 2010; Fig. 7c) to ~75 m (Rohling et al., 2014; Fig. 6e). Within LEM studies, this makes the possible resulting geometry of a terrace sequence highly uncertain. It is also striking that for slowly uplifting sequences, noise in a SL curve can have a major impact on sequence morphology already with values of 4 m amplitude (Fig. 7), which is small in comparison to meltwater pulses and Heinrich events (Fig. 1c). As such, SL noise appears to be an important fostering parameter for the geometry of a terrace sequence, especially for slowly uplifting sequences, yet difficult to quantify.

Our results align with those of Trenhaile (2002, 2014), with more wave erosion resulting in a relatively wider sequence post-MPT, although in our models we show that increasing noise may decrease the relative contrast in sequence width pre- and post-MPT (Fig. 7). In natural examples there appears to be a variation in such relative widths, as well as in the prominence of the cliff separating the two (Fig. 2). Given enough sites, perhaps the influence of changing climate cycles, noise, uplift rates and initial slopes for full Quaternary sequences can be deciphered, but given the larger number of variables we suspect this would result in many non-unique solutions.



The observed contrast in pre- and post-MPT morphology is possibly due to the specific arrangement of SL noise components, with the Rohling et al. (2014) curve quantifying those most successfully (Figs. 6, 9b). Alternatively, cliff diffusion provides an easy explanation to the first-order morphological difference before and after the MPT. Smaller amplitude, shorter period SL fluctuations form smaller cliffs and narrower terraces, which diffuse easier into wide rasas (Fig. 6, 9b), and obscure the original sequence morphology. Comparing these two mechanisms, the main difference in resulting sequence morphology lies in the slope of pre-MPT rasas in comparison to post-MPT terraces: whereas the rasa formed using the Rohling et al. (2014) curve has a similar slope to the terraces, a diffused series of narrow, small cliff terraces will result in a rasa with a steeper slope than the post-MPT terraces (Fig. 9b). In natural sites the rasa slope can be easily compared to the terrace slope to distinguish the two mechanisms. In our four examples (Fig. 2c) only the Mancora (Peru) sequence has a clearly steeper pre-MPT slope, pointing towards the importance of SL-noise arrangement. To verify this hypothesis requires a thorough morphological analysis of more long-lasting, slowly uplifting sites.

Our models show that sequences with faster uplift rates, like Xylokastro, are much less sensitive to noise than slowly uplifting sequences (Figs. 7 and 8). Considering the natural availability of terrace sequences there appears to be a trade-off by which the faster uplifting sequences tend to represent shorter time-intervals (Pedoja et al., 2014). Following the terrace degradation models of Anderson et al., (1999), an easy explanation would be the influence of river incision, with higher uplift rates implying a faster base-level drop and more aggressive channel incision. In the case of Xylokastro this appears to be true; although rapid uplift seems to have taken place since ~1 Ma (Armijo et al., 1996; Nixon et al., 2016; Fernandez-Blanco et al., 2019), the associated river incision has removed terraces older than ~240ka (Fig. 4a). Following our modeling results for Xylokastro, the MPT, and relative influence of SL noise, we conclude that within the spectrum of naturally observed sequences, the shorter-timescale, faster uplifting sites offer more potential to reconstruct first order SL variations using LEMs.

### *5.3 Sea-level curves in marine terrace analysis: recommendations and perspectives*

Selecting an appropriate SL curve within marine terrace analysis is not trivial (Caputo, 2007), and can have a significant influence on uplift rate estimates (e.g. Pedoja et al. 2018a,b). Based on our case studies, we can put forward some general recommendations. It is of primary importance to check a variety of SL curves, whether using LEMs or statistical methods. Assessing the reliability of different types of SL curves requires a thorough evaluation of assumptions and uncertainties that are difficult to quantify. Yet, within each type of SL curve it is easier to select the most appropriate curve: logical and justifiable choices can be made based on resolution of the original data, time-span, and spatial variability the SL curve is based on. For some well-studied highstands like MIS 5e and MIS 11c (e.g. Dutton et al., 2015), direct estimates from RSL compilations provide additional estimates. GIA-corrections can improve local RSL predictions, but at latitudes relatively far from ice-sheets, like Xylokastro, these are probably of secondary importance. Using LEMs has advantages over more classical methods, as they take into account the full geometry

of a terrace sequence and full shape of a SL curve. Testing several SL curves through LEMs allows for an assessment of (1) possible ages for undated marine terraces, (2) physical parameters involved in the formation of marine terrace sequences, and (3) the (types of) SL curves which are the most consistent with the geometry of the studied sequences. Marine terrace sequences that are relatively young and rapidly uplifting appear more promising for LEM studies than older, slower uplifting sequences for several reasons. Firstly, the assumptions that longer timescale (>1 Ma) SL curves rely on are highly uncertain. Secondly, for lower uplift rates the sequence morphology becomes more sensitive to SL noise like meltwater pulses, Heinrich events and local GIA-effects, and thirdly, variations of uplift rates and dynamic topography are more likely to occur over longer timescales. For the MPT case study we find that Rohling et al. (2014) RSL curve best reproduces observed sequence morphologies. For our Xylokastro case study, the RMS-misfits indicate that the best-fitting SL curves are two SL curves based on coral data for 0-125 ka, and four GMSL curves based on globally distributed datasets for 0-240 ka (Fig. 9a). These inferences are based on one sequence of 13 terraces, and thus the SL curves that have lower misfits here might not have elsewhere, but to overcome this bias similar analyses can be applied to many locations worldwide (Fig. 2a; Pedoja et al., 2014). The rapidly increasing availability of high-resolution topography, like the Xylokastro DSM used in this study, is a crucial development for such a comparison, and could allow for a global perspective on best-fitting SL curves from marine terraces. One step further would be to use LEMs to directly reconstruct GMSL, through the inversion of terrace sequence morphology with a probabilistic approach. Simultaneous analysis of both uplifting and subsiding sequences could constrain both interglacial and glacial SL, and provide unique new SL constraints.

## 6 Conclusion

For our objective of understanding how SL curves affect modeled marine terrace sequences, we draw the following conclusions from our Xylokastro, MPT and noise case studies:

1. Using a LEM to reconstruct marine terrace sequences and testing several different SL curves allows for a detailed analysis of possible terrace ages, model parameters and the SL-curves that best reproduce a sequence, while utilizing the full geometry of a terrace sequence.
2. Comparing the modeled and observed terrace geometries of rapidly uplifting sequences, the RMS misfit provides an indication how good the first order shape of a SL curve is for that particular site, whereas the number of matched terraces depends on the resolution of a SL curve. For the Xylokastro case study we find lowest RMS misfits for coral based SL-curves over ~125 ka, and SL-curves based on globally distributed datasets for ~240 ka, whereas high-resolution hydraulic model curves result in the most matched terraces on both timescales.
3. GIA-corrections can provide improvements for modeling terrace sequences, but are relatively minor in comparison to differences amongst SL curves at latitudes like Xylokastro or lower.
4. The contrasting terrace sequence morphologies pre- and post-MPT can either be the result of cliff diffusion and/or the specific noise arrangement within a

- SL curve. The dip of the slope with respect to the terraces can be used in nature to distinguish between both mechanisms.
5. Older, slower uplifting sequences are generally less favorable for LEM modeling, because SL-curves over  $>1$  Ma timescales are increasingly uncertain, terrace morphology is more sensitive to SL-noise for low uplift rates, and uplift rates are less likely to be continuous because of tectonic and/or dynamic topography variations.
  6. Young, rapidly uplifting sequences with well-preserved morphologies provide an opportunity of reconstructing past SL through LEM modeling.

## Acknowledgments

GdG, DFB, RA and RL acknowledge funding from the People Programme (Marie Skłodowska-Curie Actions) of the European Union's Seventh Framework Programme under the ITN project ALerT (Grant FP7-PEOPLE-2013-ITN number 607996). GdG also acknowledges a postdoctoral grant from the Centre National d'Etudes Spatiales (CNES, France). DM acknowledges financial support from the Millennium Nucleus The Seismic Cycle Along Subduction Zones funded by the Millennium Scientific Initiative (ICM) of the Chilean Government (grant 1150321) and Chilean National Fund for Development of Science and Technology (FONDECYT; grant 1181479). JJM acknowledges financial support from DFG grant JA 2860/1-1. We thank Arthur Delorme for assistance in producing the DSM, Riccardo Caputo for SL curve data, Stephanie Bates for her spectral analysis code, and Marco Meschis and Jennifer Robertson for fruitful discussions. Numerical computations for the DSM were performed on the S-CAPAD platform, Institut de Physique du Globe de Paris (IPGP), France.

## Data

Pleiades satellite imagery was obtained through the ISIS and Tosca programs of the CNES under an academic license and is not for open distribution. On request, we will provide the DSM calculated from this imagery to any academic researcher who gets approval from CNES (contact [isis-pleiades@cnes.fr](mailto:isis-pleiades@cnes.fr) quoting this paper, [lacassin@ipgp.fr](mailto:lacassin@ipgp.fr) in copy). The altimeter products were produced and distributed by Aviso+ (<https://www.aviso.altimetry.fr/>), as part of the Ssalto ground processing segment. The MATLAB® code used for the landscape evolution modelling is available upon request through [Julius.Jara@geo.uni-potsdam.de](mailto:Julius.Jara@geo.uni-potsdam.de). Additional data can be found in the supplementary information.

## Declarations of interest

None

## References

- Anderson, R. S., Densmore, A. L., & Ellis, M. A. (1999). The generation and degradation of marine terraces. *Basin Research*, 11(1), 7–19.

- Andrews, J. T. (1998). Abrupt changes (Heinrich events) in late Quaternary North Atlantic marine environments: a history and review of data and concepts. *Journal of Quaternary Science: Published for the Quaternary Research Association*, 13(1), 3–16.
- Armijo, R., Meyer, B., King, G. C. P., Rigo, A., & Papanastassiou, D. (1996). Quaternary evolution of the Corinth Rift and its implications for the Late Cenozoic evolution of the Aegean. *Geophysical Journal International*, 126(1), 11–53.
- Austermann, J., Mitrovica, J. X., Huybers, P., & Rovere, A. (2017). Detection of a dynamic topography signal in last interglacial sea-level records. *Science Advances*, 3(7). <https://doi.org/10.1126/sciadv.1700457>
- Authemayou, C., Pedoja, K., Heddar, A., Molliex, S., Boudiaf, A., Ghaleb, B., ... Nexer, M. (2017). Coastal uplift west of Algiers (Algeria): pre- and post-Messinian sequences of marine terraces and rasas and their associated drainage pattern. *International Journal of Earth Sciences*, 106(1), 19–41.
- Bard, E., Hamelin, B., Arnold, M., Montaggioni, L., Cabioch, G., Faure, G., & Rougerie, F. (1996). Deglacial sea-level record from Tahiti corals and the timing of global meltwater discharge. *Nature*, 382(6588), 241–244.
- Bard, E., Hamelin, B., & Fairbanks, R. G. (1990). U-Th ages obtained by mass spectrometry in corals from Barbados: sea level during the past 130,000 years. *Nature*, 346(6283), 456–458.
- Bates, S. L., Siddall, M., & Waelbroeck, C. (2014). Hydrographic variations in deep ocean temperature over the mid-Pleistocene transition. *Quaternary Science Reviews*, 88, 147–158.
- Bintanja, R., & van de Wal, R. S. W. (2008). North American ice-sheet dynamics and the onset of 100,000-year glacial cycles. *Nature*, 454(7206), 869–872.
- Bintanja, R., van de Wal, R. S. W., & Oerlemans, J. (2005). Modelled atmospheric temperatures and global sea levels over the past million years. *Nature*, 437(7055), 125–128.
- Bloom, A. L. (1967). Pleistocene Shorelines: A New Test of Isostasy. *GSA Bulletin*, 78(12), 1477–1494.
- Bloom, A. L. (1971). Glacial-eustatic and isostatic controls of sea level since the last glaciation. *Late Cenozoic Glacial Ages*, 355–379.
- Bowles, C. J., & Cowgill, E. (2012). Discovering marine terraces using airborne LiDAR along the Mendocino-Sonoma coast, northern California. *Geosphere*, 8(2), 386–402.
- Bradley, W. C. (1958). Submarine abrasion and wave-cut platforms. *GSA Bulletin*, 69(8), 967–974.
- Braithwaite, C. J. R. (2016). Coral-reef records of Quaternary changes in climate and sea-level. *Earth-Science Reviews*, 156, 137–154.
- Caputo, R. (2007). Sea-level curves: Perplexities of an end-user in morphotectonic applications. *Global and Planetary Change*, 57(3), 417–423.

- Caputo, R., Bianca, M., & D'Onofrio, R. (2010). Ionian marine terraces of southern Italy: Insights into the Quaternary tectonic evolution of the area. *Tectonics*, 29(4), TC4005.
- Chappell, J. (1974). *Geology of Coral Terraces, Huon Peninsula, New Guinea: A Study of Quaternary Tectonic Movements and Sea-Level Changes*. GSA Bulletin, 85(4), 553–570.
- Chappell, J., & Shackleton, N. J. (1986). Oxygen isotopes and sea level. *Nature*, 324(6093), 137–140.
- Clark, P. U., Archer, D., Pollard, D., Blum, J. D., Rial, J. A., Brovkin, V., ... Roy, M. (2006). The middle Pleistocene transition: characteristics, mechanisms, and implications for long-term changes in atmospheric pCO<sub>2</sub>. *Quaternary Science Reviews*, 25(23), 3150–3184.
- Collier, R. E. L., Leeder, M. R., Rowe, P. J., & Atkinson, T. C. (1992). Rates of tectonic uplift in the Corinth and Megara Basins, central Greece. *Tectonics*, 11(6), 1159–1167.
- Conrad, C. P., & Husson, L. (2009). Influence of dynamic topography on sea level and its rate of change. *Lithosphere*, 1(2), 110–120.
- Creveling, J. R., Mitrovica, J. X., Hay, C. C., Austermann, J., & Kopp, R. E. (2015). Revisiting tectonic corrections applied to Pleistocene sea-level highstands. *Quaternary Science Reviews*, 111, 72–80.
- De Boer, B., Van de Wal, R., Bintanja, R., Lourens, L. J., & Tuenter, E. (2010). Cenozoic global ice-volume and temperature simulations with 1-D ice-sheet models forced by benthic 18O records. *Annals of Glaciology*, 51(55), 23–33.
- De Gelder, G., Fernández-Blanco, D., Lacassin, R., Armijo, R., Delorme, A., Jara-Muñoz, J., & Melnick, D. (2015). Corinth terraces re-visited: Improved paleoshoreline determination using Pleiades-DEMs. *Geotectonic Research*, 97, 12–14.
- De Gelder, G., Fernández-Blanco, D., Melnick, D., Duclaux, G., Bell, R. E., Jara-Muñoz, J., ... Lacassin, R. (2019). Lithospheric flexure and rheology determined by climate cycle markers in the Corinth Rift. *Scientific Reports*, 9(1), 4260.
- Dia, A. N., Cohen, A. S., O'Nions, R. K., & Jackson, J. A. (1997). Rates of uplift investigated through 230 Th dating in the Gulf of Corinth (Greece). *Chemical Geology*, 138(3), 171–184.
- Dufaure, J.-J., & Zamanis, A. (1979). Un vieux problème géomorphologique: les niveaux bordiers au sud du Golfe de Corinthe (An old geomorphological problem: the levels developed on the southern border of the gulf of Corinth). *Bulletin de l'Association de Géographes Français*, 56(464), 341–350.
- Dutton, A., Carlson, A. E., Long, A. J., Milne, G. A., Clark, P. U., DeConto, R., ... Raymo, M. E. (2015). Sea-level rise due to polar ice-sheet mass loss during past warm periods. *Science*, 349(6244), aaa4019.

- Dwyer, G. S., Cronin, T. M., Baker, P. A., Raymo, M. E., Buzas, J. S., & Corrige, T. (1995). North Atlantic deepwater temperature change during late Pliocene and late Quaternary climatic cycles. *Science*, (270), 1347–1351.
- Elderfield, H., Ferretti, P., Greaves, M., Crowhurst, S., McCave, I. N., Hodell, D., & Piotrowski, A. M. (2012). Evolution of ocean temperature and ice volume through the mid-Pleistocene climate transition. *Science*, 337(6095), 704–709.
- Fairbanks, R. G. (1989). A 17,000-year glacio-eustatic sea level record: influence of glacial melting rates on the Younger Dryas event and deep-ocean circulation. *Nature*, 342(6250), 637–642.
- Fernández-Blanco, D., de Gelder, G., Lacassin, R., & Armijo, R. (2019). A new crustal fault formed the modern Corinth Rift. Retrieved from <https://eartharxiv.org/c6mey/>
- Grant, K. M., Rohling, E. J., Ramsey, C. B., Cheng, H., Edwards, R. L., Florindo, F., ... Williams, F. (2014). Sea-level variability over five glacial cycles. *Nature Communications*, 5, 5076.
- Hearty, P. J., Hollin, J. T., Neumann, A. C., O’Leary, M. J., & McCulloch, M. (2007). Global sea-level fluctuations during the Last Interglaciation (MIS 5e). *Quaternary Science Reviews*, 26(17), 2090–2112.
- Henry, H., Regard, V., Pedoja, K., Husson, L., Martinod, J., Witt, C., & Heuret, A. (2014). Upper Pleistocene uplifted shorelines as tracers of (local rather than global) subduction dynamics. *Journal of Geodynamics*, 78, 8–20.
- Hibbert, F. D., Rohling, E. J., Dutton, A., Williams, F. H., Chutcharavan, P. M., Zhao, C., & Tamisiea, M. E. (2016). Coral indicators of past sea-level change: A global repository of U-series dated benchmarks. *Quaternary Science Reviews*, 145, 1–56.
- Husson, L., Pastier, A.-M., Pedoja, K., Elliot, M., Paillard, D., Authemayou, C., ... Cahyarini, S. Y. (2018). Reef Carbonate Productivity During Quaternary Sea Level Oscillations. *Geochemistry, Geophysics, Geosystems*, 19(4), 1148–1164.
- Jara-Muñoz, J., Melnick, D., & Strecker, M. R. (2016). TerraceM: A MATLAB® tool to analyze marine and lacustrine terraces using high-resolution topography. *Geosphere*, 12(1), 176–195.
- Jara-Muñoz, J., Melnick, D., Zambrano, P., Rietbrock, A., González, J., Argandoña, B., & Strecker, M. R. (2017). Quantifying offshore forearc deformation and splay-fault slip using drowned Pleistocene shorelines, Arauco Bay, Chile. *Journal of Geophysical Research, [Solid Earth]*, 122(6), 4529–4558.
- Johnston, P. (1993). The effect of spatially non-uniform water loads on prediction of sea-level change. *Geophysical Journal International*, 114(3), 615–634.
- Keraudren, B., & Sorel, D. (1987). The terraces of Corinth (Greece) — A detailed record of eustatic sea-level variations during the last 500,000 years. *Marine Geology*, 77(1), 99–107.
- Lajoie, K. R. (1986). Coastal tectonics. *Active Tectonics*.

- Lambeck, K. (1995). Late Pleistocene and Holocene sea-level change in Greece and south-western Turkey: a separation of eustatic, isostatic and tectonic contributions. *Geophysical Journal International*, 122(3), 1022–1044.
- Lambeck, K., & Chappell, J. (2001). Sea level change through the last glacial cycle. *Science*, 292(5517), 679–686.
- Lambeck, K., Esat, T. M., & Potter, E.-K. (2002). Links between climate and sea levels for the past three million years. *Nature*, 419(6903), 199–206.
- Lambeck, K., Purcell, A., & Dutton, A. (2012). The anatomy of interglacial sea levels: The relationship between sea levels and ice volumes during the Last Interglacial. *Earth and Planetary Science Letters*, 315, 4–11.
- Lambeck, K., Purcell, A., Funder, S., Kjær, K., Larsen, E., & Möller, P. (2006). Constraints on the Late Saalian to early Middle Weichselian ice sheet of Eurasia from field data and rebound modelling. *Boreas*, 35(3), 539–575.
- Lambeck, K., Purcell, A., Johnston, P., Nakada, M., & Yokoyama, Y. (2003). Water-load definition in the glacio-hydro-isostatic sea-level equation. *Quaternary Science Reviews*, 22(2), 309–318.
- Lambeck, K., Purcell, A., Zhao, J., & Svensson, N.-O. (2010). The Scandinavian Ice Sheet: from MIS 4 to the end of the Last Glacial Maximum. *Boreas*, 39(2), 410–435.
- Lambeck, K., Purcell, A., & Zhao, S. (2017). The North American Late Wisconsin ice sheet and mantle viscosity from glacial rebound analyses. *Quaternary Science Reviews*, 158, 172–210.
- Lambeck, K., Rouby, H., Purcell, A., Sun, Y., & Sambridge, M. (2014). Sea level and global ice volumes from the Last Glacial Maximum to the Holocene. *Proceedings of the National Academy of Sciences of the United States of America*, 111(43), 15296–15303.
- Lea, D. W., Martin, P. A., Pak, D. K., & Spero, H. J. (2002). Reconstructing a 350ky history of sea level using planktonic Mg/Ca and oxygen isotope records from a Cocos Ridge core. *Quaternary Science Reviews*, 21(1), 283–293.
- Leeder, M. R., Portman, C., Andrews, J. E., Collier, R. E. L., Finch, E., Gawthorpe, R. L., ... Rowe, P. (2005). Normal faulting and crustal deformation, Alkyonides Gulf and Perachora peninsula, eastern Gulf of Corinth rift, Greece. *Journal of the Geological Society*, 162(3), 549–561.
- Lisiecki, L. E., & Raymo, M. E. (2005). A Pliocene-Pleistocene stack of 57 globally distributed benthic  $\delta^{18}\text{O}$  records. *Paleoceanography*, 20(PA1003).
- Ludwig, K. R., Szabo, B. J., Moore, J. G., & Simmons, K. R. (1991). Crustal subsidence rate off Hawaii determined from  $^{234}\text{U}/^{238}\text{U}$  ages of drowned coral reefs. *Geology*, 19(2), 171–174.
- Melnick, D. (2016). Rise of the central Andean coast by earthquakes straddling the Moho. *Nature Geoscience*, 9(5), 401–407.

- Merritts, D., & Bull, W. B. (1989). Interpreting Quaternary uplift rates at the Mendocino triple junction, northern California, from uplifted marine terraces. *Geology*, 17(11), 1020–1024.
- Moretti, I., Lykousis, V., Sakellariou, D., Reynaud, J.-Y., Benziane, B., & Prinzhofer, A. (2004). Sedimentation and subsidence rate in the Gulf of Corinth: what we learn from the Marion Dufresne's long-piston coring. *Comptes Rendus: Geoscience*, 336(4), 291–299.
- Murray-Wallace, C. V., & Woodroffe, C. D. (2014). *Quaternary Sea-Level Changes: A Global Perspective*. Cambridge University Press.
- Nakada, M., & Lambeck, K. (1989). Late Pleistocene and Holocene sea-level change in the Australian region and mantle rheology. *Geophysical Journal International*, 96(3), 497–517.
- Nixon, C. W., McNeill, L. C., Bull, J. M., Bell, R. E., Gawthorpe, R. L., Henstock, T. J., ... Kranis, H. (2016). Rapid spatiotemporal variations in rift structure during development of the Corinth Rift, central Greece. *Tectonics*, 35(5), 2015TC004026.
- Pedoja, K., Husson, L., Bezos, A., Pastier, A.-M., Imran, A. M., Arias-Ruiz, C., ... Choblet, G. (2018). On the long-lasting sequences of coral reef terraces from SE Sulawesi (Indonesia): Distribution, formation, and global significance. *Quaternary Science Reviews*, 188, 37–57.
- Pedoja, K., Husson, L., Johnson, M. E., Melnick, D., Witt, C., Pochat, S., ... Garestier, F. (2014). Coastal staircase sequences reflecting sea-level oscillations and tectonic uplift during the Quaternary and Neogene. *Earth-Science Reviews*, 132, 13–38.
- Pedoja, K., Husson, L., Regard, V., Cobbold, P. R., Ostanciaux, E., Johnson, M. E., ... Delcaillau, B. (2011). Relative sea-level fall since the last interglacial stage: Are coasts uplifting worldwide? *Earth-Science Reviews*, 108(1), 1–15.
- Pedoja, K., Jara-Muñoz, J., De Gelder, G., Robertson, J., Meschis, M., Fernandez-Blanco, D., ... Pinel, B. (2018). Neogene-Quaternary slow coastal uplift of Western Europe through the perspective of sequences of strandlines from the Cotentin Peninsula (Normandy, France). *Geomorphology*, 303, 338–356.
- Pedoja, K., Ortlieb, L., Dumont, J. F., Lamothe, M., Ghaleb, B., Auclair, M., & Labrousse, B. (2006). Quaternary coastal uplift along the Talara Arc (Ecuador, Northern Peru) from new marine terrace data. *Marine Geology*, 228(1), 73–91.
- Perissoratis, C., Piper, D. J. W., & Lykousis, V. (2000). Alternating marine and lacustrine sedimentation during late Quaternary in the Gulf of Corinth rift basin, central Greece. *Marine Geology*, 167(3–4), 391–411.
- Pierini, F., Demarchi, B., Turner, J., & Penkman, K. (2016). Pecten as a new substrate for IcPD dating: The quaternary raised beaches in the Gulf of Corinth, Greece. *Quaternary Geochronology*, 31, 40–52.
- Quartau, R., Trenhaile, A. S., Mitchell, N. C., & Tempera, F. (2010). Development of volcanic insular shelves: Insights from observations and modelling of Faial Island in the Azores Archipelago. *Marine Geology*, 275(1), 66–83.



- Railsback, L. B., Gibbard, P. L., Head, M. J., Voarintsoa, N. R. G., & Toucanne, S. (2015). An optimized scheme of lettered marine isotope substages for the last 1.0 million years, and the climatostratigraphic nature of isotope stages and substages. *Quaternary Science Reviews*, 111, 94–106.
- Ramalho, R., Helffrich, G., Cosca, M., Vance, D., Hoffmann, D., & Schmidt, D. N. (2010). Episodic swell growth inferred from variable uplift of the Cape Verde hotspot islands. *Nature Geoscience*, 3, 774.
- Raymo, M. E., & Mitrovica, J. X. (2012). Collapse of polar ice sheets during the stage 11 interglacial. *Nature*, 483(7390), 453–456.
- Regard, V., Pedoja, K., De La Torre, I., Saillard, M., Cortés-Aranda, J., & Nexer, M. (2017). Geometrical trends within sequences of Pleistocene marine terraces: selected examples from California, Peru, Chile and New-Zealand. *Zeitschrift Fur Geomorphologie*, 61(1), 53–73.
- Roberts, G. P., Meschis, M., Houghton, S., Underwood, C., & Briant, R. M. (2013). The implications of revised Quaternary palaeoshoreline chronologies for the rates of active extension and uplift in the upper plate of subduction zones. *Quaternary Science Reviews*, 78, 169–187.
- Robertson, J., Meschis, M., Roberts, G. P., Ganas, A., & Gheorghiu, D. M. (2019). Temporally Constant Quaternary Uplift Rates and Their Relationship With Extensional Upper-Plate Faults in South Crete (Greece), Constrained With 36 Cl Cosmogenic Exposure Dating. *Tectonics*, 38(4), 1189–1222.
- Rohling, E. J., Foster, G. L., Grant, K. M., Marino, G., Roberts, A. P., Tamisiea, M. E., & Williams, F. (2014). Sea-level and deep-sea-temperature variability over the past 5.3 million years. *Nature*, 508(7497), 477–482.
- Rohling, E. J., Grant, K., Bolshaw, M., Roberts, A. P., Siddall, M., Hemleben, C., & Kucera, M. (2009). Antarctic temperature and global sea level closely coupled over the past five glacial cycles. *Nature Geoscience*, 2(7), 500.
- Rowley, D. B., Forte, A. M., Moucha, R., Mitrovica, J. X., Simmons, N. A., & Grand, S. P. (2013). Dynamic topography change of the eastern United States since 3 million years ago. *Science*, 340(6140), 1560–1563.
- Sarr, A.-C., Husson, L., Sepulchre, P., Pastier, A.-M., Pedoja, K., Elliot, M., ... Susilohadi. (2019). Subsiding Sundaland: REPLY. *Geology*, 47(7), e470–e470.
- Shackleton, N. J. (2000). The 100,000-year ice-Age cycle identified and found to lag temperature, carbon dioxide, and orbital eccentricity. *Science*, 289(5486), 1897–1902.
- Shakun, J. D., Lea, D. W., Lisiecki, L. E., & Raymo, M. E. (2015). An 800-kyr record of global surface ocean  $\delta^{18}O$  and implications for ice volume-temperature coupling. *Earth and Planetary Science Letters*, 426, 58–68.
- Shaw, B., Ambraseys, N. N., England, P. C., Floyd, M. A., Gorman, G. J., Higham, T., ... Piggott, M. D. (2008). Eastern Mediterranean tectonics and tsunami hazard inferred from the AD 365 earthquake. *Nature Geoscience*, 1(4), 268.

- Siddall, M., Hönisch, B., Waelbroeck, C., & Huybers, P. (2010). Changes in deep Pacific temperature during the mid-Pleistocene transition and Quaternary. *Quaternary Science Reviews*, 29(1), 170–181.
- Siddall, M., Rohling, E. J., Almogi-Labin, A., Hemleben, C., Meischner, D., Schmelzer, I., & Smeed, D. A. (2003). Sea-level fluctuations during the last glacial cycle. *Nature*, 423(6942), 853–858.
- Simms, A. R., Rouby, H., & Lambeck, K. (2016). Marine terraces and rates of vertical tectonic motion: The importance of glacio-isostatic adjustment along the Pacific coast of central North America. *GSA Bulletin*, 128(1-2), 81–93.
- Sosdian, S., & Rosenthal, Y. (2009). Deep-sea temperature and ice volume changes across the Pliocene-Pleistocene climate transitions. *Science*, 325(5938), 306–310.
- Spratt, R. M., & Lisiecki, L. E. (2016). A Late Pleistocene sea level stack. *Climate of the Past*, 12(4), 1079.
- Strobl, M., Hetzel, R., Fassoulas, C., & Kubik, P. W. (2014). A long-term rock uplift rate for eastern Crete and geodynamic implications for the Hellenic subduction zone. *Journal of Geodynamics*, 78, 21–31.
- Sunamura, T. (1992). *Geomorphology of rocky coasts* (Vol. 3). John Wiley & Son Ltd.
- Trenhaile, A. (2014). Modelling the effect of Pliocene--Quaternary changes in sea level on stable and tectonically active land masses. *Earth Surface Processes and Landforms*, 39(9), 1221–1235.
- Trenhaile, A. S. (2002). Modeling the development of marine terraces on tectonically mobile rock coasts. *Marine Geology*, 185(3), 341–361.
- Waelbroeck, C., Labeyrie, L., Michel, E., Duplessy, J. C., McManus, J. F., Lambeck, K., ... Labracherie, M. (2002). Sea-level and deep water temperature changes derived from benthic foraminifera isotopic records. *Quaternary Science Reviews*, 21(1), 295–305.
- Walcott, R. I. (1972). Past Sea Levels, Eustasy and Deformation of the Earth. *Quaternary Research*, 2(1), 1–14.
- Walker, R. T., Telfer, M., Kahle, R. L., Dee, M. W., Kahle, B., Schwenninger, J.-L., ... Watts, A. B. (2016). Rapid mantle-driven uplift along the Angolan margin in the late Quaternary. *Nature Geoscience*, 9, 909.
- Ward, C. M. (1988). Marine terraces of the Waitutu district and their relation to the late Cenozoic tectonics of the southern Fiordland region, New Zealand. *Journal of the Royal Society of New Zealand*, 18(1), 1–28.
- Yamato, P., Husson, L., Becker, T. W., & Pedoja, K. (2013). Passive margins getting squeezed in the mantle convection vice. *Tectonics*, 32(6), 2013TC003375.
- Yildirim, C., Melnick, D., Ballato, P., Schildgen, T. F., Echtler, H., Erginal, A. E., ... Strecker, M. R. (2013). Differential uplift along the northern margin of the Central Anatolian Plateau: inferences from marine terraces. *Quaternary Science Reviews*, 81, 12–28.

Zeuner, F. E. (1952). Pleistocene Shore-lines. *Geologische Rundschau: Zeitschrift Fur Allgemeine Geologie*, 40(1), 39–50.