

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29

---

This manuscript is currently undergoing peer-review. Please note that the manuscript is yet to be formally accepted for publication. Subsequent versions of this manuscript may have slightly different content. If accepted, the final version of this manuscript will be available via the ‘Peer-reviewed Publication DOI’ link on the right-hand side of this webpage. Please feel free to contact any of the authors. We look forward to your feedback.

---

30 **CONTROLS OF BASEMENT FABRIC ON RIFT COUPLING AND DEVELOPMENT**  
31 **OF NORMAL FAULT GEOMETRIES: INSIGHTS FROM THE RUKWA – NORTH**  
32 **MALAWI RIFT**  
33  
34  
35  
36  
37

38 **Erin Heilman<sup>1</sup>**

39 **Folarin Kolawole<sup>2</sup>**

40 **Estella A. Atekwana<sup>3\*</sup>**

41 **Micah Mayle<sup>1</sup>**

42 **Mohamed G. Abdelsalam<sup>1</sup>**

43  
44  
45  
46 *<sup>1</sup>Boone Pickens School of Geology*  
47 *Oklahoma State University*  
48 *Stillwater, Oklahoma, USA*

49  
50 *<sup>2</sup>ConocoPhillips School of Geology & Geophysics*  
51 *University of Oklahoma*  
52 *Norman, Oklahoma, USA*

53  
54 *<sup>3</sup>Department of Geological Sciences*  
55 *College of Earth, Ocean, and Environment*  
56 *University of Delaware*  
57 *Newark, Delaware, USA*

58  
59 *\*Corresponding author email: atekwana@udel.edu*  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71

72 August 2018

73 **Highlights**

- 74 • To the SW, newfound strike-slip fault links the Rukwa and North Malawi Rift (RNMRS)
- 75 • To the NE, RNMRS border faults, intervening faults and volcanic centers are colinear
- 76 • RNMRS border faults and transfer structures align with pre-existing basement fabrics
- 77 • Basement fabrics guide the development of normal fault geometries and rift bifurcation
- 78 • Basement fabrics facilitate the coupling of the RMRS border faults and transfer structures

79

80

81 **ABSTRACT**

82 The Rukwa Rift and North Malawi Rift Segments (RNMRS) both define a major rift-oblique  
83 segment of the East African Rift System (EARS), and although the two young rifts show colinear  
84 approaching geometries, they are often regarded as discrete rifts due to the presence of the  
85 intervening Mbozi Block uplift located in-between. This problem has been complicated by the  
86 dominance of the Rungwe volcanic features along the northeastern boundary of the Mbozi Block  
87 and lack of distinct normal faults along the southwestern boundary of the block. Here, we  
88 investigate the coupling of discrete rift segments during the onset of continental rifting,  
89 modulated by the control of pre-existing basement fabrics on the development of the border fault  
90 geometries and linkage across the intra-rift transfer zone. We utilized the Shuttle Radar  
91 Topography Mission Digital Elevation Models (SRTM-DEM) to investigate the morphological  
92 architecture of the rift domains; and aeromagnetic data and SRTM-DEM to assess the  
93 relationships between the rift structures and the pre-existing basement fabric (in plan-view). Our  
94 results show that the present-day morphology of the RNMRS is characterized by along-rift  
95 alternation of rift shoulder polarity, characteristic of coupled rift segments. Careful interpretation

96 of filtered aeromagnetic maps along the northeastern and southwestern boundaries of the  
97 RNMRS reveal striking alignment of the rift-bounding faults with colinear NW-SE-trending pre-  
98 existing basement fabrics. We find that rift coupling along the northeastern boundary of the  
99 Mbozi Block transfer zone is accommodated by magmatism utilizing pre-existing fault systems,  
100 whereas, coupling along the southwestern boundary is accommodated by a new-found dextral  
101 strike-slip fault. Additionally, we show how the configuration of the pre-existing basement  
102 fabrics may influence the development of rectilinear or curvilinear normal fault geometries  
103 (plan-view) along the rifts, and the formation of basin-scale rift bifurcation around basement  
104 inter-rift transfer zones. In summary, we suggest that the structural continuation of the boundary  
105 faults along the RNMRS, and their alignment with colinear basement fabrics demonstrate the  
106 influence of structural inheritance on the coupling and amalgamation of approaching rift  
107 segments.

108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118

119 **1. INTRODUCTION**

120 Pre-existing basement fabrics are often major facilitators of continental rifting environments.  
121 Mechanically, they represent areas of structural weakness that can become reactivated and allow  
122 rifts to propagate preferentially along them (e.g., Daly et al., 1989). Several studies have  
123 documented the relationships between rift faults and the pre-existing basement fabrics (e.g.,  
124 Wheeler and Karson, 1989; Kinabo et al., 2007, 2008; Taylor et al., 2011; Phillips et al., 2016;  
125 Kolawole et al., 2018; Siuda et al., 2018). Further, recent studies have assessed the 3-  
126 dimensional relationship between pre-existing basement thrusts and intra-rift normal faults,  
127 revealing the control of the pre-existing basement structures on the nucleation and strain  
128 distribution along the normal faults (e.g., Collanega et al., 2018). In the Cenozoic East African  
129 Rift System (EARS), which is divided into an Eastern, a Western and a Southwestern Branch  
130 (Fig. 1A), several zones of well-developed basement fabric influence rifting. One of the best  
131 recently-documented examples highlighting the influence of the Precambrian basement shear  
132 zones on rifting in eastern Africa is the role of the Mwembeshi Shear Zone on the development  
133 of the Luangwa Rift (Fig. 1A; Sarafian et al., 2018). It was demonstrated that the Mwembeshi  
134 Shear Zone acted as lithospheric conduit for fluids to migrate up the lithosphere, thus facilitating  
135 the weakening and subsequent initiation of the Luangwa rift in the Permo-Triassic.

136         However, the relationship between rift segments along the western branch of the EARS  
137 and the Precambrian shear zones is rather complex and warrants detailed and careful  
138 examination. For example, the Precambrian NW-trending Aswa shear zone resulted in the  
139 termination (rather than facilitation) of the northeastward propagation of the Albertine-Rhino  
140 graben which represents the northern-most segment of the Western Branch (Katumwehe et al.,  
141 2015) (Fig. 1A). On a basin scale, previous studies have also shown that pre-existing basement

142 shear zones can influence the localization of fault development (e.g., Phillips et al., 2016;  
143 Kolawole et al., 2018), and in fact control fault segmentation and across-basin strain transfer at  
144 later stages of rift development (e.g., Muirhead and Kattenhorn, 2018).

145         The border faults along large continental rift systems, e.g., the EARS, are typically ~100  
146 km long (e.g., Foster et al., 1997; Lao-Davila et al., 2015), and the development of such large  
147 normal faults with complex segment linkage styles are yet to be fully understood (e.g., Fossen  
148 and Rotevatn, 2016; Gawthorpe et al., 2003; Rotevatn et al., 2018). However, since the  
149 interactions between the large normal faults within juvenile extensional tectonic settings lead to  
150 the systematic coupling of rift segments across transfer zones (e.g., Corti, 2012), border fault  
151 segmentation, geometries and continuity between rift segments can provide insight into the  
152 larger process of coupling between the segments of a rift system.

153         In this study, we focus on the Rukwa-North Malawi segment of the EARS, which is a  
154 major rift-oblique segment of the rift system and serves as the central segment of the system. For  
155 simplicity, we here-in refer to the Rukwa - Northern Malawi Rift segment of the East African  
156 Rift as the “RNMRS”. This segment is composed of the Rukwa Rift basin, the North Malawi  
157 Rift basin and the Mbozi Block which represents the accommodation zone between the two rifts  
158 (Fig. 1B). We address the longstanding question of the role of long-lived pre-existing basement  
159 structures in the development of the trends and geometries of rift-bounding faults, leading to  
160 subsequent coupling of individual rift segments during the onset of continental rifting. We  
161 demonstrate that there is continuous structural connectivity along the boundaries of the RNMRS,  
162 modulated by reactivation of the Precambrian metamorphic fabrics, and show that the  
163 characteristic plan-view geometries of the rift-bounding faults are modulated by the  
164 configuration of the basement fabric.

165 **2. GEOLOGIC SETTING**

166 *2.1. The Precambrian Domains*

167 The Rukwa-North Malawi Rift Segment is located within the NW-trending Paleoproterozoic  
168 Ubendian orogenic belt which is sandwiched between the Archean Tanzanian craton in the  
169 northeast and the Bangweulu cratonic block to the southwest (Figs. 2A-B; Fritz et al., 2013). The  
170 Ubendian Belt is composed of different Precambrian terranes bounded by steep shear zones  
171 (Delvaux et al., 2012). These terranes contain granulite-facies metamorphic rocks (2100-2025  
172 Ma), amphibolite-facies metamorphic rocks and granitoids (1960-1800 Ma) that have undergone  
173 dextral strike-slip shearing and granitic plutons (1090-1120 Ma), (Fritz et al., 2013). The  
174 Paleoproterozoic Usagaran orogenic belt that extends NE-SW perpendicular to the Ubendian  
175 orogenic belt in southern Tanzania, is composed of eclogites (~2000 Ma), volcano-sedimentary  
176 cover with some low-grade metamorphism (~1920 Ma), and granitoids and granitoid gneisses  
177 (1900-1730 Ma) (Fritz et al., 2013). The Usagaran and Ubendian orogenic belts resulted from  
178 collision with the Tanzania craton, where the Usagaran orogenic belt was thrust onto the craton  
179 and the Ubendian orogenic belt was accreted along the craton's margin because of strike-slip  
180 motion (Daly, 1988; Lenoir et al., 1994). The Ubendian Belt was reactivated several times, first  
181 at ~1860 Ma as a shear zone, then again at ~800 Ma and was subsequently reactivated about  
182 every 200 Ma (Lenoir et al., 1994). Mapped shear zones along the Ubendian Belt include the  
183 ~600 km long and ~30 km-wide Mughese Shear Zone, the Mtose Shear Zone, and the Chisi  
184 Shear Zone (Fig. 2; Daly, 1988; Schenk et al., 2006; Delvaux et al., 2012). Recent studies have  
185 highlighted the role of the Mughese Shear Zone in fault development and the distribution of  
186 seismicity in Northern Malawi Rift Basin (Dawson et al. 2017; Kolawole et al., 2018).

187

188 2.2. *The Rukwa Rift, Northern Malawi Rift, and the Mbozi Block accommodation zone (RNMRS)*  
189 The RNMRS evolved during the Permo-Triassic episode of rifting that affected southern and  
190 eastern Africa, also known as Karoo rifting (Chorowicz, 2005). Outcrops of Karoo sediments  
191 have been mapped along the southern end of the Rukwa Rift (Figure 2) and the northern section  
192 of the North Malawi Rift Basin (Kilembe and Rosendahl, 1992). These sediments lie  
193 unconformably over the Precambrian basement and consist mainly of sandstone, shale, and coal  
194 and thicken towards the border faults providing evidence for reactivation of synthetic faulting in  
195 the Permo-Triassic (Morley, 1992; Delvaux et al., 1992).

196 Cenozoic rifting began in the Upper Miocene, characterized by normal faulting and basin  
197 subsidence with the diagenesis of Red Sandstones and Lake Bed Sediments (Delvaux and  
198 Hanon, 1991). Additional subsidence occurred after the deposition of these packages, and in  
199 different directions, evidenced by the drag orientations of sediment packages on the faults  
200 (Kilembe and Rosendahl, 1992). The Cenozoic rifting featured the reactivation of older faults as  
201 seen in seismic profiles of the Rukwa Rift in which the faults are mostly contiguous from Karoo  
202 sediments to Red Sandstones to Lake Beds (Kervyn et al., 2006). The present-day tectonic  
203 activity in the RNMRS consists of limited volcanic eruptions, minor seismicity in the Mbozi  
204 Block region, and continued sedimentation in the Rukwa and North Malawi basins (Delvaux and  
205 Hanon, 1991).

206 The present-day architecture of the RNMRS consists of the Rukwa Rift to the northwest  
207 of the segment, with the Lupa Fault (generally considered the border fault) bounding it to the  
208 northeast and the Ufipa Fault to the SW (Fig. 1B). Towards the southeastern end of the Rukwa  
209 Rift. It bifurcates around the Mbozi Block forming the Songwe Trough (ST) to the northeast and  
210 the Musangano Trough (MT) to the southwest. The Mbozi Block transitions into the North



211 Malawi Rift (also known as the North Basin) which represents the southeastern end of the  
212 RNMRS and it consists of a half-graben, bounded to the northeast by the Livingstone Fault (Fig.  
213 1B). The Mbozi Block is referred to as an accommodation zone because it is thought to  
214 accommodate and transfer relative strain between the Rukwa and North Malawi Rift Basins  
215 (Delvaux and Hanon, 1993)

216 The Mbozi is a mass of Precambrian basement that is composed of Meta-basites and  
217 intermediate granulites and quartzites of the Mbozi Terrane (Daly, 1988) and is bounded to the  
218 southwestern by the Mughese Shear Zone (Fig. 2B), and overlain on the northeast by the  
219 volcanic deposits of the Rungwe Volcanic Province (RVP). The RVP is a ~1500 km<sup>2</sup> area of  
220 volcanic rocks and structures that evolved ca. 9 Ma (e.g., Fontijn et al., 2012), and a strong  
221 tectonic control on the localization of volcanic centers have been inferred (e.g., Fontijn et al.,  
222 2010, 2012). To the northeast of the RVP, a poorly defined NE-trending rift basin occurs, known  
223 as the “Usangu Basin”, where Permo-Triassic to Recent sedimentary rocks overlie the  
224 Precambrian basement (Mbede, 2002).

225 The crustal thickness beneath the Rukwa Rift is ~37.5 km (Kim et al., 2009), but varies  
226 between ~33 km and ~39 km along the rift shoulder (Ufipa Plateau) (Hodgson et al., 2017), and  
227 increases slightly to 41.1–42.1 km in the northwestern-most part of the rift where the rifting is  
228 minimal. These suggest that overall, the crustal thinning beneath the rift has been minimal but  
229 may be slightly more beneath the Songwe Trough (Njinju et al., 2018). Camelbeeck and Iranga  
230 (1996) estimated ~42 km crustal thickness beneath the Songwe Trough. Whereas, an average  
231 crustal thickness of 39 km has been estimated for the Rungwe Volcanic Province, ~37-39 km for  
232 the North Malawi Rift, and 38-42 km for the Proterozoic terrains surrounding the North Malawi  
233 Rift (Borrego et al., 2016; Njinju et al., 2018).

### 234 *2.3. Kinematics of the RNMRS*

235 The mode of opening of the Rukwa Rift – Northern Malawi Rift segment is controversial.  
236 Overall, two models have been proposed. One of the models advocates for orthogonal rifting due  
237 to the dominance of NW-trending pre-existing basement fabric in the region, which resulted in  
238 the rotation of the E-W directed regional extension into NE-SW, thus producing NW-striking  
239 normal faults (e.g., Delvaux et al., 1992; Morley, 2010; Delvaux et al., 2012). The other model  
240 argues for oblique extension primarily due to the obliqueness of the rift segment to the E-W  
241 directed extension, thus resulting in the development of NW-trending dextral strike-slip faults  
242 (e.g., Chorowicz, 1989; Daly et al., 1989; Wheeler and Karson, 1994; Kervyn et al., 2006;  
243 Mortimer et al., 2007). Both models are based on observations from only the Rukwa Rift and  
244 North Malawi Rifts, and did not consider the kinematics of brittle structures along the Mbozi  
245 Block accommodation zone.

246

## 247 **3.0 MATERIAL AND METHODS**

248 In this study, we carried out detailed mapping of exposed and buried fault segments within the  
249 RNMRS. We utilized Shuttle Radar Topography Mission (SRTM) Digital Elevation Model  
250 (DEM) to locate surface expressions of faults; and filtered aeromagnetic data to map the plan-  
251 view trace of basement faults and metamorphic fabrics.

252

### 253 *3.1. SRTM DEM Data*

254 We extracted topographical profiles along the length of the rift from the SRTM DEM data to  
255 investigate surface morphology of the rift segments which could provide insight into the  
256 evolving nature of the rift architecture along the RNMRS.

### 257 3.2. Aeromagnetic Data

258 We combined three separate aeromagnetic surveys consisting of data acquired over northeastern  
259 Zambia, southern Tanzania and Northern Malawi. The Tanzania survey was collected between  
260 1977-1980 with flight height of 200 m and a flight line spacing of 1 km. The Zambia survey was  
261 collected between 1973-1976 with a flight height of 150 m and a flight line spacing of 800-1000  
262 m. The Malawi survey was carried out in 2013 with a flight height of 80 m, a flight line spacing  
263 of 250 m. Before merging the three surveys, we first corrected for the skewness of the data by  
264 reducing each of them to the magnetic pole (RTP). The RTP correction normalizes the magnetic  
265 field to the magnetic field at the pole so the anomalies retain their correct strike and shape,  
266 allowing the magnetic data to be interpreted as geologic structures (Baranov, 1957; Silva, 1986).  
267 Afterwards, we applied upward-continuation (Henderson and Zeitz, 1949) of 120 m to the RTP-  
268 corrected Malawi data, 50 m to the Zambia data in order to mathematically normalize the three  
269 datasets to a 200 m observational height. We then merged the three surveys into a single  
270 aeromagnetic grid file. Further, we applied the vertical derivative (VDR) filter to the merged data  
271 in order to enhance magnetic gradients associated with possible basement faults and basement  
272 metamorphic fabrics (Salem et al., 2008).

273 The vertical derivative edge filter has been very effective in the mapping of plan-view  
274 trace of buried active faults from aeromagnetic data in different parts of the EARS (Kinabo et al.,  
275 2007, 2008; Kolawole et al., 2017, 2018). Excluding the Rungwe Volcanic Province (RVP;  
276 Figure 3) where volcanic materials overlie the crystalline basement, there is no information on  
277 the presence of basaltic rocks along the fault segments interpreted in this study. Therefore, we  
278 assume induced magnetization as the primary source of magnetization, except in the Rungwe  
279 Volcanic Province (RVP) where volcanic deposits are present. However, we do not have

280 information on remanent magnetization in the RVP at this time. Due to the higher spatial  
281 resolution of the Malawi aeromagnetic data (62 m grid cell size) compared to those covering  
282 Zambia (225 m grid cell size) and Tanzania (250 m grid cell size), the magnetic anomalies are  
283 most significantly better resolved in the Malawi part of the filtered aeromagnetic maps.

284

## 285 **4.0 RESULTS**

### 286 *4.1. Variation in rift morphology from topographic profiles*

287 We examined fifteen rift-orthogonal topographic profiles (spaced at 40 km) along the RNMRS  
288 (Fig. 1B) to understand the overall along-strike variation in rift morphology. The investigation of  
289 rift dynamics by careful analyses of the topographic structures has provided important  
290 information on the subsurface architecture of rift systems (e.g., Pik et al., 2008; Wichura et al.,  
291 2011; Lao-Davila et al., 2015). Our morphological assessments focus on the variation in the  
292 scarp-heights (relief above the surface) of exposed normal faults along the RMNRS. This  
293 provides the minimum estimate of the relative vertical displacements of the faults at the point of  
294 assessment, hence the term ‘exposed minimum vertical displacement’ (EMVD) (Lao-Davila et  
295 al., 2015).

296 Profile 1 (Fig. 3), obtained at the northern tip of the Rukwa Rift shows no pronounced  
297 fault scarps, and the slight topographic high between Lake Tanganyika and Ufipa Fault  
298 represents the northernmost tip of the Ufipa Plateau. In Profile 2, the surface morphology of the  
299 rift shows sharp topographic gradients bounding the Rukwa Rift Valley. These topographic  
300 gradients correspond to the scarps of the Ufipa Fault (600 m) and the Lupa Fault (750 m). In this  
301 northern part of the Rukwa Rift, the Ufipa and Lupa Faults have comparable scarp heights, thus  
302 illustrating a typical graben structure. In Profile 3, the rift structure changes into a half-graben

303 surface morphology with the Ufipa Fault having a significantly higher escarpment than the Lupa  
304 Fault (~900 m difference). Profiles 4 to 7 show the same half-graben morphology for the Rukwa  
305 Rift as in Profile 3; however, we observe that the Ufipa Fault scarp is much higher than the  
306 topography of the Ufipa Plateau to its west along Profile 4.

307         Along Profiles 6 and 7, we observe that the Ufipa Fault scarp is lower than in the  
308 northern profiles (Profiles 1-5) and that the Lupa Fault scarp is also higher than in the northern  
309 profiles. In Profile 7, the Rukwa Rift splits into two basins separated by the Mbozi Block such  
310 that the west basin (Musangano Trough) is bound by the Ufipa Fault to the southwest, and the  
311 east basin (Songwe Trough) is bound by the Lupa Fault to the northeast. Profiles 7 to 9 shows a  
312 continuous southeastward decrease in the scarp heights of the Ufipa and Lupa Faults; and  
313 although the Ufipa Fault scarp is still visible south of the Mbozi Block along Profile 9, the Lupa  
314 Fault scarp is significantly diminished. Profile 9 shows a gentle topographic transition from the  
315 Musangano Trough to the Mbozi Block, but to the northeast of the Mbozi Block, the topography  
316 spikes abruptly, representing the northern limits of the Rungwe Volcanic Province (RVP). In  
317 Profile 10, the Ufipa Fault bounds what appears to be the southernmost extent of the Musangano  
318 Trough, while the uplifted Mbozi – RVP domain dominates the terrain and drops off into the  
319 Usangu Trough to the northeast. Profile 11 transects the northernmost tip of the North Malawi  
320 Rift, where the RVP (bounded to the northeast by the Livingstone Fault scarp) represents the  
321 most dominant structure in the terrain and the entire topography of the Mbozi Block and areas to  
322 its southwest are relatively lower.

323         Profiles 12 and 13 illustrate half-graben morphologies for the North Malawi Rift (North  
324 Basin) with the Livingstone Fault (northeast bounding fault) dominating the topography. Profile  
325 14 transects the transfer zone between the North and Usisya Basins of the Malawi Rift showing

326 both the Livingstone border fault to the northeast and the Nyika Plateau to the southwest. Profile  
327 15 which transects the Usisya Basin describes a half-graben surface morphology but in which the  
328 rift bounding fault is located on the southwest.

329

#### 330 *4.2. Aeromagnetic lineaments and basement fabric*

331 The RTP merged (Fig. 4A) and edge-enhanced (Fig. 4B) aeromagnetic maps over the RNMRS  
332 provide a continuous plan-view image of the basement structures along the rift segment.  
333 Although the Malawi part of the merged data has the highest resolution, the moderate resolution  
334 of the Tanzania and Zambia parts of the data allows for considerable delineation of the trends of  
335 magnetic anomalies. Overall, the areas of basement exposures exhibit high amplitude, high  
336 frequency and short wavelength magnetic anomalies that delineate lineaments of interpretable  
337 trends (e.g., Kolawole et al., 2018). The high frequency lineaments can be easily observed on the  
338 rift shoulders of the Rukwa and North Malawi Rifts (Fig. 4B), and are commonly truncated at the  
339 rift margins by the rift-bounding faults (black arrows in Fig. 4B).

340         Within the rift valleys where sedimentary rocks overlie the deeply-buried basement  
341 rocks, the detailed magnetic fabric of the basement becomes suppressed such that gradients in  
342 the magnetic data could correspond to fault offset within the magnetic source (e.g., Grauch and  
343 Hudson, 2007, 2011; Kolawole et al., 2018) or remnants of the suppressed magnetic foliation of  
344 the source (Kolawole et al., 2018). In the study area, the magnetic anomalies within the rift  
345 basins are dominated by relatively lower amplitude, longer wavelength and lower frequency  
346 anomalies.

347

348

### 349 4.3. SRTM DEM Fault trends and Aeromagnetic lineaments

350 We compare the along-axis geometry of the rift-bounding faults (from SRTM DEM) with the  
351 metamorphic fabric (in plan view) of the host basement rocks along the rift shoulders (from  
352 filtered aeromagnetic data) (Figs. 5-11). Figures 5-9 focus on the southwestern boundary of the  
353 RNMRS, consisting of the Ufipa Fault of the Rukwa Rift, the southwestern boundary of the  
354 Mbozi Block and the southwestern boundary faults of the North Malawi Rift. Whereas, Figures  
355 10 and 11 focus on the northeastern boundary of the RNMRS, consisting of the Lupa Fault of  
356 Rukwa Rift, the northeastern boundary of the Mbozi Block (i.e. the Rungwe Volcanic Province)  
357 and the Livingstone Fault of the North Malawi Rift. We constrain our identification of the  
358 Precambrian terranes and strike of their fabrics with previous field studies of the Precambrian  
359 basement along the RNMRS (e.g., Daly, 1988; Wheeler and Karson, 1989; Lenoir et al., 1994;  
360 Theunissen et al., 1996; Boven et al., 1999; Fernandez-Alonso et al., 2001; Ring et al., 2002;  
361 Schenk et al., 2006; Delvaux et al., 2012, Lawley et al., 2013; Kolawole et al., 2018). Overall,  
362 within the study area, we observe that the basement fabrics exhibit two styles, (1) discrete fabrics  
363 which include isolated magnetic lineaments of strong amplitude; and (2) distributed fabrics  
364 which encompass fabric sets of multiple medium-to-low amplitude magnetic lineaments as  
365 distributed fabrics.

366

#### 367 4.3.1. Southwestern boundary of the RNMRS

368 In the northernmost part of SW Rukwa Rift (Chisi area) (Figs. 5A-C), we observe curvilinear  
369 fault geometries that follow Precambrian fabric (e.g., Ufipa, Chisi, Kalambo and Kanda Faults).  
370 In the Chisi area, the Rukwa Rift border fault consists of the Northern Ufipa Fault segment and  
371 the Chisi Fault segment. The tip of the Northern Ufipa Fault segment terminates against the

372 WNW-ESE Chisi Fault at a high angle (Fig. 5A). The Chisi Fault aligns with a strong WNW-  
373 ESE magnetic-high lineament (discrete fabric) (Fig. 5B-C) known as the Chisi Shear Zone,  
374 which extends eastward into the rift basin (Theunissen et al., 1996; Boven et al., 1999; Schenk et  
375 al., 2006; Fig. 2B). Also, the truncated Ufipa Fault segment aligns with the Ufipa Terrane  
376 basement fabrics (distributed fabrics). The North Ufipa Fault and Chisi Fault link at a very high  
377 angle to form a salient that point basinward (the Chisi Salient).

378 In the central part of the Ufipa Fault (Figs. 6A-C), the fault segments also exhibit  
379 curvilinear geometries and the hard linkage of major fault segments occur at high angles  
380 (e.g., Kwera relay ramp). The Kwera relay ramp is the largest relay zone along the Ufipa Fault.  
381 To the north of the Kwera relay ramp, the fault trends parallel to a magnetic lineament (discrete  
382 fabric) that is located to its east, whereas, to the west of this fault segment, the basement is  
383 characterized by two cross-cutting sets (NNW-SSE and NW-SE) of distributed fabrics. The  
384 NNW-SSE set represent the fabrics of the Ufipa Terrane, but the origin NW-SE is unknown at  
385 this time. The Ufipa Fault segments appear to follow the NNW-SSE basement fabrics but side-  
386 steps by means of short fault segments that align with the NW-SE fabric set. To the south of the  
387 relay ramp, the Ufipa Fault strikes parallel to the Mughese Shear Zone fabric which is colinear  
388 with the NNW-SSE set.

389 Towards the southern part of the Ufipa Fault, in Figures 7A-C, we observe that  
390 the Ufipa Fault segments exhibit rectilinear geometries such that it is difficult to delineate fault  
391 bends that could correspond to breached relay ramps. In addition, we observe a stronger  
392 alignment of the Ufipa Fault segments with the Mughese Shear Zone fabric within the area. To  
393 the east of the Ufipa Fault, a fault that strikes parallel to the Ufipa Fault separates  
394 the Musangano Trough from the Mbozi Block. The Mbozi basement is characterized by



395 metamorphic fabrics that strike WNW-ESE to NW-SE, oblique to the fault but in which subtle  
396 bends in the fault trend align with the basement fabric.

397         In the southern part of the Ufipa Fault (Figs. 8A-C), the fault is characterized by two  
398 segments. One of the segments represents a rectilinear southward continuation of the Ufipa Fault  
399 and is bounded to the east by a linear ridge that separates it from the Musangano Trough (see  
400 “Mughese Fault” segment at Tunduma in Figs. 8A-C). The other segment splays away from the  
401 former and delineates a curvilinear geometry that looks like those in the northern and central  
402 segments of the Ufipa Fault; this curvilinear segment bounds the southernmost part of  
403 the Musangano Trough (see “Ufipa Fault” in Figs. 8A-C). Around Kaseye town (see Fig. 8A),  
404 the basement is partially buried and more deeply buried around Chitipa to the south. The  
405 continuation of the Ufipa Fault is only evident in the aeromagnetic data as a strong magnetic-low  
406 lineament that is bounded by bands of magnetic-high lineaments (see magnetic fabrics  
407 around Kaseye town in Fig. 8B). Within the Kaseye area, the NW-striking Mughese Shear Zone  
408 fabrics are truncated by a discrete N-S trending magnetic-high lineament (see area within purple  
409 rectangle in Fig 8C). The magnetic-low lineament that projects as a southward continuation of  
410 the Ufipa Fault persists along the Mughese Shear Zone fabric into the Karonga area of the North  
411 Basin of Malawi Rift where the Mughese Shear Zone is abruptly truncated by the Karonga Fault  
412 (KF in Fig. 8C). The Mbozi Terrane fabric strike NW-SE, at low angles to the trend of the  
413 Mughese Shear Zone.

414         Detailed analyses of the onshore faults along the hinge zone of the Northern Malawi Rift  
415 (Karonga area) are well documented in Kolawole et al. (2018). Major onshore hinge zone faults  
416 include the Karonga Fault (KF), St. Mary Fault (SMF), Kaporo Fault (KPF), Lupaso Fault (LF),  
417 Katesula Fault (KTF) and the Mbiri Fault (Figs. 9A-C). Although the Karonga Fault cuts the

418 basement fabric, several basement-rooted buried fault segments along the rift margin align with  
419 the fabric of the Mughese Shear Zone (Kolawole et al., 2018). It is interesting to note that  
420 the Mbiri Fault, the longest fault (and potentially has the largest throw?) along the rift margin is  
421 sub-parallel to the strike of the continuation of the Ufipa Fault into the area (“Mughese Fault” in  
422 Fig. 9C). Although segments of the Mbiri Fault appear to align with the Mughese Shear  
423 Zone fabric, it is not clear if the fault reactivated the southwestern boundary of the shear zone or  
424 if the fault partially aligns with the fabric of the shear zone. However, we observe that for the  
425 most part, the side-steps along the Mbiri Fault align with the basement fabric.

426

#### 427 *4.3.2. Northeastern boundary of the RNMRS*

428 In the northern part of the Lupa Fault, the fault trend defines long, rectilinear segments (Fig.  
429 10A) and aligns with the Katuma Terrane, similar to those observed in the southern part of  
430 the Ufipa Fault where faults align with the trend of the Ufipa Terrane (Fig. 7). Although, the E-  
431 W trending fabric of the Usagaran Orogenic Belt dominates the aeromagnetic data along the  
432 northeastern Rukwa Rift shoulder (Fig. 10B), we also observe that closer to the Lupa Fault scarp,  
433 there are some lineaments that align with the fault. To the southwest of the Lupa Fault (within  
434 the rift valley), a strong magnetic-high lineament which correspond to the Chisi Shear Zone (see  
435 onshore continuation and outcrop of the shear zone in Fig. 5), strike sub-parallel to the trend of  
436 the Lupa Fault, but deviates more significantly southwards (Figs. 4B and 10).

437         The southern part of the Lupa Fault (between the Katuma Terrane and the Rungwe  
438 Volcanic Province) is characterized by curvilinear fault segments (Fig. 10 and 11). The Lupa  
439 Terrane fabrics are oriented WNW-ESE, oblique to the Lupa Fault trend (Figs. 11A-C).  
440 However, subtle steps along the curvilinear fault segments align with the Lupa Terrane fabrics.

441 In the Rungwe Volcanic Province (RVP), the magnetic lineaments strike NW-SE (Fig.  
442 11B), similar to those of the Mbozi Terrane (8B and C), and are truncated to the west by a strong  
443 NNW-striking magnetic high lineament (white arrows in Fig. 11B). Furthermore, we observe  
444 that the volcanic centers align with NNW-striking magnetic gradient superimposed on the NW-  
445 striking fabrics (black arrows in Fig. 11B). South of the Rungwe Volcanic Province, segments of  
446 the Livingstone Fault system describe rectilinear geometries and align with strong magnetic-high  
447 lineaments (shear zone?) in the Upangwa Terrane.

448

## 449 **5.0 DISCUSSION**

### 450 *5.1. Rift architecture*

451 The present morphology of the RNMRS reflects the result of multiple episodes of rifting that  
452 have affected this part of eastern Africa i.e. a Permo-Triassic rifting episode (Karoo), a  
453 Cretaceous episode and the ongoing Cenozoic rifting episode (e.g., Castaing, 1991; Morley et al.,  
454 1992). The Rukwa Rift has been described as both a graben (e.g., Zhao et al., 1997) and half-  
455 graben (e.g., Kilembe and Rosendahl, 1992) bounded by the oppositely-dipping Ufipa and Lupa  
456 normal faults. The Lupa Fault is commonly regarded as the major border fault of the Rukwa Rift  
457 due to its larger throw of ~7 km (Peirce and Lipkov, 1988) relative to the Ufipa Fault (Fig. 12);  
458 however, it has also been shown that the Cretaceous and Cenozoic sediments in the rift thicken  
459 towards both faults (Fig. 12; Morley et al., 1992; Zhao et al., 1997). In this study, our  
460 topographical assessments (Fig. 3, profiles 1-7) show that the scarp height of the Ufipa Fault  
461 consistently exceeds that of the Lupa Fault through-out the rift segment, thus suggesting  
462 significant footwall uplift along the Ufipa Fault.

463           The significantly lower scarp-height of the Lupa Fault suggests either that it has not been  
464 very active in Cenozoic times or that it has been heavily eroded since the cessation of Mesozoic  
465 rifting. However, based on these observations, we interpret that in the earlier rifting episodes  
466 (especially in the Permo-Triassic Karoo episode), the Lupa Fault played the role of the major  
467 border fault such that the basin defines a typical half-graben geometry; but in the present  
468 Cenozoic phase of rifting, the Ufipa Fault appear to have been preferentially accommodating  
469 more strain. Therefore, we suggest that the Rukwa Rift is possibly transitioning into an  
470 asymmetric-graben geometry. An asymmetric-graben is a graben with EMVD greater on one  
471 border fault compared to the other one, such that the basin polarity has shifted to the fault with  
472 the greater 'exposed minimum vertical displacement' (EMVD) (Lao-Davila et al., 2015). The  
473 implications of the present Rukwa Rift morphology on the Lupa Fault may reflect temporal and  
474 spatial migration of strain accommodation from a previously dominant border fault into another  
475 one that has been previously less-dominant. This could possibly be explained by the Scholz and  
476 Contreras (1998) suggestion that when a rift-bounding fault attains some limiting offset, motion  
477 on the fault will cease, and strain will be transferred to a new fault.

478

## 479 *5.2. The Southwestern Boundary of the RNMRS and relationships with Precambrian Basement*

### 480 *Fabric*

481 In the Chisi area, the Northern Ufipa Fault segment terminates against the Chisi Shear Zone  
482 along which the Chisi Fault segment developed. The termination of the Ufipa Fault at its  
483 intersection with this shear zone exemplifies one of the roles of pre-existing basement structures  
484 as temporal and/or spatial mechanical 'barriers' that arrest and delimit the continuous lateral  
485 propagation of a fault. Several studies on fracture propagation have demonstrated that fractures

486 are principally bifurcated, blunted, and/or arrested when they intersect discontinuities, stress  
487 barriers and/or rock layers of significantly-contrasting mechanical properties along their path of  
488 propagation (e.g., Helgeson and Aydin, 1991; Gudmundsson and Brenner, 2001; Zhang et al.,  
489 2007; Zhang and Jeffrey, 2008). Other examples of normal fault termination at long-lived  
490 basement shear zones include the case of the Albertine-Rhino Graben terminating at the Aswa  
491 Shear Zone (Katumwehe et al., 2015) and the Okavango Rift border faults against the Sekaka  
492 Shear Zone (Modisi et al., 2000). We suggest that both the reactivation of the Chisi Shear Zone  
493 into the Chisi Fault and termination of fault segments at the shear zone demonstrate the strong  
494 influence of the Chisi Shear Zone on the development of this part of the Rukwa Rift.

495         Although the northern segments of the Ufipa Fault align with the basement fabrics (Fig.  
496 5B-C) and the southern segments show even stronger alignment with the NNW-SSE fabrics of  
497 the Ufipa Terrane and/or Mughese Shear Zone (Fig. 7), we find that the central segments of the  
498 fault show only partial alignment with this basement fabric (Fig. 6A-C). These observations  
499 imply that, although the Ufipa Fault is thought to have largely propagated along the Mughese  
500 Shear Zone (Fig. 2; Delvaux et al., 2012), there is stronger control of the Mughese Shear Zone  
501 and Ufipa Terrane fabrics on the fault development along its northern and southern segments  
502 than in the central part. We suggest that the partial control of these NNW-SSE fabrics on the  
503 central Ufipa Fault segments is due to the occurrence of a NW-SE basement fabric set on the  
504 Mughese Shear Zone and Ufipa Terrane fabric (Fig. 6). This may also explain the localization of  
505 the largest relay zone along the Ufipa Fault at its central segment.

506         The Tunduma-Kaseye area of Malawi, through the Misuku Mountains (Fig. 8A)  
507 constitute the southern boundary of the Mbozi Block. Although, the SRTM DEM shows that the  
508 sub-aerial expression of the Ufipa Fault dies out roughly mid-way between Tunduma and

509 Kaseye, the filtered aeromagnetic data reveals that in the subsurface, the fault continues across  
510 the Kaseye-Chitipa area as a distinct magnetic-low lineament that bounds the Mughese Shear  
511 Zone to the south and runs southeastwards into the Karonga area (Fig. 8B). Upon closer  
512 observation of the continuation of this strong magnetic-low lineament in the Kaseye-Chitipa  
513 area, we find that it cuts-across and offsets a N-S striking magnetic-high lineament which  
514 extends 80 km southwards into the Permo-Triassic Luangwa Rift in Zambia. We interpret this N-  
515 S lineament as a mafic dike that is possibly related to one of the earlier (Triassic or Cretaceous)  
516 episodes of rifting which are known to have been associated with extensive late-stage diking  
517 events in the Luangwa Rift (e.g., Van de Velde and De Waele, 1998) and Shire Graben in  
518 southern Malawi (Castaing, 1991). Sedimentary deposits in the Kaseye-Chitipa area resulted in  
519 the burial and lack of sub-aerial exposure of this structure, thus making this study the first  
520 revelation of its existence in Northwestern Malawi.

521 Our analyses of the geometry of the interpreted dike structure (Fig. 13A), here in referred  
522 to as the Chitipa Dike, shows a distinct difference in the geometry of the structure to the north  
523 where it is cut by the strong magnetic-low lineament (continuation of the Ufipa Fault) and to the  
524 south (farther away from the fault intersection). South of Chitipa (Fig. 13B), we observe that the  
525 north-trending dike describes a consistent left-stepping geometry which diminishes across the  
526 Chitipa town location and continues with a more rectilinear geometry northwards into the  
527 Mughese Shear Zone area. This side-stepping geometry is typical of vertical sheet intrusions and  
528 are related to either magma intrusion into pre-existing stepped joint systems (e.g., Baer, 1991) or  
529 near-surface stress rotations during magma intrusion (e.g., Fossen, 2010). In this study, the  
530 coincidence and strike of the dike-steps along the NW- to NNW- striking basement fabric may in

531 fact suggest the possible influence of pre-existing basement fabric on the stepping geometries of  
532 dike intrusions during their emplacement in host metamorphic rocks (Fig. 13B).

533 Further north, where the dike is cut by the continuation of the Ufipa Fault, the filtered  
534 aeromagnetic data shows the dike exhibiting consistent right-lateral offsets across NW-SE  
535 magnetic gradients (Fig. 13C). This clear distinction in the structural style of the dike south of  
536 Chitipa and in the north across the continuation of the Ufipa Fault, suggest that the contrasting  
537 structural styles are associated with different geological processes. In a summary, we interpret  
538 that along the southern boundary of the Mbozi Block, the continuation of the Ufipa Fault, which  
539 itself aligns with the Mughese Shear Zone fabric (Fig. 8B-C), is a right-lateral strike-slip fault  
540 that displaced a N-S trending Mesozoic (?) dike intrusion (Fig. 13C). We further interpret that  
541 this strike-slip fault reactivated the Precambrian Mughese Shear Zone at some time post-  
542 Cretaceous (i.e. related to the present Cenozoic rifting phase), and therefore refer to it as the  
543 “Mughese Fault”. This interpretation is further supported by the change in the morphological  
544 expression of the Ufipa Fault from a typical single-scarp style into a narrow linear valley-ridge  
545 style in the Tunduma area in the SRTM DEM (see “linear ridge” in Fig. 8A). The observed  
546 linear valley-ridge geomorphology is typical of active strike-slip fault zones (e.g., McCalpin et  
547 al., 2009). In addition, previous field studies in the area (Delvaux et al., 2012) observed strike-  
548 slip displacement on rock outcrops at Tunduma and Mbozi Quarry (“Q” in Fig. 8A). Using the  
549 Chitipa Dike as a strain marker, we observe that the offsets increase northwards across multiple  
550 splays of the Mughese Fault and estimate that the displacement is maximum at the major  
551 Mughese Fault trace (northernmost extent of the strike-slip fault zone). Since mafic intrusions  
552 often produce magnetic anomalies larger than the actual size of the sources, it is practically  
553 impossible to estimate the true cumulative strike-slip displacement from the aeromagnetic data.

554 However, based on the lateral dike separation across the fault on our aeromagnetic data, we  
555 estimate a minimum of 500 m strike-slip displacement along the Mughese Fault (Fig. 13C). In  
556 Figure 14, we present a conceptual model that summarizes our interpretation of the Chitipa Dike  
557 geometry and the interaction of the Mughese Fault with the dike.

558 In Tunduma area (Fig. 8A), the presence of discrete breaching of the linear ridge by  
559 stream channels suggest that the strike-slip displacement along the Mughese Fault is most-likely  
560 a short-lived event that occurred at some time in the past during the development of the RNMRS.  
561 The curvilinear fault scarp adjacent to the Mughese Fault at Tunduma shows single-scarp  
562 morphology (not linear-ridge morphology) typical of normal faults as seen on the other segments  
563 of the Ufipa Fault. Therefore, we interpret this curvilinear fault scarp as a possible old segment  
564 of the Ufipa Fault that was ‘pirated’ by the Mughese strike-slip fault; thus, suggesting a phase in  
565 which the Ufipa Fault accommodated strike-slip displacement.

566 Along the southwest margin of the North Malawi Rift (Karonga area), the Mughese Fault  
567 diffuses into a zone of wide-spread faulting where the southeast-ward bend of the Mughese  
568 Shear Zone controls the development of the normal faults and recent seismicity along the basin  
569 hinge margin (Fig. 9A-C) (Kolawole et al., 2018; Dawson et al., 2018). Further south of Karonga  
570 town, the NNW-striking Mbiri Fault is the dominant fault structure in terms of its length (and  
571 displacement?) along this part of the hinge zone of the North Malawi Rift. We do not observe  
572 any direct spatial connectivity between the Mughese Fault and the Mbiri Fault. However, based  
573 on the sub-parallel geometry of both faults and the structural dominance of the Mbiri Fault in the  
574 area, we suggest that the Mbiri Fault could possibly represent a continuation of the Mughese  
575 Fault into the hinge zone of the Malawi Rift North Basin. It is also important to note that the  
576 Mbiri Fault is synthetic to the Livingstone border Fault. Following the observations and



577 interpretations above, we suggest that there exists a well-developed continuous connectivity of  
578 rift-related structures along the southwestern boundary of the RNMRS, facilitated by the extent  
579 of the Mughese Shear Zone and the Ufipa Terrane. Interestingly, the linking of the oppositely-  
580 dipping Ufipa and Mbiri Faults by the Mughese strike-slip fault describes a structure that,  
581 overall, is similar to that of the Morley et al. (1990) convergent-approaching normal fault  
582 system.

583

### 584 *5.3. The Northeastern Boundary of the RNMRS and relationships with Precambrian Basement* 585 *Fabric*

586 The northern segment of the Lupa Fault exhibits clear alignment with the trend of the Katuma  
587 Terrane (Fig. 10) and with a few interpretable magnetic lineaments (likely due to low resolution  
588 of the aeromagnetic data). We also note that the Lupa Fault is sub-parallel to the Chisi Shear  
589 Zone in this area. Farther south of the Lupa Fault (Kapalala-Kanga area; Figs. 10 A-C), the fault  
590 segments occur at a high angle to the Usagaran Belt and Lupa Terrane fabrics, thus indicating an  
591 apparent lack of control of pre-existing basement fabric on the propagation of the southern Lupa  
592 Fault, except in the coincidence of the fault steps with the trend of the basement fabrics.

593         The southern segment of the Lupa Fault transitions into the Rungwe Volcanic Province  
594 (RVP) where surficial cover of volcanic deposits obscures the southward continuation of the  
595 Lupa Fault (Fig. 11A). In addition, the presence of mafic volcanic deposits in the RVP (e.g.,  
596 Fontijn et al., 2012) makes it difficult to make a reliable interpretation of the magnetic fabric of  
597 the underlying Precambrian basement (Fig. 11B). However, we find that a distinct magnetic-high  
598 lineament aligns with the Mbaka Fault surface trace (white arrows in Figs. 11A and 11B). Also,  
599 the distribution of volcanic centers in the RVP show alignment with both the Mbaka Fault trace

600 and a subtle curvilinear gradient (black arrows in Fig. 11B). Further south, the curvilinear  
601 gradient connects with the Livingstone Fault, the northeastern border fault of the North Malawi  
602 Rift. This curvilinear gradient coincides with the location and extents of the so-called “Ngozi-  
603 Rungwe Line” of Fontijn et al. (2010), described as a buried fault system that served as a conduit  
604 for magmatic fluids to migrate to the surface volcanic vents. Therefore, although it is possible  
605 that the magnetic anomalies in this area are affected by remanent magnetization from the  
606 volcanic deposits, we interpret that this aeromagnetic gradient provides a possible subsurface  
607 evidence of the fault system (Ngozi-Rungwe Line) that connects the Lupa Fault and the  
608 Livingstone Fault across the RVP. The filtered aeromagnetic data and previous field studies  
609 (e.g., Wheeler and Karson, 1989) shows that the Livingstone Fault segments align with and  
610 reactivated the fabric of the Upangwa Terrane (Figs. 11B-C). The observations and  
611 interpretations above suggest that there exists a well-developed continuous connectivity of rift-  
612 related structures along the northeastern boundary of the RNMRS. However, the relationship  
613 between the basement fabric and the buried faults beneath the volcanic deposits remains unclear.

614

#### 615 *5.4. Implications for Rift Development*

##### 616 *5.4.1. Rift Coupling*

617 In the Rukwa Rift, the substantial dominance of the Ufipa Fault rift shoulder over that of the  
618 Lupa Fault may imply that the Ufipa Fault is the present-day active border fault of the Rukwa  
619 Rift. This proposition may be supported by the hypocentral location of the 1994 Mw5.9 Rukwa  
620 earthquake and its aftershocks with epicentral location in the northern part of the rift (Fig. 12;  
621 Zhao et al., 1997). The nodal planes of the earthquake focal mechanism solution are broadly  
622 consistent with the orientation of both the Lupa and Ufipa faults, and relative position of

623 aftershocks to the main shock is well determined (Fig. 12; Zhao et al., 1997). Considering the  
624 uncertainty range of the earthquakes, the spatial distribution of the aftershocks relative to the  
625 main shock delineates a sub-horizontal fault zone that most fits the subsurface projection of the  
626 Ufipa Fault (Fig. 12). In addition, field investigations of the Kwera relay ramp (see Fig. 6 for  
627 location) revealed features that indicate recent activity along the Ufipa Fault (Delvaux et al.,  
628 2012). Camelbeeck and Iranga (1996) observed several lower crustal seismicity in the Rukwa  
629 Rift with most of the events clustering beneath the Songwe Trough and the Rungwe Volcanic  
630 Province (southern parts of the rift; Fig. 1B). The locations of the clusters suggest activity along  
631 the southern Lupa Fault and Mbeya Range Fault. However, since the scarp height (and throw?)  
632 of the Ufipa Fault decreases southwards (Fig. 3), and the throw on the Lupa Fault increases  
633 southwards (and rapidly along the Songwe Trough) (Morley et al., 1992), we infer that the  
634 present border fault role of the Ufipa Fault excludes the southermost parts of the Rukwa Rift.

635         It has also been observed that the early stage of continental rifting is typically  
636 characterized by the development of along-axis alternating polarity of rift segments, rift border  
637 faults, uplifted rift flanks (e.g., Bosworth, 1985; Rosendahl, 1987; Hayward and Ebinger, 1996;  
638 Lao-Davila et al., 2015). The zones of polarity changes (transfer/accommodation zones) serve to  
639 transfer extensional strain between the rift segments and link the border faults which often have  
640 variable structural styles and geometries (e.g., Morley et al., 1990; Wilson, 1999). Within young  
641 continental rift settings, interactions between these large border faults lead to the systematic  
642 coupling of border faults and rift segments across the transfer zones, and subsequent growth of  
643 the rift system (e.g., Corti, 2012). Along the RNMRS, the alternating location of the of rift  
644 shoulder uplift (SW in Rukwa Rift and NE in North Malawi Rift), typical of coupled rift  
645 segments, suggest that the RNMRS can be considered a coupled rift segment. Further, our study

646 here shows that there is in fact, continuous structural continuation along the northeastern and  
647 southwestern margins of the RNMRS, typical of a coupled rift segment. Although, studies in the  
648 EARS and illustrations of its rift segments had always assumed this to be true, we hereby  
649 provide evidence supporting it, for the first time. In the West Antarctic rift system, the  
650 localization of recent volcanism along transverse structures within an accommodation zone (the  
651 Discovery accommodation zone) suggests active structural interactions between the flanking rift  
652 segments (Wilson, 1999). Therefore, we further suggest that the focusing of Neogene volcanism  
653 (e.g., Fontijn et al., 2010, 2012) along the northeastern boundary faults of the Mbozi  
654 accommodation zone (Fig. 15A) may be indicative of the ongoing coupling of the Rukwa and  
655 North Malawi Rift's northeastern border faults.

656

#### 657 *5.4.2. Rift Kinematics*

658 Several studies have suggested that the development of the RNMRS has been dominated by  
659 dextral strike-slip kinematics (e.g., Chorowicz, 1989; Daly et al., 1989; Wheeler and Karson,  
660 1994; Kervyn et al., 2006; Mortimer et al., 2007). However, analyses of fault architecture, fault-  
661 kinematics, paleostress and present-day earthquake focal mechanism solution in the Rukwa Rift  
662 show that the present-architecture of the rift largely developed within a pure extensional setting  
663 with extension direction orthogonal to the trend of the RNMRS (Morley, 2010; Delvaux et al.,  
664 2012). Furthermore, Delvaux et al. (2012) observed dextral strike-slip faulting along the fault  
665 systems bounding the Rukwa Rift, but concluded that the strike-slip event was transitory and was  
666 associated with an early Mesozoic transpressional event that resulted in the inversion of Karoo  
667 sediments. In this study, we observe the existence of a well-defined strike-slip fault bounding the  
668 SW margin of the Mbozi Block that reactivated the Precambrian Mughese Shear Zone. We also

669 observe lack of present-day activity along the strike-slip fault, and that this fault displaced a  
670 buried mafic dike with at least 500 m of dextral offset.

671         In the absence of chronological data on the mapped dike, we posit that the dike is most-  
672 likely associated with the widespread late-Karoo dike swarms observed in the Luangwa Rift and  
673 Shire Graben (southern Malawi) (Castaing, 1991; Van de Velde and De Waele, 1998). We refer  
674 to this strike-slip fault as the “Mughese Fault”, and the buried dike as the “Chitipa Dike”. The  
675 Chitipa Dike, presented for the first time in this work, may constitute the most excellent record  
676 of strike-slip kinematics along the RNMRS. Although, our results agree with Delvaux et al.  
677 (2012) in that the strike-slip faulting along the RNMRS was short-lived, we suggest that future  
678 geochronological analyses of this intrusion may provide the most-reliable constraint on the  
679 timing of strike-slip faulting event. It is also interesting to note that if the Mughese strike-slip  
680 fault is post-Karoo, its development represents a late reactivation of the Mughese Shear Zone in  
681 the evolution of the RNMRS. However, late reactivation of rift-oblique basement shear zones is  
682 not uncommon in rift basins (e.g., Muirhead and Kattenhorn, 2018).

683         Following the considerations above, we present cartoons of the RNMRS, illustrating the  
684 continuous structural connectivity along the northeast and southwest boundaries guided by the  
685 basement fabrics (Fig. 15B), and possible subsurface geometries and interactions of the domain-  
686 bounding structures (Fig. 15B-E). The inferred dominance of the Ufipa Fault in the northern and  
687 central parts of the Rukwa Rift (Fig. 12) suggest possible truncation of the Lupa Fault at depth,  
688 such that the load of the basin hanging wall block is being carried by the Ufipa Fault (Fig. 15B).  
689 However, seismic data is needed to confirm this interpretation. We illustrate a possible spatial  
690 relationship between the RVP magma pathways and the Mbozi Block bounding faults in Figure  
691 15C. In Figure 15D-E, we show a generalized basin geometry and flip in border fault polarity

692 from the Malawi Rift North Basin to the Usisya Basin. Overall, we posit that, along the  
693 northeastern boundary of the Mbozi Block transfer zone strain is accommodated by magmatism  
694 utilizing pre-existing fault systems, whereas, along the southwestern boundary, strain is  
695 accommodated by dextral strike-slip faulting.

696

#### 697 *5.5. Control of basement fabrics on normal fault geometries*

698 Overall, along the border faults of the Rukwa Rift, we find that strongly-curvilinear normal fault  
699 geometries (in plan-view) occur in three distinct settings. One, in areas where the basement  
700 fabrics describe high curvatures ( $>15^\circ$ ) (e.g., in the Chisi area, northern segment of the Lupa  
701 Fault; Fig. 5). Second, in areas of superposition of discordant sets of basement fabrics, in which  
702 the overall fault strike is parallel to one of the sets and relay ramp breach-faults follow the other  
703 fabric set that is oblique to the overall fault strike (e.g., central segment of Ufipa Fault; Fig. 6).  
704 Third, in areas where the faults propagate at high-angle to the strike of the basement fabrics (e.g.,  
705 in the Kapalala-Kanga area, southern segment of the Lupa Fault; Fig. 10). In general, along the  
706 RNMRS, we find that in areas where the basement fabrics show low curvatures ( $<10^\circ$ ), the fault  
707 segments tend to describe long, rectilinear geometries with narrow or almost unidentifiable  
708 breached-relay zones (e.g., Southern Ufipa Fault, Fig. 7; Mughese Fault, Fig. 8; Livingstone  
709 Fault, Fig. 11). We suggest that the control of rectilinear ( $<10^\circ$  curvature) basement fabrics on  
710 the propagation of normal faults may result in the development of rectilinear fault segments with  
711 greater likelihood of occurrence of narrow relay ramps and tip-to-tip fault linkage.

712 Curvilinear normal faults have been observed at various scales and at different  
713 extensional tectonic settings (Fossen and Rotevatn, 2016). However, the first order curvilinear  
714 normal faults are typically characterized by segment boundaries with salients (cusps) that plunge

715 basin-ward e.g., Salt-lake salient and Transverse Mountain salients of the Wasatch Fault in Utah  
716 (Fig. 16A), and the Gullfaks salient in the North Sea (Fig. 16B) (Fossen and Rotevatn, 2016).  
717 Also, the reported curvilinear faults show characteristic basin-ward concave geometries.  
718 However, the South Oquirrh Mountains normal fault zone, although curvilinear (Wu and Bruhn,  
719 1994), exhibits striking basin-ward convex geometry in which the cusps point into the footwall  
720 of the fault (Fig. 16A). Wu and Bruhn (1994) suggested that convex curvilinear geometry of the  
721 South Oquirrh Mountains normal fault zone developed by sequential propagation of the fault into  
722 its foot-wall, guided by linkage across smaller en-echelon faults created by the lateral shear  
723 components at the tips of the propagating fault. Here, in the Rukwa Rift, we observe both  
724 convex- and concave-curvilinear normal fault geometries along the Ufipa and Lupa Faults. The  
725 central Ufipa Fault (Fig. 6) and southern Lupa Fault (Kapalala-Mwambani area in Fig. 10A-C)  
726 exhibit convex-curvilinear normal fault geometries, whereas the northern Ufipa Fault and the  
727 Chisi Fault demonstrate concave-curvilinear fault geometries (linked at the Chisi salient) (Fig.  
728 5). Also, we find striking similarities between the concave- and convex-curvilinear fault  
729 geometries of the Rukwa Rift (Fig. 16C) with those in the southern Malawi Rift (Fig. 16D), the  
730 Jurassic sedimentary sequence of the northern North Sea rift (Gullfaks area) (Fig. 16B), and the  
731 Provo-Salt Lake City area (Fig. 16A). As show in Figure 16D, the segments of the Bilila-  
732 Mtakataka Fault in southern Malawi Rift present excellent examples of convex-curvilinear  
733 normal fault geometries (Jackson and Blekinsop, 1997). Recent studies on the relationships  
734 between the basement fabrics and the Bilila-Mtakataka Fault segments (Johnson et al., 2018;  
735 Hodge et al., 2018) show that some of the segments appear to align with the distributed basement  
736 fabrics, while others cut across the basement fabrics.

737           Based on the observations above, we present conceptual models for the control of  
738 various configurations of pre-existing basement fabrics on the development of curvilinear normal  
739 fault plan-view geometries (Fig. 17). We show how discrete and distributed basement fabrics and  
740 combinations of the two categories of fabrics can influence the plan-view geometry of normal  
741 faults. However, there is need to better understand (1) the influence of the basement fabrics on  
742 the geometries of curvilinear normal faults in 3-dimensions, (2) the influence of the extension  
743 direction on the development of the observed curvilinear normal faults in areas where the  
744 basement fabric present mechanical anisotropy.

745           Although, Fossen and Rotevatn (2016) provide evidence of subsidiary short-cut faulting  
746 across a salient, suggesting an impending evolution of concave-curvilinear faults into rectilinear  
747 faults, it is not yet clear if the model applies to convex-curvilinear normal faults since the two  
748 styles of faults are geometrically different. However, the synthetic Bilila-Mtakataka and the  
749 Chirobwe-Ncheu Faults (Fig. 16D) in southern Malawi Rift may provide some insight. Since the  
750 Chirobwe-Ncheu Fault is older than the Bilila-Mtakataka Fault (Jackson and Blekinsop, 1997),  
751 the rectilinear geometry of the Chirobwe-Ncheu Fault suggests that the apices of convex  
752 curvilinear segments may likewise eventually get breached to form more-rectilinear fault  
753 segments. Conversely, we observe the opposite of this model along the Lupa Fault, where the  
754 more-rectilinear northern Lupa Fault has accommodated much less strain compared to its  
755 curvilinear southern segment which has the most strain within the Rukwa Rift (Morley et al.,  
756 1992). Therefore, we suggest that although the model of temporal progression from a curvilinear  
757 to rectilinear fault geometry may apply to some large normal faults, it may not apply to others.

758

759



## 760 5.6. Rift Bifurcation

761           The basin scale splaying of the Rukwa Rift around the Mbozi Block into the Musangano  
762 and Songwe Troughs (Fig. 1B) obviously represents a smaller-scale of rift bifurcation when  
763 compared to the continental scale bifurcation of rift systems around microplates. Examples of  
764 such continental-scale rift bifurcation include the branching of the East African Rift around the  
765 Tanzania microplate into the western and eastern branches (Fig. 1A; e.g., Rosendahl, B.R., 1987;  
766 Versfelt and Rosendahl, 1989), the Red Sea Rift around the Sinai microplate into the Gulf of  
767 Suez Rift and Dead Sea Transform, and the South Atlantic Rift around the Sergipe microplate  
768 into the Tucano-Recôncavo Rift and Sergipe-Alagoas Transform (e.g., Szatmari and Milani,  
769 1999) (Figs. 18A-C). Based on scale distinction, we therefore refer to the continental rift-system  
770 scale bifurcation as first (1<sup>st</sup>) order rift bifurcation (Figs. 18A-C) and the rift-basin scale  
771 bifurcation as second (2<sup>nd</sup>) order rift bifurcation (Fig. 18D-F). Similar to the Rukwa Rift, 2<sup>nd</sup>  
772 order rift bifurcations are common along the East African Rift System. Examples include the  
773 Southern Malawi Rift bifurcation around the Shire Horst, the Shire Graben bifurcation around  
774 the Namalambo Horst, and the Albertine Graben bifurcation around the Rwenzori Block (Fig.  
775 18B; e.g., Castaing, 1991; Koehn et al., 2008; Lao-Davila et al., 2015; Xue et al., 2017).

776           Regardless of scale, numerical models have showed that inherited structural  
777 heterogeneity and lateral strength variations are key controls on rift bifurcation (e.g., Brune and  
778 Autin, 2013; Brune et al., 2017). However, it appears that 1<sup>st</sup> order bifurcations commonly occur  
779 at the tip of pre-existing microcratonic blocks along the path of propagation of a continental rift  
780 system (e.g., Fig. 18A). Also, 2<sup>nd</sup> order bifurcations appear to occur at the transfer zones between  
781 approaching rift segments, possibly due to a high tendency for the development of interfingering  
782 fault blocks in the transfer zones between colinear approaching rift segments (Morley, 1995) and

783 lateral rotation of the trapped blocks in the transfer zones between overlapping approaching rift  
784 segments (Koehn et al., 2008).

785         It is possible that the bifurcation of the Rukwa Rift around the Mbozi Block is related to  
786 the location of the block within the transfer zone of the colinear approaching Rukwa and North  
787 Malawi Rift segments. However, we suggest that the basement fabrics of the Ubendian Belt  
788 around the Mbozi Block transfer zone could be playing a complementary role in facilitating and  
789 guiding the intra-rift bifurcation of the Rukwa Rift around the block. Our filtered aeromagnetic  
790 map (Fig. 4B) shows that the block is dominated by a WNW-ESE and N-S fabrics. According to  
791 Daly (1988), the Meta-basites and intermediate granulites and quartzites of the Mbozi Block are  
792 characterized by lineations that trend NE-SW. These observations, however, suggest that the  
793 Mbozi Block fabrics strike at oblique to high-angles to the trend of the colinear fabrics of the  
794 basement terranes bounding the Mbozi Block (Katuma-Upangwa Terranes to the northeast, and  
795 those of the Ufipa Terrane to the southwest) and the main rift-bounding faults (the Lupa, Ufipa,  
796 Livingstone and Mughese Faults) (Fig. 15A). Therefore, we suggest that the colinear fabrics of  
797 the basement terranes surrounding the Mbozi Block which are already controlling the  
798 propagation and linking of the rift-bounding faults may be playing a significant role in guiding  
799 the bifurcation of the Rukwa Rift around the Mbozi Block (Figs. 4B, 7, 8 and 11).

800

## 801 **6.0 CONCLUSIONS**

802         Our topographic analyses of the morphology of the RNMRS and detailed study of the  
803 relationships between the pre-existing basement fabric and rift-related faults provide, for the first  
804 time, evidence supporting the coupling of the Rukwa and North Malawi Rift. Our topographic  
805 analyses in the Rukwa Rift suggest that the Ufipa Fault is the present-day active border fault of

806 the rift. We find that the Ufipa fault is the dominant topographic feature in the northern part of  
807 the RNMRS and diminishes as it encounters the Mbozi block where it becomes a strike slip fault  
808 (Mughese Fault), at which point the border fault polarity flips and the Livingstone Fault is the  
809 dominant fault in the southern part of the RNMRS.

810 Further, we demonstrate the continuity structures along the northeastern and  
811 southwestern margins of the RNMRS. We show that this structural connectivity across the  
812 Mbozi Block transfer zone between the rifts is guided by the pre-existing Precambrian terrane  
813 fabrics and associated shear zones. We show that the coupling of the RNMRS along the  
814 northeastern boundary of the Mbozi Block transfer zone is accommodated by magmatism along  
815 the linking faults, whereas, coupling along the southwestern boundary is accommodated by  
816 strike-slip faulting. Overall, we suggest that the continuation of the boundary faults along the  
817 RNMRS, and their alignment with colinear Precambrian basement fabric and shear zones  
818 indicate the influence of the pre-existing basement structures on the coupling and amalgamation  
819 of approaching colinear rift segments. On the basin-scale bifurcation of the Rukwa Rift, we infer  
820 that the discordance of the basement fabrics within the Mbozi Block transfer zone to those of the  
821 basement terranes bounding it may have facilitated the development of intra-rift bifurcation of  
822 the rift around the transfer zone.

823 Furthermore, we show the influence of pre-existing basement fabrics on the development  
824 of the RNMRS as evidenced in the geometry, termination and kinematics of the rift-bounding  
825 fault segments. Our observations suggest that curvilinear normal fault geometries developed in  
826 areas where the basement fabrics are either curvilinear, composed of superposed sets of  
827 differently-orientated fabrics, or not favorably oriented to the extensional stress field. Whereas,  
828 long, rectilinear fault geometries with narrow or almost unidentifiable breached-relay zones

829 developed in areas where the pre-existing basement fabrics are roughly rectilinear, suggesting  
830 the greater likelihood of occurrence of tip-to-tip linkage of fault segments.

831 Finally, we present the existence of a buried Pre-Cenozoic strike-slip-faulted mafic dike,  
832 which we suggest is potentially the most excellent record of strike-slip kinematics along the  
833 RNMRS. We further suggest that future geochronological analyses of this intrusion may provide  
834 the most-reliable constraint on the timing of the controversial strike-slip faulting event along the  
835 RNMRS.

836

### 837 **Acknowledgements**

838 This work was partially supported by the National Science Foundation (NSF) grant EAR10-  
839 09988 (awarded to E.A. Atekwana). Also, we acknowledge the Oklahoma State University  
840 (OSU) for the Niblack Scholarship awarded to Erin Heilman. We thank the Geological Survey  
841 Department of Malawi for allowing us to purchase the 2013 aeromagnetic data used in this study.  
842 We also thank the Zambia Geological Survey for allowing us to purchase the aeromagnetic data  
843 used in this study. We thank South African Development Community (SADC) for providing the  
844 Tanzania aeromagnetic data used in this study. This is the Oklahoma State University Boone  
845 Pickens School of Geology contribution no. 2018-.

846

847 **References**

- 848 Baer, G., 1991. Mechanisms of dike propagation in layered rocks and in massive, porous  
849 sedimentary rocks. *Journal of Geophysical Research: Solid Earth*, 96(B7), pp.11911-  
850 11929.
- 851 Baranov, V., 1957. A new method for interpretation of aeromagnetic maps: pseudo-gravimetric  
852 anomalies. *Geophysics*, 22(2), pp.359-382.
- 853 Borrego, D.J., 2016. Crustal Structure of the Rungwe Volcanic Province and Region  
854 Surrounding the Northern Lake Malawi Rift Basin. MSc Thesis, Pennsylvania State  
855 University.
- 856 Bosworth, W., 1985. Geometry of propagating continental rifts. *Nature*, 316(6029), p.625.
- 857 Boven, A., Theunissen, K., Sklyarov, E., Klerkx, J., Melnikov, A., Mruma, A. and Punzalan, L.,  
858 1999. Timing of exhumation of a high-pressure mafic granulite terrane of the  
859 Paleoproterozoic Ubende belt (West Tanzania). *Precambrian Research*, 93(1), pp.119-  
860 137.
- 861 Brune, S. and Autin, J., 2013. The rift to break-up evolution of the Gulf of Aden: Insights from  
862 3D numerical lithospheric-scale modelling. *Tectonophysics*, 607, pp.65-79.
- 863 Brune, S., Corti, G. and Ranalli, G., 2017. Controls of inherited lithospheric heterogeneity on rift  
864 linkage: Numerical and analog models of interaction between the Kenyan and Ethiopian  
865 rifts across the Turkana depression. *Tectonics*, 36(9), pp.1767-1786.
- 866 Camelbeeck, T. and Iranga, M.D., 1996. Deep crustal earthquakes and active faults along the  
867 Rukwa trough, eastern Africa. *Geophysical Journal International*, 124(2), pp.612-630.
- 868 Castaing, C., 1991. Post-Pan-African tectonic evolution of South Malawi in relation to the  
869 Karroo and recent East African rift systems. *Tectonophysics*, 191(1-2), pp.55-73.

870 Chorowicz, J., 1989. Transfer and transform fault zones in continental rifts: examples in the  
871 Afro-Arabian rift system. Implications of crust breaking. *Journal of African Earth*  
872 *Sciences (and the Middle East)*, 8(2-4), pp.203-214.

873 Chorowicz, J., 2005. The east African rift system. *Journal of African Earth Sciences*, 43(1),  
874 pp.379-410.

875 Collanega, L., Bell, R., Coleman, A.J., Lenhart, A. and Breda, A., 2018. How do intra-basement  
876 fabrics influence normal fault growth? Insights from the Taranaki Basin, offshore New  
877 Zealand. EarthArxiv, DOI: 10.31223/osf.io/8rn9u

878 Corti, G., 2012. Evolution and characteristics of continental rifting: Analog modeling-inspired  
879 view and comparison with examples from the East African Rift System. *Tectonophysics*,  
880 522, pp.1-33.

881 Daly, M.C., 1988. Crustal Shear Zones in Central Africa - A kinematic approach to Proterozoic  
882 Tectonics. *Episodes*, 11(1), pp.5-11.

883 Daly, M.C., Chorowicz, J. and Fairhead, J.D., 1989. Rift basin evolution in Africa: the influence  
884 of reactivated steep basement shear zones. *Geological Society, London, Special*  
885 *Publications*, 44(1), pp.309-334.

886 Dawson, S.M., Laó-Dávila, D.A., Atekwana, E.A. and Abdelsalam, M.G., 2018. The influence  
887 of the Precambrian Mughese Shear Zone structures on strain accommodation in the  
888 northern Malawi Rift. *Tectonophysics*, 722, pp.53-68.

889 Delvaux, D., and Hanon, M., 1991. Neotectonics of the Mbeya area, SW Tanzania. *Annual*  
890 *report of the Royal Museum of Central Africa, Department of Geology and*  
891 *Mineralogy*, 1992, pp.87-97.

892 Delvaux, D., Kervyn, F., Macheyeke, A.S. and Temu, E.B., 2012. Geodynamic significance of  
893 the TRM segment in the East African Rift (W-Tanzania): Active tectonics and paleostress  
894 in the Ufipa plateau and Rukwa basin. *Journal of Structural Geology*, 37, pp.161-180.

895 Delvaux, D., Levi, K., Kajara, R. and Sarota, J., 1992. Cenozoic paleostress and kinematic  
896 evolution of the Rukwa–North Malawi rift valley (East African Rift System). *Bulletin des*  
897 *Centres de Recherche Exploration-Production ElfAquitaine*, 16, pp.383-406.

898 Fernandez-Alonso, M., Delvaux, D., Klerkx, J. and Theunissen, K., 2001. Structural link  
899 between Tanganyika-and Rukwa-rift basins at Karema-Nkamba (Tanzania): basement  
900 structural control and recent evolution. *Mus. Roy. Afr. Centr., Tervuren (Belgique), Dép.*  
901 *Géol. Min., Rap. Ann*, pp.91-100.

902 Fontijn, K., Delvaux, D., Ernst, G.G., Kervyn, M., Mbede, E. and Jacobs, P., 2010. Tectonic  
903 control over active volcanism at a range of scales: case of the Rungwe Volcanic  
904 Province, SW Tanzania; and hazard implications. *Journal of African Earth Sciences*,  
905 58(5), pp.764-777.

906 Fontijn, K., Williamson, D., Mbede, E. and Ernst, G.G., 2012. The Rungwe Volcanic Province,  
907 Tanzania—A volcanological review. *Journal of African Earth Sciences*, 63, pp.12-31.

908 Fossen, H., 2010. *Structural geology*. Cambridge University Press.

909 Fossen, H. and Rotevatn, A., 2016. Fault linkage and relay structures in extensional settings—A  
910 review. *Earth-Science Reviews*, 154, pp.14-28.

911 Foster, A., Ebinger, C., Mbede, E. and Rex, D., 1997. Tectonic development of the northern  
912 Tanzanian sector of the East African Rift System. *Journal of the Geological*  
913 *Society*, 154(4), pp.689-700.

914 Fritz, H., Abdelsalam, M., Ali, K.A., Bingen, B., Collins, A.S., Fowler, A.R., Ghebreab, W.,  
915 Hauzenberger, C.A., Johnson, P.R., Kusky, T.M. and Macey, P., 2013. Orogen styles in  
916 the East African Orogen: a review of the Neoproterozoic to Cambrian tectonic  
917 evolution. *Journal of African Earth Sciences*, 86, pp.65-106.

918 Gawthorpe, R.L., Jackson, C.A.L., Young, M.J., Sharp, I.R., Moustafa, A.R. and Leppard, C.W.,  
919 2003. Normal fault growth, displacement localisation and the evolution of normal fault  
920 populations: the Hammam Faraun fault block, Suez rift, Egypt. *Journal of Structural*  
921 *Geology*, 25(6), 883-895.

922 Grauch, V.J.S. and Hudson, M.R., 2007. Guides to understanding the aeromagnetic expression of  
923 faults in sedimentary basins: Lessons learned from the central Rio Grande rift, New  
924 Mexico. *Geosphere*, 3(6), pp.596-623.

925 Grauch, V.J.S. and Hudson, M.R., 2011. Aeromagnetic anomalies over faulted strata. *The*  
926 *Leading Edge*, 30(11), pp.1242-1252.

927 Gudmundsson, A. and Brenner, S.L., 2001. How hydrofractures become arrested. *Terra*  
928 *Nova*, 13(6), pp.456-462.

929 Hayward, N.J. and Ebinger, C.J., 1996. Variations in the along-axis segmentation of the Afar  
930 Rift system. *Tectonics*, 15(2), pp.244-257.

931 Helgeson, D.E. and Aydin, A., 1991. Characteristics of joint propagation across layer interfaces  
932 in sedimentary rocks. *Journal of Structural Geology*, 13(8), pp.897-911.

933 Henderson, R.G. and Zietz, I., 1949. The upward continuation of anomalies in total magnetic  
934 intensity fields. *Geophysics*, 14(4), pp. 517-534.



935 Hodge, M., Fagereng, Å., Biggs, J., and Mdala, H., 2018.. Controls on early-rift geometry: New  
936 perspectives from the Bilila-Mtakataka fault, Malawi. *Geophysical Research Letters*, 45,  
937 3896–3905.

938 Hodgson, I., Illsley-Kemp, F., Gallacher, R.J., Keir, D., Ebinger, C.J. and Mtelela, K., 2017.  
939 Crustal structure at a young continental rift: A receiver function study from the  
940 Tanganyika Rift. *Tectonics*, 36(12), pp.2806-2822.

941 Jackson, J. and Blenkinsop, T., 1997. The Bilila-Mtakataka fault in Malaŵi: An active, 100-km  
942 long, normal fault segment in thick seismogenic crust. *Tectonics*, 16(1), pp.137-150.

943 Johnson, S., Mendez, K., Beresh, S.C.M., Mynatt, W.G., Elifritz, E.A., Laó-Dávila, D.A.,  
944 Atekwana, E.A., Abdelsalam, M.G., Chindandali, P.R.N., Chisenga, C. and Gondwe, S.,  
945 2017.. The Relationships of Subparallel Synthetic Faults and Pre-existing Structures in  
946 the Central Malawi Rift. *AGU Fall Meeting poster number T22C-03*.

947 Katumwehe, A.B., Abdelsalam, M.G. and Atekwana, E.A., 2015. The role of pre-existing  
948 Precambrian structures in rift evolution: The Albertine and Rhino grabens,  
949 Uganda. *Tectonophysics*, 646, pp.117-129.

950 Kervyn, F., Ayub, S., Kajara, R., Kanza, E. and Temu, B., 2006. Evidence of recent faulting in  
951 the Rukwa rift (West Tanzania) based on radar interferometric DEMs. *Journal of African*  
952 *Earth Sciences*, 44(2), pp.151-168.

953 Kilembe, E.A. and Rosendahl, B.R., 1992. Structure and stratigraphy of the Rukwa  
954 rift. *Tectonophysics*, 209(1-4), pp.143-158.

955 Kim, S., Nyblade, A.A. and Baag, C.E., 2009. Crustal velocity structure of the Rukwa Rift in the  
956 western branch of the East African Rift system. *South African Journal of Geology*, 112(3-  
957 4), pp.251-260.

958 Kinabo, B.D., Atekwana, E.A., Hogan, J.P., Modisi, M.P., Wheaton, D.D. and Kampunzu, A.B.,  
959 2007. Early structural development of the Okavango rift zone, NW Botswana. *Journal of*  
960 *African Earth Sciences*, 48(2), pp.125-136.

961 Kinabo, B.D., Hogan, J.P., Atekwana, E.A., Abdelsalam, M.G. and Modisi, M.P., 2008. Fault  
962 growth and propagation during incipient continental rifting: Insights from a combined  
963 aeromagnetic and Shuttle Radar Topography Mission digital elevation model  
964 investigation of the Okavango Rift Zone, northwest Botswana. *Tectonics*, 27(3).

965 Koehn, D., Aanyu, K., Haines, S. and Sachau, T., 2008. Rift nucleation, rift propagation and the  
966 creation of basement micro-plates within active rifts. *Tectonophysics*, 458(1-4), pp.105-  
967 116.

968 Kolawole, F., Atekwana, E.A., Laó-Dávila, D.A., Abdelsalam, M.G., Chindandali, P.R., Salima,  
969 J. and Kalindekafe, L., 2018. Active deformation of Malawi Rift's North Basin hinge  
970 zone modulated by reactivation of pre-existing Precambrian shear zone fabric. *Tectonics*,  
971 37, pp.683–704.

972 Lao-Davila, D.A., Al-Salmi, H.S., Abdelsalam, M.G. and Atekwana, E.A., 2015. Hierarchical  
973 segmentation of the Malawi Rift: The influence of inherited lithospheric heterogeneity  
974 and kinematics in the evolution of continental rifts. *Tectonics*, 34(12), pp.2399-2417.

975 Lawley, C.J., Selby, D., Condon, D.J., Horstwood, M., Millar, I., Crowley, Q. and Imber, J.,  
976 2013. Litho-geochemistry, geochronology and geodynamic setting of the Lupa Terrane,  
977 Tanzania: implications for the extent of the Archean Tanzanian Craton. *Precambrian*  
978 *Research*, 231, pp.174-193.

979 Lenoir, J.L., Liégeois, J.P., Theunissen, K. and Klerkx, J., 1994. The Palaeoproterozoic  
980 Ubendian shear belt in Tanzania: geochronology and structure. *Journal of African Earth*  
981 *Sciences*, 19(3), pp.169-184.

982 Mbede, E.I., 2002. Interpretation of reflection seismic data from the Usangu Basin, East African  
983 Rift System. *Tanzania Journal of Science*, 28(1), pp.83-97.

984 McCalpin, J.P., Rockwell, T.K. and Weldon II, R.J., 2009. Paleoseismology of Strike-Slip  
985 Tectonic Environments. *International Geophysics*, 95, pp.421-496.

986 Modisi, M.P., Atekwana, E.A., Kampunzu, A.B. and Ngwisanyi, T.H., 2000. Rift kinematics  
987 during the incipient stages of continental extension: Evidence from the nascent Okavango  
988 rift basin, northwest Botswana. *Geology*, 28(10), pp.939-942.

989 Morley, C.K., 1995. Developments in the structural geology of rifts over the last decade and their  
990 impact on hydrocarbon exploration. *Geological Society, London, Special*  
991 *Publications*, 80(1), pp.1-32.

992 Morley, C.K., 2010. Stress re-orientation along zones of weak fabrics in rifts: An explanation  
993 for pure extension in 'oblique' rift segments?. *Earth and Planetary Science*  
994 *Letters*, 297(3), pp.667-673.

995 Morley, C.K., Cunningham, S.M., Harper, R.M. and Wescott, W.A., 1992. Geology and  
996 geophysics of the Rukwa rift, East Africa. *Tectonics*, 11(1), pp.69-81.

997 Morley, C.K., Nelson, R.A., Patton, T.L. and Munn, S.G., 1990. Transfer zones in the East  
998 African rift system and their relevance to hydrocarbon exploration in rifts (1). *AAPG*  
999 *Bulletin*, 74(8), pp.1234-1253.

1000 Mortimer, E., Paton, D.A., Scholz, C.A., Strecker, M.R. and Blisniuk, P., 2007. Orthogonal to  
1001 oblique rifting: effect of rift basin orientation in the evolution of the North basin, Malawi  
1002 Rift, East Africa. *Basin Research*, 19(3), pp.393-407.

1003 Muirhead, J.D. and Kattenhorn, S.A., 2018. Activation of preexisting transverse structures in an  
1004 evolving magmatic rift in East Africa. *Journal of Structural Geology*, 106, pp.1-18.

1005 Njinju, E.A., Atekwana, E.A., Stamps, D.S., Abdelsalam, M.G., Atekwana, E.A., Mickus, K.L.,  
1006 Kolawole, F. and Nyalugwe, V., 2018. Lithospheric Structure of the Malawi Rift:  
1007 Implications for Rifting Processes in Magma Poor Rift Systems. *EarthArXiv*, DOI:  
1008 10.31223/osf.io/83qd9

1009 Peirce, J.W. and Lipkov, L., 1988. Structural interpretation of the Rukwa rift, Tanzania.  
1010 *Geophysics*, 53(6), pp.824-836.

1011 Phillips, T.B., Jackson, C.A., Bell, R.E., Duffy, O.B. and Fossen, H., 2016. Reactivation of  
1012 intrabasement structures during rifting: A case study from offshore southern  
1013 Norway. *Journal of Structural Geology*, 91, pp.54-73.

1014 Pik, R., Marty, B., Carignan, J., Yirgu, G. and Ayalew, T., 2008. Timing of East African Rift  
1015 development in southern Ethiopia: Implication for mantle plume activity and evolution of  
1016 topography. *Geology*, 36(2), pp.167-170.

1017 Ring, U., Kröner, A., Buchwaldt, R., Toulkeridis, T. and Layer, P.W., 2002. Shear-zone patterns  
1018 and eclogite-facies metamorphism in the Mozambique belt of northern Malawi, east-  
1019 central Africa: implications for the assembly of Gondwana. *Precambrian*  
1020 *Research*, 116(1), pp.19-56.

1021 Rosendahl, B.R., 1987. Architecture of continental rifts with special reference to East  
1022 Africa. *Annual Review of Earth and Planetary Sciences*, 15(1), pp.445-503.

- 1023 Rotevatn, A., Jackson, C.A.L., Tvedt, A.B.M., Bell, R. and Blækkan, I., 2018. How do normal  
1024 faults grow?. EarthArXiv.
- 1025 Salem, A., Williams, S., Fairhead, J.D., Ravat, D. and Smith, R., 2007. Tilt-depth method: A  
1026 simple depth estimation method using first-order magnetic derivatives. *The Leading  
1027 Edge*, 26(12), pp.1502-1505.
- 1028 Sarafian, E., Evans, R.L., Abdelsalam, M.G., Atekwana, E., Elsenbeck, J., Jones, A.G. and  
1029 Chikambwe, E., 2018. Imaging Precambrian lithospheric structure in Zambia using  
1030 electromagnetic methods. *Gondwana Research*, 54, pp.38-49.
- 1031 Schenk, V., Boniface, N., Loose, D., Alkmim, F.F. and Noce, C.M., 2006. Paleoproterozoic  
1032 subduction zones at the margins of the Tanzania and Congo Cratons: evidence from  
1033 eclogites with MORB-type chemistry in the Usagaran–Ubendian Belts of Tanzania and  
1034 the Nyong complex of Cameroon. *The Paleoproterozoic Record of the São Francisco  
1035 Craton*, pp.102-103.
- 1036 Scholz, C.H. and Contreras, J.C., 1998. Mechanics of continental rift  
1037 architecture. *Geology*, 26(11), pp.967-970.
- 1038 Silva, J.B., 1986. Reduction to the pole as an inverse problem and its application to low-latitude  
1039 anomalies. *Geophysics*, 51(2), pp.369-382.
- 1040 Siuda, K., Magee, C., Bell, R., Jackson, C.A.L. and Collanega, L., 2018. Pre-existing basement  
1041 thrusts influence rifting in the Taranaki Basin, New Zealand. EarthArXiv.
- 1042 Szatmari, P. and Milani, E.J., 1999. Microplate rotation in northeast Brazil during South Atlantic  
1043 rifting: Analogies with the Sinai microplate. *Geology*, 27(12), pp.1115-1118.

- 1044 Taylor, B., Weiss, J.R., Goodliffe, A.M., Sachpazi, M., Laigle, M. and Hirn, A., 2011. The  
1045 structures, stratigraphy and evolution of the Gulf of Corinth rift, Greece. *Geophysical*  
1046 *Journal International*, 185(3), pp.1189-1219.
- 1047 Theunissen, K., Klerkx, J., Melnikov, A. and Mruma, A., 1996. Mechanisms of inheritance of  
1048 rift faulting in the western branch of the East African Rift, Tanzania. *Tectonics*, 15(4),  
1049 pp.776-790.
- 1050 Van de Velde, P., and De Waele, B., 1998. Geology of the Mupamadzi River area. Explanation  
1051 of degree sheet 1231, SW quarter, Report No. 105, *Geological Survey Department*,  
1052 *Republic of Zambia*.
- 1053 Versfelt, J. and Rosendahl, B.R., 1989. Relationships between pre-rift structure and rift  
1054 architecture in Lakes Tanganyika and Malawi, East Africa. *Nature*, 337(6205), p.354.
- 1055 Wheeler, W.H. and Karson, J.A., 1989. Structure and kinematics of the Livingstone Mountains  
1056 border fault zone, Nyasa (Malawi) Rift, southwestern Tanzania. *Journal of African Earth*  
1057 *Sciences (and the Middle East)*, 8(2-4), pp.393-413.
- 1058 Wheeler, W.H. and Karson, J.A., 1994. Extension and subsidence adjacent to a "weak"  
1059 continental transform: An example from the Rukwa rift, East Africa. *Geology*, 22(7),  
1060 pp.625-628.
- 1061 Wichura, H., Bousquet, R., Oberhänsli, R., Strecker, M.R. and Trauth, M.H., 2011. The Mid-  
1062 Miocene East African Plateau: a pre-rift topographic model inferred from the  
1063 emplacement of the phonolitic Yatta lava flow, Kenya. *Geological Society, London*,  
1064 *Special Publications*, 357(1), pp.285-300.
- 1065 Wilson, T.J., 1999. Cenozoic structural segmentation of the Transantarctic Mountains rift flank  
1066 in southern Victoria Land. *Global and Planetary Change*, 23(1-4), pp.105-127.

1067 Wu, D. and Bruhn, R.L., 1994. Geometry and kinematics of active normal faults, South Oquirrh  
1068 Mountains, Utah: implication for fault growth. *Journal of Structural Geology*, 16(8),  
1069 pp.1061-1075.

1070 Xue, L., Gani, N.D. and Abdelsalam, M.G., 2017. Geomorphologic proxies for bedrock rivers:  
1071 A case study from the Rwenzori Mountains, East African Rift system. *Geomorphology*,  
1072 285, pp.374-398.

1073 Zhang, X. and Jeffrey, R.G., 2008. Reinitiation or termination of fluid-driven fractures at  
1074 frictional bedding interfaces. *Journal of Geophysical Research: Solid Earth*, 113(B8).

1075 Zhang, X., Jeffrey, R.G. and Thiercelin, M., 2007. Deflection and propagation of fluid-driven  
1076 fractures at frictional bedding interfaces: a numerical investigation. *Journal of Structural*  
1077 *Geology*, 29(3), pp.396-410.

1078 Zhao, M., Langston, C.A., Nyblade, A.A. and Owens, T.J., 1997. Lower-crustal rifting in the  
1079 Rukwa graben, East Africa. *Geophysical Journal International*, 129(2), pp.412-420.

1080

1081

1082

1083

1084

1085

1086

1087


1088

1089

1090

1091

1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102



1103

1104 Fig.1. (A) Topographic map of the East African Rift System, showing the segments of the rift  
1105 system and the location of the Rukwa-North Malawi Rift Segment (RNMRS) (black square). (B)  
1106 Digital Elevation Model (DEM) of the RNMRS showing the different domains. Solid black lines  
1107 represent topographic profile lines shown in Figure 3. ARR= Albertine-Rhino Rift, ASZ= Aswa



1108 Shear Zone, MT= Musangano Trough, MwSZ= Mwembeshi Shear Zone, RVP= Rungwe  
 1109 Volcanic Province, ST= Songwe Trough.

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126


1127

1128

1129

1130

1131

















1132 Fig. 2. (A) Precambrian domains around the Tanzanian Craton (modified after Theunissen et al.,  
 1133 1996; Boven et al., 1999; Fernandez-Alonso et al., 2001; Schenk et al., 2006). 1 = Archean  
 1134 craton; 2 = Paleoproterozoic Usagaran and Ubendian orogenic belts; 3 = Mozambique orogenic  
 1135 belt; 4 = western limit of Pan-African influence; 5 = Meso and/or Neoproterozoic sediments  
 1136 (only for the Ubendian Belt region); 6 = Phanerozoic volcanics and sedimentary rocks. (B)  
 1137 Regional geological map of southwest Tanzania showing the terrane structure of the Ubendian  
 1138 orogenic belt within which the Rukwa-Malawi Rift Segment (RNMR) developed.

1139

1140

1141



- Normal Fault
- Cenozoic Sediments
- Rungwe Volcanics
- Precambrian Basement

1142  
1143  
1144  
1145  
1146  
1147  
1148  
  
1149  
  
1150  
  
1151  
  
1152  
  
1153  
  
1154  
  
1155  
  
1156  
  
1157  
  
1158  
  
1159  
  
1160  
  
1161  
  
1162  
  
1163  
  
1164  
  
1165  
  
1166  
  
1167

Fig. 3. Topographic profiles across the Rukwa - North Malawi Rift segment. Profile numbers correspond to the profiles in Figure 1B. KLF = Kalambo Fault, KF = Kanda Fault, KSF = Kasanga Fault, LF = Lupa Fault, LVF = Livingstone Fault, MB = Mbozi Block, MBF = Mbiri Fault, MF = Mbaka Fault, MT = Musangano Trough, RVP = Rungwe Volcanic Province, ST = Songwe Trough, UF = Ufipa Fault, USF = Usisya Fault.



1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210




Fig. 4. (A) The reduced to pole Total Magnetic Intensity (RTP-TMI) map of the Rukwa-North Malawi Rift area, draped over the topographic digital elevation model of SW Tanzania. (B) The 1<sup>st</sup> vertical derivative map of the RTP-TMI map of the Rukwa-North Malawi Rift area draped over the topographic digital elevation model.

1211  
 1212  
 1213  
 1214  
 1215  
 1216  
 1217  
 1218  
 1219  
 1220  
 1221  
 1222  
 1223  
 1224  
 1225  
 1226  
 1227  
 1228  
 1229  
 1230  
 1231  
 1232  
 1233  
 1234  
 1235  
 1236  
 1237  
 1238  
 1239  
 1240  
 1241  
 1242  
 1243  
 1244  
 1245  
 1246  
 1247  
 1248  
 1249  
 1250




Fig. 5. Northernmost part of the Ufipa Fault, the SW boundary fault of the Rukwa Rift. Topographic digital elevation model in the top panel (A), the vertical derivative of the magnetic data in the middle panel (B), and a structural interpretation in the bottom panel (C).



1251  
 1252  
 1253  
 1254  
 1255  
 1256  
 1257  
 1258  
 1259  
 1260  
 1261  
 1262  
 1263  
 1264  
 1265  
 1266  
 1267  
 1268  
 1269  
 1270  
 1271  
 1272  
 1273  
 1274  
 1275  
 1276  
 1277  
 1278




Fig. 6. Central segment of the Ufipa Fault, SW boundary of the Rukwa Rift. Topographic digital elevation model in the top panel (A), the vertical derivative of the magnetic data in the middle panel (B), and a structural interpretation in the bottom panel (C).

1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322




Fig. 7. The Musangano Trough part of the SW boundary of the Rukwa Rift. Topographic digital elevation model in the top panel (A), the vertical derivative of the magnetic data in the middle panel (B), and a structural interpretation in the bottom panel (C).



1323  
 1324  
 1325  
 1326  
 1327  
 1328  
 1329  
 1330  
 1331  
 1332  
 1333  
 1334  
 1335  
 1336  
 1337  
 1338  
 1339  
 1340  
 1341  
 1342  
 1343  
 1344  
 1345  
 1346  
 1347  
 1348  
 1349  
 1350  
 1351  
 1352  
 1353




Fig. 8. The SW margin of the Mbozi Block. Topographic digital elevation model in the top panel (A), the vertical derivative of the magnetic data in the middle panel (B), and a structural interpretation in the bottom panel (C).



1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399




Fig. 9. The SW margin of the North Malawi Rift Basin (Karonga area). Topographic digital elevation model in the top panel (A), the vertical derivative of the magnetic data in the middle panel (B), and a structural interpretation in the bottom panel (C). MSZ = Mughese Shear Zone, KF = Karonga Fault. Fault and basement fabric interpretations from Kolawole et al. (2018).

1400  
 1401  
 1402  
 1403  
 1404  
 1405  
 1406  
 1407  
 1408  
 1409  
 1410  
 1411  
 1412  
 1413  
 1414  
 1415  
 1416  
 1417  
 1418  
 1419  
 1420  
 1421  
 1422  
 1423  
 1424  
 1425  
 1426  
 1427  
 1428  
 1429  
 1430  
 1431  
 1432  
 1433  
 1434  
 1435  
 1436  
 1437  
 1438  
 1439  
 1440  
 1441  
 1442  
 1443  
 1444  
 1445





Fig. 10. The northern section of the Lupa Fault, the NE boundary fault of the Rukwa Rift. Topographic digital elevation model in the top panel (A), the vertical derivative of the magnetic data in the middle panel (B), and a structural interpretation in the bottom panel (C).





1447  
 1448  
 1449  
 1450  
 1451  
 1452  
 1453  
 1454  
 1455  
 1456  
 1457  
 1458  
 1459  
 1460  
 1461  
 1462

Fig. 11. The southern section of the Lupa Fault, the NE margin of the Mbozi Block (Rungwe Volcanic Province) and a section of the Livingstone Fault (NE border fault if the North Malawi Rift). Topographic digital elevation model in the top panel (A), the vertical derivative of the magnetic data in the middle panel (B), and a structural interpretation in the bottom panel (C). KVC = Kyejo Volcanic Center; NVC = Ngozi Volcanic Center; RVC = Rungwe Volcanic Center. Volcanic centers from Fontijn et al. (2010, 2012).

1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508




Fig. 12. Generalized geometrical relations of stratigraphic units and normal faults of the Rukwa rift, showing thickening of Cenozoic sediments towards both the Lupa and Ufipa border faults (modified after Zhao et al., 1997).



1509  
 1510  
 1511  
 1512  
 1513  
 1514  
 1515  
 1516  
 1517  
 1518  
 1519  
 1520  
 1521  
 1522  
 1523  
 1524  
 1525  
 1526  
 1527  
 1528  
 1529  
 1530  
 1531  
 1532  
 1533  
 1534  
 1535  
 1536  
 1537  
 1538  
 1539  
 1540  
 1541  
 1542  
 1543  
 1544  
 1545  
 1546  
 1547  
 1548  
 1549  
 1550  
 1551  
 1552  
 1553  
 1554




Fig. 13. (A) 1<sup>st</sup> vertical derivative of the aeromagnetic data covering the Kaseye-Chitipa area along the SW margin of the Mbozi Block (see location in Fig. 8C). (B) Close-up of the central and southern segments of a buried mafic dike (the “Chitipa Dike”) showing side-stepping segments that coincide with pre-existing basement fabric. (C) Close-up of the northern segment of the dike showing right-lateral offsets by the continuation of the Ufipa Fault. This strike-slip fault segment is here-in referred to as the Mughese Fault.

1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600





Fig. 14. 3-dimensional (3-D) conceptual model of the geometry of the interpreted dike (the "Chitipa Dike") and its interactions with the pre-existing basement fabric and the post-emplacement Mugheze strike-slip fault offset.



1602

1603

1604

1605

1606

1607

1608

1609


1610

1611

1612

Fig. 15. (A) Generalized cartoon (map view) of the Rukwa-North Malawi Rift Segment (RNMRS) illustrating the continuous structural connectivity along the northeast and southwest boundaries, guided by the basement fabric. Black solid line = fault, blue solid line = basement fabric. RVP = Rungwe Volcanic Province. (B-E) Cross-section cartoons across the segments of the RNMRS, illustrating the possible subsurface geometries and interactions of the domain-bounding structures. LLF = Livingstone-Lupa Fault.





1613  
 1614  
 1615  
 1616  
 1617  
 1618  
 1619  
 1620  
 1621  
 1622  
 1623  
 1624  
 1625  
 1626  
 1627  
 1628  
 1629  
 1630  
 1631  
 1632  
 1633  
 1634  
 1635  
 1636  
 1637  
 1638  
 1639  
 1640  
 1641  
 1642  
 1643  
 1644  
 1645  
 1646  
 1647

Fig. 16. (A-B) The Wasatch Fault in the Salt Lake area, Utah, and the first-order faults in the northern North Sea (base Cretaceous unconformity) showing concave-curvilinear fault geometries (modified after Fossen and Rotevatn, 2016). (C) Strikingly similar Concave-curvilinear fault geometry occurs in the northern Ufipa Fault and Chisi Fault in the Rukwa Rift. However, the Central Ufipa Fault shows convex-curvilinear fault geometry. (D) The Bilila-Mtakataka Fault also shows excellent convex curvilinear fault geometries.







Fig. 17. Models illustrating the control of varying configurations of pre-existing basement fabrics on the development of concave- and convex-curvilinear normal fault plan-view geometries. These models are based on the observations from the Rukwa – North Malawi Rift Segment (this study).

1678

1679

1680  
 1681  
 1682  
 1683  
 1684  
 1685  
 1686  
 1687  
 1688  
 1689  
 1690  
 1691  
 1692  
 1693  
 1694  
 1695  
 1696  
 1697  
 1698  
 1699  
 1700  
 1701  
 1702  
 1703  
 1704



1705 Fig. 18. Examples of system-scale rift bifurcation in the (A) East African Rift System, (B) Red  
 1706 Red Sea Rift and (C) South Atlantic Rift; and examples of basin-scale rift bifurcation in the (D)  
 1707 Rukwa Rift (see Fig. 18A for location), (E) southern Malawi Rift and (F) Albertine Rift (see Fig.  
 1708 18A for location). These examples show that the 2<sup>nd</sup> order bifurcations are integral components  
 1709 of inter-rift transfer zones where the coupling of rift segments take place.