
This paper is a non-peer reviewed manuscript submitted to EarthArXiv. The paper will be published on the websites of the International Association of Hydrogeologists (IAH) and the IAH Commission on Groundwater Quality. Subsequent versions of this manuscript may have slightly different content. We welcome feedback and comments, which can be sent to the authors or to info@iah.org.

International Association of Hydrogeologists Technical Insight Paper Series:

Antimicrobial Resistance – A Growing Global Groundwater Challenge

T.Boving^{1*}, C.Hernández-Zepeda², D.Lapworth³, M.Lenczewski⁴, B.P.Vellanki⁵

¹University of Rhode Island, Department of Geosciences, Kingston, RI, USA

²Centro de Investigación Científica de Yucatán, Mérida, Yucatán, México

³British Geological Survey, Nottingham, United Kingdom

⁴Northern Illinois University, Department of Earth, Atmosphere and Environment, DeKalb, IL, USA

⁵Indian Institute of Technology, Roorkee, India

**Corresponding author: tboving@gmail.com*

@iahgroundwater

Non-peer reviewed preprint submitted to EarthArXiv

Abstract

Antimicrobials are composed of medications such as antibiotics and antivirals which are widely used to ensure human and animal health. However, inappropriate use of antimicrobials is a major driver of antimicrobial resistance. AMR arises from complex interactions between humans, animals, microbial organisms, medicines, wildlife and the environment. Once in the wastewater stream, antibiotics are only partially removed by most wastewater treatment plants and are frequently detected in both effluent water and sludges. Bacteria can develop antibiotic resistance through several mechanisms such as genetic mutations or by acquiring antibiotic resistance genes (ARGs) from other bacteria via horizontal gene transfer. Based on studies worldwide, hundreds of ARGs have been detected in various aquatic systems, posing significant environmental and public health concerns. Once in the environment, ARGs can disseminate within bacterial populations from which some species could re-enter human and animal systems via contaminated water, food, or recreational activities. This environmental reservoir of resistance undermines the effectiveness of antibiotic therapies, complicates the treatment of infectious diseases, and increases the risk of outbreaks caused by multidrug-resistant pathogens. The current strategy of preventing AMR is to protect aquatic environments from pharmaceutical contamination, especially antibiotics. But because of the difficulty and expense of detecting compounds associated with AMR, most countries do not have national, legally enforceable water quality standards. Further developments in affordable screening and treatment solutions will be necessary to tackle this insidious contamination.



Technical Insight Series

Antimicrobial Resistance – A Growing Global Groundwater Challenge



International Association
of Hydrogeologists

Thick foam carpeting the Yamuna River downstream from a sewage channel in New Delhi, India (T.Boving URI)

Antimicrobials have been used for decades and are an essential part of modern medicine. Antimicrobials are composed of medications such as antibiotics, antifungals, antivirals, anthelmintics, etc., widely used to ensure human and animal health. However, inappropriate use of antimicrobials is known to be a major driver of antimicrobial resistance (AMR)¹. AMR arises from complex interactions between humans, animals, microbial organisms, medicines, wildlife and the environment (Figure 1).

A recent study estimated the proportion of inappropriate antibiotic use globally to be on average 30% (~13,000,000 kg) equivalent to the annual consumption of antibiotics in China². AMR is a chronic public health problem globally, set to account for 10 million deaths each year by 2050³.

Animal usage of antibiotics varies considerably, for example in the USA it accounts for around 70% of medically important antibiotic sales, while in the UK this figure is around 30%, indeed across Europe between 2011-2018 there was a 33% reduction in the sales of antibiotics for livestock⁴. However, in Asia, the far East and Oceania, antibiotic use per unit livestock remains around 4 times higher than Europe⁴.

To manage the threat of AMR a 'One Health' Approach is required which integrates efforts across multiple sectors and disciplines. Rapid advances in genomic surveillance for AMR has enabled monitoring of persistence and abundance of AMR genes and mutations within microbial organisms⁵ and forms a central tool for monitoring AMR spread globally.




Figure 1: Environmental releases and pathways of chemical compounds that can cause antimicrobial resistance (AMR). (AI Generated. Design by B.P. Vellanki IIT-R; M. Lenczewski NIU)

Antimicrobial Pharmaceuticals and Their Detection in Waters

Most modern antibiotics are synthetic derivatives of various natural compounds and encompass a diverse range of chemical structures. Of the many antibiotics on the market, ciprofloxacin, tetracycline, sulfamethoxazole, and amoxicillin are among the most

frequently detected antibiotic residues in rivers, groundwater, and wastewater^{6,7}. Notably, most antibiotics were discovered decades ago with very few new ones developed since.

The global surge in antibiotic consumption for both medical and agricultural applications has inadvertently introduced these bioactive compounds into aquatic environments⁸. Importantly, antibiotics are not completely metabolized when administered to humans and animals. In the case of antibiotics administered to animals, approximately 90% are excreted in original form or as metabolites⁹. Once in the wastewater stream, antibiotics are incompletely removed by most wastewater treatment plants (WWTPs) and are frequently detected in both effluent water and sludges. Sulfamethoxazole, ciprofloxacin, and trimethoprim were among the most persistent antimicrobics, with overall aqueous removal efficiencies typically below 40%. Untreated municipal sewage may contain as much as 10 $\mu\text{g L}^{-1}$ antibiotics, whereas typical antibiotic concentrations in treated municipal sewage range from non-detects to approximately 1 $\mu\text{g L}^{-1}$ ¹⁰. When AMR causing compounds enter the groundwater, they pose a high risk to human health, as indicated in a study of private wells¹¹. A study of groundwater in China showed that 47 antibiotics, mainly sulfonamides and fluoroquinolones, have been detected and that these antibiotics in groundwater will induce the production of resistance genes, causing ecological harm¹².

Antibiotic Resistance Genes (ARG)

WWTPs are hot spots for the transfer and proliferation of AMR bacteria in the environment²⁶. Especially during biological treatment of domestic sewage, conditions are conducive for the emergence of AMR bacteria via horizontal gene transfer. A recent study showed that full scale WWTP in a metropolitan city in India could not remove the clinically relevant antimicrobial resistant genes (AMG) from wastewater and anaerobically digested sludge¹³. Another study revealed that the agricultural soils amended with sewage sludge had higher abundance of ARGs than normal agricultural soils¹⁴. Elevated levels of AMR-indicating class 1 integron-integrase gene, *int11*, and a wide range of emerging contaminants in urban surface waters and lower levels were found in groundwaters in Bangaluru, India¹⁵. Risk quotients indicated potential AMR development caused by high concentrations of azithromycin, fluconazole, and sulfanilamide in surface waters due to sewage inflows.

Bacteria can develop antibiotic resistance through several mechanisms¹⁶. These traits in antibiotic-resistant bacteria (ARB) can arise through genetic mutations or by acquiring antibiotic resistance genes (ARGs) from other bacteria via horizontal gene transfer (HGT), occurring mainly through conjugation, transformation, and transduction.

Urban water systems (Figure 2) contribute to the dissemination of resistance determinants between bacterial species, thereby making aquatic environments reservoirs and routes that facilitate the spread of ARGs and the emergence of antibiotic-resistant pathogens¹⁷. To

date, hundreds of ARGs have been detected in various aquatic systems, posing significant environmental and public health concerns ¹⁸.

Box 1

The Potomac River sewage spill, 2026

The 2026 Potomac River sewage spill was a massive raw sewage release caused by a major sewer line collapse north of Washington, D.C. Estimates put the spill at roughly 240–300 million gallons of raw sewage discharged into the river, one of the largest wastewater spills in U.S. history. Persistently elevated *E. coli* and *Staphylococcus aureus*, including antibiotic-resistant strain of this bacterium known as MRSA, in river water and nearby soils weeks after the break were reported, with contamination detected several miles downstream. Because many first-line antibiotics no longer work, MRSA infections can be harder to treat, may require stronger or intravenous drugs.

AMR indicators in groundwater and surface water in urban India

Urban groundwater and surface water in the South Indian city of Bengaluru was investigated for antimicrobial resistance (AMR) indicators and emerging organic contaminants (EOC) ¹⁵. Out of 125 EOCs detected in this study, 62 were present in groundwaters and a number of these compounds could be linked directly to distinct recharge sources. Risk quotients were computed to evaluate the level of risk associated with the concentrations of EOCs to contribute to AMR development. Of the nine detected antimicrobials that could have RQs calculated, three were found at concentrations posing a risk of AMR development azithromycin, fluconazole and sulfanilamide. The ubiquitous detection of the sweetener sucralose indicated recent groundwater recharge and a contribution of imported Cauvery River water for recharge. This study highlights the need for monitoring and water protection and the role of EOCs as potential drivers of AMR.



Figure 2: Untreated sewage is discharging in a channel that drains wastewater into the Yamuna River in New Delhi, India. Once released into the surface water, compounds like antimicrobial pharmaceuticals can migrate into the groundwater (T. Boving – URI)

Once in the environment, ARGs can disseminate within bacterial populations from which some species could re-enter human and animal systems via contaminated water, food, or recreational activities (Figure 1). This environmental reservoir of resistance undermines the effectiveness of antibiotic therapies, complicates the treatment of infectious diseases, and increases the risk of outbreaks caused by multidrug-resistant pathogens ^{17, 19, 20}.

Regulations

The current strategy of preventing AMR is to protect aquatic environments from pharmaceutical contamination, especially antibiotics. However, monitoring pharmaceuticals is insufficient to fully assess the threat associated with AMR. Rather, additional indicators are needed, such as monitoring the presence of antibiotic-resistant bacteria and analyzing the transfer of resistance genes ²¹. But because of the difficulty and expense of detecting compounds associated with AMR, most countries do not have national, legally enforceable water quality standards.

In the United States, the Environmental Protection Agency (USEPA) issues a list of unregulated drinking water contaminants that are known or anticipated to occur in public water systems but currently are not subject to national primary drinking water regulations. Compounds used in pharmaceuticals or byproducts of their use, such as estrogenic hormones (estradiol, estriol, estrone) and antibiotics (erythromycin), were proposed but were not considered for further regulations when the 4th Contaminant Candidate List (CCL) was finalized in 2021 ²². The current CCL5 list includes 81 contaminants or contaminant groups; none are antibiotics.

In the European Union (EU), the Water Framework Directive (WFD) is the main tool to reduce the emissions of priority substances that harm the aquatic environment. Updated lists of these substances have been published since 2001 but pharmaceuticals were not included in any of these lists ²¹. In 2013, the EU established the first Surface Water Watch List, which, like the EPA's CCL, lists compounds suspected of posing a risk to aquatic environments but lack sufficient data to be included in the WFD. Updated every 4 years, the Watch List includes several antibiotics, hormones and other drugs.

The Indian Government proposed a bill on pollution limits for antibiotics at manufacturing sites. The proposed limits were based on measured concentrations in treated effluents. However, the bill passed in 2020 without antibiotic discharge limits due to critiques primarily from manufacturers ²³.

The AMR Industry Alliance, an industry group headquartered in Switzerland, formulated Antibiotic Manufacturing Standard (AMS), which is a set of minimum requirements for evaluating risks in antibiotic supply chains to prevent the release of antibiotic residues into the environment and mitigate AMR ²⁴. The voluntary standard requires that the manufacturer

of an antibiotic must have an effective environmental management system and that the antibiotic's Predicted No-Effect Concentrations (PNEC), or the level at which a substance will not have an adverse effect on its environment, are met²⁴. The PNEC below which AMR is not likely to develop ranges from about 0.1 to 2.0 $\mu\text{g L}^{-1}$ for many antibiotics⁹. For comparison, typical antibiotic concentrations in untreated hospital effluent and industrially polluted surface water can be orders of magnitude higher than PNEC and even can reach the minimal inhibitory concentration (which is the lowest concentration of an antimicrobial drug that prevents the visible growth of a microorganism in a laboratory setting¹⁰).



Figure 3: As part of a complex mixture of pharmaceuticals and personal care products (PPCPs), antimicrobial compounds are being released into the environment via many pathways where they can pollute groundwater and surface water bodies. Sampling and testing of wastewater in the field is a first step towards studying the environmental impacts of AMR. Here, students from IIT Roorkee in India studying an artificial lake fed by partially treated wastewater. (T. Boving – URI).

Future Outlook

AMR and the presence of pharmaceuticals in general is a major groundwater quality issue in the coming decades. The ubiquitous distribution of AMR around the globe has the potential to significantly impact the quality and availability of water resources in many regions. The pervasive presence of AMR in the environment raises concerns about impacts to ecosystem and human health. Further developments in affordable screening and treatment solutions will be necessary to tackle this insidious contamination.

Further Information

The global reach of the AMR problem is reflected in numerous guidance documents prepared by national and international organizations. In response to growing concerns of AMR the World Health Organization launched Global Antimicrobial Resistance and Use Surveillance (GLASS) in 2015, which includes annual reporting on AMR and antimicrobial consumption data. In 2024, the WHO published the Guidance for Responsible and Sustainable Manufacturing of Antibiotics²³ in which it provides a framework for manufacturers to reduce antibiotic pollution from their facilities by setting human health-based targets. It also established a risk management process for wastewater and solid waste. The United Nation's General Assembly High-Level Meeting on Antimicrobial Resistance²⁵ is an international effort to address AMR. This initiative responds to drug-resistant infections know no borders, meaning no single country can respond to AMR alone. However, it must be noted that the water quality regulations of many countries so far pay little attention to the growing AMR problem. Drivers of AMR in groundwater and groundwater dependent ecosystems clearly remains an understudied area of research and there is a need for more detailed studies to fill this knowledge gap.

References

- ¹ Goossens H, Ferech M, Vander Stichele R, et al. Outpatient antibiotic use in Europe and association with resistance: a crossnational database study. *Lancet* 2005;365:579–87. [https://doi.org/10.1016/S0140-6736\(05\)17907-0](https://doi.org/10.1016/S0140-6736(05)17907-0)
- ² Mulchandani, R., Tiseo, K., Nandi, A., Klein, E., Gandra, S., Laxminarayan, R., Van Boeckel, T. (2025). Global trends in inappropriate use of antibiotics, 2000-2021: scoping review and prevalence estimates. *BMJ public health*, 3(1), e002411. <https://doi.org/10.1136/bmjph-2024-002411>
- ³ Tang, K.W.K., Millar, B.C. and Moore, J.E., 2023. Antimicrobial resistance (AMR). *British journal of biomedical science*, 80, p.11387. <https://doi.org/10.3389/bjbs.2023.11387>
- ⁴ Singh, B., Bhat, A. and Ravi, K., 2024. Antibiotics misuse and antimicrobial resistance development in agriculture: a global challenge. *Environment & Health*, 2(9), pp.618-622. <https://pubs.acs.org/doi/full/10.1021/envhealth.4c00094>
- ⁵ Djordjevic, S.P., Jarocki, V.M., Seemann, T., Cummins, M.L., Watt, A.E., Drigo, B., Wyrsh, E.R., Reid, C.J., Donner, E. and Howden, B.P., 2024. Genomic surveillance for antimicrobial resistance—a One Health perspective. *Nature Reviews Genetics*, 25(2), pp.142-157. <https://www.nature.com/articles/s41576-023-00649-y>

- ⁶Githinji, L. J. M., Musey, M. K., Ankumah, R. O. (2011). Evaluation of the fate of ciprofloxacin and amoxicillin in domestic wastewater. *Water, Air, and Soil Pollution*, 219(1–4), 191–201. <https://doi.org/10.1007/s11270-010-0697-1>.
- ⁷ Samrot, A. V., Wilson, S., Preeth, R. S. S., Prakash, P., Sathiyasree, M., Saigeetha, S., Shobana, N., Pachiyappan, S., & Rajesh, V. V. (2023). Sources of antibiotic contamination in wastewater and approaches to their removal—An overview. *Sustainability*, 15(16). <https://doi.org/10.3390/su1516>
- ⁸ Boving T.B., Dixit D., Dutta A., 2026. Photocatalytic Treatment of Antibiotic Compounds in Wastewater Streams. In: *Technologies for Soil and Water Pollution Remediation*. Ed: Uthandi S., Li P., Elumalai V., Boving T., Elsevier. DOI: <https://doi.org/10.1016/B978-0-443-36731-1.00001-7>
- ⁹ Adhikari S., Lee H.H., Kim D.-H., 2024. Primary' antibiotics in wastewater treatment plants. *iScience*, Volume 27, Issue 9, 110789, <https://doi.org/10.1016/j.isci.2024.110789> .
- ¹⁰ Larsson DGJ, Flach CF, 2022. Antibiotic resistance in the environment. *Nat Rev Microbiol*. 2022 May;20(5):257-269. doi: 10.1038/s41579-021-00649-x.
- ¹¹ Alawi, M., Smyth, C., Drissner, D. et al., 2024. Private and well drinking water are reservoirs for antimicrobial resistant bacteria. *npj Antimicrob Resist* 2,7. <https://doi.org/10.1038/s44259-024-00024-9>
- ¹² Zeng H., Li J., Zhao W., Xu J., Xu H., Li D., Zhang J., 2022. The Current Status and Prevention of Antibiotic Pollution in Groundwater in China. *Int J Environ Res Public Health*. 7;19(18):11256. doi: 10.3390/ijerph191811256 .
- ¹³ Dixit, S., Vijayan, J., Singh, H., Nambi, I. M., Renganathan, T., Ravikrishna, R., & Singh, G. (2025). Removal of antibiotic resistance genes and co-selectors in a full-scale sewage treatment plant during droughts and floods. *Environmental Science: Water Research & Technology*, 11(3), 702-713.
- ¹⁴ Hung, W. C., Miao, Y., Truong, N., Jones, A., Mahendra, S., & Jay, J. (2022). Tracking antibiotic resistance through the environment near a biosolid spreading ground: resistome changes, distribution, and metal (loid) co-selection. *Science of the Total Environment*, 823, 153570.
- ¹⁵ Brauns B., Chandra S., Civil W., Lapworth D.J., MacDonald A.M., McKenzie A.A., Read D.S., Sekhar M., Singer A.C., Thankachan A., Tipper H.J., 2024. Presence of emerging organic contaminants and microbial indicators in surface water and groundwater in urban India, *Environmental Pollution*, 362, <https://doi.org/10.1016/j.envpol.2024.124983> .
- ¹⁶ Taylor N.G.H, Verner-Jeffreys D.W., Baker-Austin C., 2011. Aquatic systems: maintaining, mixing and mobilising antimicrobial resistance? *Trends in Ecology & Evolution*, 26/6, <https://doi.org/10.1016/j.tree.2011.03.004>.

- ¹⁷ Karkman A., Do T.T., Walsh F., Virta M.P.J., (2018) Antibiotic-Resistance Genes in Waste Water, *Trends in Microbiology*, 26(3):220-228 <https://doi.org/10.1016/j.tim.2017.09.005>.
- ¹⁸ Zhang X.-X., Zhang T., Fang H.H.P., 2009. Antibiotic resistance genes in water environment. *Mini Review. Appl. Microbiol. Biotechnol.* 82:397–414, DOI 10.1007/s00253-008-1829-z
- ¹⁹ Berendonk, T.U., Manaia, C.M., Merlin, C., Fatta-Kassinos, D., Cytryn, E., Walsh, F., Bürgmann, H., Sørum, H., Norström, M., Pons, M. N., Kreuzinger, N., Huovinen, P., Stefani, S., Schwartz, T., Kisand, V., Baquero, F., & Martinez, J. L. (2015). Tackling antibiotic resistance: the environmental framework. *Nature reviews. Microbiology*, 13(5), 310–317. <https://doi.org/10.1038/nrmicro3439>
- ²⁰ Manaia, C., Aga, D., Cytryn, E., Gaze, W., Graham, D., Guo, J., Leonard, A., Liguán, L., Murray, A., Nunes, O., Rodríguez-Mozaz, S. Topp, E., Zhang, T. (2022). The Complex Interplay Between Antibiotic Resistance and Pharmaceutical and Personal Care Products in the Environment. *Environmental Toxicology and Chemistry*. 43. 10.1002/etc.5555. Mulchandani R, et al. *BMJ Public Health* 2025;3:e002411. doi: 10.1136/bmjph-2024-002411
- ²¹ Rogowska J, Gałęzowska G, Zimmermann A, 2024. Challenges and Current Trends in Preventing Antimicrobial Resistance in EU Water Law Context. *Antibiotics (Basel)*. 2024 Dec 31;14(1):18. doi: 10.3390/antibiotics14010018
- ²² USEPA 2021. Announcement of Final Regulatory Determinations for Contaminants on the Fourth Drinking Water Contaminant Candidate List. A Rule by the Environmental Protection Agency; Federal Register 40 CFR Part 141. [EPA-HQ-OW-2019-0583; FRL-10019-70-OW] RIN 2040-AF93 Pages 12272-12291. Currently available at: <https://www.federalregister.gov/documents/2021/03/03/2021-04184/announcement-of-final-regulatory-determinations-for-contaminants-on-the-fourth-drinking-water>
- ²³ WHO 2024. Guidance on wastewater and solid waste management for manufacturing of antibiotics. 3 September 2024. Currently available at: <https://www.who.int/publications/i/item/9789240097254> 12639.
- ²⁴ AMR 2025. Antibiotic Manufacturing Standard: Minimizing risk of developing antibiotic resistance and aquatic ecotoxicity in the environment resulting from the manufacturing of human antibiotics. Currently available at: <https://www.amrindustryalliance.org/antibiotic-manufacturing-standard/>
- ²⁵ UN 2024. Multi-stakeholder Hearing on Antimicrobial Resistance. Currently available at: <https://www.un.org/pga/78/multi-stakeholder-hearing-on-antimicrobial-resistance/>

²⁶ Kunhikannan S, Thomas CJ, Franks AE, Mahadevaiah S, Kumar S, Petrovski S. Environmental hotspots for antibiotic resistance genes. *Microbiologyopen*. 2021 Jun;10(3):e1197. doi: 10.1002/mbo3.1197. PMID: 34180594; PMCID: PMC8123917.

T.Boving, C.Hernández-Zepeda,
D.Lapworth, M.Lenczewski, B.P.Vellanki

IAH Groundwater Quality Commission

This series is designed both to inform professionals in other sectors of key interactions with groundwater resources and hydrogeological science, and to guide IAH members in their outreach to related sectors.