

Abiotic Oxygen and the Limits of Earth Analogy in Exoplanet Biosignature Interpretation

Alexandre Vaillancourt

tilex3r.av@gmail.com

Domain: Geochemistry · Atmospheric Evolution · Philosophy of Science · Exobiology

Working Paper — August 2026 NOT PEER REVIEWED

Abstract

The detection of atmospheric oxygen on a rocky exoplanet is widely regarded as a strong indicator of biological activity. This expectation derives from a single observational precedent: Earth. We argue that this extrapolation conflates two epistemologically distinct categories of knowledge — universal physical laws, which apply to any rocky planet under any conditions, and contingent terrestrial chemistry, which reflects the specific geochemical boundary conditions of one planet around one star. Through a process-by-process flux analysis of four independently documented abiotic O_2 production mechanisms on Archean Earth — UV-C photolysis coupled to hydrodynamic H_2 escape, SO_2 photodissociation (directly evidenced by the sulfur mass-independent fractionation record), silicate mechanochemistry, and terrestrial plasma electrolysis — we demonstrate that their combined production ($\sim 6 \times 10^{11} - 6 \times 10^{12} \text{ mol } O_2 \text{ yr}^{-1}$) is of comparable magnitude to the total Archean reductive sink capacity ($\sim 5.1 \times 10^{12} \text{ mol } O_2\text{-eq. yr}^{-1}$). This near-equivalence means that the boundary between abiotic and biological oxygen cannot be resolved from Earth's own geochemical record with current data — an epistemological result with direct consequences for exoplanet life detection. We propose a formal framework distinguishing physically universal constraints from geochemically contingent parameters, and argue that biosignature interpretation must be grounded in the former before observational data from JWST and future missions can be evaluated without circular reasoning.

1. Introduction

The modern biosignature framework rests on a sample of one. Every molecule proposed as a potential indicator of life beyond Earth — O_2 , O_3 , CH_4 , N_2O , dimethyl sulfide — derives its status from its role in Earth's biosphere. This is a necessary starting point, but it carries an implicit epistemological commitment that is rarely examined: the assumption that what biology does on Earth, it does in qualitatively similar ways elsewhere, under conditions that may differ by orders of magnitude in temperature, pressure, stellar flux, atmospheric composition, and geological history.

This paper examines that assumption through the specific case of atmospheric oxygen. O_2 has long been considered among the most compelling potential biosignatures, on the grounds that no known abiotic process produces it in quantities sufficient to dominate a planetary atmosphere. Recent work has significantly eroded this position: (Wordsworth & Pierrehumbert, 2014) demonstrated abiotic O_2 accumulation in CO_2 -rich atmospheres; (Krissansen-Totton et al., 2021) showed that waterworld planets can develop Earth-like O_2 inventories through hydrodynamic H_2 escape alone; and a growing body of experimental work documents abiotic O_2 production through mineral chemistry (Stone et al., 2022), (He et al., 2023) and SO_2 photodissociation (Farquhar et al., 2000), (Wogelius & Walther, 2022).

The response of the biosignature community to these findings has been to identify and catalog abiotic false-positive scenarios, then to propose contextual discriminants — additional observations intended to rule out abiotic explanations and confirm biological ones. This approach implicitly accepts terrestrial chemical parameters as universal reference values, and asks only how to navigate their ambiguities.

We propose a more fundamental critique: the terrestrial chemical framework is not universal. It is the product of a specific set of geochemical boundary conditions that obtained on one planet, around one star, with one particular tectonic regime, atmospheric composition, and oceanic chemistry. Applying its quantitative results to other planetary systems is not an extrapolation of physical laws — it is an extrapolation of contingent

geochemical outcomes. The distinction matters enormously for the interpretation of atmospheric data from JWST and future observatories.

This paper does not apply the proposed framework to a specific exoplanet target. This is deliberate. The quantitative parameters required to establish a physically universal abiotic baseline for any given planet — stellar UV flux, planetary mass, atmospheric composition, tectonic state — are not derivable from terrestrial analogy. They must be observed directly for each target. The present contribution establishes the analytical framework and its epistemological grounding; its application awaits observational data of sufficient quality from JWST and future missions.

This paper is organized as follows. Section 2 presents a rigorous quantitative analysis of abiotic O₂ production on Archean Earth, with explicit uncertainty bounds. Section 3 formalizes the distinction between universal physical constraints and contingent terrestrial chemistry. Section 4 develops the implications for exoplanet biosignature interpretation and proposes falsifiable predictions.

2. Abiotic O₂ Production on Archean Earth: A Constrained Flux Analysis

We analyze four abiotic O₂ production mechanisms with independent geochemical or experimental support, and present quantified flux ranges with explicit uncertainty bounds. Where a mechanism cannot be quantitatively constrained with current data, it is identified as such and excluded from the numerical balance, with the specific experimental gap noted.

2.1 Archean Boundary Conditions



A qualitative distinction from the modern system governs all flux estimates. In the modern atmosphere (21% O₂), any O* radical produced by photochemistry immediately encounters abundant ambient O₂ and recombines — the net yield is negligible. In the Archean anoxic atmosphere (<10⁻⁶ PAL O₂), O* has no recombination partner. The sign of the net O₂ flux from any dissociative process is therefore a function of ambient O₂ concentration — a qualitative difference in mechanism, not merely a quantitative scaling. Table 1 summarizes the boundary conditions that determine the magnitude of each production flux.

Table 1. Key Archean boundary conditions relevant to abiotic O₂ production (~3.0 Ga). PAL = present atmospheric level.

Parameter	Archean (~3.0 Ga)	Modern	Primary consequence for O ₂ yield
Atmospheric O ₂	<10 ⁻⁶ PAL	0.21 atm	No O* recombination with ambient O ₂
Atmospheric CO ₂	~0.1–0.3 bar	4×10 ⁻⁴ bar	Enhanced photodissociation substrate
Volcanic SO ₂	~10–100 ppmv	<1 ppbv	Continuous SO ₂ photolysis source
UV-C flux (100–280 nm)	~10× modern (no O ₃ shield)	~0 at surface	Direct photolysis of H ₂ O and SO ₂
EUV/X flux	3–10× modern (young Sun)	1×	Enhanced H ₂ ionization and escape
Tectonic activity	3–5× modern	1×	Continuous silicate fracturing
H ₂ escape rate	~10 ¹² –10 ¹³ mol yr ⁻¹	~4×10 ¹⁰ mol yr ⁻¹	Irreversible O ₂ accumulation ratchet

2.2 Mechanism P1: UV-C Photolysis of H₂O and Hydrodynamic H₂ Escape

In the absence of an ozone shield, UV-C radiation (100–280 nm) from the young Sun photodissociated atmospheric water vapor and CO₂ continuously. The governing reactions are:



UV photolysis and hydrodynamic H₂ escape are treated as a single integrated flux. The H₂ escape rate provides the most direct and best-constrained quantitative handle on net irreversible O₂ production: each mole of H₂ lost to space renders $\frac{1}{2}$ mol of O₂ permanently non-recombinant (Eq. 2.3). (Catling et al., 2001) estimated the pre-GOE H₂ escape flux at $\sim 10^{12}$ – 10^{13} mol H₂ yr⁻¹ — a value independently supported by xenon isotope fractionation records, which document a sustained Archean H₂ escape rate 2–3 orders of magnitude above the post-GOE value of $\sim 4 \times 10^{10}$ mol H₂ yr⁻¹ (Zahnle et al., 2019):

$$P_1 = \frac{1}{2} \cdot F(\text{H}_2) [\text{mol O}_2 \text{ yr}^{-1}] \quad (2.3)$$





$$\text{Lower bound: } F(\text{H}_2)_{\text{escape}} = 10^{12} \text{ mol yr}^{-1} \rightarrow P_1 = 5 \times 10^{11} \text{ mol O}_2 \text{ yr}^{-1}$$

$$\text{Upper bound: } F(\text{H}_2)_{\text{escape}} = 10^{13} \text{ mol yr}^{-1} \rightarrow P_1 = 5 \times 10^{12} \text{ mol O}_2 \text{ yr}^{-1}$$

P₁ estimate: $\sim 5 \times 10^{11}$ – 5×10^{12} mol O₂ yr⁻¹. This is the best-constrained single production mechanism, anchored to independent isotopic evidence. Cumulative total: $\sim 5 \times 10^{11}$ – 5×10^{12} mol O₂ yr⁻¹.

2.3 Mechanism P2: SO₂ Photodissociation — The MIF-S Constraint

This mechanism holds a unique evidential status: it is the only abiotic O₂ production pathway with a direct, continuously preserved geochemical signature in the rock record. The sulfur mass-independent fractionation (MIF-S) signal — anomalous $\Delta^{33}\text{S}$ and $\Delta^{36}\text{S}$ values in sediments older than ~ 2.4 Ga — was established by (Farquhar et al., 2000) as the product of UV photolysis of volcanic SO₂ in an anoxic, ozone-free atmosphere. (Pavlov & Kasting, 2002) constrained atmospheric O₂ at $< 10^{-5}$ PAL from MIF-S preservation requirements, simultaneously confirming the anoxic condition and the continuous SO₂ photolysis flux. The disappearance of MIF-S at ~ 2.4 Ga marks the GOE with sub-million-year precision.



The elemental sulfur produced (Eqs. 2.5–2.6) polymerizes into S₈ aerosols and deposits — it exits the cycle permanently. The O produced by Eqs. 2.4–2.5 has no ambient O₂ to recombine with, yielding a net positive O₂ flux per mole of SO₂ photolyzed. An ionic pathway for direct O₂ production from SO₂ double photoionization has been additionally documented (Wogelius & Walther, 2022). The volcanic SO₂ flux to the Archean atmosphere is estimated at ~ 1 – 10 Tmol SO₂ yr⁻¹ (Catling & Kasting, 2017). Applying a photolysis efficiency ϵ of 1–10% — consistent with Archean UV-C flux and SO₂ column density estimates — yields:

$$P_2 = \epsilon \cdot F(\text{SO}_2) [\text{mol O}_2 \text{ yr}^{-1}] \quad (2.8)$$




$$\text{Lower bound: } \epsilon = 0.01, F(\text{SO}_2) = 10^{12} \text{ mol yr}^{-1} \rightarrow P_2 = 10^{10} \text{ mol O}_2 \text{ yr}^{-1}$$

$$\text{Upper bound: } \epsilon = 0.10, F(\text{SO}_2) = 10^{13} \text{ mol yr}^{-1} \rightarrow P_2 = 10^{12} \text{ mol O}_2 \text{ yr}^{-1}$$

P₂ estimate: $\sim 10^{10}$ – 10^{12} mol O₂ yr⁻¹. The MIF-S record provides direct observational evidence that this mechanism operated continuously throughout the Archean. The primary uncertainty is the photolysis efficiency ϵ , which requires experimental determination under Archean UV-C intensity and SO₂ column density conditions. Cumulative total: $\sim 5 \times 10^{11}$ – 6×10^{12} mol O₂ yr⁻¹.

2.4 Mechanism P3: Silicate Mechanochemistry

(Stone et al., 2022) demonstrated experimentally that fault movements in hot underground fractures produce H₂O₂ and molecular O₂ under entirely anoxic conditions. (He et al., 2023) confirmed this mineral-based pathway through systematic quartz abrasion experiments. The governing reactions are:



(Zhang et al., 2023) estimated a global Mesoarchean ROS production flux of $\sim 3.9\text{--}6.7 \times 10^8$ mol oxidants yr^{-1} from continental silicate weathering. Extending to the oceanic ultramafic crust — which represents 3–5× the continental surface area and was continuously renewed at 3–5× modern tectonic rates — and applying the stoichiometric conversion of Eq. 2.11:

Lower bound: Continental estimate alone (Zhang et al., 2023): $\sim 10^8$ mol O_2 yr^{-1}

Upper bound: Extended to oceanic crust at 5× tectonic activity: $\sim 10^{10}$ mol O_2 yr^{-1}

P₃ estimate: $\sim 10^8\text{--}10^{10}$ mol O_2 yr^{-1} . This mechanism is individually minor in total flux but carries a distinct biological significance: the H_2O_2 produced in turbulent subaqueous environments was directly available to proto-photosynthetic microorganisms, a point developed in Section 3.3. Cumulative total: $\sim 6 \times 10^{11}\text{--}6 \times 10^{12}$ mol O_2 yr^{-1} .

2.5 Mechanism P4: Terrestrial Lightning — An Unresolved Contribution

Electrical discharges in an anoxic $\text{CO}_2/\text{H}_2\text{O}$ atmosphere constitute a physically plausible additional source of O^* . We exclude this mechanism from the quantitative analysis for two reasons.



First, the available yield data do not apply to natural lightning. The most widely cited NO_x yield per flash (Schumann & Huntrieser, 2007) derives from controlled laboratory plasma discharge experiments. The kinetics of a natural lightning channel — involving a return stroke at $\sim 30,000$ K followed by millisecond-timescale cooling and complex channel chemistry — are not reproduced in laboratory plasma chambers. Laboratory yields cannot be directly applied to natural lightning.

Second, lightning in a nitrogen-bearing atmosphere functions primarily as a nitric acid generator. Electrical discharges dissociate N_2 and O_2 to produce NO , which is rapidly oxidized to NO_2 and then to HNO_3 via reaction with H_2O . The net O_2 yield per flash is effectively zero. In an Archean atmosphere dominated by CO_2 and H_2O , the nitrogen fixation pathway is absent, but the plasma kinetics of $\text{CO}_2/\text{H}_2\text{O}$ dissociation under natural lightning discharge conditions have not been experimentally characterized. The O^* yield per flash cannot be estimated by analogy without introducing an unquantified systematic error.

The net contribution of terrestrial lightning to Archean O_2 production is therefore currently unresolvable. Its quantitative characterization — specifically, laboratory measurement of O_2 yield per joule of discharge energy in $\text{CO}_2/\text{H}_2\text{O}$ mixtures at Archean atmospheric pressures — is identified as a research priority.

2.6 Reductive Sink Capacity and Net Flux Balance

Table 2 presents the principal Archean reductive sink fluxes. A critical methodological point governs the dominant sink: the oceanic Fe^{2+} system was not a static depleting reservoir but a dynamically renewed flux, sustained by continuous hydrothermal input from mid-ocean ridge and off-axis vent systems. (Thompson et al., 2019) calculated that hydrothermal and weathering Fe^{2+} inputs were sufficient to sustain peak banded iron formation (BIF) deposition at ~ 4.5 Tmol Fe yr^{-1} . The O_2 demand of this dynamic sink is calculated from stoichiometry:



Applying Eq. 2.12 to the peak Fe^{2+} flux of 4.5 Tmol Fe yr^{-1} yields an O_2 demand of $\sim 1.1 \times 10^{12}$ mol O_2 yr^{-1} .

Table 2. Archean reductive sink fluxes and O_2 -equivalent demand ($\sim 3.0\text{--}2.4$ Ga).

Sink mechanism	Governing equation	O_2 -eq. flux (mol yr^{-1})	Source
Fe^{2+} oceanic oxidation (dynamic BIF flux)	Equations 2.12–2.13	$\sim 1.1 \times 10^{12}$	(Thompson et al., 2019)
Serpentinization of	$\text{Fe}^{2+} + \text{H}_2\text{O} \rightarrow \text{Fe}^{3+} + \text{H}_2$	$\sim 2.0 \times 10^{12}$	(Meier et al., 2021)

komatiites			
Volcanic reductant gases (H ₂ S, SO ₂ , H ₂)	$\text{H}_2\text{S} + \frac{3}{2} \text{O}_2 \rightarrow \text{SO}_2 + \text{H}_2\text{O}$	$\sim 1.0 \times 10^{12}$	(Kump, 2001)
Crustal FeS ₂ weathering	$\text{FeS}_2 + \frac{7}{4} \text{O}_2 + \text{H}_2\text{O} \rightarrow \text{Fe}^{2+} + 2\text{SO}_4^{2-}$	$\sim 0.5 \times 10^{12}$	(Kasting, 2013)
Atmospheric CH ₄ oxidation	$\text{CH}_4 + 2\text{O}_2 \rightarrow \text{CO}_2 + 2\text{H}_2\text{O}$	$\sim 0.5 \times 10^{12}$	(Kharecha et al., 2005)
Total sink capacity		$\sim 5.1 \times 10^{12}$	

Table 3. Abiotic O₂ reductive sink fluxes by mechanism.

Mechanism	Lower bound (mol O ₂ yr ⁻¹)	Upper bound (mol O ₂ yr ⁻¹)	Evidential basis
P1 — UV photolysis + H ₂ escape	$\sim 5 \times 10^{11}$	$\sim 5 \times 10^{12}$	Xe isotope fractionation; H ₂ escape models
P2 — SO ₂ photodissociation (MIF-S)	$\sim 10^{10}$	$\sim 10^{12}$	Archean sedimentary MIF-S record
P3 — Silicate mechanochemistry	$\sim 10^8$	$\sim 10^{10}$	Experimental (Stone et al., 2022)
P4 — Terrestrial lightning	—	—	Unresolved — plasma kinetics uncharacterized
Total abiotic (P1+P2+P3)	$\sim 6 \times 10^{11}$	$\sim 6 \times 10^{12}$	
Total sink capacity	$\sim 5.1 \times 10^{12}$	$\sim 5.1 \times 10^{12}$	
Net O ₂ flux	-4.5×10^{12} (deficit)	$+9 \times 10^{11}$ (marginal surplus)	

The result of Table 3 must be stated with precision: the combined abiotic O₂ production from three quantitatively constrained mechanisms brackets the total Archean reductive sink capacity. The lower bound yields a deficit; the upper bound yields a marginal surplus. Neither is demonstrably correct with current data. This near-equivalence is the central empirical finding of this paper — not a quantitative failure, but an epistemological result whose implications are developed in Section 3.

3. Universal Physical Laws versus Contingent Geochemistry: A Necessary Distinction

3.1 A Category Error in the Biosignature Literature

The result of Section 2 — that abiotic O₂ production on Earth cannot be cleanly distinguished from biological production using currently available geochemical data — reveals a structural problem in the biosignature literature. Current frameworks implicitly treat two distinct categories of knowledge as equivalent:

Physically universal constraints are statements that follow from physical laws applicable to any rocky planet: the Hoyle resonance ($^{12}\text{C} + ^4\text{He} \rightarrow ^{16}\text{O} + \gamma$, $E_{\text{res}} = 7.65 \text{ MeV}$) guarantees oxygen-rich lithospheres wherever stars produce heavy elements; UV photolysis of H₂O is thermodynamically inevitable on any water-bearing planet orbiting a UV-active star without an ozone shield; hydrodynamic H₂ escape is a function of planetary mass and temperature, independent of atmospheric composition. These constraints apply universally — to any rocky planet we observe.

Contingent geochemical parameters are quantities that reflect the specific combination of boundary conditions that obtained on Earth: a particular atmospheric CO₂ and SO₂ content, a particular tectonic renewal

rate, a particular oceanic Fe^{2+} concentration, a particular volcanic reductant flux. The quantitative flux estimates of Section 2 — both production rates and sink capacities — belong to this category. They are not derivable from physical laws alone; they require the specific Archean boundary conditions of Table 1, themselves only partially constrained.

This distinction is not terminological. The quantitative result of Section 2 — that abiotic and biological O_2 fluxes bracket the same order of magnitude on the only inhabited planet accessible to direct study — makes the distinction empirically consequential. If the boundary cannot be resolved on Earth with Earth's own data, it cannot be resolved on a planet observed at interstellar distances by inference from Earth's geochemical parameters. The category error consists in applying contingent geochemical parameters — derived from one planet under one set of historical conditions — to other planetary systems as though they constituted universal physical law.

3.2 The $n = 1$ Problem and Its Epistemological Consequences

Exobiology operates with a sample of one inhabited planet. The implications of this constraint have not been drawn fully. From Earth alone, we cannot distinguish between:

(a) Universal properties of life: biology, wherever it exists, produces O_2 as a thermodynamic consequence of oxygenic photosynthesis, in quantities that systematically exceed abiotic production under any planetary boundary conditions.

(b) Contingent properties of life on Earth: Earth biology produces O_2 in the observed quantities because of the specific evolutionary trajectory of cyanobacteria under Archean geochemical conditions — conditions that may not be replicated on any other known planet.

The analysis of Section 2 demonstrates that this distinction cannot be resolved from Earth's geochemical record: abiotic and biological contributions to Archean O_2 are of comparable magnitude and cannot be cleanly separated with current data. If we cannot resolve this on the only inhabited planet accessible to direct study, we cannot resolve it on a planet observed at interstellar distances through a telescope.

3.3 A Causal Implication: Abiotic Oxidants as Selective Pressure

There is a further consequence of this analysis that deserves recognition. If abiotic O_2 and H_2O_2 production is a physically universal consequence of UV irradiation and silicate chemistry on wet rocky planets, it may also constitute a universal selective pressure driving the evolution of oxidant-utilizing metabolisms. Hao et al. (2021, *Nature Communications*) demonstrated experimentally that abiotic H_2O_2 and O_2 from silicate abrasion in turbulent subaqueous environments created the oxidant gradients that selectively disadvantaged anaerobic microorganisms — concluding that this abiotic supply "created an evolutionary impetus for the origin of oxygenic photosynthesis." Under this interpretation, cyanobacteria did not invent O_2 production from scratch in a fully reduced environment; they evolved to exploit an oxidant gradient that already existed.

This reframes the relationship between abiotic geochemistry and biology. Rather than treating biological O_2 production as a discrete event that superseded an abiotic baseline, the evidence suggests a continuum: abiotic oxidants → selective pressure → biological amplification → atmospheric oxygenation. The GOE, under this framework, represents not the origin of O_2 production but the point at which the combined abiotic and biological flux permanently exceeded the renewal capacity of geochemical reductive sinks.

4. Implications for Exoplanet Biosignature Interpretation

4.1 Rebuilding the Framework on Universal Constraints

A biosignature framework grounded in physically universal constraints rather than contingent terrestrial chemistry would have the following properties:

- Universal baseline first. For any target planet, the first step in atmospheric interpretation is to establish what O_2 and related gases would be expected from photolysis, escape, and mineral chemistry alone — given the directly observable parameters of that planet: stellar UV flux, planetary mass and escape velocity, atmospheric composition, and estimated tectonic state. This baseline will differ for every target and cannot be assumed from terrestrial analogy.

- Biological signal as departure from universal baseline. The diagnostic signature of biology is not O_2 above an absolute threshold derived from Earth, but an atmospheric state that cannot be explained by the universal baseline alone — a departure that requires an additional source. The classical $CH_4 + O_2$ disequilibrium biosignature (Krissansen-Totton, Olson & Catling, 2018) approximates this principle, though its application requires that the universal baseline be established individually for each target planet.
- Observational context as prerequisite, not supplement. Stellar UV flux, planetary mass, tectonic indicators, and atmospheric composition are not optional contextual supplements to biosignature detection; they are the inputs required to compute the universal baseline. A detection without this context is not a weak detection — it is an uninterpretable signal.

4.2 Testable Predictions

Prediction 1. Any tectonically active rocky planet with liquid water, a CO_2/SO_2 -bearing atmosphere, and a host star with sufficient UV-C output will develop a measurable abiotic O_2 signal, independently of biological activity. The magnitude of this signal will scale with planetary age, tectonic rate, stellar UV flux, and atmospheric composition. This prediction is falsifiable with sufficient spectral sensitivity from JWST or its successors.

Note on quantitative thresholds: The absence of precise numerical thresholds in Prediction 1 is not a weakness of the framework — it is a direct consequence of its central argument. The scaling parameters governing abiotic O_2 production (stellar UV flux, tectonic rate, atmospheric SO_2 abundance, planetary escape velocity) are not derivable from terrestrial analogy; they must be measured directly for each target planet. Assigning a universal threshold derived from Earth's boundary conditions would reproduce the category error this framework is designed to correct. The prediction is falsifiable in principle; its quantitative form requires planet-specific observational inputs that are unavailable prior to observation.

Prediction 2. The presence of sulfur mass-independent fractionation (MIF-S) in a planetary atmosphere — detectable in principle through S isotopologue spectroscopy — would constitute direct evidence of ongoing SO_2 photolysis in an anoxic environment. Its co-detection with O_2 would confirm abiotic O_2 production without biological amplification. Its disappearance from a planet's spectral record would mark an analog of the GOE — the onset of sustained atmospheric oxygenation.

Prediction 3 (falsification). A rocky, water-bearing planet with confirmed active tectonics, a CO_2/SO_2 atmosphere, a UV-active host star, and an age exceeding 3 Ga that shows no detectable O_2 or O_3 would require revision of the universal physical baseline presented here. Such an observation would imply either an unexpectedly efficient sink system or boundary conditions outside the range of current models.

5. Limitations and Research Priorities

Flux uncertainty. The flux estimates of Section 2 span one to two orders of magnitude. This uncertainty is genuine and reflects the incomplete constraint of Archean boundary conditions. Narrowing these bounds — particularly the SO_2 photolysis efficiency ϵ and the tectonic scaling of P3 — requires targeted experimental work under Archean UV and pressure conditions.

Lightning plasma kinetics. The O^* yield of natural lightning discharge in CO_2/H_2O mixtures at Archean atmospheric pressures has not been experimentally measured. Laboratory determination of this parameter would directly constrain P4 and potentially alter the quantitative balance of Table 3.

Operational classification of universal vs. contingent parameters. The distinction introduced in Section 3 requires a systematic classification of existing biosignature components — identifying which are grounded in physically universal constraints and which in contingent terrestrial chemistry. This classification exercise is beyond the scope of the present paper and constitutes a necessary foundational contribution to the field.

Selective pressure hypothesis. (Hao et al., 2021) demonstrate that abiotic oxidants could have created selective pressure for the evolution of oxygenic photosynthesis; they do not demonstrate that this pressure was necessary. Whether oxygenic photosynthesis could have evolved in a fully anoxic environment remains an open question.

These limitations do not invalidate the central argument. They define the research agenda required to place it on a fully quantitative foundation.

6. Conclusion

Atmospheric oxygen has long been treated as a strong indicator of biological activity on exoplanets, on the grounds that no abiotic process produces it in sufficient quantities to dominate a planetary atmosphere. This paper has challenged that position on two counts.

First, a pole-by-pole flux analysis of three quantitatively constrained abiotic O₂ production mechanisms on Archean Earth — UV-C photolysis coupled to hydrodynamic H₂ escape, SO₂ photodissociation directly evidenced by the sulfur mass-independent fractionation record, and silicate mechanochemistry — yields a combined production of $\sim 6 \times 10^{11} - 6 \times 10^{12} \text{ mol O}_2 \text{ yr}^{-1}$. This brackets the total documented Archean reductive sink capacity of $\sim 5.1 \times 10^{12} \text{ mol O}_2\text{-eq. yr}^{-1}$. The mathematical result is:

$$\sum_i P_i \sim 6 \times 10^{11} - 6 \times 10^{12} \text{ mol O}_2 \text{ yr}^{-1} \quad \sum_j S_j \sim 5.1 \times 10^{12} \text{ mol O}_2\text{-eq. yr}^{-1} \quad (6.1)$$

The near-equivalence of Eq. 6.1 means that the boundary between abiotic and biological oxygen cannot be resolved from Earth's own geochemical record with current data. This is not a quantitative failure — it is an epistemological result. It demonstrates that even for the only inhabited planet accessible to direct study, we cannot cleanly separate the abiotic and biological contributions to atmospheric O₂.

Second, this result exposes a structural category error in current biosignature frameworks: the conflation of physically universal constraints — which follow from physical laws applicable to any rocky planet — with contingent geochemical parameters, which reflect the specific boundary conditions of one planet under one set of historical conditions. Applying the quantitative outputs of terrestrial geochemistry to other planetary systems as though they constituted universal physical law introduces an unquantified systematic bias into life detection.

The implication for exoplanet observation is direct: the interpretation of any atmospheric signal as a biosignature requires that a physically universal abiotic baseline be established first — derived from the directly observable parameters of the target planet, not assumed from terrestrial analogy. Before that baseline is established, a detection is not evidence of life. It is an unresolved signal.

This paper does not claim that life is rare or that O₂ is uninformative. It claims that the framework currently used to interpret O₂ as a biosignature has been built on a foundation that conflates the universal and the contingent — and that correcting this conflation is a necessary condition for meaningful life detection in the era of JWST and beyond.

Acknowledgements

The author thanks Claude.ai (Anthropic) and Gemini (Google DeepMind) for their contribution to the clarification of certain conceptual ideas and editorial review. All scientific claims, interpretations, and conclusions are the sole responsibility of the author.

- Catling, D.C., Zahnle, K.J., & McKay, C.P. (2001). Biogenic methane, hydrogen escape, and the irreversible oxidation of early Earth. *Science*, 293(5531), 839–843. <https://doi.org/10.1126/science.1061976>
- Catling, D.C., & Kasting, J.F. (2017). *Atmospheric Evolution on Inhabited and Lifeless Worlds*. Cambridge University Press. <https://doi.org/10.1017/9781139020558>
- Farquhar, J., Bao, H., & Thiemens, M.H. (2000). Atmospheric influence of Earth's earliest sulfur cycle. *Science*, 289(5480), 756–758. <https://doi.org/10.1126/science.289.5480.756>
- Hao, J., Knoll, A.H., Huang, F., Schieber, J., Hazen, R.M., & Daniel, I. (2021). An abiotic source of Archean hydrogen peroxide and oxygen that pre-dates oxygenic photosynthesis. *Nature Communications*, 12, 6611. <https://doi.org/10.1038/s41467-021-26916-2>
- He, H., Wu, X., Zhu, J., et al. (2023). A mineral-based origin of Earth's initial hydrogen peroxide and molecular oxygen. *Proceedings of the National Academy of Sciences*, 120(13), e2221984120. <https://doi.org/10.1073/pnas.2221984120>
- Hoyle, F. 1954, *ApJS*, 1, 121. <https://doi.org/10.1086/190005> *Astrophysical Journal Supplement Series*, volume 1, page 121, 1954.
- Kasting, J.F. (2013). What caused the rise of atmospheric O₂? *Chemical Geology*, 362, 13–25. <https://doi.org/10.1016/j.chemgeo.2013.05.039>
- Kharecha, P., Kasting, J.F., & Siefert, J.L. (2005). A coupled atmosphere–ecosystem model of the early Archean Earth. *Geobiology*, 3(2), 53–76. <https://doi.org/10.1111/j.1472-4669.2005.00049.x>
- Krissansen-Totton, J., Fortney, J.J., Nimmo, F., & Wogan, N. (2021). Oxygen false positives on habitable zone planets around sun-like stars. *AGU Advances*, 2, e2020AV000294. <https://doi.org/10.1029/2020AV000294>
- Krissansen-Totton, J., Olson, S., & Catling, D.C. (2018). Disequilibrium biosignatures over Earth history and implications for detecting exoplanet life. *Science Advances*, 4(1), eaao5747. <https://doi.org/10.1126/sciadv.aao5747>
- Kump, L.R. (2001). Rise of atmospheric oxygen and the "upside-down" Archean mantle. *Geochemistry, Geophysics, Geosystems*, 2(1). <https://doi.org/10.1029/2000GC000114>
- Meier, D.V.P., et al. (2021). Decreasing extents of Archean serpentinization contributed to the rise of an oxidized atmosphere. *Nature Communications*, 12, 7341. <https://doi.org/10.1038/s41467-021-27541-9>
- Pavlov, A.A., & Kasting, J.F. (2002). Mass-independent fractionation of sulfur isotopes in Archean sediments: strong evidence for an anoxic Archean atmosphere. *Astrobiology*, 2(1), 27–41. <https://doi.org/10.1089/153110702753621321>
- Schumann, U., & Huntrieser, H. (2007). The global lightning-induced nitrogen oxides source. *Atmospheric Chemistry and Physics*, 7, 3823–3907. <https://doi.org/10.5194/acp-7-3823-2007>
- Stone, J., Edgar, J.O., Gould, J.A., & Telling, J. (2022). Tectonically driven oxidant production in the hot biosphere. *Nature Communications*, 13, 4529. <https://doi.org/10.1038/s41467-022-32129-y>
- Thompson, K.J., et al. (2019). Photoferrotrophy, deposition of banded iron formations, and methane production in Archean oceans. *Science Advances*, 5(11), eaav2869. <https://doi.org/10.1126/sciadv.aav2869>
- Wogelius, R.A., & Walther, J.V. (2022). Abiotic molecular oxygen production — ionic pathway from sulfur dioxide. *Science Advances*, 8, abq5411. <https://doi.org/10.1126/sciadv.abq5411>
- Wordsworth, R., & Pierrehumbert, R. (2014). Abiotic oxygen-dominated atmospheres on terrestrial habitable zone planets. *The Astrophysical Journal Letters*, 785, L20. <https://doi.org/10.1088/2041-8205/785/2/L20>
- Zahnle, K.J., Gacesa, M., & Catling, D.C. (2019). Strange messenger: A new history of hydrogen on Earth, as told by Xenon. *Geochimica et Cosmochimica Acta*, 244, 56–85. <https://doi.org/10.1016/j.gca.2018.09.017>
- Zhang, S., et al. (2023). Geodynamic oxidation of Archean terrestrial surfaces. *Communications Earth & Environment*, 4, 179. <https://doi.org/10.1038/s43247-023-00789-3>