

## Acknowledgements

The author would like to thank Dr. Tarmo Remmel for his helpful comments and discussions regarding this work.

## Bibliography

- R. S. Bivand (2000) Using the R statistical data analysis language on GRASS 5.0 GIS database files. *Computers & Geosciences* 26: 1043-1052.
- A. Brimicombe (2003) *GIS, environmental modelling and engineering*. New York, USA: Taylor & Francis
- J. M. Chen, J. Liu, J. Chilar, M.L. Goulden (1999) Daily canopy photosynthesis model through temporal and spatial scaling for remote sensing applications. *Ecological Modelling* 124: 99-119.
- G. D. Farquhar, S. von Caemmerer, J.A. Berry (1980) A biochemical model of photosynthetic CO<sub>2</sub> assimilation in leaves of C<sub>3</sub> species. *Planta* 149: 78-90.
- P. G. Jarvis (1976) The interpretation of the variations in leaf water potential and stomatal conductance found in canopies in the field. *Philosophical Transactions Of The Royal Society Of London*: 593-610.
- J. Liu, J. M. Chen, J. Chilar, W. M. Park (1997) A process-based Boreal Ecosystem Productivity Simulator using remote sensing inputs. *Remote Sensing of Environment* 62: 158-175.
- J. Liu, J. M. Chen, J. Chilar, W. Chen (1999) Net primary productivity distribution in the BOREAS region from a process model using satellite and surface data. *Journal of Geophysical Research-Atmospheres* 104: 27735-27754.
- J. Liu, J. M. Chen, J. Chilar, W. Chen (2002) Net primary productivity mapped for Canada at 1-km resolution. *Global Ecology and Biogeography* 11: 115-129.
- J. L. Monteith (1965) Evaporation and environment. in *Proceedings of the 19th Symposium of the Society for Experimental Biology* New York, USA: Cambridge University Press.
- S. W. Running, J. C. Coughlan (1988) A general model of forest ecosystem processes for regional applications. 1. Hydrologic balance, canopy gas-exchange and primary production processes. *Ecological Modelling* 42: 125-154.
- M. S. Wigmosta, L. W. Vail, D. P. Lettenmaier (1994) A distributed hydrology-vegetation model for complex terrain. *Water Resources Research* 30: 1665-1679.

Oliver Sonnentag  
 Department of Geography  
 University of Toronto  
 Toronto, ON, Canada  
[oliver.sonnentag \(at\) gmail.com](mailto:oliver.sonnentag@gmail.com)

# r.roughness – a new tool for morphometric analysis in GRASS

by Carlos Henrique Grohmann

## Introduction

This article briefly describes `r.roughness`, a shell script written to calculate the surface roughness of raster surfaces. The method is based on Hobson (1972), where roughness is defined as the ratio between surface and plan area of square cells. This script will create square sub-regions with size defined by the user; in each sub-region, the real and planar areas will be calculated by `r.surf.area`, and the results (points at the centre of sub-regions) will be interpolated with `v.surf.rst`. The user also can set the tension and smooth parameters of interpolation.

## Surface Roughness

Surface roughness (or topographical roughness) was first introduced as a morphometric parameter by

Stone and Dugundji (1965) and Hobson (1967, 1972). To Hobson (1972), one possible way to calculate it is the ratio between surface (real) area and flat (plan) area of square cells; in this approach, flat surfaces would present values close to 1, whilst in irregular ones the ratio shows a curvilinear relationship which asymptotically approaches infinity as the real areas increases.

Day (1979) describes surface roughness as the expression of non-systematic variability of the topographic surface, and used the dispersion of vector normals to surface plans as a roughness indicator to discriminate tropical karst stiles.

Ferrari et al. (1998) argue that surfaces with distinct characteristics can present the same roughness value, due the existence of interactions between the number and magnitude of terrain irregularities. Grohmann (2004), considers this method useful for morphological characterisation since it is mainly related with the shape of land-forms and not its elevation; thus, tectonically tilted areas have their expression shown, while it could be masked in a hypsometric map, as consequence of altimetric variations.




Figure 1: Study area.




Figure 2: Surface roughness map, grid=500m.

The method has been applied in studies related to morphology of lake bottoms (Hakanson, 1974), as a discriminant of karstic areas (Day, 1979; Karmann et al., 1996; Ferrari et al., 1998), for structural compartmentation of sedimentary basins basement (Grohmann et al., 2005), in morphometric analysis of alkaline massifs (Roldan et al., 2006; Grohmann et al., 2007), in structural analysis of strike-slip shear zones (Steiner et al., 2006) and for macro-geomorphological compartmentation (Grohmann & Riccomini, 2006).

## Usage and Examples

The script has five options: `map`, `grid`, `rough`, `tension` and `smooth`. `map` stands for the input raster surface and is the only required option. `grid` is the size of the sub-regions in which roughness will be calculated; the default value is 1000m. `rough` is the name for the output map; if a name is not provided, it will be set to `input_map_name.roughness.grid_size`. `tension` and `smooth` will be used by `v.surf.rst` for interpolation of roughness values; the default values are `tension=40` and `smooth=0.1`.

The examples presented are from an area located in southeastern Brazil, southern region of São Paulo State. Local geology consists of NE-SW trending metapelitic and metacalcareous rocks where karstic landscapes developed over the carbonatic rocks, with altimetric differences up to 700m between non-carbonatic (pelitic, psamitic and granitic) crests and karstic valley bottoms. NW-SE trending dikes cut across the area and have a strong influence on geomorphological development (Fig. 1). Surface roughness was calculated for grid sizes of 500, 1000 and

2000m.

With a grid size of 500m (Fig. 2), a good correlation with land-forms can be seen. Higher roughness values are related with the Bethary River valley, developed over a NW-SE trending dike. Also, karstic valleys have smaller roughness values than non-carbonatic crests. The general picture of the features present in (Fig. 2) can be seen in the map for grid size of 1000m (Fig. 3), although is not possible to individualise the answer from each carbonatic unit. A grid of 2000m (Fig. 4) does not give much information, indicating that land-forms within this area cannot be well described with a wavelength this large.

## Concluding Remarks

Surface roughness is a useful parameter for morphological compartmentation. `r.roughness` is a shell script that automises the process, but users must be aware that it uses `r.surf.area` to calculate both real and planar area for each grid cell (sub-regions) and that raster resolution plays an important role on area estimations.

The script is available through GRASS Wiki site <sup>3</sup> in two versions: `r.roughness` for GRASS 6.1+ and `r.roughness60` for GRASS 6.0.x.

## Acknowledgements

This work was supported by FAPESP grant 04/06260-5 (C.H.G. Doctor's Degree Fellowship). The author is thankful to Tomás Senabre (Spain) for his feedback which helped to improve the script usability.

<sup>3</sup>[http://grass.gdf-hannover.de/wiki/GRASS\\_AddOns](http://grass.gdf-hannover.de/wiki/GRASS_AddOns)




Figure 3: Surface roughness map, grid=1000m.




Figure 4: Surface roughness map, grid=2000m.



## Bibliography

- M. J. Day (1979) Surface roughness as a discriminator of tropical karst styles *Zeitschrift für Geomorphologie*, Suppl.-Bd.32:1-8.
- J. A. Ferrari, S. T. Hiruma, I. Karmann (1998) Caracterização morfométrica de uma superfície cárstica do Vale do Ribeira, São Paulo (Núcleo Caboclos - PETAR) *Revista do Instituto Geológico*, 19:9-17.
- C. H. Grohmann (2004) Morphometric analysis in Geographic Information Systems: applications of free software GRASS and R *Computers & Geosciences*, 30:1055-1067. <http://dx.doi.org/10.1016/j.cageo.2004.08.002>
- C. H. Grohmann, C. Riccomini, F. Paula e Silva and H. K. Chang (2005) Compartimentação tectônica da Bacia Bauru no centro-oeste do Estado de São Paulo *In: X Simpósio Nacional de Estudos Tectônicos / IV International Symposium on Tectonics, Curitiba-PR*, Boletim de Resumos Expandidos:25-27.
- C. H. Grohmann and C. Riccomini (2006) Análise regional de rugosidade de relevo *In: XLIII Congresso Brasileiro de Geologia, Aracaju-SE* Anais:97.
- C. H. Grohmann, C. Riccomini and F. M. Alves (2007) SRTM-based morphotectonic analysis of the Poços de Caldas Alkaline Massif, southeastern Brazil *Computers & Geosciences (in press)*. <http://dx.doi.org/10.1016/j.cageo.2006.05.002>
- L. Hakanson (1974) A mathematical model for establishing numerical values of topographical roughness for lake bottoms *Geografiska Annaler. Series A, Physical Geography*, 56:183-200.
- R. D. Hobson (1967) FORTRAN IV programs to determine the surface roughness in topography for the CDC 3400 computer *Computer Contribution 14, State Geol. Survey Kansas*, 28p.
- R. D. Hobson (1972) Surface roughness in topography: quantitative approach *In: Chorley, R.J., 1972. Spatial analysis in geomorphology*, 225-245.
- I. Karmann, R. F. Pereira and J. A. Ferrari (1996) Índice de rugosidade: parâmetro morfométrico da intensidade de relevo. Exemplo do carste da bacia do Rio Una, Bahia *In: XXXIX Congresso Brasileiro de Geologia, Salvador*, Anais:575-579.
- L. F. Roldan, S. S. Steiner, C. H. Grohmann and R. Machado (2006) Análise morfométrica da região do Domo de Lages, SC *In: XLIII Congresso Brasileiro de Geologia, Aracaju-SE* Anais:266.
- S. S. Steiner, R. Machado and C. H. Grohmann, C. H. (2006) Análise morfométrica da estrutura divergente na região do Rio Paraíba do Sul, Rio de Janeiro *In: XLIII Congresso Brasileiro de Geologia, Aracaju-SE* Anais:97.
- R. O. Stone and J. Dugundji (1965) A study of microrelief – its mapping, classification and quantification by means of a Fourier analysis *Engineering Geology*, 1:89-187.

Carlos Henrique Grohmann  
Institute of Geosciences  
University of São Paulo, Brazil  
<http://www.igc.usp.br>  
guano AT usp br