

How EU policies could reduce nutrient pollution in European inland and coastal waters?

Grizzetti B.^{*1}, Vigiak O.¹, Udias A.¹, Aloe A.², Zanni M.², Bouraoui F.¹, Pistocchi A.¹, Dorati C.², Friedland R.¹, De Roo A.¹, Benitez Sanz C.³, Leip A.¹, Bielza M.⁴

¹European Commission Joint Research Centre (JRC), Ispra (VA), Italy

²Arhs Developments, Italia

³Emgrisa, Madrid, Spain

⁴Seidor, Barcelona, Spain

*Corresponding author bruna.grizzetti@ec.europa.eu

This is a non-peer reviewed preprint of an article that was subsequently accepted for publication in *Global Environmental Change*. Please cite the published article which is freely available at <https://doi.org/10.1016/j.gloenvcha.2021.102281>

Abstract

Intensive agriculture and densely populated areas represent major sources of nutrient pollution for European inland and coastal waters, altering the aquatic ecosystems and affecting their capacity to provide ecosystem services and support economic activities. Ambitious water policies are in place in the European Union (EU) for protecting and restoring aquatic ecosystems under the Water Framework Directive and the Marine Strategy Framework Directive. This research quantified the current pressures of point and diffuse nitrogen and phosphorus emissions to European fresh and coastal waters (2005-2012), and analysed the effects of three policy scenarios of nutrient reduction: 1) the application of measures currently planned in the Rural Development Programmes and under the Urban Waste Water Treatment Directive (UWWTD); 2) the full implementation of the UWWTD and the absence of derogations in the Nitrates Directive; 3) high reduction of nutrient, using best technologies in wastewaters treatment and optimal fertilisation in agriculture. The results of the study show that for the period 2005-2012, the nitrogen load to European seas was 3.3-4.1 TgN/y and the phosphorus load was 0.26-0.30 TgP/y. EU policy measures could decrease the nutrient export to the seas up to 14% for nitrogen and 20% for phosphorus, improving the ecological status of rivers and lakes, but widening the nutrient unbalance in coastal ecosystems, affecting eutrophication. Further nutrient reductions could be possible by a combination of measures especially in the agricultural sector. The respective contribution of nutrient sources and expected changes differ per European regional seas. The study highlights the advantages of adopting a nexus thinking when addressing the land-sea dynamics, checking the coherence of measures taken under different policies.

Keywords

Nutrient pollution, policy scenarios, EU Water Framework Directive, EU Marine Strategy Framework Directive.

1. Introduction

In Europe, intensive agricultural practices together with high population density represent important sources of nutrients for fresh and coastal waters (Sutton et al., 2011). Nutrient pollution is one of the major pressures on European aquatic ecosystems altering their condition (Grizzetti et al., 2017). At present in the EU more than half of water bodies are not in good ecological status, with nutrient being one of the major causes of degradation (Poikane et al., 2019a). Many marine ecosystems suffer from hypoxia and eutrophication (Diaz and Rosenberg, 2008; Romero et al., 2013). In estuaries and coastal waters the increase of nutrient availability from riverine loads foster primary productivity causing the phenomenon of eutrophication (Howarth et al., 2011). In particular, the imbalance of nitrogen and phosphorus over silica can be responsible for the proliferation of harmful algal blooms (HABs) (Billen and Garnier, 2007). Eutrophication impairs water quality and alters the condition and functioning of fresh and marine ecosystems, affecting their capacity to supply key ecosystem services and sustain economic activities, such as water purification, coastal protection, lifecycle maintenance, drinking water, fishing, shellfish farming, recreation and tourism (Culhane et al., 2019; Grizzetti et al., 2016; Liqueste et al., 2016; Piroddi et al., 2017).

Ambitious water policies are in place in the European Union (EU) for protecting and restoring freshwater and marine ecosystems under the Water Framework Directive (WFD, 2000/60/EC) and the Marine Strategy Framework Directive (MSFD, 2008/56/EC). The goal of the WFD is to achieve a Good Ecological Status for all water bodies in the EU, including river, lakes, transitional and coastal waters. Under the WFD, Member States develop River Basin Management Plans (RBMP) for the protection and restoration of aquatic ecosystems and the sustainable use of water resources. Measures adopted under the Nitrates Directive (91/676/EEC) and the Common Agricultural Policy (CAP), to reduce nutrient water pollution from agriculture, as well as actions taken under the Urban Waste Water Treatment Directive (UWWTD, 91/271/EEC) contribute to the objectives of the WFD. Similarly, the MSFD aims at achieving Good Environmental Status (GES) of all EU marine waters, protecting biodiversity and resilience of marine ecosystems, and promoting their sustainable use. The GES is described by 11 qualitative descriptors and nutrient pollution directly or indirectly influences several of them concerning biodiversity, non-indigenous species, fish population, reproduction, eutrophication and sea floor integrity (Descriptors 1-6). European seas are also protected under four international Conventions: the Helsinki Convention on the Baltic Sea (HELCOM, 2020), the OSPAR Convention on the North-East Atlantic (OSPAR, 2020), the Barcelona Convention on the Mediterranean (UNEP, 2020) and the Bucharest Convention on the Black Sea (Black Sea Commission, 2020).

The provision of ecosystems services depend on the condition of aquatic ecosystems (Grizzetti et al., 2019). Current pressures and future changes, such as climate changes, can further degrade the status of aquatic ecosystems (Jonkers et al., 2019; Lajaunie-Salla et al., 2018; Macias et al., 2018) and the resilience of water resources (Zampieri et al., 2019). Modelling tools can be useful to study the impacts of future scenarios, policy measures and climate changes at the regional and continental scale (Arheimer et al., 2012; Bartosova et al., 2019; Bouraoui et al., 2014; Ludwig et al., 2010; Seitzinger et al., 2010), and to

check the coherence of different policy targets, for instance between water (WFD and MSFD) and agricultural (Common Agricultural Policy, CAP) policies. The assessment of policy scenarios for Europe requires a flexible and spatially detailed analysis to account for the wide climatic, hydrological and socio-economic gradients, as well as the use of consistent data across many different countries.

The objective of this study was to quantify the current nitrogen and phosphorus pressures on European fresh and coastal waters and to assess the effects of policy scenarios to reduce nutrient pollution with progressive levels of ambition. In particular, the study aimed at developing a spatial analysis for inland and coastal waters to fit different level of policy intervention from the water body to the river basin and the regional sea, including the most recent data available across European countries by standard data reporting flows.

Key questions of the analysis were: ‘Would the current measures in place ensure the achievement of the WFD goal of good ecological status for water bodies and MSFD target of good environmental status for the marine ecosystems?’ and ‘How far different scenarios of nutrient pressures reduction could contribute to the improvements of ecological status of freshwaters and the reduction of nutrient export to the sea?’ To address these questions we applied a new version of the model GREEN (Grizzetti et al., 2012). The model GREEN has been used in previous studies for assessing the nutrient loads to the European seas, the nitrogen retention in European freshwaters, and for scenario analysis (Bouraoui et al., 2014; Grizzetti et al., 2012; La Notte et al., 2017; Leip et al., 2015; Malagó et al., 2019).

2. Materials and methods

2.1 Overview of the assessment

The model GREEN (Geospatial Regression Equation for European Nutrient losses) (Grizzetti et al., 2012, 2008) was applied to assess the nitrogen and phosphorus pressures on European fresh and coastal waters. In specific, nutrients’ annual load and average concentration at different points of the river network and at the river outlets to the sea were estimated, as well as the contribution of different diffuse and point sources to the total load.

The spatial extent of the analysis covered all river basins draining into European seas and whose waters fall completely or in part within EU28 countries (Figure 1). EU overseas territories were not included in the study. The assessment was carried out for the period 2005-2012 (with a focus on 2012) for which best available data for model input and calibration were available at the European scale.

Nitrogen and phosphorus inputs from different sources in river basins were estimated for all Europe to describe the current nutrient pressures on waters (reference scenario). Then, three scenarios were developed to simulate the effect of policies with progressive levels of ambition to reduce nutrient pressures. The scenarios on nutrients were associated to concurrent measures to alleviate water scarcity (De Roo et al., 2020).

Finally, the results of the scenarios were analysed in view of the EU water policy (Water Framework Directive) target of good ecological status for all rivers, lakes and transitional waters in the EU and the good environmental status for marine waters.




Figure 1. Extent covered by the analysis: European river basins up to the red line. Colours indicate the river basins per European Regional Seas. An example of the catchments and river network delineation is provided in the upper right corner.

2.2 The model GREEN

The model GREEN (Grizzetti et al., 2012) includes a geospatial data model for Europe (geo-database), where data are linked to the hydrological structure of the river network (developed in ESRI environment), and routines to model nitrogen and phosphorus flow in the river basin according to different pathways (developed in SQL server environment).

The geospatial data model is based on the CCM2 model (Vogt et al., 2007) and includes the Ecrin data for lakes (EEA, 2012). Europe is divided in ~1 million catchments of 7 km² average size. Each catchment has an elementary river stretch, except in coastal areas, where river stretch might be absent.

The model distinguishes two major pathways to represent the fate of nutrients: diffuse sources that undergo a retention in the land phase (basin retention) before reaching the stream, and point sources that are directly discharged into surface waters. Once in the river all sources are reduced by the in-stream retention (river retention). Diffuse sources include nutrient from mineral fertilisers, manure application, nitrogen crop and soil fixation, nitrogen atmospheric deposition, background losses (only for phosphorus) and inputs from scattered dwellings, i.e. isolated houses that are not connected to sewerage systems. Point sources consist of urban and industrial wastewater discharges. Nutrient input from the different sources and the basin and river retention are simulated in each small catchment and routed through the river network. Basin retention is modelled as a decay function proportional to the inverse of the total annual precipitation in the catchment; river retention is estimated as a decay function proportional to the river length, considered as a proxy for water residence time. In addition, lake retention is simulated as a function of lakes residence time and average depth. Two parameters are calibrated at the European scale,

a basin retention coefficient (*basinCoeff*) and a river retention coefficient (*riverCoeff*). In the model, inputs from agricultural sources are reduced by the basin and then the river retention, while point sources are reduced only by the river retention. Input from scattered dwellings is considered to be halved before reaching the stream. Nitrogen atmospheric deposition and phosphorus background losses are split into two parts, i.e. inputs to agricultural land undergo the basin retention (which include also the crop uptake), while in all other areas they are reduced by a fixed rate, derived from the literature, before entering into the stream (Model equations are provided in SuppMat S1).

Model input and output data are organised in the geo-database at the spatial resolution of catchments. The model results are aggregated per Functional Elementary Catchments (FECs) (EEA, 2012), containing on average 2-3 catchments), per river basins and River Basin Districts (RBD), according to the WFD (EEA, 2019a), and per basins draining into marine regions, according to the MFSD (EEA, 2018a).

For the present study a new version of the model GREEN was developed. The major changes from the previous version (Grizzetti et al., 2012) are the higher spatial resolution of the model application (sub-catchments of 7 km² average size), the inclusion of all coastal areas (also in absence of streams), the adoption of Corine Land Cover maps for land cover allocation, and a new setup of nutrient input data corresponding to the most recent European data publicly available. In addition, the model was calibrated using a different dataset and for a more recent period (2008-2012).

2.3 Input of nutrient sources and model parameterisation

The allocation of different land cover type in each catchment was based on the Corine Land Cover map (CLC, 2012) (grid at 100 m resolution, available for year 2000, 2006 and 2012 for Europe). For areas falling outside the CLC, land cover data was taken from the Climate Change Initiative Land Cover map (ESA, 2017) (global grid at 300 m resolution, yearly maps from 1992 to 2015). Time series on total agricultural area and fertiliser application per major crops and grassland were provided by the model CAPRI at the administrative unit level NUTS2 (European Commission, 2018) (yearly values from 1984 to 2013, estimated considering year 2012 as base year). The CAPRI model covers EU28, Norway, and non-EU countries in the Balkans. Information on fertiliser applications were also retrieved from FAO (FAOSTAT, 2020), for European countries not covered by CAPRI.

Two annual time series of nitrogen and phosphorus mineral and manure fertiliser maps from 1995 to 2015 were developed for Europe based on two different methods. The first adopts the CLC for the spatial location of crops and the CAPRI model for the information on fertilisers and utilised agricultural area by crop type. The second considers the extent and location of agricultural area provided by ESA and the total fertilisation rate reported by FAO per country. The first approach offers higher details on crop location and fertilisation, and the utilised agricultural area is consistent with data reported by national statistics of EU28 countries (as CAPRI is based on EUROSTAT). The second method provides average nutrient fertilisation rates per country (without distinguishing crop types) in agricultural areas detected by satellite images, and can be applied to all European countries (also those not covered by CAPRI) and at the global scale. In order to develop nitrogen and phosphorus fertiliser maps for all river basins draining in European seas the results of the two methods were combined, completing the areas (countries) not covered by the first method by the second one. The model CAPRI provides also the nitrogen fixation by crops. In addition biological nitrogen fixation in soils of 5 kgN/ha and phosphorus background sources of 0.15 kgP/ha were considered as in the previous version of the model. Annual total nitrogen atmospheric deposition was computed using the data from the model EMEP (EMEP, 2020).

Nitrogen and phosphorus inputs from domestic waste were retrieved from Vigiak et al. (Vigiak et al., 2020), which used the most recent information reported by EU countries under the Urban Waste Water Treatment Directive. Further, industrial discharges were retrieved from the European Pollutant Release and Transfer Register (E-PRTR) (EEA, 2018b) and official statistics available in EUROSTAT and WHO for non EU countries. The dataset is based on the spatial analysis of different sources and pathways of human and industrial waste to surface waters, including the location and level of treatment of each treatment plants (Vigiak et al. 2019).

While annual time series were available for nutrient inputs from agriculture and nitrogen atmospheric deposition, inputs from scattered dwellings and point sources discharges were available only for the year 2012 and applied as constant for the period 2005-2012.

In each catchment the stream length was based on the CCM2 model and the time series of annual precipitation was computed from EFAS-Meteo (Arnal et al., 2019; Ntegeka et al., 2013). The water flow estimated by the model LISFLOOD (De Roo et al., 2020) was included in the analysis, by interpolating the average annual water flow, provided by the model on a 1 km² grid resolution, at each catchment outlet of the geospatial data model.

2.4 Model calibration and validation

The calibration of the two model parameters (basinCoeff and riverCoeff) was performed over the period 2008-2012 using the data publicly available in the WaterBase v14 (EEA, 2020) on mean annual nitrogen and phosphorus concentrations. In total 9335 observations for total nitrogen and 13890 observations for total phosphorus were used for the calibration, corresponding to 3685 and 4163 different locations in Europe for nitrogen and phosphorus, respectively (additional information in SuppMat S2). The period 2008-2012 was selected for the calibration as the quality of the data and their spatial coverage (despite not homogeneous across Europe) were better than in previous years. In addition, using the whole period for calibration was necessary to cover different hydrological years (in the studied area the average rainfall in the period 2008-2012 ranged between 739-881 mm). The annual nitrogen and phosphorus “observed” loads used for calibration were computed multiplying the concentrations times the annual water flow estimated by the model LISFLOOD for all the points in the river network where observed data were available.

The calibration was carried out using a routine designed specifically for this purpose using R (R Core Team, 2014), including several optimization algorithms (Simulated annealing, Genetic Algorithm, etc.). It was found that the application of a Markov Chain Monte Carlo method (Kuczera and Parent, 1998) was the most efficient when the number of evaluations performed in the calibration process was less than 500 (which is desirable for practical purposes given the computation time requirements for each evaluation). Several goodness of fit measures were considered for the optimal parameter selection, among others: Nash-Sutcliffe Efficiency (NSE), Percent Bias (pbias), Relative Index of Agreement (rd), Relative Nash-Sutcliffe efficiency (rNSE) (Krause et al., 2005). The calibration was run both with actual values and log10 values to compare the effects of large nutrient loads (generally in downstream catchments) with small loads (generally in upstream catchments). The final parameters selection was based on the Nash-Sutcliffe Efficiency (NSE) (Nash and Sutcliffe, 1970) statistic computed on the no-log simulation.

The model was run from 2005 to 2012. The verification of the model estimation was conducted comparing the nutrients load estimated by the model GREEN at the outlet of the 50 largest rivers in Europe with data

available from various independent sources in the literature. Correlation was computed for the whole data set as well as for the different data sources. The linear fit between reported and modelled freshwater runoff and nutrient loads was computed, assuming the intersect at 0.

2.5 Scenarios development

The scenarios of reduction of nutrient pollution in inland and coastal waters were designed with the specific intent of providing support to the EU water policy and with a progressive increasing level of ambition. Three scenarios were developed, including both a reduction of urban point sources and diffuse agricultural pollution (Table 1). The first scenario, called Business As Usual Scenario (BAU), was meant to represent the current level of investment in water protection from nutrient pollution foreseen by actual planning instruments in EU Member States, notably, investments in upgrading urban waste water treatment plants under the UWWTD Art.17 (Benitez Sanz et al., 2018), and investments in measures to protect water quality under the Rural Development Programme (RDP) 4b priority. In the latter, the effectiveness of the mix of measures was assumed to be 10% reduction of the nitrogen and phosphorus entering the water system (Sarteel et al., 2016). This reduction was applied in all catchments, but only considering the fraction of Utilised Agricultural Area (UAA) that is covered by RDP 4b priority in the NUTS2, as the location of UAA subject to RDP 4b priority is unknown. The second scenario, called Nutrient Scenario (NUTR), intended to represent a full or enhanced implementation of two EU Directives, the UWWTD for collecting and treating wastewater from urban agglomerations, and the Nitrates Directive to protect freshwater from agricultural pollution. In the NUTR scenario all wastewater treatments in the EU were set compliant with the requirement of the UWWTD (Pistocchi et al., 2019) and nitrogen application in agricultural fields was limited to maximum 170 kgN/ha in all Nitrates Vulnerable Zones in the EU irrespectively of the presence of Derogations (phosphorus in manure application was reduced proportionally). Finally the third scenario, called High Technically Feasible Reduction scenario (MTFR), considered that all wastewaters in the EU are treated with the maximum level of nutrient reduction currently possible (correspondent to a tertiary treatment with an enhanced reduction of phosphorus), while mineral fertiliser are reduced according to an optimal fertilisation. The latter was simulated keeping the nitrogen surplus (difference between nitrogen input and output in the agricultural field) to a minimum, which was set to 10% of the output. The amount of mineral nitrogen to be reduced per each EU country was estimated using the data of nitrogen balance (surplus) reported by EUROSTAT for year 2012 and considering that only 70% of the current manure application could be used by plants. The country specific reduction of mineral nitrogen was applied distributing the reduction rate according to a weight that accounts for catchments with higher application of mineral+manure fertilisers. The corresponding reduction of mineral phosphorus was estimated.

Each scenario of nutrient reduction was associated to a corresponding scenario of measures for preventing water scarcity implemented in the LISFLOOD model (De Roo et al., 2020). The BAU and NUTR scenarios were associated with the implementation of measures on water quantity (increase of water use efficiency in irrigation and in domestic usage, changing cooling water requirements due to change in energy demand and energy mix, and implementation of wastewater re-use for irrigation), corresponding to the level of current investments foreseen in River Basin Management Plans of the WFD. Differently, the MTFR scenario was associated to a higher level of implementation of the same measures.

Table 1. Description of the scenarios. The scenarios for water quantity are described in De Roo et al. (2020).

Scenario acronym	Water quantity (model LISFLOOD)	Water quality – Point pollution (model GREEN)	Water quality – Diffuse pollution (model GREEN)
REF	Situation in 2005-2012	Situation in 2012	Situation in 2005-2012
BAU (Business As Usual)	Change in irrigation efficiency, urban water usage efficiency, water re-use, and water use changes due to energy demand changes, under the current investments in River Basin Management Plants of the WFD	Nutrient reductions related to the current investments in EU under the Art.17 of the Urban Waste Water Treatment Directive (Council Directive 91/271/EEC).	Nutrient reductions due to the measures funded under the Rural Development Programme 4b priority (spatial information on the measures for implementing the Nitrates Directive were not available).
NUTR (Nutrient Reduction)	<u>As BAU</u>	Full implementation of the Urban Waste Water Treatment Directive (Council Directive 91/271/EEC) in EU.	Maximum application of manure nitrogen is limited to 170 kgN/ha in areas draining the Nitrates Vulnerable Zones (NVZ), irrespective of the presence of Derogations. The corresponding decrease of phosphorus in manure application is considered.
MTFR (High Technically Feasible Reduction)	Change in irrigation efficiency, urban water usage efficiency, water re-use, and water use changes due to energy demand changes under the Maximum Technical Feasibility scenario	All urban waste water treatment plants are upgraded to the highest treatment level.	Scenario of optimal fertilization. The nitrogen surplus on agricultural soils is set to 10% of the reported output. The corresponding reduction of mineral phosphorus is considered.

2.6 Scenarios analysis and link to ecological targets

Annual nutrient load to the sea and concentration in freshwater estimated by the model GREEN for the three scenarios (BAU, NUTR, MTFR) were compared with the values of the reference scenario (REF) for year 2012. This year was chosen as the most representative of average hydrological conditions (average rainfall of 823 mm, compared to the range of 739-881 mm for the period 2008-2012) and input data for this year were the most accurate. The analysis was conducted at different levels of aggregation: 1) Europe (extent covered by the modelling), 2) MSFD regional sea basins, and 3) river basins, which are of interest for the WFD, as River Basin Districts are composed of one or several river basins (results not shown). The share of different nutrient input sources to the river basin was evaluated as well as their contribution to the total load in rivers. In addition, the predicted changes in nutrient concentration and N:P ratio were analysed.

The potential impact of the three scenarios on achieving the target of good ecological status established by the WFD was assessed by spatially linking the model predictions with the location and ecological conditions of water bodies reported by the EU Member States under the WFD. For this purpose, the most recent data on the ecological status of lakes, rivers and transitional waters, referring to second River Basin Management Plans, 2010/2016 (EEA, 2019b), were overlaid with the geospatial data model of GREEN. The WISE dataset includes information on water bodies' delineation, and on the ecological status and main impact types observed. Each WISE water body was assigned to one or multiple catchments in the GREEN geospatial model according to its location and spatial geometry. A one to one correspondence between WISE water bodies and the GREEN geospatial model was not possible as the delineation of water bodies depends on decisions of River Basin District Authorities in the different EU countries, while the catchments and river network delineation in the GREEN geospatial model are derived from consistent topographic information at the European scale (CCM2 model). The water bodies area (for lakes) and length (for rivers and transitional waters) falling in each catchment of the geospatial model were computed, as one water body could lay in one or many catchments. The nutrient concentration estimated by the model GREEN for each WISE water body was computed as area-weighted or length-weighted average across the CCM2 catchments involved.

The spatial association allowed to compute nutrient concentration estimated by the model GREEN in the WISE water bodies and derive the distribution of nutrient concentration for each ecological status classes. A value derived from the average nitrogen and phosphorus concentration in water bodies in Good Ecological Status in the baseline simulation (REF) was used as threshold for scenarios comparison. These thresholds were computed for rivers, lakes, and transitional waters bodies considering (i) the whole EU and (ii) a subset of water bodies where 'Nutrient Pollution' impact was reported (25% of the total water bodies). The potential impacts of the scenarios on the ecological status was evaluated considering the percentage of lakes area and rivers length that was below the threshold in the different scenarios.

3. Results

3.1 Model calibration and validation

The results of model calibration indicated a good agreement between observed and estimated nutrient loads, with NSE of 0.96 and 0.75 for nitrogen and phosphorus, respectively (SuppMat S3).

The freshwater runoff estimated by the LISFLOOD model and total nutrient loads estimated by the model GREEN at the outlet of the 50 largest European rivers showed a good agreement with the values reported by other data sources and the literature (Figure 2 and SuppMat S4). Correlation coefficients of model estimates with loads reported by HELCOM and OSPAR Conventions are 0.85 for nitrogen and 0.90 for phosphorus; the correlation with data found in the literature (listed in SuppMat S4) is 0.94 for nitrogen and 0.84 for phosphorus. The gradients of the linear fitting are almost all around 1, indicating a good fit between reported and modelled values. Overall, the model results with respect to the nitrogen loads are better than for the phosphorus ones. Beside the good overall agreement, for some individual rivers severe deviations occur, e.g. freshwater runoff of river Scheldt is two magnitudes below the reported values, while nutrient loads fit well. On the other hand, for some rivers (e.g. Thames, Humber) GREEN provides the nutrient loads directly at the outlet to the sea, while observations are collected upstream.




Figure 2. Comparison between annual Total Nitrogen loads reported by different sources and estimated by the GREEN model (tN/y) shown on a log-log scale. The sources of reported values are indicated by the dot colours and are detailed in SuppMat S4).

3.2 Scenarios analysis

3.2.1 Changes in nutrient inputs

In Europe (region covered by the study) for the reference year 2012 the total input of nitrogen was estimated at 27.6 TgN (23.3 TgN in EU28) of which 71% from mineral and manure fertilisers, 17% from atmospheric deposition, 8% from biological natural fixation (in crops and soils), 3.5% from human and industrial waste wasters and 0.5% from scattered dwellings. The total input of phosphorus was estimated at 4.1 TgP (3.5 TgP in EU28), of which 93% from mineral and manure fertilisers, 5% from human and industrial waste wasters, 2.3% from natural background and 0.5% from scattered dwellings. On average point nutrient sources represented only 3-5% of the total input to the river basin system, but locally they could reach much higher shares.

Overall, in the BAU, NUTR and MTRF scenario compared to the REF scenario, the total nitrogen input was reduced by 1%, 5% and 18% respectively, and the total phosphorus input was reduced by 1%, 10% and 14%, respectively. In particular, for EU28 the MTRF scenario foresaw a reduction of 21% of nitrogen input (-24% point sources and -25% agricultural sources) and 16% of phosphorus input (-49% point sources and -16% agricultural sources). When looking at regional differences, the BAU scenario resulted in limited reduction (1-2%) in all regions for both nitrogen and phosphorus, while the NUTR scenario involved large changes in some regions, such as in the basins draining into the Greater North Sea characterised by intensive agriculture and livestock production. The MTRF scenario foresaw reductions between 13-37%

for total nitrogen and 9-32% for total phosphorus input, with substantial changes in the Bay of Biscay and Iberian Coast and in the Mediterranean region, especially by abating point sources (SuppMat S5).

3.2.2 Changes in nutrient loads to the sea

For the period 2005-2012, the model estimated that the total nitrogen load to European seas was 3.3-4.1 TgN/y (3.9 TgN/y in 2012) and the total phosphorus load was 0.26-0.30 TgP/y (0.29 TgP/y in 2012) (Figure 3). Considering the most recent year 2012, which was an average year for precipitation, the scenarios BAU, NUTR and MTRF bring about a reduction of nitrogen load of 2%, 6% and 14%, respectively, and a reduction of phosphorus load of 3%, 8%, 20%, respectively. The changes are related mainly to a reduction of agricultural sources for nitrogen and to an improvement of wastewater discharges for phosphorus.




Figure 3. Total nitrogen (left) and phosphorus (right) annual load to European seas estimated by the GREEN model (ton/y) for the period 2005-2012. Colours indicate the relative contribution of major sources: point sources (red), scattered dwellings (orange), agriculture (green) and background (P) or atmospheric deposition (N) (blue).

The scenarios produce different effects regionally (Figure 4 and 5, tables provided in SuppMat S6). The BAU scenario involves quite limited reductions for both nitrogen and phosphorus load (1%-3%) in almost all regional seas except in the Baltic Sea, where improvements are related to investments in upgrading waste water treatments, and in the Adriatic Sea (mainly for phosphorus abatement). The NUTR scenario benefits especially the Greater North Sea (-9% N load, -14% P load) for reduction of nutrients from agricultural sources, and the Central and Western Mediterranean (up to -11% for N load and -13% for P load) and the Black Sea (-6% N load, -12% P load) for improvements in wastewater treatments and agricultural sources. The highest decline of nutrient loads to European seas are obtained by the MTRF scenario, with reductions of nitrogen ranging from around 10% in the Baltic and Danube regions, to about 15% in the Greater North Sea, Celtic sea, Bay of Biscay and Iberian Coast, and 20% and more in the Central and Western Mediterranean. These improvements are related to both upgrading of wastewater treatment plants and reduction of mineral fertiliser application in the Mediterranean and Balkan regions, and mainly due to a reduction of mineral fertilisers in the Great North and Baltic regions. Concerning phosphorus, the MTRF foresaw a decrease of 9% in the Baltic Sea, around 20% in the Greater North Sea, Celtic Sea and Black Sea, and a drop of 45-47% in the Central and Western Mediterranean. Also, in this case changes are associated to reduction in the use of mineral fertiliser (in the Mediterranean and Baltic region) and to the abatement of point sources (in the Mediterranean and Bay of Biscay and Iberian Coast).





Figure 4. Total nitrogen load to different regional seas estimated by the GREEN model (ton/y) for the reference (REF) and the BAU, NUTR and MTRF scenarios (Table 1). Colours indicate the relative contribution of major sources: mineral nitrogen (MinN), manure nitrogen (ManN), atmospheric deposition (AtmN), crop fixation (FixN), soil fixation (SoilN), scattered dwellings (SdN) and point sources (PsN).



2 Figure 5. Total phosphorus load to different regional seas estimated by the GREEN model (ton/y) for the reference (REF) and the BAU, NUTR and MTR scenarios (Table 1). Colours
 3 indicate the relative contribution of major sources: mineral phosphorus (MinP), manure phosphorus (ManP), background (BgP), scattered dwellings (SdP) and point sources (PsP).

3.2.3 Changes in nutrient concentration and N:P ratio

The nutrient load and concentration in the scenarios depend on the combined effects of measures to combat water scarcity and actions to curb nutrient pollution. The measures implemented for water saving generally slightly increase the water flow in rivers (dilution effect for nutrients), especially in the Mediterranean region and in some irrigated basins in Western Europe, while in Eastern Europe future investments in water use are foreseen to produce a small decrease of the water flow (concentration effect for nutrients).

Mean concentrations of both nitrogen and phosphorus decrease under the three scenarios. On average, in inland waters the reduction is higher for phosphorus (up to -18% in MTRF) than for nitrogen (up to -11% in MTRF). The measures affect mainly polluted areas, with decreases observed especially in streams with concentrations above the 75th percentile. The decline in concentration is more prominent at the outlets to the sea than in inland waters, with small variation under the BAU scenario and changes of about -10% and -20% in the NUTR and the MTRF scenario respectively (SuppMat S7).

The ratio between nitrogen and phosphorus (N:P) in inland waters varies greatly across Europe, according to local conditions and nutrient sources. The effect of the scenarios on the respective proportion of the nutrient can be very different. At the sea outlets, the BAU does not substantially alter the N:P ratio, as by construction the scenario involves the same reduction of nitrogen and phosphorus in agriculture field and limited improvements in wastewater treatment plants. Conversely, overall the N:P ratio increases under the NUTR scenario and decreases under the MTRF scenario. This is related to the fact that the N:P ratio in the manure is different from that of the mineral fertilisation, with the NUTR scenario limiting manure and the MTRF tackling mineral fertilisation (SuppMat S7).

3.3 Link to the ecological status of water bodies

The changes in nutrient concentration and N:P ratio have implications on the ecological condition and biological components of the aquatic ecosystems. The average concentration of nitrogen and phosphorus estimated by the model GREEN for the WISE water bodies for the different class of ecological status is shown in Table 2. Considering only WISE water bodies that are currently reported in good ecological status, for year 2012 (baseline) the model estimates an average concentration of 1.13 mgN/l and 0.08 mgP/l for lakes, 2.52mgN/l and 0.16 mgP/l for rivers, and 3.88 mgN/l and 0.44 mgP/l for transitional waters. Class average nutrient concentration increases for water bodies from high to more degraded status. It also takes high values in water bodies for which the ecological status is not reported.

Based on the average concentration found by modelling for water bodies in good ecological status, the threshold of 2.0 mgN/l for nitrogen and 0.1 mgP/l for phosphorus were set to compare the potential effect of the three scenarios for reaching the policy target of good ecological status (Table 3). Overall, the predicted improvement concerns only a limited fraction of inland waters, with only an additional 1% of rivers length and 3-4% of lakes and transitional waters area being below the thresholds in the MTRF scenario. These fractions slightly increase when restricting the analysis to water bodies for which the “nutrient pollution” impact is reported in the WISE data (i.e. 2% of rivers length and 4-5% of lakes and transitional waters area).

Table 2. Concentration estimated by the model GREEN per water body type and class of ecological status. Values refer to the baseline scenario (REF) in year 2012. Concentrations are weighted by the fraction of segment length or polygon area of the WISE water bodies falling in each catchment of the GREEN spatial data model. The values discussed in the text are reported in bold.

	Class of Ecological Status	Total segment length (km)	Total polygon area (km ²)	Mean nitrogen concentration (mgN/l)		Mean phosphorus concentration (mgP/l)	
				(weighted by length)	(weighted by area)	(weighted by length)	(weighted by area)
Lakes							
	High	-	56948	-	0.80	-	0.09
	Good	-	160056	-	1.13	-	0.08
	Moderate	-	283399	-	1.75	-	0.12
	Poor	-	111154	-	2.76	-	0.15
	Bad	-	15004	-	24.31	-	1.28
	Unknown	-	5392	-	3.47	-	0.20
Rivers							
	High	764081	5159	0.81	3.17	0.07	0.31
	Good	2551093	12467	2.52	3.97	0.16	0.38
	Moderate	5012752	19176	4.63	4.67	0.25	0.42
	Poor	2287402	3634	8.05	4.41	0.41	0.42
	Bad	808669	632	10.72	19.92	0.60	0.74
	Unknown	232524	648	4.52	6.33	0.26	0.42
Transitional waters							
	High	8	118	-	0.72	-	0.06
	Good	74	3494	13.19	3.88	0.55	0.44
	Moderate	56	12102	1.27	9.16	0.08	1.17
	Poor	21	7628	2.75	14.26	0.18	1.17
	Bad	-	9830	-	12.46	-	1.23
	Unknown	-	424	-	42.03	-	2.77

Table 3. Fraction of rivers length, lakes area and transitional waters area of WISE water bodies that is below a threshold concentration of 2.0 mgN/l and 0.1 mgP/l under the reference and BAU, NUTR and MTFR scenarios (Table 1). (*) Compute on a subset of water bodies reporting 'nutrient pollution' impact in WISE data.

Scenario	Nitrogen concentration THRESHOLD < 2.0 (mgN/l)		Phosphorus concentration THRESHOLD < 0.1 (mgP/l)	
		(*)		(*)
Lakes				
REF	0.87	<i>0.68</i>	0.88	<i>0.70</i>
BAU	0.87	<i>0.68</i>	0.88	<i>0.70</i>
NUTR	0.88	<i>0.69</i>	0.88	<i>0.71</i>
MTFR	0.88	<i>0.70</i>	0.89	<i>0.72</i>
Rivers				
REF	0.43	<i>0.30</i>	0.46	<i>0.33</i>
BAU	0.44	<i>0.31</i>	0.46	<i>0.34</i>
NUTR	0.44	<i>0.31</i>	0.49	<i>0.37</i>
MTFR	0.47	<i>0.34</i>	0.49	<i>0.37</i>
Transitional waters				
REF	0.08	<i>0.10</i>	0.07	<i>0.10</i>
BAU	0.09	<i>0.10</i>	0.07	<i>0.10</i>
NUTR	0.09	<i>0.10</i>	0.08	<i>0.10</i>
MTFR	0.11	<i>0.12</i>	0.11	<i>0.14</i>

4. Discussion

4.1 Modelling and scenarios limitations

Assessing nutrient pressures and the effect of policy scenarios on fresh and coastal waters at the European scale is challenging. Europe is a wide continent with a large variability of soil, climatic, hydrological and ecological features, and much diversified landscapes and agronomic production systems. Nutrient pressures on water from agriculture or point sources and remediation measures in place vary greatly according to the regional socio-economic conditions. To address the complexity of nutrient pollution in European waters, special attention was dedicated to the spatial representation of the hydrological system and coastal catchments, and to use the most updated and homogeneous data on nutrient pressures at the European scale. Both aspects are crucial for assessing policy scenarios, indeed the consistency and transparency of nutrient sources are necessary to ensure a sound representation of policy actions, and the quantification of their impacts from the single water bodies, to the river basins and marine regions is useful to support different levels of policy intervention. The high spatial resolution of model input and output (catchments of $\sim 7\text{km}^2$) afforded a detailed representation of nutrient sources and a good link to water bodies delineated by Member States under the WDF (WISE data). A specific effort was dedicated to the inclusion of coastal catchments, to improve the representation of nutrient sources in these areas, which are generally highly populated with nutrient discharges directly into the sea. Nevertheless, the model has some structural limitations, as it provides a simplified representation of nutrient processes and pathways in the aquatic environment. It does not include feedback processes such as the effect of irrigation on fertiliser input or the pollution legacy in groundwater. The model provides annual estimations of nitrogen and phosphorus loads and their inter-annual variation but for ecological impacts seasonal distribution is also relevant.

In the assessment, nutrient concentrations result from the ratio between nutrient loads estimated by the model GREEN and water flow produced by the model LISFLOOD. In some catchments, concentrations can take high values, for examples in upstream dry areas where modelled water flow is low, or in presence of large point sources discharges, with high share of artificial water circulation, which is not well captured by the model GREEN (water abstractions and return water is taken into consideration in the computation of water flow in the model LISFLOOD).

Data availability also imposed some constraints. Observed data for model calibration were limited to the information available by the European Environment Agency (EEA), which concerned only annual nutrient concentration (without associated information on water flow) and were not homogeneously distributed across Europe. In future model calibration, statistical techniques, such as bootstrap sampling, could be used to balance the influence of heterogeneous distribution of gauging points, so that upstream and downstream stations could be proportionally represented. With regard to nutrient input, urban wastewater discharges were estimated only for year 2012 and kept constant in the simulation, disregarding the changes that have occurred between 2005 and 2012. This limits loads simulations but has no relevance for the scenarios analysis.

All the scenarios considered nutrient reductions in both point and diffuse agricultural sources. The BAU scenario envisaged the application of measures limiting nutrient losses to water according to the current investments under the Urban Waste Water Treatment Directive (Art.17) and Rural Development Programmes (priority 4b). An average 10% effectiveness of measure for agricultural losses was considered in the modelling based on the study of (Sarteel et al., 2016), but more regional variability is possible. It is

important to note that in the BAU scenario the effect of additional measures in place under Action Programmes of the Nitrates Directive or other schemes could not be taken into consideration for the lack of consistent and quantitative information across Europe. More knowledge in this regard would benefit future assessments. The NUTR scenario focused on the reduction of manure in Nitrates Vulnerable Zones, independently from the presence of Derogations. Information on the manure was available at the NUT2 level from the model CAPRI and was then spatialized using the Corine Land Cover map (as explained in the Section 2.3), but no information was available on the way manure is treated and stored, which can influence the impact on water pollution. The MTFR scenario minimised the use of mineral fertilisers, recycling almost all manure produced as fertiliser, but not reducing the current livestock production. The optimisation was computed using nitrogen balance at the country scale, as regional values were not available, but this might not be fully representative for countries such as Italy that have a strong gradient in fertiliser inputs across the country.

4.2 Nutrients loads to the European seas

According to the estimations of the BAU scenario, the current investments by the EU Member States for reducing point and diffuse nutrient pollution might result in little improvements of water quality (only 2-3% reduction of the nutrient load to the seas). There is still a potential for nutrient recovery in wastewaters that could be tapped into, especially in the Danube and the Adriatic regions. Also, in many EU countries measures to reduce nutrient losses to water could be extended, as they currently cover only a small fraction of the agricultural land (according to the budget allocation under priority 4b in Rural Development Programmes).

The NUTR and MTFR scenarios represent the possible effects of more ambitious investments. Concerning wastewater, significant nutrient reductions could be obtained in the Atlantic coasts and in the Mediterranean and Danube regions, with improvements possible also beyond the implementation of the UWWTD. However, a significant reduction of nutrient loads to the seas would only be possible with important cuts to mineral fertiliser applications, if livestock production remains unchanged (MTFR scenario), or through a lower production or a different management of manure (NUTR scenario). This could be achieved applying optimal fertilisation techniques across all Europe, adopting a better synergy between crop and livestock production and/or reducing the livestock intensity. Importantly, in the three scenarios analysed in this study, the reduction of nutrient from point sources corresponds to progressive upgrading of wastewater treatment plants, while the decrease from agricultural sources corresponds to three different strategies that could be combined, yielding to a larger nutrient decline. In addition, measures to reduce nitrogen input from atmospheric deposition were not considered, neither the model includes a feedback mechanism to simulate the positive effect of agricultural measures on limiting nitrogen emission/deposition from the atmosphere.

4.3 Eutrophication and policy targets

Many European seas and coastal areas suffer from problems of hypoxia and eutrophication related to high nutrient loads from the draining basins (Diaz and Rosenberg, 2008; Ménesguen and Lacroix, 2018), with large areas affected in the Baltic Sea (Meier et al., 2019, 2018; Skogen et al., 2014), Greater Northern Sea (Garnier et al., 2019; Lancelot et al., 2011; Passy et al., 2016; Thieu et al., 2010), Mediterranean Sea (Colella et al., 2016; Cozzi et al., 2018; Cozzi and Giani, 2011; Karydis and Kitsiou, 2012; Macias et al., 2018), and Black Sea (Kudryavtseva et al., 2019; Lancelot et al., 2002). Eutrophication is caused by excess

of nitrogen and phosphorus over silica (Billen and Garnier, 2007; Howarth et al., 2011). Generally, nitrogen controls the eutrophication in coastal marine ecosystems (Howarth and Marino, 2006), while phosphorus is considered the limiting nutrient in freshwaters (Elser et al., 2007). The impact of riverine nutrient inputs on eutrophication depend on the receiving waters. In coastal marine ecosystems shelf orography, morphology, water circulation, turbidity, light and salinity, all influence the eutrophication process (Carstensen et al., 2019; Viaroli et al., 2015). For example, the marine region under the freshwater influence is larger for the Po and the Danube than for the Ebro and Rhone rivers because of the shelf orography. These rivers are characterized by an excess of dissolved inorganic nitrogen and silica over phosphate, which is sometimes compensate by the bioavailability of organic phosphorus (Cozzi et al., 2018). Anthropogenic activities have increased the amount of nutrients delivered to the aquatic environment but at the same time have also altered their balance. Similarly, measures to reduce nutrient pollution can create or exacerbate nutrient imbalance in the receiving coastal waters (Howarth and Marino, 2006). For instance, in south-eastern Europe the potential risk for coastal ecosystems eutrophication in relation to nitrogen and phosphorus has changed between the 1990s and the 2000s (Romero et al., 2013). For all these reasons, an important aspect to be look at in the scenarios is the expected changes of the N:P ratio in the nutrient load at the sea. All the scenarios indicate a higher abatement of phosphorus loads than nitrogen loads for the whole area of study (for example -20% P and -14% N in the MTFR scenario), but the impacts are very site specific and should be analysed case by case.

The analysis on the potential change in the ecological status of rivers, lakes and transitional waters suggests that an ambitious scenario (such as the MTFR) would only slightly increase the fraction of river length and water body area in good ecological status, with better effects in regions under high nutrient pressures. However, it has to be noted that the thresholds chosen in the analysis were very conservative, and an average concentration of the 'good ecological status' class rather than the boundary between good and moderate class was adopted for the computation. Also, in the case of freshwaters, the vulnerability of the ecosystem to nutrient enrichment strongly depends on the water body type. Across Europe, good-moderate total nitrogen threshold concentrations of 0.25-4.00 mgN/l (median 1.0 mgN/l) and total phosphorus threshold concentrations of 5-500 µgP/l (median 27.5 µgP/l) were reported per lakes, and good-moderate total nitrogen threshold concentrations of 0.25-35 mgN/l (median 2.5 mgN/l) and total phosphorus threshold concentrations of 8-660 µgP/l (median 100 µgP/l) were reported per rivers (Poikane et al., 2019a), indicating that rivers and lakes with nutrient concentrations higher than the thresholds adopted in the present analysis could also reach good ecological status. In addition, the scenarios results clearly show that the reduction in nutrient concentration mainly concerns concentrations in the 75th percentile, which indicates that ecological condition might improve in many degraded water bodies even without dropping their concentration below the chosen thresholds. Interestingly, the model GREEN captured the distribution of nutrient concentration per water body type and per class of ecological status, reporting increasing average concentrations from lakes, to river and transitional waters for both nitrogen and phosphorus, which generally corresponds to the characteristics observed for these aquatic ecosystems (Poikane et al., 2019a).

The scenarios analysed in this study were conceived to test possible effects of current and future policy actions in the EU. To curb eutrophication in fresh and coastal waters specific targets for nitrogen and phosphorus should be based on the local ecosystem condition, considering the inland sources of nutrient. The results of this study could help in this sense, allowing to link basin specific riverine nutrients per source to nutrient targets of fresh and coastal waters established under the WFD and MFSD (for nutrient targets

see (Poikane et al., 2019b; Salas Herrero et al., 2019)), contemporary checking the coherence of the freshwater and marine policy objectives.

5. Conclusions

The results of the study show that current investments in the EU countries for limiting point and diffuse nutrient pollution might result in a mild reduction of the nutrient load to the European seas, while more ambitious measures could decrease nutrient export to the sea up to 14% for nitrogen and 20% for phosphorus. Further reductions could be possible by a combination of measures especially in the agricultural sector. Importantly, future actions could widen the unbalance between nitrogen and phosphorus in receiving waters, affecting the aquatic ecosystems. In Europe regional differences and ecosystems specificity are present and need to be taken into consideration in the analysis of pressures and impacts, as well as when setting nutrient restoration targets both for freshwater and coastal ecosystems.

The study provides a picture of the major nitrogen and phosphorus sources in European river basins and the consequent pressures on fresh and coastal waters, adopting homogeneous data reported by EU and non-EU countries through standard and/or official data reporting flows. The analysis addressed the needs of model simplicity, data transparency, high spatial resolution and hydrological consistency that allow linking the riverine nutrient sources and possible measures to the ecological targets of the current EU water policies, WFD and MFS, also checking their coherence with other sectoral policies, such as the agricultural policy.

The scenarios analysis highlighted the advantages of adopting a nexus thinking when addressing the land-sea dynamics, with measures taken in several sectors under different policies evaluated together. The scenarios implied measures for water quantity, such as irrigation efficiency and water for energy cooling, and for water quality, including actions under the UWWTP Directive, the Nitrates Directive and the CAP. The relevance of adopting a holistic approach will be key to meet the ambitious policy goals of the new European Green Deal on Zero Pollution, Farm to Fork and Biodiversity Strategy, with the objective of protecting aquatic ecosystems, ensuring their resilience and their sustainable use.

Acknowledgements

The authors declare no conflict of interest.

References

- Arheimer, B., Dahné, J., Donnelly, C., 2012. Climate Change Impact on Riverine Nutrient Load and Land-Based Remedial Measures of the Baltic Sea Action Plan. *Ambio* 41, 600–612.
<https://doi.org/10.1007/s13280-012-0323-0>
- Arnal, L., Asp, S.-S., Baugh, C., Roo, A. de, Disperati, J., Dottori, F., Garcia, R., GarciaPadilla, M., Gelati, E., Gomes, G., Kalas, M., Krzeminski, B., Latini, M., Lorini, V., Mazzetti, C., Mikulickova, M., Muraro, D., Prudhomme, C., Rauthe-Schöch, A., Rehfeldt, K., Sa, P., Ziese, M., 2019. EFAS upgrade for the extended model domain – technical documentation, EUR 29323 EN, Publications Office of the European Union, Luxembourg, ISBN 978-92-79-92881-9, doi:10.2760/806324, JRC111610.
- Bartosova, A., Capell, R., Olesen, J.E., Jabloun, M., Refsgaard, J.C., Donnelly, C., Hyytiäinen, K., Pihlainen, S., Zandersen, M., Arheimer, B., 2019. Future socioeconomic conditions may have a larger impact than climate change on nutrient loads to the Baltic Sea. *Ambio* 48, 1325–1336.
<https://doi.org/10.1007/s13280-019-01243-5>
- Benitez Sanz, C., Wolters, H., Martí, B., Mora, B., 2018. EU Water and Marine Measures Data base. Deliverable to Task B2 of the BLUE2 project “Study on EU integrated policy assessment for the freshwater and marine environment, on the economic benefits of EU water policy and on the costs of its non- implementatio.
- Billen, G., Garnier, J., 2007. River basin nutrient delivery to the coastal sea: Assessing its potential to sustain new production of non-siliceous algae. *Mar. Chem.* 106, 148–160.
<https://doi.org/https://doi.org/10.1016/j.marchem.2006.12.017>
- Black Sea Commission, 2020. Bucharest Convention.
- Bouraoui, F., Thieu, V., Grizzetti, B., Britz, W., Bidoglio, G., 2014. Scenario analysis for nutrient emission reduction in the European inland waters. *Environ. Res. Lett.* 9. <https://doi.org/10.1088/1748-9326/9/12/125007>
- Carstensen, J., Conley, D.J., Almroth-Rosell, E., Asmala, E., Bonsdorff, E., Fleming-Lehtinen, V., Gustafsson, B.G., Gustafsson, C., Heiskanen, A.S., Janas, U., Norkko, A., Slomp, C., Villnäs, A., Voss, M., Zilius, M., 2019. Factors regulating the coastal nutrient filter in the Baltic Sea. *Ambio*.
<https://doi.org/10.1007/s13280-019-01282-y>
- CLC, 2012. CORINE Land Cover Map Analysis.
- Colella, S., Falcini, F., Rinaldi, E., Sammartino, M., Santoleri, R., 2016. Mediterranean Ocean Colour Chlorophyll Trends. *PLoS One* 11, e0155756.
- Cozzi, S., Giani, M., 2011. River water and nutrient discharges in the Northern Adriatic Sea: Current importance and long term changes. *Cont. Shelf Res.* 31, 1881–1893.
<https://doi.org/https://doi.org/10.1016/j.csr.2011.08.010>
- Cozzi, S., Ibáñez, C., Lazar, L., Raimbault, P., Giani, M., 2018. Flow regime and nutrient-loading trends from the largest South European watersheds: Implications for the productivity of mediterranean and Black Sea’s Coastal Areas. *Water (Switzerland)* 11. <https://doi.org/10.3390/w11010001>
- Culhane, F., Teixeira, H., Nogueira, A.J.A., Borgwardt, F., Trauner, D., Lillebø, A., Piet, G., Kuemmerlen, M., McDonald, H., O’Higgins, T., Barbosa, A.L., van der Wal, J.T., Iglesias-Campos, A., Arevalo-Torres, J., Barbière, J., Robinson, L.A., 2019. Risk to the supply of ecosystem services across aquatic

- ecosystems. *Sci. Total Environ.* 660, 611–621.
<https://doi.org/https://doi.org/10.1016/j.scitotenv.2018.12.346>
- De Roo, A., Bisselink, B., Guenther, S., Gelati, E., Adamovic, M., 2020. Assessing the effects of water saving measures on Europe's water resources; BLUE2 project – Freshwater quantity. JRC Technical Report.
- Diaz, R.J., Rosenberg, R., 2008. Spreading dead zones and consequences for marine ecosystems. *Science* (80-.). 321, 926–929. <https://doi.org/10.1126/science.1156401>
- EEA, 2020. Waterbase - Water Quantity v14.
- EEA, 2019a. WISE WFD Reference Spatial Datasets reported under Water Framework Directive 2016 - PUBLIC VERSION - version 1.3, Apr. 2019.
- EEA, 2019b. WISE Water Framework Directive Database.
- EEA, 2018a. Marine regions and subregions under the Marine Strategy Framework Directive.
- EEA, 2018b. E-PRTR.
- EEA, 2012. European catchments and Rivers network system (Ecrins).
- Elser, J.J., Bracken, M.E.S., Cleland, E.E., Gruner, D.S., Harpole, W.S., Hillebrand, H., Ngai, J.T., Seabloom, E.W., Shurin, J.B., Smith, J.E., 2007. Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecol. Lett.* 10, 1135–1142.
<https://doi.org/10.1111/j.1461-0248.2007.01113.x>
- EMEP, 2020. Nitrogen deposition data.
- ESA, 2017. Land Cover CCI Product User Guide Version 2. Tech. Rep.
- European Commission, 2018. Strategic plans, financing, management and monitoring of CAP, common organization of markets - Impact Assessment Part 3. Commission Staff Working Document SWD(2018) 301.
- FAOSTAT, 2020. Fertilizers data.
- Garnier, J., Riou, P., Le Gendre, R., Ramarson, A., Billen, G., Cugier, P., Schapira, M., Théry, S., Thieu, V., Ménesguen, A., 2019. Managing the agri-food system of watersheds to combat coastal eutrophication: A land-to-sea modelling approach to the french coastal English channel. *Geosci.* 9.
<https://doi.org/10.3390/geosciences9100441>
- Grizzetti, B., Bouraoui, F., Aloe, A., 2012. Changes of nitrogen and phosphorus loads to European seas. *Glob. Chang. Biol.* 18. <https://doi.org/10.1111/j.1365-2486.2011.02576.x>
- Grizzetti, B., Bouraoui, F., De Marsily, G., 2008. Assessing nitrogen pressures on European surface water. *Global Biogeochem. Cycles* 22. <https://doi.org/10.1029/2007GB003085>
- Grizzetti, B., Lanzanova, D., Liqueste, C., Reynaud, A., Cardoso, A.C., 2016. Assessing water ecosystem services for water resource management. *Environ. Sci. Policy* 61.
<https://doi.org/10.1016/j.envsci.2016.04.008>
- Grizzetti, B., Liqueste, C., Pistocchi, A., Vigiak, O., Zulian, G., Bouraoui, F., De Roo, A., Cardoso, A.C., 2019. Relationship between ecological condition and ecosystem services in European rivers, lakes and

- coastal waters. *Sci. Total Environ.* 671, 452–465.
<https://doi.org/https://doi.org/10.1016/j.scitotenv.2019.03.155>
- Grizzetti, B., Pistocchi, A., Liqueste, C., Udias, A., Bouraoui, F., Van De Bund, W., 2017. Human pressures and ecological status of European rivers. *Sci. Rep.* 7. <https://doi.org/10.1038/s41598-017-00324-3>
- HELCOM, 2020. Helsinki Convention.
- Howarth, R., Chan, F., Conley, D.J., Garnier, J., Doney, S.C., Marino, R., Billen, G., 2011. Coupled biogeochemical cycles: eutrophication and hypoxia in temperate estuaries and coastal marine ecosystems. *Front. Ecol. Environ.* 9, 18–26. <https://doi.org/10.1890/100008>
- Howarth, R.W., Marino, R., 2006. Nitrogen as the limiting nutrient for eutrophication in coastal marine ecosystems: Evolving views over three decades. *Limnol. Oceanogr.* 51, 364–376.
https://doi.org/10.4319/lo.2006.51.1_part_2.0364
- Jonkers, L., Hillebrand, H., Kucera, M., 2019. Global change drives modern plankton communities away from the pre-industrial state. *Nature*. <https://doi.org/10.1038/s41586-019-1230-3>
- Karydis, M., Kitsiou, D., 2012. Eutrophication and environmental policy in the Mediterranean Sea: a review. *Environ. Monit. Assess.* 184, 4931–4984. <https://doi.org/10.1007/s10661-011-2313-2>
- Krause, P., Boyle, D.P., Bäse, F., 2005. Comparison of different efficiency criteria for hydrological model assessment. *Adv. Geosci.* 5, 89–97. <https://doi.org/10.5194/adgeo-5-89-2005>
- Kuczera, G., Parent, E., 1998. Monte Carlo assessment of parameter uncertainty in conceptual catchment models: The Metropolis algorithm. *J. Hydrol.* 211, 69–85.
- Kudryavtseva, E., Aleksandrov, S., Bukanova, T., Dmitrieva, O., Rusanov, I., 2019. Relationship between seasonal variations of primary production, abiotic factors and phytoplankton composition in the coastal zone of the south-eastern part of the Baltic Sea. *Reg. Stud. Mar. Sci.* 32.
<https://doi.org/10.1016/j.rsma.2019.100862>
- La Notte, A., Maes, J., Dalmazzone, S., Crossman, N.D., Grizzetti, B., Bidoglio, G., 2017. Physical and monetary ecosystem service accounts for Europe: A case study for in-stream nitrogen retention. *Ecosyst. Serv.* 23. <https://doi.org/10.1016/j.ecoser.2016.11.002>
- Lajaunie-Salla, K., Sottolichio, A., Schmidt, S., Litrico, X., Binet, G., Abril, G., 2018. Future intensification of summer hypoxia in the tidal Garonne River (SW France) simulated by a coupled hydro sedimentary-biogeochemical model. *Environ. Sci. Pollut. Res.* 25, 31957–31970.
<https://doi.org/10.1007/s11356-018-3035-6>
- Lancelot, C., Staneva, J., van Eeckhout, D., Beckers, J.-M., Stanev, E., 2002. Modelling the Danube-influenced North-western Continental Shelf of the Black Sea. II: Ecosystem Response to Changes in Nutrient Delivery by the Danube River after its Damming in 1972. *Estuar. Coast. Shelf Sci.* 54, 473–499. <https://doi.org/https://doi.org/10.1006/ecss.2000.0659>
- Lancelot, C., Thieu, V., Polard, A., Garnier, J., Billen, G., Hecq, W., Gypens, N., 2011. Cost assessment and ecological effectiveness of nutrient reduction options for mitigating Phaeocystis colony blooms in the Southern North Sea: An integrated modeling approach. *Sci. Total Environ.* 409, 2179–2191.
<https://doi.org/https://doi.org/10.1016/j.scitotenv.2011.02.023>
- Leip, A., Billen, G., Garnier, J., Grizzetti, B., Lassaletta, L., Reis, S., Simpson, D., Sutton, M.A., De Vries, W.,

- Weiss, F., Westhoek, H., 2015. Impacts of European livestock production: Nitrogen, sulphur, phosphorus and greenhouse gas emissions, land-use, water eutrophication and biodiversity. *Environ. Res. Lett.* 10. <https://doi.org/10.1088/1748-9326/10/11/115004>
- Liquete, C., Piroddi, C., Macías, D., Druon, J.N., Zulian, G., 2016. Ecosystem services sustainability in the Mediterranean Sea: Assessment of status and trends using multiple modelling approaches. *Sci. Rep.* 6. <https://doi.org/10.1038/srep34162>
- Ludwig, W., Bouwman, A.F., Dumont, E., Lespinas, F., 2010. Water and nutrient fluxes from major Mediterranean and Black Sea rivers: Past and future trends and their implications for the basin-scale budgets. *Global Biogeochem. Cycles* 24. <https://doi.org/10.1029/2009GB003594>
- Macias, D., Garcia-Gorriz, E., Stips, A., 2018. Major fertilization sources and mechanisms for Mediterranean Sea coastal ecosystems. *Limnol. Oceanogr.* 63, 897–914. <https://doi.org/10.1002/lno.10677>
- Malagó, A., Bouraoui, F., Grizzetti, B., De Roo, A., 2019. Modelling nutrient fluxes into the Mediterranean Sea. *J. Hydrol. Reg. Stud.* 22. <https://doi.org/10.1016/j.ejrh.2019.01.004>
- Meier, H.E.M., Edman, M., Eilola, K., Placke, M., Neumann, T., Andersson, H.C., Brunnabend, S.-E., Dieterich, C., Frauen, C., Friedland, R., Gröger, M., Gustafsson, B.G., Gustafsson, E., Isaev, A., Kniebusch, M., Kuznetsov, I., Müller-Karulis, B., Naumann, M., Omstedt, A., Ryabchenko, V., Saraiva, S., Savchuk, O.P., 2019. Assessment of Uncertainties in Scenario Simulations of Biogeochemical Cycles in the Baltic Sea. *Front. Mar. Sci.* .
- Meier, H.E.M., Edman, M.K., Eilola, K.J., Placke, M., Neumann, T., Andersson, H.C., Brunnabend, S.-E., Dieterich, C., Frauen, C., Friedland, R., Saraiva, S., Savchuk, O.P., 2018. Assessment of eutrophication abatement scenarios for the baltic sea by multi-model ensemble simulations. *Front. Mar. Sci.* 5. <https://doi.org/10.3389/fmars.2018.00440>
- Ménesguen, A., Lacroix, G., 2018. Modelling the marine eutrophication: A review. *Sci. Total Environ.* 636, 339–354. <https://doi.org/10.1016/J.SCITOTENV.2018.04.183>
- Nash, J.E., Sutcliffe, J.V., 1970. River flow forecasting through conceptual models. Part 1: a discussion of principles. *J. Hydrol.* 10, 282–290.
- Ntegeka, V., Salamon, P., Gomes, G., Sint, H., Lorini, V., Zambrano-Bigiarini, M., Thielen, J., 2013. EFAS-Meteo: A European daily high-resolution gridded meteorological data set for 1990 – 2011. EUR --- Scientific and Technical Research series ---ISSN 1831-9424 (online) Luxembourg: Publications Office of the European Union.
- OSPAR, 2020. OSPAR Convention.
- Passy, P., Le Gendre, R., Garnier, J., Cugier, P., Callens, J., Paris, F., Billen, G., Riou, P., Romero, E., 2016. Eutrophication modelling chain for improved management strategies to prevent algal blooms in the Bay of Seine. *Mar. Ecol. Prog. Ser.* 543, 107–125. <https://doi.org/10.3354/meps11533>
- Piroddi, C., Coll, M., Liquete, C., Macias, D., Greer, K., Buszowski, J., Steenbeek, J., Danovaro, R., Christensen, V., 2017. Historical changes of the Mediterranean Sea ecosystem: Modelling the role and impact of primary productivity and fisheries changes over time. *Sci. Rep.* 7. <https://doi.org/10.1038/srep44491>
- Pistocchi, A., Dorati, C., Grizzetti, B., Udias, A., Vigiak, O., Zanni, M., 2019. Water quality in Europe:

- effects of the Urban Wastewater Treatment Directive. A retrospective and scenario analysis of Dir. 91/271/EEC, EUR 30003 EN, Publications Office of the European Union, Luxembourg, ISBN 978-92-76-11263-1, doi:10.2760/303163, JRC115.
- Poikane, S., Kelly, M.G., Salas Herrero, F., Pitt, J.-A., Jarvie, H.P., Claussen, U., Leujak, W., Lyche Solheim, A., Teixeira, H., Phillips, G., 2019a. Nutrient criteria for surface waters under the European Water Framework Directive: Current state-of-the-art, challenges and future outlook. *Sci. Total Environ.* 695, 133888. <https://doi.org/https://doi.org/10.1016/j.scitotenv.2019.133888>
- Poikane, S., Phillips, G., Birk, S., Free, G., Kelly, M.G., Willby, N.J., 2019b. Deriving nutrient criteria to support 'good' ecological status in European lakes: An empirically based approach to linking ecology and management. *Sci. Total Environ.* 650, 2074–2084. <https://doi.org/https://doi.org/10.1016/j.scitotenv.2018.09.350>
- R Core Team, 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Romero, E., Garnier, J., Lassaletta, L., Billen, G., Le Gendre, R., Riou, P., Cugier, P., 2013. Large-scale patterns of river inputs in southwestern Europe: seasonal and interannual variations and potential eutrophication effects at the coastal zone. *Biogeochemistry* 113, 481–505. <https://doi.org/10.1007/s10533-012-9778-0>
- Salas Herrero, F., Teixeira, H., Poikane, S., 2019. A Novel Approach for Deriving Nutrient Criteria to Support Good Ecological Status: Application to Coastal and Transitional Waters and Indications for Use. *Front. Mar. Sci.* .
- Sarteel, M., Tostivint, C., Landowski, A., Basset, C., Muehmel, K., Lockwood, S., Ding, H., Oudet, N., Mudgal, S., Cherrier, V., Grebot, B., Naumann, S., Dooley, E., Lukat, E., Freluh-Larsen, A., Wunder, S., Carter, M.S., Ambus, P., Provololo, G., Koeijer, T.D., Linderhof, V., Michels, R., 2016. Resource efficiency in practice: closing mineral cycles : final report. - Ebook. - Luxembourg : European Commission, Directorate-General for the Environment, 2016. - ISBN 9789279582387.
- Seitzinger, S.P., Mayorga, E., Bouwman, A.F., Kroeze, C., Beusen, A.H.W., Billen, G., Van Drecht, G., Dumont, E., Fekete, B.M., Garnier, J., Garnier, J., Harrison, J.A., 2010. Global river nutrient export: A scenario analysis of past and future trends. *Global Biogeochem. Cycles* 24. <https://doi.org/10.1029/2009GB003587>
- Skogen, M.D., Eilola, K., Hansen, J.L.S., Meier, H.E.M., Molchanov, M.S., Ryabchenko, V.A., 2014. Eutrophication status of the North Sea, Skagerrak, Kattegat and the Baltic Sea in present and future climates: A model study. *J. Mar. Syst.* 132, 174–184. <https://doi.org/https://doi.org/10.1016/j.jmarsys.2014.02.004>
- Sutton, M., Howard, C., Erisman, J.W., Billen, G., Bleeker, A., Greenfelt, P., van Grinsven, H., Grizzetti, B., 2011. *The European Nitrogen Assessment*. Cambridge University Press, Cambridge.
- Thieu, V., Garnier, J., Billen, G., 2010. Assessing the effect of nutrient mitigation measures in the watersheds of the Southern Bight of the North Sea. *Sci. Total Environ.* 408, 1245–1255. <https://doi.org/https://doi.org/10.1016/j.scitotenv.2009.12.031>
- UNEP, 2020. Barcelona Convention.
- Viaroli, P., Nizzoli, D., Pinardi, M., Soana, E., Bartoli, M., 2015. Eutrophication of the Mediterranean Sea: a watershed—cascading aquatic filter approach. *Rend. Lincei* 26, 13–23.

<https://doi.org/10.1007/s12210-014-0364-3>

Vigiak, O., Grizzetti, B., Zanni, M., Aloe, A., Dorati, C., Bouraoui, F., Pistocchi, A., 2020. Domestic waste emissions to European waters in the 2010s. *Sci. Data* 7. <https://doi.org/10.1038/s41597-020-0367-0>

Vogt, J., Soille, P., de Jager, A., et al, 2007. A pan-European River and Catchment Database. EC-JRC Report EUR 22920 EN. Luxembourg.

Zampieri, M., Grizzetti, B., Meroni, M., Scoccimarro, E., Vrieling, A., Naumann, G., Toreti, A., 2019. Annual green water resources and vegetation resilience indicators: Definitions, mutual relationships, and future climate projections. *Remote Sens.* 11. <https://doi.org/10.3390/rs11222708>

Supplementary Material

S1. Model GREEN equations

For each catchment i in the geo-data model of GREEN the nutrient load L_i is estimated by the **general equation**:

$$L_i = (1 - Lret_i) * (DS_i * (1 - Bret_i) + PS_i + U_i) * (1 - Rret_i) \quad (\text{Equation S1})$$

Where:

L = Nutrient load at the catchment outlet (ton/yr)

DS = Nutrient diffuse sources in the catchment (ton/yr)

PS = Nutrient point sources in the catchment (ton/yr)

U = Nutrient load from upstream catchemnts (ton/yr)

$Lret$ = Lake retention (fraction)

$Bret$ = Basin retention (fraction)

$Rret$ = River retention (fraction)

$$Bret_i = 1 - \exp(-\mathit{basinCoeff} * \text{Inverse of precipitation}_i) \quad (\text{Equation S2})$$

$$Rret_i = 1 - \exp(-\mathit{riverCoeff} * \text{River length}_i) \quad (\text{Equation S3})$$

In the Equation S2 and S3 the inverse of precipitation and the river length are scaled by maximum scaling (Frank and Todeschini, 1994).

The retention occurring in lakes ($Lret$) was computed according to Kronvang et al. (2004), as follows:

$$Lret = 1 - 1/[1+(7.3/z)*RT] \quad (\text{for nitrogen}) \quad (\text{Equation S4})$$

$$Lret = 1 - 1/[1+(26/z)*RT] \quad (\text{for phosphorus}) \quad (\text{Equation S5})$$

Where:

z = average lake depth (m),

RT = hydraulic residence time (yr).

The average lake depth and hydraulic residence time were obtained from HydroLAKES database (<https://www.hydrosheds.org/pages/hydrolakes>, Messenger et al. 2016).

In **GREEN nitrogen model**, for each catchment i the total nitrogen load L_i is estimated by the equation:

$$L_i = (1 - R_{ret,i}) * [(MinN_i + ManN_i + FixN_i + SoilN_i + (1 - FF_i) * AtmN_i) * (1 - R_{ret,i}) + 0.38 * FF_i * AtmN_i + 0.5 * SdN_i + PsN_i + U_i] * (1 - R_{ret,i}) \quad (\text{Equation S6})$$

Where:

MinN = Nitrogen mineral fertilisers (ton/yr)

ManN = Nitrogen in manure fertilisers (ton/yr)

FixN = Nitrogen fixation by leguminous crops and fodder (ton/yr)

SoilN = Nitrogen fixation by bacteria in soils (ton/yr)

AtmN = Nitrogen deposition from atmosphere (ton/yr)

SdN = Nitrogen input from scattered dwellings (ton/yr)

PsN = Nitrogen input from point sources (ton/yr)

U = Nitrogen load from upstream catchments (ton/yr)

FF = Non-agricultural land cover in the catchment (fraction)

Input from scattered dwelling (SD) are estimated to be reduced by 50% before entering the river

Background losses for nitrogen are estimated as $0.38 * FF * AtmN$. For an atmospheric deposition of 10 kgN/ha this corresponds to a background of 3.8 kgN/ha (in line with the values reported by HELCOM, 2003).

In **GREEN phosphorus model**, for each catchment i the total phosphorus load L_i is estimated by the equation:

$$L_i = (1-Rret_i) * [(MinP_i + ManP_i + (1-FF_i)*BgP_i)*(1-Bret_i) + FF_i*BgP_i + 0.5*SdP_i + PsP_i + U_i] * (1-Rret_i) \quad \text{(Equation S7)}$$

MinP = Phosphorus mineral fertilisers (ton/yr)

ManP = Phosphorus in manure fertilisers (ton/yr)

BgP = Phosphorus background losses (ton/yr)

SdP = Phosphorus input from scattered dwellings (ton/yr)

PsP = Phosphorus input from point sources (ton/yr)

U = Phosphorus load from upstream catchments (ton/yr)

FF = Non-agricultural land cover in the catchment (fraction)

Input from scattered dwelling (SD) are estimated to be reduced by 50% before entering the river.

Background losses for phosphorus are estimated at 0.15 kgP/ha (in line with the values reported by HELCOM, 2003).

References

Frank, I.E. and Todeschini, R., 1994. The data analysis handbook. Elsevier, Amsterdam, the Neatherlands.

Kronvang, B., Hezlar, J., Boers, P., Jansen, J. P., Behrendt, H., Anderson, T., Arheimer, B., Venohr, M., Hoffmann, C.C., 2004. Nutrient retention handbook. Software manual for EUROHARP NUTRET and scientific review on nutrient retention, no. 9-2004 in EUROHARP report, Oslo, <http://www.euroharp.org>.

HELCOM, Helsinki Commission, 2003. Fourth Baltic Sea Load Compilation. Balt. Sea Environ. Proc. No. 93. 189pp.

Messenger, M.L., Lehner, B., Grill, G., Nedeva, I., Schmitt, O., 2016: Estimating the volume and age of water stored in global lakes using a geo-statistical approach. Nature Communications: 13603. doi: 10.1038/ncomms13603

S2. Number of observations for model calibration

Table S2.1 Number and average area of catchments in the geo-spatial model of GREEN

Number of catchments	Total Area (km ²)	Average catchment area (km ²)
950472	6327575	6.66

Table S2.2 Number of observed nitrogen and phosphorus concentration per year available for the calibration of the model GREEN.

Year	Number observations Total Nitrogen	Number observations Total Phosphorus
2008	2492	2837
2009	2018	2186
2010	1564	2894
2011	1578	2941
2012	1683	3032


Table S2.3 Number of observed nitrogen and phosphorus concentration per country per year available for the calibration of the model GREEN.

Country	N. catchments	Area (km ²)	Number observations Total Nitrogen					Number observations Total Phosphorus				
			2008	2009	2010	2011	2012	2008	2009	2010	2011	2012
AD	71	93	0	0	0	0	0	0	0	0	0	0
AL	2761	5676	0	0	0	0	0	17	4	4	3	3
AT	7354	16768	3	3	3	3	2	40	45	45	39	39
BA	3340	10222	2	11	9	11	14	10	11	9	11	14
BE	481	6149	39	40	39	66	66	39	40	39	66	66
BG	3308	22219	32	29	36	52	54	2	31	32	52	55
BY	1956	17624	0	0	0	0	0	0	0	0	0	0
CH	3979	8286	6	0	6	48	48	8	7	8	88	87
CY	179	1850	12	15	0	0	0	12	15	17	15	18
CZ	2032	15784	56	0	0	0	0	57	1	1	1	1

DE	7394	71457	176	171	177	183	152	179	199	209	207	190
DK	493	8631	19	19	19	18	18	19	19	19	18	18
EE	707	9094	44	44	45	41	43	44	44	45	41	43
ES	17119	99576	166	178	184	0	80	197	169	246	87	92
FI	9388	67878	93	87	85	84	85	93	88	86	85	85
FR	17879	109755	1000	862	1	2	2	1001	854	1013	1015	1015
GB	5586	48865	147	157	216	206	176	154	161	233	227	199
GG	3	14	0	0	0	0	0	0	0	0	0	0
GR	8894	26311	0	0	0	0	0	43	0	0	0	0
HR	2453	11271	17	16	15	15	16	17	16	15	15	16
HU	1123	18519	7	6	5	5	5	8	8	6	5	5
IE	1697	13962	0	0	12	12	15	0	0	20	17	25
IM	14	114	0	0	0	0	0	0	0	0	0	0
IT	18650	59997	350	89	416	462	430	484	90	500	553	537
JE	2	24	0	0	0	0	0	0	0	0	0	0
KS	410	2172	0	0	0	0	0	19	19	18	18	18
LI	8	34	0	0	0	0	0	2	2	2	2	0
LT	1181	13132	30	30	32	35	32	30	30	32	35	32
LU	63	514	0	0	0	3	3	3	3	3	3	3
LV	1013	12848	26	23	2	2	2	26	23	2	2	2
MD	225	2930	0	0	0	0	0	0	0	0	0	0
ME	1315	2857	0	0	0	0	0	0	0	0	0	0
MK	1555	5052	0	0	0	0	0	0	0	0	0	0
MT	5	63	0	0	0	0	0	0	0	0	0	0
NL	216	6918	10	10	10	4	4	10	10	10	4	4
NO	11763	64782	29	29	29	28	48	29	29	29	28	49
PL	5107	62372	23	21	5	81	170	24	20	5	81	168
PT	2815	17709	14	12	14	3	10	17	29	30	9	34
RO	6092	47769	48	49	58	64	65	71	71	70	67	69
RS	1971	15539	34	43	44	43	32	43	43	44	43	32
RU	19148	142092	1	1	1	1	2	1	1	1	1	2
SE	9271	89903	46	44	76	74	74	75	74	76	74	74
SI	1331	4060	8	6	9	11	4	8	6	8	10	5
SK	2009	9735	53	22	15	20	29	54	23	16	18	30
SM	2	13	0	0	0	0	0	0	0	0	0	0
TR	6563	102218	0	0	0	0	0	0	0	0	0	0
UA	948	9492	1	1	1	1	2	1	1	1	1	2


S3. Model calibration

Figure S3.1 GREEN model calibration: Nitrogen loads



Period of calibration	Number iterations	basinCoeff interval	riverCoeff interval	sdCoeff	basinCoeff	riverCoeff	NSE (noLOG)
2008-2012	500	30-50	0.005-0.1	0.666667	35.8880	0.0083	0.9614

Figure S3.2 GREEN model calibration: Phosphorus loads



Period of calibration	Number iterations	basinCoeff interval	riverCoeff interval	sdCoeff	basinCoeff	riverCoeff	NSE (noLOG)
2008-2012	500	40-75	0.005-0.1	0.714286	58.1965	0.0350	0.7483

S4. Model validation

Figure S4.1 The maps of the 50 European rivers (with the largest nitrogen load) where freshwater runoff and nutrient loads entering the European seas from the reference simulation of the model GREEN were compared with a variety of observed data from independent sources.




Table S4.1 Sources of data of water runoff and nutrient loads used to compared the results of the model GREEN.

Reference	Web link
Global Runoff Data Centre (GRDC)	https://www.bafg.de/GRDC/EN/01_GRDC/grdc_node.html
Global Nutrient Export from WaterSheds 2 (NEWS 2; (Mayorga et al., 2010))	https://doi.org/10.1016/j.envsoft.2010.01.007
European Environment Agency (EEA)	https://www.eea.europa.eu/data-and-maps/data/waterbase-rivers-10
Helsinki Commission (HELCOM; HELCOM (2015):	https://helcom.fi/wp-content/uploads/2019/08/BSEP145_Lowres.pdf
HELCOM (2018)	https://helcom.fi/wp-content/uploads/2019/12/BSEP163.pdf
HELCOM Map and Data Service	http://maps.helcom.fi/website/mapservice/
OSPAR Data and Information Management System	https://odims.ospar.org/
UK National River Flow Archive	https://nrfa.ceh.ac.uk/data/search
Hydro-Data provided by the Irish Office of Public Works	http://waterlevel.ie/hydro-data/list.html#
OSPAR Intersessional Correspondence Group on Eutrophication Modelling (ICG-EMO; provided by Sonja van Leeuwen (NIOZ, pers. comm.), extension of Lenhart et al. (2010)	https://doi.org/10.1016/j.jmarsys.2009.12.014

Table S4.2. Literature sources to compare freshwater runoff and nutrient loads of selected rivers with the results of the model GREEN.

Reference	River	Web link
(Cozzi et al., 2018)	Danube, Ebro, Po, Rhone	doi:10.3390/w11010001
(Friedland et al., 2019)	Oder	https://doi.org/10.3389/fmars.2018.00521
(Hartmann et al., 2011)	Rhine	https://doi.org/10.1007/s10201-010-0322-4
(Hesse and Krysanova, 2016)	Elbe	https://doi.org/10.3390/w8020040
(Howden et al., 2010)	Thames	DOI: 10.1002/hyp.7835
(Karydis and Kitsiou, 2012)	Adige, Drin, Ebro, Po, Rhone, Tiber	DOI 10.1007/s10661-011-2313-2
(Kauppila and Koskiaho, 2003)	Kemijoki	https://doi.org/10.2166/nh.2003.0004
(Lajaunie-Salla et al., 2018)	Garonne	https://doi.org/10.1007/s11356-018-3035-6

(Lassaletta et al., 2012)	Ebro	doi:10.5194/bg-9-57-2012
(Ludwig et al., 2010)	Adige, Arno, Danube, Ebro, Po, Rhone	doi:10.1029/2009GB003594
(Ménesguen et al., 2019)	Dordogne, Garonne, Loire, Seine	https://doi.org/10.1016/j.ocemod.2018.11.002
(Minaudo et al., 2015)	Loire	doi:10.5194/bg-12-2549-2015
(Mockler et al., 2017)	Shannon	http://dx.doi.org/10.1016/j.scitotenv.2017.05.186
(Passy et al., 2013)	Scheldt, Seine, Somme	http://dx.doi.org/10.1016/j.jmarsys.2013.05.005
(Passy et al., 2016)	Seine	doi: 10.3354/meps11533
(Petus et al., 2014)	Adour	http://dx.doi.org/10.1016/j.csr.2013.11.011
(Tockner et al., 2009)	Danube, Daugava, Duoro, Ebro, Elbe, Loire, Nemunas, Neva, Oder, Rhine, Rhone, Vistula	Tockner, Klement, Urs Uehlinger, and Christopher T. Robinson. Rivers of Europe. Academic Press, 2009.
(Radach and Pätsch, 2007)	Elbe, Ems, Rhine, Weser	DOI: 10.1007/BF02782968
(Romero et al., 2013)	Adige, Adour, Arno, Dordogne, Duoro, Ebro, Garonne, Loire, Po, Rhone, Scheldt, Seine, Somme, Tagus, Tiber, Vilaine	DOI 10.1007/s10533-012-9778-0
(Skarbøvik et al., 2014)	Black Drin	DOI:10.2298/ABS1402667S
(Thieu et al., 2010)	Scheldt, Seine, Somme	doi:10.1016/j.scitotenv.2009.12.031
(Valsecchi et al., 2015)	Adige, Arno, Po	http://dx.doi.org/10.1016/j.chemosphere.2014.07.044
(Vybernaite-Lubiene et al., 2018)	Nemunas	doi:10.3390/w10091178
(Ylöstalo et al., 2016)	Neva	http://dx.doi.org/10.1016/j.marchem.2016.07.004

Figure S4.2 Comparison of reported freshwater runoff with GREEN model [m³ s⁻¹] (derived from the model LISFLOOD), shown on a log-log-scale, color-coded are the different sources for the reported values.




Figure S4.3 Comparison of reported annual Total Phosphorus load with GREEN model [t a-1], shown on a log-log-scale, color-coded are the different sources for the reported values.




Table S4.3 Correlation coefficients (R2) and gradients of a linear fit (a) for freshwater runoff and nutrient loads between GREEN model and different data sources (listed in Table S4.1 and Table S4.2). The correlation was computed for the whole data set as well as for the different data sources separately. The linear fit between reported and modeled freshwater runoff and nutrient loads was computed, assuming the intersect at 0. *Data sources from the literature per river basin are provided in Table S4.2.

	Freshwater Runoff		Total Nitrogen load		Total Phosphorus load	
	R2	a	R2	a	R2	a
All data	0.96	1.079	0.90	0.961	0.49	0.565
All data without NEWS2					0.80	0.993
NEWS2 (global database)	0.95	1.079	0.91	1.084	0.72	0.371
GRDC (Global Runoff Database)	0.97	1.086				
OSPAR ICG EMO	0.89	1.174	0.74	1.085	0.63	1.078
Reported by HELCOM or OSPAR	0.90	1.210	0.85	1.020	0.90	1.146
EEA data			0.93	0.969	0.75	1.006
Data from the literature*	0.96	1.059	0.94	0.894	0.84	0.968

References

- Cozzi, S., Ibáñez, C., Lazar, L., Raimbault, P., Giani, M., 2018. Flow regime and nutrient-loading trends from the largest South European watersheds: Implications for the productivity of mediterranean and Black Sea's Coastal Areas. *Water (Switzerland)* 11. <https://doi.org/10.3390/w11010001>
- Friedland, R., Schernewski, G., Gräwe, U., Greipsland, I., Palazzo, D., Pastuszek, M., 2019. Managing eutrophication in the Szczecin (Oder) lagoon-development, present state and future perspectives. *Front. Mar. Sci.* 5. <https://doi.org/10.3389/fmars.2018.00521>
- Hartmann, J., Levy, J., Kempe, S., 2011. Increasing dissolved silica trends in the Rhine River: An effect of recovery from high P loads? *Limnology* 12, 63–73. <https://doi.org/10.1007/s10201-010-0322-4>
- Hesse, C., Krysanova, V., 2016. Modeling climate and management change impacts on water quality and in-stream processes in the Elbe river basin. *Water (Switzerland)* 8. <https://doi.org/10.3390/w8020040>
- Howden, N.J.K., Burt, T.P., Worrall, F., Whelan, M.J., Bierozza, M., 2010. Nitrate concentrations and fluxes in the River Thames over 140 years (1868-2008): Are increases irreversible? *Hydrol. Process.* 24, 2657–2662. <https://doi.org/10.1002/hyp.7835>
- Karydis, M., Kitsiou, D., 2012. Eutrophication and environmental policy in the Mediterranean Sea: a review. *Environ. Monit. Assess.* 184, 4931–4984. <https://doi.org/10.1007/s10661-011-2313-2>
- Kaupilla, P., Koskiaho, J., 2003. Evaluation of Annual Loads of Nutrients and Suspended Solids in Baltic Rivers. *Hydrol. Res.* 34, 203–220. <https://doi.org/10.2166/nh.2003.0004>

- Lajaunie-Salla, K., Sottolichio, A., Schmidt, S., Litrico, X., Binet, G., Abril, G., 2018. Future intensification of summer hypoxia in the tidal Garonne River (SW France) simulated by a coupled hydro sedimentary-biogeochemical model. *Environ. Sci. Pollut. Res.* 25, 31957–31970. <https://doi.org/10.1007/s11356-018-3035-6>
- Lassaletta, L., Romero, E., Billen, G., Garnier, J., García-Gómez, H., Rovira, J.V., 2012. Spatialized N budgets in a large agricultural Mediterranean watershed: High loading and low transfer. *Biogeosciences* 9, 57–70. <https://doi.org/10.5194/bg-9-57-2012>
- Ludwig, W., Bouwman, A.F., Dumont, E., Lespinas, F., 2010. Water and nutrient fluxes from major Mediterranean and Black Sea rivers: Past and future trends and their implications for the basin-scale budgets. *Global Biogeochem. Cycles* 24. <https://doi.org/10.1029/2009GB003594>
- Mayorga, E., Seitzinger, S.P., Harrison, J.A., Dumont, E., Beusen, A.H.W., Bouwman, A.F., Fekete, B.M., Kroeze, C., Van Drecht, G., 2010. Global Nutrient Export from WaterSheds 2 (NEWS 2): Model development and implementation. *Environ. Model. Softw.* 25, 837–853. <https://doi.org/10.1016/j.envsoft.2010.01.007>
- Ménesguen, A., Dussauze, M., Dumas, F., Thouvenin, B., Garnier, V., Lecornu, F., Répécaud, M., 2019. Ecological model of the Bay of Biscay and English Channel shelf for environmental status assessment part 1: Nutrients, phytoplankton and oxygen. *Ocean Model.* 133, 56–78. <https://doi.org/10.1016/j.oceanmod.2018.11.002>
- Minaudo, C., Meybeck, M., Moatar, F., Gassama, N., Curie, F., 2015. Eutrophication mitigation in rivers: 30 years of trends in spatial and seasonal patterns of biogeochemistry of the Loire River (1980–2012). *Biogeosciences* 12, 2549–2563. <https://doi.org/10.5194/bg-12-2549-2015>
- Mockler, E.M., Deakin, J., Archbold, M., Gill, L., Daly, D., Bruen, M., 2017. Sources of nitrogen and phosphorus emissions to Irish rivers and coastal waters: Estimates from a nutrient load apportionment framework. *Sci. Total Environ.* 601–602, 326–339. <https://doi.org/10.1016/j.scitotenv.2017.05.186>
- Passy, P., Gypens, N., Billen, G., Garnier, J., Thieu, V., Rousseau, V., Callens, J., Parent, J.-Y., Lancelot, C., 2013. A model reconstruction of riverine nutrient fluxes and eutrophication in the Belgian Coastal Zone since 1984. *J. Mar. Syst.* 128, 106–122. <https://doi.org/10.1016/j.jmarsys.2013.05.005>
- Passy, P., Le Gendre, R., Garnier, J., Cugier, P., Callens, J., Paris, F., Billen, G., Riou, P., Romero, E., 2016. Eutrophication modelling chain for improved management strategies to prevent algal blooms in the Bay of Seine. *Mar. Ecol. Prog. Ser.* 543, 107–125. <https://doi.org/10.3354/meps11533>
- Petus, C., Marieu, V., Novoa, S., Chust, G., Bruneau, N., Froidefond, J.-M., 2014. Monitoring spatio-temporal variability of the Adour River turbid plume (Bay of Biscay, France) with MODIS 250-m imagery. *Cont. Shelf Res.* 74, 35–49. <https://doi.org/10.1016/j.csr.2013.11.011>
- Radach, G., Pätsch, J., 2007. Variability of continental riverine freshwater and nutrient inputs into the North Sea for the years 1977–2000 and its consequences for the assessment of eutrophication. *Estuaries and Coasts* 30, 66–81. <https://doi.org/10.1007/BF02782968>
- Romero, E., Garnier, J., Lassaletta, L., Billen, G., Le Gendre, R., Riou, P., Cugier, P., 2013. Large-scale patterns of river inputs in southwestern Europe: seasonal and interannual variations and potential eutrophication effects at the coastal zone. *Biogeochemistry* 113, 481–505. <https://doi.org/10.1007/s10533-012-9778-0>

- Skarbøvik, E., Perović, A., Shumka, S., Nagothu, U.S., 2014. Nutrient inputs, trophic status and water management challenges in the transboundary Lake Skadar/Shkodra, Western Balkans. *Arch. Biol. Sci.* 66, 667–681.
- Thieu, V., Garnier, J., Billen, G., 2010. Assessing the effect of nutrient mitigation measures in the watersheds of the Southern Bight of the North Sea. *Sci. Total Environ.* 408, 1245–1255. <https://doi.org/https://doi.org/10.1016/j.scitotenv.2009.12.031>
- Tockner, K., Uehlinger, U., Robinson, C.T., 2009. *Rivers of Europe*. Academic Press.
- Valsecchi, S., Rusconi, M., Mazzoni, M., Viviano, G., Pagnotta, R., Zaghi, C., Serrini, G., Polesello, S., 2015. Occurrence and sources of perfluoroalkyl acids in Italian river basins. *Chemosphere* 129, 126–134. <https://doi.org/https://doi.org/10.1016/j.chemosphere.2014.07.044>
- Vybernaite-Lubiene, I., Zilius, M., Saltyte-Vaisiauske, L., Bartoli, M., 2018. Recent Trends (2012–2016) of N, Si, and P Export from the Nemunas River Watershed: Loads, Unbalanced Stoichiometry, and Threats for Downstream Aquatic Ecosystems. *Water* . <https://doi.org/10.3390/w10091178>
- Ylöstalo, P., Seppälä, J., Kaitala, S., Maunula, P., Simis, S., 2016. Loadings of dissolved organic matter and nutrients from the Neva River into the Gulf of Finland – Biogeochemical composition and spatial distribution within the salinity gradient. *Mar. Chem.* 186, 58–71. <https://doi.org/https://doi.org/10.1016/j.marchem.2016.07.004>

S5. Nutrient inland input under different scenarios

Table S5.1 Nitrogen inland input per regional seas (per source and scenario).

Nitrogen inland input per regional seas (per source and scenario)								
Scenario	Atmospheric deposition	Scattered dwellings	Point sources	% of change	Agriculture (Min, Man, BNF)	% of change	Total Input	% of change
	(ton/y)	(ton/y)	(ton/y)		(ton/y)		(ton/y)	
Baltic								
REF	989390	33773	90578		3365835		4479576	
BAU	971834	33730	88588	-2	3312010	-2	4406162	-2
NUTR	989390	32746	86179	-5	3298474	-2	4406788	-2
MTFR	989390	32746	83618	-8	2618555	-22	3724308	-17
Greater North Sea								
REF	1120137	15805	245245		6284123		7665310	
BAU	1108643	15805	245200	0	6217190	-1	7586839	-1
NUTR	1120137	13161	235980	-4	5422629	-14	6791907	-11
MTFR	1120137	13161	214615	-12	4857025	-23	6204937	-19
Celtic Sea								
REF	151298	1755	55090		1635782		1843925	
BAU	148476	1755	52685	-4	1604992	-2	1807908	-2
NUTR	151298	1253	48805	-11	1592667	-3	1794024	-3
MTFR	151298	1253	37532	-32	1367870	-16	1557953	-16
Bay of Biscay and Iberian Coast								
REF	444798	8008	79653		2970126		3502585	
BAU	439683	7947	75373	-5	2934104	-1	3457107	-1
NUTR	444798	7908	65736	-17	2963165	0	3481606	-1
MTFR	444798	7908	55760	-30	2313345	-22	2821811	-19
Black Sea								
REF	959768	38907	178796		3820915		4998386	
BAU	947052	38405	151675	-15	3782809	-1	4919940	-2
NUTR	959768	36305	143400	-20	3724705	-3	4864178	-3
MTFR	959768	36305	141393	-21	3212028	-16	4349494	-13
Aegean-Levantine Sea								
REF	213160	14123	54715		969524		1251522	
BAU	211992	14123	52772	-4	963854	-1	1242740	-1
NUTR	213160	13011	52447	-4	967552	0	1246170	0
MTFR	213160	13011	52277	-4	813571	-16	1092019	-13
Ionian Sea and Central Med Sea								
REF	40239	4939	15504		238346		299028	
BAU	39425	4933	14911	-4	233376	-2	292645	-2
NUTR	40239	3229	11218	-28	235555	-1	290241	-3
MTFR	40239	3229	10441	-33	135060	-43	188969	-37
Adriatic Sea								
REF	346900	11802	50053		884785		1293540	
BAU	343854	11665	46745	-7	876638	-1	1278902	-1
NUTR	346900	9682	40931	-18	848085	-4	1245597	-4
MTFR	346900	9682	39632	-21	534385	-40	930598	-28
Western Med Sea								
REF	313139	6087	98436		1262171		1679833	
BAU	310041	6053	96028	-2	1249788	-1	1661910	-1
NUTR	313139	5072	81316	-17	1224325	-3	1623852	-3
MTFR	313139	5072	65512	-33	837689	-34	1221412	-27

Table S5.2 Phosphorus inland input per regional seas (per source and scenario).

Phosphorus inland input per regional seas (per source and scenario)								
	Background	Scattered dwellings	Point sources		Agriculture (Min, Man)		Total Input	
Scenario	(ton/y)	(ton/y)	(ton/y)	% of change	(ton/y)	% of change	(ton/y)	% of change
Baltic								
REF	24788	7790	12768		530411		575758	
BAU	24788	7777	12095	-5	521432	-2	566092	-2
NUTR	24788	7538	11268	-12	507270	-4	550864	-4
MTFR	24788	7538	10329	-19	442224	-17	484879	-16
Greater North Sea								
REF	14315	2995	32323		1033242		1082874	
BAU	14315	2995	32300	0	1022322	-1	1071932	-1
NUTR	14315	2478	31229	-3	738886	-28	786908	-27
MTFR	14315	2474	23205	-28	938456	-9	978450	-10
Celtic Sea								
REF	2929	301	9281		290097		302608	
BAU	2929	301	8867	-4	284694	-2	296792	-2
NUTR	2929	241	9153	-1	279416	-4	291739	-4
MTFR	2929	213	4441	-52	267602	-8	275185	-9
Bay of Biscay and Iberian Coast								
REF	9909	1735	15884		566851		594379	
BAU	9909	1720	14494	-9	559545	-1	585669	-1
NUTR	9909	1714	14125	-11	564704	0	590451	-1
MTFR	9909	1710	7482	-53	456045	-20	475146	-20
Black Sea								
REF	16204	7965	28502		616570		669240	
BAU	16204	7868	23064	-19	609207	-1	656342	-2
NUTR	16204	7409	23618	-17	576299	-7	623529	-7
MTFR	16204	7353	20102	-29	564287	-8	607946	-9
Aegean-Levantine Sea								
REF	5517	2897	12236		178358		199008	
BAU	5517	2897	11747	-4	177509	0	197670	-1
NUTR	5517	2655	11760	-4	177890	0	197821	-1
MTFR	5517	2646	9845	-20	159501	-11	177509	-11
Ionian Sea and Central Med Sea								
REF	1214	810	2441		37761		42226	
BAU	1214	809	2346	-4	36989	-2	41357	-2
NUTR	1214	525	2296	-6	36806	-3	40841	-3
MTFR	1214	522	815	-67	26250	-30	28801	-32
Adriatic Sea								
REF	3572	2321	7881		202626		216401	
BAU	3572	2294	6741	-14	200848	-1	213455	-1
NUTR	3572	1956	6128	-22	186650	-8	198306	-8
MTFR	3572	1933	3906	-50	154648	-24	164059	-24
Western Med Sea								
REF	6409	1150	17930		242760		268249	
BAU	6409	1145	17354	-3	240245	-1	265152	-1
NUTR	6409	986	16624	-7	231508	-5	255526	-5
MTFR	6409	985	6863	-62	179053	-26	193310	-28

Table S5.3 Nitrogen inland input in EU28 under REF and MTRF scenarios.

Nitrogen inland input in EU28 countries (per source and scenario MTRF)										
Country	Scenario REF (ton)					Changes in scenario MTRF compared to scenario REF (%)				
	Atmospheric deposition	Scattered dwellings	Point sources	Agriculture	Total input	Atmospheric deposition	Scattered dwellings	Point sources	Agriculture	Total input
AT	107290	0	15487	256670	379447	0	-40	0	-3	-2
BE	52005	2502	11528	382658	448693	0	0	-17	-31	-26
BG	74982	1159	10761	293765	380667	0	0	-46	-22	-19
CY	2959	673	1099	15262	19993	0	-2	-27	-26	-22
CZ	95376	4822	7546	449503	557248	0	-14	-13	-51	-41
DE	582254	3259	83122	2658311	3326944	0	-64	-1	-24	-19
DK	49490	0	9113	440232	498835	0	0	-1	-25	-22
EE	27418	1	1052	84117	112588	0	-40	-2	-24	-18
ES	241078	228	73957	1697343	2012607	0	-47	-34	-43	-37
FI	103108	2599	7931	222687	336325	0	0	-1	-29	-19
FR	603176	16655	71311	3583686	4274828	0	0	-14	-9	-8
GB	194772	744	109117	1761634	2066267	0	-52	-35	-33	-30
GR	78778	5172	8810	372337	465097	0	-25	-4	-48	-39
HR	57536	576	8780	166449	233341	0	-64	-67	-63	-47
HU	84353	3602	8389	442588	538932	0	-35	-19	-37	-31
IE	50028	1108	7810	727191	786137	0	-40	-50	0	-1
IT	373198	9510	91189	1187167	1661064	0	-47	-28	-43	-32
LT	58698	2548	2225	241232	304703	0	-10	-1	-19	-15
LU	4045	40	503	26626	31214	0	-18	-8	-45	-39
LV	45071	1545	1345	97957	145917	0	-8	-24	-19	-13
MT	109	0	655	2724	3488	0	0	-3	-45	-36
NL	74042	263	18159	590486	682951	0	0	-1	-25	-22
PL	352123	18643	39566	1594469	2004801	0	-2	-16	-28	-23
PT	33459	664	20319	186077	240520	0	0	-41	-35	-31
RO	187531	1778	51876	784563	1025747	0	-5	-53	-11	-11
SE	158667	0	12611	298413	469692	0	0	0	-14	-9
SI	27698	2160	2362	42294	74515	0	-6	-49	-26	-16
SK	47480	4453	4133	161768	217834	0	-16	-21	-26	-20
Total EU28	3766724	84702	680757	18768211	23300394	0	-15	-24	-25	-21

Table S5.4 Phosphorus inland input in EU28 under REF and MTR scenarios.

Phosphorus inland input in EU28 countries (per source and scenario MTR)										
Country	Scenario REF (ton)					Changes in scenario MTR compared to scenario REF (%)				
	Background	Scattered dwellings	Point sources	Agriculture	Total input	Background	Scattered dwellings	Point sources	Agriculture	Total input
AT	1258	0	1265	61169	63692	0	-42	-1	-2	-2
BE	461	380	1480	87678	89999	0	0	-25	-7	-7
BG	1667	209	1623	33197	36695	0	0	-67	-16	-17
CY	139	217	313	4132	4801	0	-2	-42	-21	-21
CZ	1184	1043	1032	47709	50968	0	-16	-30	-26	-25
DE	5360	513	7198	445985	459056	0	-74	-7	-10	-10
DK	647	0	883	64753	66283	0	0	-1	-5	-5
EE	682	0	113	10571	11366	0	-42	-3	-20	-19
ES	7470	52	15338	370825	393686	0	-52	-59	-35	-35
FI	5090	416	523	31831	37860	0	0	-3	-15	-12
FR	8234	3588	11947	591610	615379	0	0	-43	-5	-6
GB	3665	165	20048	276421	300299	0	-59	-56	-18	-20
GR	1974	1075	2936	57175	63161	0	-27	-67	-39	-39
HR	845	179	2631	23647	27303	0	-74	-83	-46	-49
HU	1389	761	1200	58865	62215	0	-38	-42	-23	-23
IE	1047	161	1035	141523	143767	0	-46	-70	0	-1
IT	4502	1425	12642	250632	269200	0	-52	-59	-26	-27
LT	985	435	208	33332	34960	0	-12	-2	-15	-14
LU	39	6	45	4910	4999	0	-21	-17	-18	-18
LV	964	286	176	16800	18226	0	-9	-39	-17	-16
MT	5	0	274	646	924	0	0	-50	-21	-30
NL	519	37	1679	116145	118380	0	0	-6	-4	-4
PL	4678	5277	6790	319049	335795	0	-2	-34	-20	-20
PT	1329	150	4477	43652	49608	0	0	-55	-29	-30
RO	3583	290	8341	127108	139323	0	-5	-65	-7	-11
SE	6743	0	1043	42355	50141	0	0	0	-7	-6
SI	305	346	356	12801	13807	0	-7	-64	-15	-16
SK	730	888	540	15396	17555	0	-18	-46	-16	-16
Total EU28	65494	17898	106137	3289917	3479447	0	-14	-49	-16	-16

S6. Nutrient loads to European regional seas under different scenarios

Table S6.1 Nitrogen load at sea outlets (per source and scenario) estimated by the model GREEN.

Nitrogen load at sea outlets (per source and scenario)								
Scenario	Atmospheric deposition (ton/y)	Scattered dwellings (ton/y)	Point sources (ton/y)	% of change	Agriculture (Min, Man, BNF) (ton/y)	% of change	Total load (ton/y)	% of change
Baltic								
REF	203992	20171	83561		217699		525423	
BAU	200051	20145	81745	-2	213669	-2	515608	-2
NUTR	203963	19530	79532	-5	213140	-2	516165	-2
MTFR	203962	19530	77214	-8	171710	-21	472416	-10
Greater North Sea								
REF	226159	9815	236788		655866		1128627	
BAU	223834	9815	236746	0	648476	-1	1118870	-1
NUTR	226156	8137	227692	-4	564141	-14	1026126	-9
MTFR	226156	8137	206589	-13	509292	-22	950173	-16
Celtic Sea								
REF	36655	1149	54908		324813		417525	
BAU	35968	1149	52503	-4	318664	-2	408284	-2
NUTR	36654	818	48631	-11	316696	-2	402798	-4
MTFR	36654	818	37382	-32	265481	-18	340335	-18
Bay of Biscay and Iberian Coast								
REF	86320	5132	75714		209743		376908	
BAU	85332	5092	71677	-5	207585	-1	369686	-2
NUTR	86318	5066	62208	-18	209225	0	362817	-4
MTFR	86317	5066	52697	-30	178043	-15	322123	-15
Black Sea								
REF	163158	21202	149583		191089		525033	
BAU	160717	20937	125850	-16	188359	-1	495863	-6
NUTR	163157	19873	118845	-21	183064	-4	484939	-8
MTFR	163156	19873	117262	-22	162536	-15	462828	-12
Aegean-Levantine Sea								
REF	50016	9016	52978		81797		193807	
BAU	49748	9016	51103	-4	81435	0	191303	-1
NUTR	50010	8285	50782	-4	81657	0	190735	-2
MTFR	50010	8285	50619	-4	73137	-11	182051	-6
Ionian Sea and Central Med Sea								
REF	9310	3258	15343		22389		50300	
BAU	9124	3254	14754	-4	21928	-2	49061	-2
NUTR	9308	2125	11112	-28	22122	-1	44667	-11
MTFR	9305	2125	10340	-33	12504	-44	34274	-32
Adriatic Sea								
REF	81602	7588	48586		95888		233665	
BAU	80909	7497	45317	-7	95150	-1	228873	-2
NUTR	81598	6196	39670	-18	91711	-4	219174	-6
MTFR	81540	6196	38365	-21	60995	-36	187096	-20
Western Med Sea								
REF	74916	3956	96524		75350		250746	
BAU	74189	3933	94154	-2	74507	-1	246783	-2
NUTR	74913	3281	79612	-18	74324	-1	232131	-7
MTFR	74907	3281	64084	-34	58814	-22	201086	-20

Table S6.2 Phosphorus load at sea outlets (per source and scenario) estimated by the model GREEN.

Phosphorus load at sea outlets (per source and scenario)								
Scenario	Background sources (ton/y)	Scattered dwellings (ton/y)	Point sources (ton/y)	% of change	Agriculture (Min, Man) (ton/y)	% of change	Total load (ton/y)	% of change
Baltic								
REF	12342	4473	10733		6317		33865	
BAU	12341	4465	10191	-5	6178	-2	33175	-2
NUTR	12341	4320	9525	-11	5987	-5	32174	-5
MTFR	12341	4320	8772	-18	5321	-16	30755	-9
Greater North Sea								
REF	5685	1809	30196		28389		66079	
BAU	5685	1809	30176	0	28062	-1	65731	-1
NUTR	5685	1485	29186	-3	20438	-28	56793	-14
MTFR	5685	1482	21449	-29	25879	-9	54495	-18
Celtic Sea								
REF	1085	210	9215		21487		31997	
BAU	1085	210	8802	-4	21078	-2	31175	-3
NUTR	1085	168	9089	-1	20764	-3	31105	-3
MTFR	1085	148	4405	-52	19461	-9	25099	-22
Bay of Biscay and Iberian Coast								
REF	3774	1123	14598		9264		28759	
BAU	3774	1113	13381	-8	9163	-1	27431	-5
NUTR	3774	1108	13020	-11	9225	0	27127	-6
MTFR	3774	1106	6775	-54	8067	-13	19722	-31
Black Sea								
REF	4551	3487	19398		5892		33328	
BAU	4551	3451	15593	-20	5775	-2	29370	-12
NUTR	4551	3287	15972	-18	5360	-9	29170	-12
MTFR	4551	3266	13704	-29	5475	-7	26995	-19
Aegean-Levantine Sea								
REF	2959	1935	11653		4384		20931	
BAU	2959	1935	11206	-4	4374	0	20474	-2
NUTR	2959	1766	11219	-4	4380	0	20323	-3
MTFR	2959	1759	9311	-20	4186	-5	18215	-13
Ionian Sea and Central Med Sea								
REF	582	569	2402		1021		4574	
BAU	582	568	2308	-4	1001	-2	4459	-3
NUTR	582	368	2260	-6	998	-2	4208	-8
MTFR	582	366	802	-67	683	-33	2432	-47
Adriatic Sea								
REF	1776	1563	7369		6650		17359	
BAU	1776	1544	6282	-15	6604	-1	16207	-7
NUTR	1776	1312	5769	-22	6170	-7	15028	-13
MTFR	1776	1295	3599	-51	5274	-21	11945	-31
Western Med Sea								
REF	3270	759	17126		3514		24669	
BAU	3270	755	16576	-3	3473	-1	24075	-2
NUTR	3270	642	15880	-7	3455	-2	23248	-6
MTFR	3270	642	6515	-62	3043	-13	13471	-45

S7. Nutrient concentration in European rivers under different scenarios

Table S7.1 Statistics on the distribution of nitrogen and phosphorus concentration and N:P ratio at all catchments outlets, estimated by the model GREEN under different scenarios.

Nitrogen all catchments

Scenario	n	mean	sd	median	trimmed	Q0.05	Q0.1	Q0.25	Q0.75	Q0.9	Q0.95
BAU	845314	2.49	18.11	1.05	1.40	0.08	0.15	0.47	2.32	4.69	7.19
NUTR	845314	2.45	17.71	1.06	1.40	0.08	0.15	0.48	2.32	4.63	7.01
MTFR	845347	2.26	21.53	0.99	1.29	0.08	0.15	0.46	2.11	4.21	6.32
REF	845312	2.54	18.19	1.07	1.42	0.08	0.15	0.48	2.36	4.77	7.33

Phosphorus all catchments

Scenario	n	mean	sd	median	trimmed	Q0.05	Q0.1	Q0.25	Q0.75	Q0.9	Q0.95
BAU	845314	0.16	2.16	0.06	0.07	0.02	0.02	0.03	0.12	0.25	0.42
NUTR	845314	0.15	2.05	0.06	0.07	0.02	0.02	0.03	0.11	0.24	0.40
MTFR	845347	0.14	2.51	0.05	0.07	0.02	0.02	0.03	0.11	0.22	0.37
REF	845312	0.17	2.19	0.06	0.08	0.02	0.02	0.03	0.12	0.25	0.43

N:P ratio

Scenario	n	mean	sd	median	trimmed	Q0.05	Q0.1	Q0.25	Q0.75	Q0.9	Q0.95
BAU	845314	18.95	16.68	16.64	17.49	3.08	5.35	11.43	23.82	33.35	40.99
NUTR	845314	19.39	16.99	16.90	17.85	3.11	5.35	11.57	24.40	34.68	42.56
MTFR	845347	18.79	16.71	16.58	17.37	3.12	5.48	11.55	23.35	32.73	40.31
REF	845312	19.06	16.71	16.70	17.59	3.12	5.39	11.45	23.98	33.70	41.23

Table S7.2 Statistics on the distribution of nitrogen and phosphorus concentration and N:P ratio at the river basins outlets, estimated by the model GREEN under different scenarios.

Nitrogen Sea Outlets

Scenario	n	mean	sd	median	trimmed	Q0.05	Q0.1	Q0.25	Q0.75	Q0.9	Q0.95
BAU	4148	4.50	18.06	1.16	1.63	0.04	0.06	0.25	2.98	6.24	15.26
NUTR	4148	4.09	15.55	1.15	1.56	0.04	0.06	0.25	2.82	5.88	14.25
MTFR	4148	3.77	14.94	0.99	1.38	0.04	0.06	0.24	2.43	5.30	12.98
REF	4148	4.57	18.25	1.19	1.66	0.04	0.06	0.25	3.07	6.42	15.61

Phosphorus Sea Outlets

Scenario	n	mean	sd	median	trimmed	Q0.05	Q0.1	Q0.25	Q0.75	Q0.9	Q0.95
BAU	4148	0.37	1.71	0.07	0.11	0.01	0.01	0.03	0.20	0.52	1.28
NUTR	4148	0.33	1.55	0.07	0.11	0.01	0.01	0.03	0.18	0.46	1.17
MTFR	4148	0.29	1.42	0.06	0.10	0.01	0.01	0.03	0.17	0.38	0.93
REF	4148	0.38	1.72	0.07	0.12	0.01	0.01	0.03	0.20	0.53	1.36

N:P Sea Outlets

Scenario	n	mean	sd	median	trimmed	Q0.05	Q0.1	Q0.25	Q0.75	Q0.9	Q0.95
BAU	4148	14.00	10.03	12.97	13.08	1.75	3.61	8.11	18.11	23.97	28.75
NUTR	4148	14.75	11.65	13.11	13.26	1.75	3.51	7.99	18.59	25.56	34.31
MTFR	4148	13.79	9.40	13.25	13.12	1.75	3.64	8.62	17.62	22.26	26.58
REF	4148	14.04	10.07	12.98	13.12	1.75	3.61	8.14	18.25	24.05	28.93

Figure S7.1 Distribution of nitrogen and phosphorus concentration and N:P ratio at the sea outlets (right) and at all catchments outlets (left), estimated by the model GREEN under different scenarios.

