

# Can hydrocarbon extraction from the crust enhance or inhibit seismicity in tectonically active regions? A statistical study in Italy

Alexander Garcia<sup>1</sup>, Licia Faenza,<sup>2</sup> Andrea Morelli,<sup>3</sup> and Ilaria Antoncecchi<sup>4,5</sup>

<sup>1</sup> Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Bologna. Via Donato Creti 12, 40128 Bologna, Italy. Email: alexander.garcia@ingv.it

<sup>2</sup> Istituto Nazionale di Geofisica e Vulcanologia, Osservatorio Nazionale Terremoti, Bologna Italy. . Email: licia.faenza@ingv.it

<sup>3</sup> Istituto Nazionale di Geofisica e Vulcanologia, Sezione di Bologna. Via Donato Creti 12, 40128 Bologna, Italy. Email: andrea.morelli@ingv.it

<sup>4</sup> Ministero di Sviluppo Economico, Direzione Generale per le Infrastrutture e la Sicurezza dei Sistemi Energetici e Geominerari, DGISSEG, Rome, Italy. Email: ilaria.antoncecchi.ext@mise.gov.it

<sup>5</sup> Ricerca sul Sistema Energetico S.p.A (RSE), Milano, Italy

## Version I: 16 March 2021.

Note: This manuscript has been submitted for publication in *Frontiers in Earth Sciences, section Solid Earth Geophysics*, and currently it is under peer-review. Therefore, subsequent versions of this manuscript may have slightly different content. If accepted, the final version of this manuscript will be available via the Peer-reviewed Publication DOI (link on the right-hand side of the EarthArXiv webpage). Feedback is welcome, so please feel free to contact any of the authors.

## SUMMARY

A number of oil- and gas-producing leases have been operating in Italy in the last decades, many of which are located in the surroundings of tectonically active regions. Identifying human-induced seismicity in areas with high levels of natural seismicity is a difficult task for which virtually any result can be a source of controversy. We implemented a large-scale analysis aiming at tracking significant departures of background seismicity from a stationary behavior around active oil and gas development leases in Italy. We analyzed seismicity rates before and after hydrocarbon peak production in 6 oil-producing and 43 gas-producing leases, and evaluate the significance of possible seismicity rate changes. In a considerable number of cases seismicity rate results stationary. None of the observed cases of seismicity rate increase after the peak production is statistically significant (at a  $s.l. = 0.05$ ). Conversely, considering cases of seismicity rate decrease after peak production, our results suggest that the seismicity rate reduction is statistically significant ( $s.l. = 0.05$ ) around one oil-producing lease (Val d'Agri, Basilicata) and around a cluster of gas-producing leases in Sicily. Our results put in evidence correlated changes between the rates of shallow seismicity and hydrocarbon production in these areas, which are then identified as hotspots requiring more detailed research; assessing actual causal relationships between these processes will require further physically-based modelling. If a physical causative link between these processes exists, then the observed seismicity rate reduction could either be due to increased seismicity during the progressive increase in production before reaching its maximum, or to an actual seismicity rate reduction after that peak. Considering that there is evidence of seismicity occurring before the start of hydrocarbon production, which contrasts with the evident reduction of events observed after the peak production, we think it likely that the seismicity inhibition is a plausible hypothesis. Using a simple model we also calculate Coulomb stress changes in planes optimally oriented for failure, and we show that under some conditions the inhibition of seismicity is feasible in at least one of these cases. We conclude that more efforts to study the mechanisms and the possible consequences of anthropogenically-driven seismicity inhibition are required.

**Key words:** Regional seismicity, Hydrocarbon production, Correlation analysis, Italy, Anthropogenic hazards, Seismicity rate changes

## 1 INTRODUCTION

The complex geological setting that characterizes the Italian peninsula is the result of different geodynamical processes closely acting in time and space; consequently, today the crust in this zone is characterized by a complex stress field with tightly spaced compressional and extensional regions (Amato & Montone 1997). The most predominant geomorphologic features in this region are the Southern Alps and the Apennines mountain chains, which are characterized by thrust-and-fold belts originated from the interaction between the European and the Adriatic-African tectonic plates (see e.g., Calamita et al. 1994; Cello & Mazzoli 1998; D'Agostino et al. 2001; Bertello et al. 2010; Handy et al. 2010; Cazzini et al. 2015; van Hinsbergen et al. 2020). Such an active and complex tectonic setting makes Italy a seismically-active region where, on average, more than 2000 seismic events with magnitude  $\geq 2.0$  are located each year by the Italian national seismic network (see e.g., the *Bollettino Sismico Italiano*, Pagliuca et al. 2020).

In recent years, the interest in the possible influence of anthropogenic activities on seismicity has significantly grown mainly because of a generalized public concern, which has pushed the development of independent scientific research to support objective policy making (e.g., Ellsworth 2013; Dahm et al. 2015; van der Voort & Vanclay 2015; GarciaAristizabal et al. 2020). The Italian territory is the scenario of a wide number of underground industrial activities such as oil and gas extraction, geothermal energy production, and gas storage, many of which have been suspected to have direct or indirect causal links with some seismic events located nearby; however, to date there are no unambiguously documented reports of damaging seismic events associated with anthropogenic activities in the country (see e.g., Braun et al. 2018). Probably, the only clear cases of seismicity linked to underground geo-resource development in Italy are the low-magnitude seismicity occurrences recorded in connection with wastewater reinjection at the Costa Molina 2 well in the High Val d'Agri, southern Italy (Valoroso et al. 2009; Improta et al. 2015), and the seismicity recorded near geothermal power plants in Tuscany (Evans et al. 2012).

Among all the anthropic activities having the potential to stimulate earthquakes to occur, the effects of fluid injection or extraction from the crust are probably the processes arising more concern, in particular the activities related with oil and gas production. In Italy, hydrocarbons are found in several oil and gas provinces, most of which are located in the Po plain, the northern Adriatic sea, the southern Apennines, and in Sicily (e.g., Bertello et al. 2010); as a consequence, these are the areas hosting most of the development leases in the country (Figure 1). According to data published by the Italian ministry of Economic Development (MISE), as of 2019 there were in total 193 development leases in the country, 127 of which are onshore and the other 66 offshore (UNMIG 2020).

Discriminating natural from induced seismicity in seismically active regions is a particularly complex task. Early attempts to discriminate induced from natural seismicity were performed, for fluid injection operations, by Davis & Fröhlich (1993), and for fluid withdrawal by Davis & Nyffenegger (1995); however, these approaches were mainly based on qualitative assessments.

More quantitative, physically-based and/or stochastic methods for discriminating natural from induced seismicity have recently been proposed in literature (a review can be found, e.g., in Grigoli et al. 2017). For example, Dahm et al. (2015) propose a quantitative probabilistic approach to discriminate induced, triggered, and natural earthquakes, calculating the probability that events have been anthropically triggered/induced from the modeling of Coulomb stress changes and a rate-and-state dependent seismicity model.

Schoenball et al. (2015) analyzed inter-event times, spatial distribution, and frequency-size distributions for natural and induced earthquakes around a geothermal field. Determining the distribution of nearest neighbor distances in a combined space-time-magnitude metric, they identify clear differences between both kinds of seismicity. For example, it is suggested that compared to natural earthquakes, induced earthquakes feature a larger population of background seismicity and nearest neighbors at large magnitude rescaled times and small magnitude rescaled distances. They argue that unlike tectonic processes, stress changes caused by anthropic underground operations occur on much smaller time scales and appear strong enough to drive small faults through several seismic cycles. As a result, it is likely to record seismicity close to previous hypocenters after short time periods.

Zhang et al. (2016) compared moment tensors of both natural and induced events in the Western Canadian Sedimentary Basin. These authors calculated full moment tensors and stress drop values for eight induced earthquakes (magnitudes between 3.2 and 4.4), as well as for a nearby M5.3 event considered as a natural earthquake. This study suggests that, first, it may be possible to discriminate between induced and natural seismicity considering region-specific attributes, as for example the focal depths (which they suggest as the most robust parameter since the induced events in their study area are significantly shallower than most of the intra-plate earthquakes in the Canadian Shield). Moreover, they found a non-negligible ( $> 25\%$ ), non double couple component for most of the induced events studied.

Zaliapin & BenZion (2016) analyzed statistical features of background and clustered subsets of earthquakes in California and in South Africa. These authors suggest that, compared to regular tectonic activity, induced seismicity in the analyzed data sets exhibit remarkable features as (i) a higher rate of background events, (ii) faster temporal offspring decay, (iii) higher rate of repeating events, (iv) larger proportion of small clusters, and (v) larger spatial separation between parent and offspring.

Discriminating human-induced from natural seismicity is therefore a difficult problem for which virtually any solution can be a source of controversy. For this reason, the evaluation of possible interactions between seismicity and hydrocarbon production, if possible, should rely on multi-disciplinary analyses such as e.g., detailed physically-based modelling complemented by sophisticated stochastic methods able to provide probabilistic assessments and to take uncertainties into account. It is worth noting however that the ways in which these interactions may occur are complex and their identification in a context characterized by high levels of naturally-occurring seismicity is not straightforward. Moreover, given the relatively high number of development leases active in Italy, performing such analyses at the national scale may be considered an intractable problem.

These reasons pushed us to explore the possibility to implement large-scale screening methods aiming at tracking measurable phenomena, such as e.g., changes in seismicity rates, that plausibly could occur if notable interactions between underground human operations and nearby seismicity sources are actually occurring in a given area. Spatial and temporal correlation between human activity and event rates is usually considered a key parameter to suspect possible relationships between seismicity and underground anthropic activity (e.g., Shapiro et al. 2007, 2010; Cesca et al. 2014; Leptokaropoulos et al. 2017; Garcia-Aristizabal 2018; Molina et al. 2020); for example, significant changes in seismicity rates with respect to background seismicity, as well as the spatial and temporal correlation of gas injection operations and seismicity were analyzed by Cesca et al. (2014) to suggest a possible case of triggered/induced seismicity near an offshore platform used for gas storage in Spain (the Casor project).

However, underground human-induced perturbations (as e.g., pore pressure variations due to fluid injections) can produce changes at large distances and/or with large temporal delays, potentially causing earthquakes to occur several kilometers away as well as months/years after the industrial operations have stopped or reached the maximum peak (e.g., Mulargia & Bizzarri 2014); likewise, natural seismicity may also occur within few kilometers from industrial sites. Therefore, in seismically active regions (such as in Italy), spatio-temporal correlations between industrial activity and significant changes in seismicity rates with respect to background activity by themselves usually do not provide irrefutable proofs of causal relationships between hydrocarbon production and seismic activity. Despite this, we argue that studying such correlations has a remarkable added value since it gives us the possibility of performing large-scale, systematic analyses of a huge amount of seismic and production data and, under the working hypotheses considered, to identify *hotspot areas* where it could be possible to perform, in a later stage, more detailed research to verify possible causal relationships.

The working hypothesis in this work starts from assuming that fluid withdrawal from the crust may induce deformations in the surroundings of the host rock; the magnitude of such deformations will depend on multiple factors such as, for example, the lithology, the structural geology, the volume and rate of fluid removed, the geomechanical features of the reservoir, and its behavior during the fluid withdrawal process, among others. The deformations may in turn alter the local stress field and, as a consequence, stimulate or inhibit seismicity in the surroundings.

We assume that when in a given region the seismicity release is driven by long-term, steady tectonic processes only, then the rate at which independent seismic events occur tends to be stationary. However, if other natural (e.g., hydrogeology: see Hainzl et al. 2006; Pintori et al. 2021) or man-made (e.g., pressurized fluid injections: see Shapiro et al. 2007; Garcia-Aristizabal 2018) processes are able to perturb the local stress field, then it is possible that the rate at which seismicity is released in that specific area can be altered. In such a case, slight deviations from stationarity could possibly be measured.

In this work we are interested in identifying significant departures of background seismicity from a stationary be-

havior around active oil and gas development leases in Italy. Moreover, we are also interested in testing whether possible changes in the rate at which seismicity is released in such areas are correlated with main changes in the hydrocarbon production patterns. The article is structured as follows: first we present the seismic and the hydrocarbon production data available for this study. Second, we present the methodological approach used in order to identify zones with possible anomalies in seismicity rates concomitant with significant changes in oil and gas production. Finally, we present the results and discuss the importance and limitations of these findings.

## 2 DATA


For this study we use a national-wide seismic catalog containing earthquake locations and magnitudes, as well as the most detailed public oil and gas production data from development leases in Italy. All the used data are freely accessible from public sources (see the *Data availability statement* section for details).

### 2.1 Seismicity

We use the seismic data collected in the publicly-available HORUS catalog (Homogenized instrumental seismic catalog, Lolli et al. 2020a,b). This is an extended instrumental seismic catalog reporting earthquake locations and magnitudes since 1960 and is continuously updated as new data is processed. An outstanding feature of this catalogue is the effort made to harmonize the event magnitudes in terms of an *equivalent homogeneous moment magnitude*,  $M_w$ .

We use the data in the time interval 1980-2017, which includes 368,258 earthquakes (Figure 1). This time interval is selected because it covers the same time window of available hydrocarbon production data; moreover, the eighties are probably the period when the national seismic network started to grow more consistently, improving as a consequence the quality of the earthquake locations and reducing the completeness magnitude ( $M_c$ ). However, this process does not evolve uniformly throughout the whole country, since the spatial and temporal distribution of new seismic stations is not spatially homogeneous. For example, Figure 2 shows the temporal evolution of the Italian National Seismic Network; the stations belong to different networks deployed for seismic monitoring in Italy (e.g., the Italian Seismic Network and the Euro-Mediterranean Network, both maintained by INGV; the Italian Strong Motion Network, managed by the National Civil Protection; the networks operated by other national institutes and universities such as the Istituto Nazionale di Oceanografia e di Geofisica Sperimentale, the University of Basilicata, the University of Genoa, the University of Trieste, the University of Bari, and other organisations in the border areas).

The completeness magnitude ( $M_c$ ) of this catalogue therefore changes in time and in space, as a consequence of different factors as e.g., changes in the number and distribution of seismic stations available for locating events, as well as the quality of the instrumentation and of the site facilities. Therefore, identifying a single completeness magnitude (in both time and space) for the full catalog results in a very high



**Figure 1.** Map showing the spatial distribution of the oil and gas production leases in Italy, as well as the epicenters of the earthquakes from the HORUS catalog used in this study.

$M_c$  value; adopting such a choice would force us to discard an important amount of seismic data, and for this reason, the  $M_c$  is rather determined locally for each site, as discussed in the Methods section.


## 2.2 Oil and gas production data

We collected the most detailed public information about hydrocarbon production in Italy (available from the Italian Ministry of Economic Development, MISE, see the *Data Availability Statement* section for details). Out of 109 active development permits (Figure 1), we got production data from 102 leases in which a total of 588 production wells are present. The available data set includes the annual oil and gas production since 1980 (volumes in thousands of cubic meters of oil or standard cubic meters of gas); the data is aggregated by development lease, that is, summing up the production/year from all the wells producing within the lease. For development leases in which production started before 1980, the total oil and gas production preceding 1980 is also available.

It is worth noting that besides the hydrocarbon production, a number of sites where underground waste water injection is performed are also present in Italy; nevertheless, the fluid injection-related data are not publicly available and for this reason these activities were not considered in this study.

As Figure 1 shows, the active development leases are mostly distributed along the Adriatic coast (onshore and offshore), as well as along the Po Valley (in the North), the central/southern Apennines and Sicily. In this study we consider the hydrocarbon production from 1980 up to 2017 (Figure 3). From the 102 development leases for which production data is available, 85 exclusively produce gas, 1 exclusively produces oil, and 16 produce both oil and gas. The development leases with the highest annual oil production in the study period are found in the Val d’Agri (Basilicata region), Villafortuna-Trecate (Piemonte region, western Po Valley), B.C 8.LF (offshore, southern Adriatic sea), and C.C 6.EO (Sicily, southern Italy) leases (Figure 3A); on the other hand, those with the highest annual gas productions in the same period are the A.C 7.AS (offshore, central Adriatic sea), D.C





**Figure 2.** Temporal evolution of the deployment of monitoring seismic stations in Italy, maintained by various Italian and international institutes and universities (see the text for details). Substantial increases in the number of stations concurred with the occurrence of different seismic sequences.

1.AG (Calabria region), A.C 2.AS and A.C. 27.EA (offshore, northern Adriatic sea, Figure 3B).


### 3 METHODS

We perform a large-scale screening of the behavior of background seismicity (i.e., the events considered independent, as described in Section 3.3) in areas around oil and gas production sites in Italy. Our goal is to attempt (i) to identify correlated changes between seismicity rates and hydrocarbon production, and (ii) to test the significance of these seismicity rate changes.

First, a pre-processing step is performed in order to identify the areas around the target leases where a reasonable amount of both seismic and production data is available. Afterwards, we proceed with the proper data analyses, namely (i) identification of independent background seismicity in the lease area, (ii) selection of background events located within a distance  $\delta x$  from the production wells, and (iii) test the significance of possible seismicity rate changes correlated with changes in hydrocarbon production. These steps are explained in the following paragraphs.

#### 3.1 Identifying development leases for the analyses

If the withdrawal of fluids from the crust affects the stress field and the seismicity release in a given zone, then a possible way to identify potential traces of such effects is to look for changes in background seismicity rates before and




**Figure 3.** Annual oil and gas produced between 1980 and 2017 in Italy from 102 development leases in Italy: (a) Oil is produced from 17 leases (1 of which exclusively produces oil); (b) Gas is produced from 101 leases (85 of which exclusively produce gas). Data available from the Italian Ministry of Economic Development (MISE).

after important changes in the rate at which fluids are extracted. Observing the time series of production data (Figure 3), it can be seen that in most of the sites, the oil production (Figure 3A) and the gas production (Figure 3B) have some outstanding features: after the production starts, it follows different paths up to a point in which it reaches maximum production. The time at which oil or gas production reaches its maximum is hereinafter called the *maximum production time*,  $t_m$ . Afterwards, for times  $t > t_m$ , the production tends to decrease (probably related to field depletion processes).

Looking in detail at the development of the time series of production data, the main change points in oil and gas production data of potential interest for the analyses in this article are the start of production and the time at which production reaches its maximum. We consider the peak of maximum hydrocarbon production as the reference point for exploring possible changes in seismicity rates (before and after the peak). Changes before and after the production start cannot be analysed because in most of the sites the time at which production initiates precede the start of the seismic data catalog.

An essential requirement to study changes in seismicity rates consists of taking a sufficiently long time window of seismic data in the time periods preceding and following the change point in production data considered for comparing seismicity rates. In general, the longer the time window, the more representative will our data set likely be. Taking into account the time window of data available for this study (1980-2017), as well as the average seismicity rates observed around the active development leases, we consider as a re-



**Figure 4.** Year at which the maximum production is reached in the available data at each development lease ( $t_m$ ). Histograms show the number of (A) oil- and (B) gas-producing leases in which  $t_m$  is reached in the respective year; the blue bars identify the period selected for the analyses (1995–2008) according to the described criteria.

requirement for including a given development lease in this analysis the availability of a complete set of seismic data for at least 10 years before and 10 years after the  $t_m$  associated with the peak production in the respective development lease.

Figure 4 shows histograms summarizing the  $t_m$  at which maximum oil (Figure 4A) and gas (Figure 4B) production are reached in all the development leases for which data are available. Adopting the minimum 10-years time window before and after  $t_m$ , and also considering the temporal completeness identified for the seismicity around the leases (see section 3.2 for details), the sites that can be reliably analyzed in this work reduce to those for which the peak production in the available time series is reached between 1995 and 2008 (i.e.,  $1995 \leq t_m \leq 2008$ ). In this way, 6 oil-producing leases and 43 gas-producing leases comply with these conditions and are selected for further analyses.

The time series of oil and gas production data from the selected development leases are shown, respectively, in Figure 5A and 5B. Taking the peak oil/gas production as a reference, the data follow different patterns that can be better observed normalizing the production by the maximum production reached in each lease and realigning the time series with respect to  $t_m$  (Figure 5C and 5D). For the sake of simplicity, we identify three main patterns (Figure 5E): pattern I (the most frequent), is when the production rate steadily increases with time up to reach its maximum; afterwards the production progressively reduces with time. Patterns II and III instead refer to cases where production increases up to remain at relatively high levels for a while, before signifi-

cantly decreasing again. In some cases the peak production is reached at the end (pattern II), and in other cases it is reached at the beginning (pattern III) of the high-production period.

### 3.2 Seismic Data Selection


We first calculate the epicentral distance between each earthquake in the catalogue and the nearest production well in a lease. This information is used to select the whole seismicity located around each lease, paying particular attention to include all possible clusters of events in time and space observed in the surroundings (i.e., to avoid including incomplete data of seismic sequences, for example). This generally means including seismicity located within a few hundred kilometers around each lease (Figure 6A). Moreover, since it is generally observed that events induced by man-made, underground operations tend to be shallower than most natural, tectonically driven events, and that induced events often occur at depths comparable to the depth of wells (see e.g., Zhang et al. 2016; Foulger et al. 2018), then for this analysis we only consider the events with depth shallower than 15km.

Finally, for each resulting (local) catalog we calculate the completeness magnitude ( $M_c$ ) using, comparatively, three methods [the Maximum Curvature, the Goodness of Fit (Wiemer & Wyss 2000), and the Modified Goodness of Fit (Leptokaropoulos et al. 2013)], that are available as open tools in the EPOS (European Plate Observing System) platform for anthropogenic hazards (IS-EPOS 2016; Orlecka-Sikora et al. 2020). In this way, for each lease we obtain a seismic catalogue covering a given time interval and is complete above a given minimum magnitude  $M_c$  (Figure 6A and 6B).

### 3.3 Declustering the seismic catalog

To perform the analyses proposed in this work we suggest to use a seismic catalog composed as much as possible by independent events, that is, events not likely triggered by earthquake interaction processes (as e.g., aftershocks). That set of independent events is hereinafter referred to as *background seismicity* (see e.g., Figure 6B).

The seismic declustering is a crucial issue in statistical seismology. In general, the term “background” (or “independent”) events refers to events that are typically related to regional tectonic activity. Triggered events occur in space-time clusters, and are associated with the occurrence of previous events (e.g., stimulated by stress perturbations); they are often referred to as “dependent events”. The distinction within a catalogue between the contribution of independent and dependent earthquakes is very complex, and each method inherently contains subjectivity (Zhuang et al. 2002). This characteristic implies that applying different declustering models to the same catalogue may generate catalogs of independent events that may differ. In seismology, several techniques have been developed to address the problem of declustering; van Stiphout et al. (2012) provides an overview of this issue, describing the pros and cons of the most popular algorithms. In this work we use the Nearest-Neighbour (NN) Clustering Analysis technique, developed by Zaliapin & Ben-Zion (2013). An outstanding advantage




**Figure 5.** Annual production of identified sites in which it is possible to perform the proposed analyses. Annual production of (A) oil and (B) gas in the selected development leases. Production data normalized by the maximum value reached in each time series and realigned respect to peak time,  $t_m$  are shown in plots C (oil) and D (gas). (E) Main temporal trends observed in production data.

of the NN algorithm is its simplicity because the link between the background events with those triggered is a metric that only concerns a distance measure in time, space and magnitude between any pair of events. The technique consists of calculating, for each earthquake  $i$  in the catalog, the distance to any other  $j$ -th earthquake subsequently occurred as:

$$\eta_{ij} = t_{ij} r_{ij}^{d_f} 10^{-bm_i} \quad (1)$$

where  $t_{ij} = t_j - t_i$  is the difference between the two occurrence times expressed in years (with  $t_{ij} > 0$ ),  $r_{ij}$  is the distance between the two hypocenters in km,  $d_f$  is the fractal dimension of the distribution of hypocenters,  $b$  is the Gutenberg-Richter  $b$ -value, and  $m_i$  is the magnitude of the  $i$  earthquake. For each earthquake  $i$ , the smaller  $\eta_{ij}$  is



**Figure 6.** Summary of the processing for selecting seismic data around a given production lease for the correlation analysis: (A) Regional seismicity around the lease, for which the  $M_c$  is evaluated; (B) Identification of background seismicity from the regional catalog (considering events above  $M_c$ ) using declustering techniques; (C) Spatial filtering by selecting events located within a distance  $\delta x$  from production wells; (D) binomial test for evaluating the significance of seismicity rate changes.

selected. Within a complete seismic catalog, the distribution of these 3-dimensional distances always shows a bimodal pattern: the first group of earthquakes has an unusually small distance and represents earthquakes that are “clustered”, whereas the second group identifies the events interpreted as “background earthquakes”, since in the considered parameter space they are characterized by greater distances from each other.

### 3.4 Evaluating the significance of seismicity rate changes

Once the background seismicity around a given lease is obtained, we select a set of seismic events for studying possible changes in seismicity rate correlated with changes in hydrocarbon production. With this aim, we set a maximum distance from the producing wells,  $\delta x$ , to define the volume enclosing the events to be included in the analysis (Figure 6C). Defining  $\delta x$  is critical since it reflects the spatial extent where the potential deformations and stress perturbations are supposed to alter the natural occurrence of seismicity. In principle it should be carefully evaluated case by case accounting for different local factors such as, e.g., the size and depth of the reservoir, the volume of fluids withdrawn from the crust, etc. If  $\delta x$  is small, there is a risk of including a small, non representative sample of events; on the other hand, if  $\delta x$  is very large, the significance of possible local changes in seismicity rates can be hidden by a large sample of regional (and presumably stationary) seismicity.

In tectonic earthquake interaction studies, the size of the area at which earthquake triggering is mostly expected to occur is usually mapped by static stress perturbation; if  $L$  is the mainshock source length (as derived from scaling relations, such as for example Wells & Coppersmith 1994), a characteristic distance in the range 1 to 3  $L$  is often suggested as a plausible distance within which triggered aftershock are expected to occur (e.g., Parsons & Velasco 2009; Tahir et al. 2012). Some authors have suggested similar scaling properties for determining characteristic distances for seismicity induced by reservoir impoundments (Grasso et al. 2019) and gas reservoirs (Grasso et al. submitted); in such cases, the characteristic distance is determined as a function of the size of the reservoir. We do not have information about the dimensions of the oil and gas reservoirs from where hydrocarbons are produced in the analyzed cases; therefore, heuristically we select events located at distances within  $\delta x = 5$  and 10 km from the production wells, which seems a reasonable and conservative choice for this study.

We look for significant seismicity rate changes before and after  $t_m$  using the binomial test proposed by Lepotokaropoulos et al. (2017). Let  $T_{pre} = [t_1, t_m]$  be the time interval identified before the maximum production peak (with  $t_m - t_1 \geq 10$  years, as defined before), and  $T_{post} = [t_m, t_2]$  the time interval identified after the maximum production peak (with  $t_2 - t_m \geq 10$  years as well, see Figure 6D for reference). Let  $n_{pre}$  be the number of events that occurred in the period  $T_{pre}$ , and  $n_{post}$  the number of events that occurred in the period  $T_{post}$ . The total number of events



in both periods is therefore  $N = n_{pre} + n_{post}$  (Figure 6D). If the seismicity rate exhibit changes that are correlated with changes in the production trend before and after the time at which the maximum production is reached ( $t_m$ ), that is, if the seismicity rate during  $T_{pre}$  is significantly different from the seismicity rate during  $T_{post}$ , then the actual division of the total number of events  $N$  in both periods into  $n_{pre}$  and  $n_{post}$  should be significantly different from the division which could be attained at random. Therefore, if we hypothesize stationary seismicity, the proposed null hypothesis,  $H_0$ , states that:

$H_0 : n_{post}$  can be obtained at random from  $N$  under probability  $P$

where  $P$  is related to the time partitions as follows:

$$P = \frac{T_{post}}{T_{pre} + T_{post}} \quad (2)$$

This hypothesis is tested by means of the binomial test (e.g., Wonnacott & Wonnacott 1977). If  $N$  events occur randomly in the interval  $[t_1, t_2]$ , this test provides (i) the probability  $p_1$  that the number of events in the interval  $[t_m, t_2]$  is less than or equal to  $n_{post}$ ,

$$p_1 = \quad (3)$$

or (ii) the probability  $p_2$  that the number of events in the interval  $[t_m, t_2]$  is greater than or equal to  $n_{post}$ ,

$$p_2 = \quad (4)$$

The binomial test assumes that each event is independent, with equal probability of occurrence in the interval  $[t_1, t_2]$ ; the null hypothesis ( $H_0$ ) is evaluated at a given significance level (e.g.  $s.l. = 0.05$ ), so that if  $p_1$  (or  $p_2$ )  $< s.l.$ , we conclude that there is evidence, with a given significance  $s.l.$ , that rate in the interval  $[t_m, t_2]$  decreased (or increased) with respect to the rate in the interval  $[t_1, t_m]$ .

## 4 RESULTS

One of the main issues faced to perform the data analysis was to concentrate our efforts in areas where both seismic and production data were sufficiently representative to avoid, as much as possible, data-driven biases in our results. For this reason, and given the time span covered by the seismic and production data available for this study, we avoided analyzing any region in which the minimum data requirements defined in the Methods section was not accomplished. In particular, the minimum length of the time window of seismic data before and after  $t_m$ , the time at which the maximum production is reached, becomes one of the main constraints, forcing us to not considering about 56% of the leases for which production data are actually available (that is, 57 out of the 102 leases). In such cases, the peak production occurred too close to the end or the beginning of the seismic catalogue, which prevents an accurate estimation of the pre- and post-  $t_m$  seismic rate. On the remaining 45 leases (39 produce gas only, 2 produce oil only, and 4 produce both gas and oil), we applied the analyses described in section 3 (Methods). The spatial distribution of the analyzed leases cover different areas of the country, including the western Po plain, the northern and central Adriatic sea, southern Apennines and Sicily.

The completeness magnitudes determined for the regional seismicity around each lease are summarized in Table 1 (for the oil-producing leases) and Table 2 (for the gas-producing leases). It is worth noting that, in general, the completeness magnitudes tend to be relatively high due to our interest in having seismic data sets as long as possible in time; in fact, the  $M_c$  has been selected for the seismicity around each lease so that completeness is ensured, at least, since 1985. The seismic catalogs, composed of nearby regional events with magnitudes above the  $M_c$  specifically determined for each study area, are then declustered using the NN clustering analysis technique (Section 3.3) to obtain a sample of background seismicity composed of independent events at regional scale.

Finally, we identify background events located within  $\delta x = 5$  km and  $\delta x = 10$  km from the production wells in a given lease, and select the events before and after the respective  $t_m$ . In this way we calculate the number of events before ( $n_{pre}$ ) and after ( $n_{post}$ ) the peak production time,  $t_m$ , as well as the respective time window lengths  $T_{pre}$  and  $T_{post}$  (see e.g., Figure 6D for reference).

Using these data, we first calculate the seismicity rate (in terms of average number of events/year) observed before and after  $t_m$  for events located within a distance  $\delta x$  from the producing wells. Comparing the seismicity rates before and after the peak production, it is possible to highlight the areas with seismicity rate variations. For example, Figure 7A and Figure 8A show the location and estimate of seismicity rate variations around, respectively, the 6 oil-producing and 43 gas-producing leases analyzed in this study when considering seismicity within  $\delta x = 5$  km from production wells; colors indicate the behavior of the seismicity rate (i.e., increase, decrease, unchanged) before and after  $t_m$ . Regarding the oil-producing leases, the seismicity rate around 3 sites results null and unchanged before and after  $t_m$  (green squares in Figure 7A; names can be seen in Table 1), in 2 cases (Masseria Verticchio and Val d'Agri) the seismicity rate before the peak production results higher than the seismicity rate in the time window after  $t_m$  (blue squares in Figure 7A), whereas in one case (Ragusa) we observe the opposite situation (that is, the seismicity rate before is lower than the seismicity rate after  $t_m$ , red squares in Figure 7A). Regarding the gas-producing leases, we observe that in 20 sites the seismicity rate does not change (18 of which have zero events before and after  $t_m$ ). These sites are represented as green squares in Figure 8A. In the remaining 23 leases (names can be seen in Table 2), a seismicity rate variation has been detected: 9 sites exhibit an increase (red squares in Figure 8A), whereas 14 sites exhibit a decrease in the rate (blue squares in Figure 8A). When considering  $\delta x = 10$  km, 1 oil-producing and 10 gas-producing leases do not exhibit seismicity rate changes, in 4 and 22 leases (oil and gas, respectively) the seismicity rate before  $t_m$  results higher than the seismicity rate after the peak production, whereas the opposite behavior is observed in 1 oil- and 11 gas-producing leases.

We then evaluate the significance of these seismicity rate variations using the binomial test described in Section 3.4. For the areas exhibiting a reduction in seismicity rate after  $t_m$ , we calculate  $p_1$  (Eq. 3) to evaluate the significance of the observed seismicity rate reduction in the time interval  $T_{post} = [t_m, t_2]$  with respect to the rate observed in the



**Table 1.** Oil-producing leases analyzed in this study. For each lease we present the estimated completeness magnitude, as well as the p values ( $p_1$  and  $p_2$ ) of the binomial test performed considering the seismicity located within a distance  $\delta x = 5$  and  $\delta x = 10$  km (the symbol – indicates cases in which both the seismicity rate in both the pre- and the post- $t_m$  time windows are zero).

Oil leases		Completeness		$\delta x = 5$ km		$\delta x = 10$ km	
n.	Lease name	$M_c$	since...	$p_1$ (lesser)	$p_2$ (greater)	$p_1$ (lesser)	$p_2$ (greater)
1	FC 2.AG	2.8	1985	–	–	$4.2 \times 10^{-1}$	
2	GIAURONE	2.1	1985	–	–	$6.1 \times 10^{-1}$	
3	MASSERIA VERTICCHIO	2.2	1985	$4.8 \times 10^{-1}$		$3.3 \times 10^{-1}$	
4	RAGUSA	1.7	1985		$4.3 \times 10^{-1}$		$4.3 \times 10^{-1}$
5	<b>VAL D'AGRI</b>	2.4	1985	$1.6 \times 10^{-5}$		$3.3 \times 10^{-8}$	
6	VILLAFORTUNA-TRECATE	1.9	1985	–	–	–	–

interval  $T_{\text{pre}} = [t_1, t_m]$ . Likewise, for the areas exhibiting an increase in seismicity rate after  $t_m$  we calculate  $p_2$  (Eq. 4) to evaluate the significance of the observed increase in the time interval  $T_{\text{post}}$  with respect to the rate observed in the interval  $T_{\text{pre}}$ . The results of these calculations, considering seismicity located within both  $\delta x = 5$  and  $\delta x = 10$  km from production wells, are presented in Table 1 for the oil-producing and in Table 2 for the gas-producing leases.

The resulting probability values ( $p_1$  and  $p_2$ ) provide an indication about how likely it is to observe the number of events  $n_{\text{post}}$  under the null hypothesis (which basically reflects what would be expected in case of stationary seismicity in the whole interval  $[t_1, t_2]$ , see Figure 6D for reference). Therefore, the lower the p value, the more unlikely is to observe  $n_{\text{post}}$  under such hypothesis (and therefore the more evidence in favor of a significant seismicity rate change). Considering for example a significance level  $s.l. = 0.05$  we find that, on the one hand, none of the cases in which an increase in seismicity rate was observed after  $t_m$  is statistically significant (considering seismicity located within both 5 and 10 km from production wells). On the other hand, regarding the cases exhibiting a seismicity rate decrease after  $t_m$ , our results suggest that the observed seismicity rate change is statistically significant for one oil-producing lease (the Val d'Agri) and two gas-producing leases (Fiumetto and Rocca Cavallo). The location of all the analyzed leases, classified by the calculated p values considering as reference a  $s.l. = 0.05$ , are shown in Figure 7B for the oil-producing and Figure 8B for the gas-producing leases. The Val d'Agri lease is located in the Basilicata region, in the Southern Apennines, whereas Fiumetto and Rocca Cavallo leases are located in Sicily, near Etna Volcano.

The leases for which the seismicity rate change is considered statistically significant are identified using events within both  $\delta x = 5$  and  $\delta x = 10$  km (see Tables 1 and 2). It is worth to note that in most of the cases the rate change (i.e., increase or decrease) is coherent for seismicity selected considering both  $\delta x$  values. However, in a few cases of data selected around some gas-producing leases (5 sites out of 43, namely A.C 28.EA, Bronte-S.Nicola, Monte Morrone, Recovato, and S. Andrea), there is a change in the observed trend of seismicity rate variation (for example, a decrease observed for the seismicity located within 5 km contrasts with an increase observed when selecting events within a 10 km distance); discrepancies in such areas are due to instabilities mainly caused by either low seismic activity (such as,

e.g., in the Po plain or offshore in the Adriatic sea) or sites close to seismically active sources (such as, e.g., very active tectonic or volcanic areas). It is worth noting however that in all these cases, as well as for cases in which no events are identified within the first 5 km, the observed rate changes are in any case not statistically significant.

## 5 DISCUSSION

In this work we monitor significant departures of background seismicity from a stationary behavior around active oil and gas development leases in Italy. We collected oil and gas production data from 102 leases in the period 1980 - 2017, and we used the seismic data from the HORUS catalog (Lolli et al. 2020b). After a close and conservative inspection of the available data, it has been possible to implement the proposed analyses in 6 oil-producing and 43 gas-producing leases in the country (including about 44% of the leases from which production data was available). We identify statistically significant seismicity rate changes (considering a  $s.l. = 0.05$ ) concomitant with outstanding changes in hydrocarbon production (i.e., before and after peak production) in one oil-producing lease, Val d'Agri (Figure 7B), and two gas-producing leases, Fiumetto and Rocca Cavallo (Figure 8B). In all these cases, the seismicity rate after the peak production results significantly lower with respect to the seismicity rate observed before.

It is worth to remind that spatial and temporal correlations between background seismicity rates and industrial activity, as the one highlighted in these areas, do not constitute an absolute proof to establish a causal relationship between hydrocarbon production and seismic activity. Therefore, we stress that the main value of this finding is to highlight these areas as *hotspot* zones deserving further detailed analyses to verify (or discard) possible causal relationships. If the oil or gas production in the Val d'Agri, Fiumetto and Rocca Cavallo leases played a role to alter shallow earthquake occurrences in these areas, then more sophisticated, physically-based studies are required to understand if the observed changes in seismicity rates are actually an observable consequence of physical mechanisms able to generate such changes. However, such analyses require access to detailed, geological, structural and geomechanical information.

Val d'Agri, located in the Basilicata region, is a particularly interesting case (Figure 9). This area hosts the largest onshore oil and gas field in Europe, where possibly-induced

**Table 2.** Gas-producing leases analyzed in this study. For each lease we present the estimated completeness magnitude, as well as the p values ( $p_1$  and  $p_2$ ) of the binomial test performed considering the seismicity located within a distance  $\delta x = 5$  and  $\delta x = 10$  km (the symbol ++ indicate cases with equal seismicity rates in both the pre- and the post- $t_m$  time windows, whereas the symbol – indicates cases in which both rates are zero).


Gas leases		Completeness		$\delta x = 5\text{km}$		$\delta x = 10\text{km}$	
n.	Lease name	$M_c$	since...	$p_1$ (lesser)	$p_2$ (greater)	$p_1$ (lesser)	$p_2$ (greater)
1	A.C 1.AG	2.2	1985		$7.0 \times 10^{-1}$		$7.0 \times 10^{-1}$
2	A.C 13.AS	2.1	1985	$6.0 \times 10^{-1}$		$4.6 \times 10^{-1}$	
3	A.C 17.AG	2.1	1985	$5.8 \times 10^{-1}$		$1.9 \times 10^{-1}$	
4	A.C 18.AG	2.2	1985	–	–	$6.7 \times 10^{-1}$	
5	A.C 21.AG	2.1	1985	–	–	$6.4 \times 10^{-1}$	
6	A.C 25.EA	2.2	1985	–	–	–	–
7	A.C 27.EA	2.2	1985		$3.7 \times 10^{-1}$	++	++
8	A.C 28.EA	2.2	1985	$5.2 \times 10^{-1}$			$4.8 \times 10^{-1}$
9	A.C 34.AG	1.9	1985	–	–	–	–
10	A.C. 6.AS	1.9	1985	–	–	$3.9 \times 10^{-1}$	
11	A.C 7.AS	2.2	1985	–	–		$4.9 \times 10^{-1}$
12	A.C 8.ME	1.9	1985	–	–	$1.8 \times 10^{-1}$	
13	B.C 13.AS	2.0	1985		$5.2 \times 10^{-1}$		$5.2 \times 10^{-1}$
14	B.C 14.AS	2.2	1985	$3.3 \times 10^{-1}$		$1.9 \times 10^{-1}$	
15	B.C 17.TO	2.3	1985	–	–	–	–
16	B.C 22.AG	2.2	1985	–	–	–	–
17	B.C 5.AS	2.0	1985	–	–	–	–
18	BRONTE - S.NICOLA	2.7	1985		$6.3 \times 10^{-1}$	$6.5 \times 10^{-2}$	
19	CASE SCHILLACI	2.7	1985	$4.8 \times 10^{-1}$		$1.6 \times 10^{-1}$	
20	CERVIA MARE	1.9	1985		$4.9 \times 10^{-1}$		$6.5 \times 10^{-1}$
21	D.C 4.AG	2.6	1985		$4.4 \times 10^{-1}$		$3.0 \times 10^{-1}$
22	F.C 2.AG	2.8	1985	–	–	$5.2 \times 10^{-1}$	
23	<b>FIUMETTO</b>	2.7	1985	$2.1 \times 10^{-2}$		$1.2 \times 10^{-2}$	
24	FORNOVO DI TARO	2.2	1985	–	–	$4.4 \times 10^{-1}$	
25	MASSERIA VERTICCHIO	2.2	1985	++	++	++	++
26	MISANO ADRIATICO	2.1	1985	$6.1 \times 10^{-1}$		$8.2 \times 10^{-2}$	
27	MONTE MORRONE	1.5	1985	$3.6 \times 10^{-1}$			$3.1 \times 10^{-1}$
28	MONTE URANO	2.0	1985	$4.3 \times 10^{-1}$		$2.5 \times 10^{-1}$	
29	MONTEARDONE	2.1	1985	++	++		$4.3 \times 10^{-1}$
30	MONTIGNANO	1.8	1985	$5.9 \times 10^{-1}$		$1.5 \times 10^{-1}$	
31	POLICORO	2.2	1985	–	–	$3.3 \times 10^{-1}$	
32	QUARTO	2.2	1985	–	–	–	–
33	RAGUSA	1.7	1985		$4.3 \times 10^{-1}$		$4.3 \times 10^{-1}$
34	RAPAGNANO	2.0	1985		$7.3 \times 10^{-2}$		$6.2 \times 10^{-2}$
35	RECOLETA	1.5	1985	–	–	$4.8 \times 10^{-1}$	
36	RECOVATO	1.6	1985		$3.3 \times 10^{-1}$	$5.9 \times 10^{-1}$	
37	<b>ROCCA CAVALLO</b>	2.7	1985	$1.6 \times 10^{-2}$		$4.5 \times 10^{-3}$	
38	S. ANDREA	1.8	1985	$4.6 \times 10^{-1}$			$9.5 \times 10^{-2}$
39	SAN MARCO	1.9	1985	$3.5 \times 10^{-1}$		$8.2 \times 10^{-2}$	
40	SORESINA	2.1	1985	–	–	–	–
41	TERTIVERI	2.4	1985	$1.1 \times 10^{-1}$		$1.1 \times 10^{-1}$	
42	TORRENTE CIGNO	2.0	1985	–	–	$5.5 \times 10^{-1}$	
43	VILLAFORTUNA-TRECATE	1.9	1985	–	–	–	–

seismicity has been detected in connection with wastewater reinjection at the Costa Molina 2 well (e.g., Stabile et al. 2014b; Improta et al. 2015; Buttinelli et al. 2016). Moreover, some authors reported clustered seismicity located to the south of the nearby Pertusillo artificial water reservoir, whose origin has been suggested to be induced by rapid water level changes of the Pertusillo impoundment (Valoroso et al. 2009, 2011; Stabile et al. 2014a).

Figure 9A shows the location of the Val d’Agri lease, the production wells, as well as the regional seismicity in the area (gray circles), the background seismicity identified using the NN clustering analysis technique (squares), and

the selected background seismicity located within a distance  $\delta x = 10$  km (yellow squares) and  $\delta x = 5$  km (red squares) from production wells. Only a few selected events are located in the southern part of the lease, close to the cluster of seismicity located south of the Pertusillo lake. Figure 9B shows the time series of annual oil and gas production from the Val d’Agri, and the plot of event times and depths of selected seismicity. What we actually observe is a clear reduction in the number of shallow ( $z \leq 15$  km) background seismic events just after the oil production in Val d’Agri reached its maximum in 2005.

For completeness, we also reproduced the study using in



**Figure 7.** Map of the pre- and post- $t_m$  seismicity rate changes (A) and p values (B) for the 6 oil-producing leases analyzed in this work considering the seismicity located within  $\delta x = 5$  km from producing wells.

alternative an ETAS-based (Epidemic-type Aftershock Sequences) method for declustering the seismic catalog (Ogata 1988, 1998); despite the events identified as background events using NN and ETAS methods may slightly differ, the output of the analysis using the ETAS-based declustering confirms the results.


A similar significant change in the seismicity rate has been detected in the surroundings of Fiumetto and Rocca Cavallo leases, which are located in Sicily, near Etna Volcano (Figure 10). These two leases are located in an area where other five gas-producing leases operate (namely Bronte-S.Nicola, Case Schillaci, Gagliano A, Gagliano B, and Samperi), forming a “cluster” of production leases distributed in a relatively small area (about 400 km<sup>2</sup>, Figure 10A and 10C). The results of the correlation analysis for Bronte-S.Nicola and Case Schillaci are also reported in Table 2 and Figure 8, whereas the last three leases were not included in this study because they did not meet the data requirements defined in Section 3.1.

Figure 10 shows the geographical location and gas production for Fiumetto and Rocca Cavallo leases, and the background seismicity identified for this zone. We also plotted the location of all the other leases in the area, as well as the gas production of the other two cases included in this study (Bronte-S.Nicola and Case Schillaci). It is worth noting that Gagliano has been producing gas for much more longer time with respect to the other four, but it was not possible to include it in the analysis since the peak production precedes the time interval considered in this study. In these plots we highlight all the background events located within 5 and 10 km from the production wells in Fiumetto (Figure 10A and 10B) and Rocca Cavallo (Figure 10C and 10D) leases.

In this context, a correlation analysis between seismicity


and hydrocarbon production for this area is particularly complex, probably requiring an integrated analysis considering all the leases together. Analyzing the results obtained considering each lease independently, similar to what is observed in Val d’Agri, we observe a decrease in the rate of shallow seismicity after the peak gas production in both Fiumetto and Rocca Cavallo (Figure 10B and 10D). A similar behavior is observed in the Case Schillaci site as well, but the p value in this case is not statistically significant at the *s.l.* adopted (neither considering  $\delta x = 5$  km nor  $\delta x = 10$  km). Finally, and as mentioned before, the results for Bronte-S.Nicola are contrasting when considering different  $\delta x$  values, as a probable effect of intense, shallow seismicity related to activity at the nearby Etna volcano. It is worth noting that the higher peaks in gas production in this area (in the time interval considered in this study) are reached in Fiumetto, Rocca Cavallo and Bronte-S.Nicola between  $\sim 1997$  and  $\sim 2005$ , a period in which the reduction in the number of events starts to be evident in the zone. However, it should be kept in mind that this area is a particularly complex case due to different factors, such as the closeness of different active leases and the proximity of a seismically active volcano, therefore any further analysis probably requires taking into account the whole cluster of leases together (considering also the influence in this particular area of possible stress perturbations caused by activity at the nearby Etna volcano).

In the areas highlighted by low p values in this study we observe that the seismicity after  $t_m$  occurs at a lower rate with respect to the seismicity occurring before the peak production. In the case of a link with fluid withdrawal, such observation could result from either an increase in seismicity rate associated with the phase preceding  $t_m$ , when the fluid withdrawal rate had an increasing trend (see e.g., Figure 5E for reference), or because seismicity is somehow inhibited as



**Figure 8.** Map of the pre- and post- $t_m$  seismicity rate changes (A) and p values (B) for the 43 gas-producing leases analyzed in this work considering the seismicity located within  $\delta x = 5$  km from producing wells.





**Figure 9.** Seismic and production data used in the analysis for the Val d'Agri lease. (A) Background seismicity, highlighting events located at distances  $\delta x = 5$  and 10km from the production wells. (B) Time series of production data and plot of the temporal occurrence and depths of seismic events


the fluid withdrawal process goes on (which results particularly evident after  $t_m$ ). An analysis of the seismicity rate before and after the start of fluid withdrawal operations would be useful to understand which of the two scenarios dominates; however, the seismic data for times preceding the start of operations is too scarce for reliable and systematic analyses. Looking however at the seismicity and production data in Figures 9 and 10, it is interesting to note that background seismic events are anyhow present before the start of production in the highlighted cases, and this observation contrasts with the apparent seismicity rate drop observed after the peak production; this observation leads us to think that the change in rate could primarily be due to a decrease in the number of earthquakes after the peak.

In an attempt to understand if deformative phenomena associated with fluid withdrawal from the crust may explain an apparent inhibition of shallow seismicity, we analyzed

a hypothetical case of deformation related to reservoir depletion and calculate the Coulomb stress changes on fault planes optimally oriented for failure (see e.g., Lin & Stein 2004; Toda et al. 2005). In practice, we assume the contraction of a sub-horizontal dike in an elastic halfspace. For this theoretical exercise we use geomechanical properties of the Val d'Agri zone taken from literature. The modeled dike has an area that roughly covers the area outlined by the production wells active in the Val d'Agri lease (that we assume as an approximate proxy of the actual areal distribution of the reservoir), and is located at a depth of  $\sim 4$  km below surface (which approximately corresponds to the depth of the reservoir in Val d'Agri, as reported in e.g., Stabile et al. 2014b).

In agreement with the regional stress field inferred for this region by Cucci et al. (2004), we assume an extensional tectonic environment (characteristic of normal faulting regimes) with  $S_V > S_{H_{\max}} > S_{H_{\min}}$ , and where the





**Figure 10.** Seismic and production data used in the analysis for (A) Fiumetto and (B) Rocca Cavallo leases. Top: map showing background seismicity, highlighting all the events located at distances  $\delta x = 5$  and 10 km from the production wells. Bottom Time series of production data and plot of the temporal occurrence and depths of seismic events. Production data and location of the nearby Bronte-S. Nicola and Case Schillaci leases are also shown for reference.

minimum horizontal stress is oriented in the direction NNE-SSW. The results of the Coulomb stress change resolved on optimally oriented planes for this setting is shown in Figure 11; cold (blue) colors in Figure 11 indicate zones with negative Coulomb stress, outlining areas in which the modeled deformation would tend to inhibit seismicity. Such results depict a simple example in which fluid withdrawal operations are allowed to induce a contraction of the reservoir; in such a case, the resulting deformation would tend to inhibit seismicity at least in the shallow crust around the area of a depleting reservoir. Therefore, under some circumstances it seems feasible to observe a reduction in seismicity rates following the period of largest hydrocarbon production.

## 6 CONCLUSIONS


In areas characterized by high levels of natural seismicity, the identification of human-induced seismicity is a difficult task for which virtually any result can be a source of controversy. In Italy, a relatively high number of oil- and gas-producing leases have been operating in the last decades, many of which are located in the surroundings of seismically active regions (e.g., Sicily, and the central and southern Apennines). Besides hydrocarbon production, underground waste water injection is also performed in different areas of the country, but these activities were not considered in this study because injection-related data are not publicly available.

Therefore, our analyses are focused on areas where fluids (oil and gas) are withdrawn from the crust. Performing

detailed, physically-based analyses at the national scale to identify cases of anthropogenic seismicity in Italy may be considered an intractable problem. Consequently, we implemented a large-scale screening procedure aiming at tracking measurable phenomena, such as, e.g., changes in seismicity rates, that plausibly could occur if notable interactions between fluid withdrawal from the crust and nearby seismicity sources are actually occurring in a given area. We stress however that spatial-temporal correlations between proxies of industrial activity and significant changes in seismicity rates do not provide proofs of causal relationships between hydrocarbon production and seismic activity; rather, studying such correlations gives us the possibility of performing large-scale, systematic analyses of a huge amount of seismic and production data, and to identify in this way *hotspot areas* where to focus more detailed research to verify, in a later stage, possible causal relationships.

In this context, we analyzed seismicity rates before and after peak production in 6 oil-producing and 43 gas-producing leases, and evaluate the significance of potential seismicity rate changes. Such cases are about the 44% of the development leases active in Italy (the other 56% were not analyzed due to data limitation constraints). The main findings resulting from this study can be summarized as follows:

- When considering the background seismicity located within 5 km from the production wells, about 50% of the oil-producing and 46% of the gas-producing leases analyzed in this study do not show any change in seismicity rate before and after the time at which the peak production is reached; in most of these cases the seismicity rate is zero (basically due to the short distance considered). The percentage of oil- and



**Figure 11.** Coulomb stress change (bar) calculated on fault planes optimally oriented for failure, considering as the source of deformation the contraction of a sub-horizontal dike located at  $\sim 4$  km depth. Both plots show a vertical cross section in the NS direction, crossed by horizontal layers located at (A) 1.5 km and (B) 5.0 km below the surface.

gas-producing leases with no observed change in the seismicity rate reduces respectively to about 17% and 23% when considering seismicity located within 10 km from production wells.

- None of the observed cases of seismicity rate increase after the hydrocarbon peak production is statistically significant (at a  $s.l. = 0.05$ ); such cases include about 17% of the oil- and 21 to 25% of the gas-producing leases analyzed in this study (depending on  $\delta x$ ). This result is obtained selecting background seismicity within both 5 and 10 km from production wells.

- Regarding the cases exhibiting a seismicity rate decrease after the hydrocarbon peak production, our results suggest that the observed seismicity rate change is statistically significant ( $s.l. = 0.05$ ) for one oil-producing lease (the Val d'Agri, in Basilicata) and two gas-producing leases (Fiumetto and Rocca Cavallo, in Sicily). These three leases, which basically correspond with two geographical areas since the later two are adjacent to each other (and to other active leases as Bronte S.Nicola, Case Schillaci, and Gagliano, all clustered in a relatively small area) can be identified as hotspots deserving future research to study whether there may exist a causal relationship between the hydrocarbon production and the observed reduction in seismicity rate following the peak production.

In conclusion, our analyses highlight areas near some oil- and gas-producing leases in Italy where the seismicity rate reduces after peak production is reached (as compared to the seismicity rate preceding it). We emphasize however that our results just put in evidence a correlated change between the rates of shallow seismicity and hydrocarbon production in these areas, and that assessing actual causal relationships between these two processes will require further detailed, physically-based research. Despite this, we argue that should a physical link exist between these processes, the observed seismicity rate reduction could either be due to increased seismicity during the progressive increase in production rate before  $t_m$ , or to actual seismicity rate reduction after  $t_m$ . This second scenario would put in evidence possible processes of seismicity inhibition. Considering that the occurrence of seismicity before the start of hydrocar-

bon production in the hotspot areas contrasts with the reduction of events observed after the peak production, we suspect that the seismicity inhibition is a plausible hypothesis in these cases. With a simple theoretical exercise based on modelling Coulomb stress changes we showed that, at least under some simplified conditions, inhibition of seismicity is actually possible. To our knowledge, such an effect of hydrocarbon production on seismicity has not been clearly documented in literature until now.

Our observations draw attention to an interesting research problem: the characteristics and implications of increased seismicity caused by anthropogenic activities (e.g., pressurized fluid injection) have so far had a prominent role in research on induced seismicity; the implications, instead, of anthropic processes potentially capable of inhibiting seismicity — in particular, on seismic hazard in seismically active regions — have received less attention, or have been neglected. We consider that more efforts to study the mechanisms and the possible consequences of anthropogenically-driven seismicity inhibition, as well as to document possible cases of such phenomenon, are required.

## CONFLICT OF INTEREST STATEMENT

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Author Contributions

AG and LF conceived the experiment, implemented the methods used to analyze the data, and interpreted the achieved results. LF performed the spatial and temporal declustering of the seismic catalog. AG performed the correlation analyses and the Coulomb stress calculations. AM contributed in the discussion of results and in writing. IA collected the industrial data and created the database of hydrocarbon production used in the study. AG and LF wrote the bulk of the text, with inputs from all the co-authors. All the authors have read and approved the manuscript.

## Funding

This study was performed with the support of Clypea, the Innovation Network for Future Energy financed by the Italian Ministry of Economic Development, Direzione Generale per le Infrastrutture e la Sicurezza dei Sistemi Energetici e Geominerari (MISE - DGISSEG).

## Data Availability Statement

The seismic data set analyzed in this study can be accessed through the HORUS (HOMogenized instRumental Seismic catalog) website: <http://horus.bo.ingv.it/> (see also Lollì et al. 2020a,b). The hydrocarbon production data are publicly available in the website of the “Direzione generale per le infrastrutture e la sicurezza dei sistemi energetici e geominerari (DGISSEG), Ufficio nazionale minerario per gli idrocarburi e le georisorse (UNMIG)”, from the Italian Ministry of Economic Development (MISE): <https://unmig.mise.gov.it/index.php/it/dati>. The ViDEPI project – Visibilit  dei dati afferenti all’attivit  di esplorazione petrolifera in Italia (Visibility of data related to mineral exploration activities in Italy, in Italian), is available at <https://www.videpi.com/>. (all sites, last accessed: February 2021)

## ACKNOWLEDGMENTS

The authors thank the invaluable support from the “Direzione generale per le infrastrutture e la sicurezza dei sistemi energetici e geominerari (DGISSEG), Ufficio nazionale minerario per gli idrocarburi e le georisorse (UNMIG)”, from the Italian Ministry of Economic Development (MISE), as well as their Innovation Network for Future Energy (Clypea). Maps were plotted using Generic Mapping Tools (GMTs) (Wessel et al. 2019, <https://www.generic-mapping-tools.org/>, last accessed January 2021). The authors thank also Anna Nardi, Antonio Rossi and Patrizia Battelli for sharing data on the history of the Italian seismic monitoring network. Any use of trade, firm, or product names is for descriptive purposes only. The results and conclusions of this work do not necessarily represent the position of MISE - DGISSEG.

## REFERENCES

Amato, A. & Montone, P., 1997. Present-day stress field and active tectonics in southern peninsular Italy, *Geophysical Journal International*, **130**(2), 519–534.

Bertello, F., Fantoni, R., Franciosi, R., Gatti, V., Ghielmi, M., & Pugliese, A., 2010. From thrust-and-fold belt to foreland: hydrocarbon occurrences in Italy, *Geological Society, London, Petroleum Geology Conference series*, **7**(1), 113126.

Braun, T., Cesca, S., Khn, D., Martirosian-Janssen, A., & Dahm, T., 2018. Anthropogenic seismicity in Italy and its relation to tectonics: State of the art and perspectives, *Anthropocene*, **21**, 80 – 94.

Buttinelli, M., Improta, L., Bagh, S., & Chiarabba, C., 2016. Inversion of inherited thrusts by wastewater injection induced seismicity at the val d’agri oilfield (Italy), *Scientific Reports*, **6**(1), 37165.

Calamita, F., Cello, G., Deiana, G., & Paltrinieri, W., 1994. Structural styles, chronology rates of deformation, and timespace relationships in the UmbriaMarche thrust system (central Apennines, Italy), *Tectonics*, **13**(4), 873–881.

Cazzini, F., Zotto, O. D., Fantoni, R., Ghielmi, M., Ronchi, P., & Scotti, P., 2015. Oil and gas in the Adriatic foreland, Italy, *Journal of Petroleum Geology*, **38**(3), 255–279.

Cello, G. & Mazzoli, S., 1998. Apennine tectonics in southern Italy: A review, *J. Geodyn.*, **27**(2), 191–211.

Cesca, S., Grigoli, F., Heimann, S., Gonzalez, A., Buforn, E., Maghsoudi, S., Blanch, E., & Dahm, T., 2014. The 2013 September–October seismic sequence offshore Spain: a case of seismicity triggered by gas injection?, *Geophysical Journal International*, **198**(2), 941–953.

Cucci, L., Pondrelli, S., Frepoli, A., Mariucci, M. T., & Moro, M., 2004. Local pattern of stress field and seismogenic sources in the Pergola-Melandro basin and the Agri valley (Southern Italy), *Geophysical Journal International*, **156**(3), 575–583.

D’Agostino, N., Giuliani, R., Mattone, M., & Bonci, L., 2001. Active crustal extension in the central Apennines (Italy) inferred from GPS measurements in the interval 1994–1999, *Geophys. Res. Lett.*, **28**(10), 2121–2124.

Dahm, T., Cesca, S., Hainzl, S., Braun, T., & Krger, F., 2015. Discrimination between induced, triggered, and natural earthquakes close to hydrocarbon reservoirs: A probabilistic approach based on the modeling of depletion-induced stress changes and seismological source parameters, *Journal of Geophysical Research: Solid Earth*, **120**(4), 2491–2509.

Davis, S. D. & Frohlich, C., 1993. Did (or will) fluid injection cause earthquakes? - criteria for a rational assessment, *Seismological Research Letters*, **64**(3–4), 207–224.

Davis, S. D. & Nyffenegger, P. A., 1995. The 9 April 1993 earthquake in south-central Texas: Was it induced by fluid withdrawal?, *Bulletin of the Seismological Society of America*, **85**(6), 1888–1895.

Ellsworth, W. L., 2013. Injection-induced earthquakes, *Science*, **341**(6142).

Evans, K. F., Zappone, A., Kraft, T., Deichmann, N., & Moia, F., 2012. A survey of the induced seismic responses to fluid injection in geothermal and CO<sub>2</sub> reservoirs in Europe, *Geothermics*, **41**, 30–54.

Foulger, G. R., Wilson, M. P., Gluyas, J. G., Julian, B. R., & Davies, R. J., 2018. Global review of human-induced earthquakes, *Earth-Science Reviews*, **178**, 438 – 514.

Garcia-Aristizabal, A., 2018. Modelling fluid-induced seismicity rates associated with fluid injections: examples related to fracture stimulations in geothermal areas, *Geophysical Journal International*, **215**(1), 471493.

Garcia-Aristizabal, A., Danesi, S., Braun, T., Anselmi, M., Zaccarelli, L., Famiani, D., & Morelli, A., 2020. Epistemic Uncertainties in Local Earthquake Locations and Implications for Managing Induced Seismicity, *Bulletin of the Seismological Society of America*, **110**(5), 2423–2440.

Grasso, J.-R., Amorese, D., & Karimov, A., 2019. Anthropogenic seismicity as aftershocks for geo-resource production? Implications for Mmax estimates (reservoir impoundment cases), *Geophysical Journal International*, **219**(2), 958–967.

Grasso, J.-R., Karimov, A., & Amorese, D., submitted. The longest seismic swarm triggered by fluid manipulations? Lacq, France, 1969–2016, *Submitted to Bulletin of the Seismological Society of America*.

Grigoli, F., Cesca, S., Priolo, E., Rinaldi, A. P., Clinton, J. F., Stabile, T. A., Dost, B., Fernandez, M. G., Wiemer, S., & Dahm, T., 2017. Current challenges in monitoring, discrimination, and management of induced seismicity related to underground industrial activities: A European perspective, *Reviews of Geophysics*, **55**(2), 310340.

Hainzl, S., Kraft, T., Wassermann, J., Igel, H., & Schmedes, E., 2006. Evidence for rainfall-triggered earthquake activity, *Geophysical Research Letters*, **33**(19).

Handy, M. R., M. Schmid, S., Bousquet, R., Kissling, E., & Bernoulli, D., 2010. Reconciling plate-tectonic reconstructions of Alpine Tethys with the geological-geophysical record of spreading and subduction in the Alps, *Earth-Science Rev.*, **102**(3–4), 121–158.

Improta, L., Valoroso, L., Piccinini, D., & Chiarabba, C., 2015. A detailed analysis of wastewater-induced seismicity in the val d’agri oil field

- (Italy), *Geophysical Research Letters*, **42**(8), 2682–2690.
- IS-EPOS, 2016. Completeness magnitude estimation [web application/source code] – <https://tcs.ah-epos.eu/>.
- Leptokaropoulos, K., Staszek, M., Lasocki, S., Martinez-Garzn, P., & Kwiatek, G., 2017. Evolution of seismicity in relation to fluid injection in the North-Western part of The Geysers geothermal field, *Geophysical Journal International*, **212**(2), 11571166.
- Leptokaropoulos, K. M., Karakostas, V. G., Papadimitriou, E. E., Adamaki, A. K., Tan, O., & İnan, S., 2013. A homogeneous earthquake catalog for western turkey and magnitude of completeness determination, *Bulletin of the Seismological Society of America*, **103**(5), 2739–2751.
- Lin, J. & Stein, R. S., 2004. Stress triggering in thrust and subduction earthquakes and stress interaction between the southern san andreas and nearby thrust and strike-slip faults, *Journal of Geophysical Research: Solid Earth*, **109**(B2).
- Lolli, B., Randazzo, D., Vannucci, G., & Gasperini, P., 2020a. The Homogenized Instrumental Seismic Catalog (HORUS) of Italy from 1960 to Present, *Seismological Research Letters*, **91**(6), 3208–3222.
- Lolli, B., Vannucci, G., Randazzo, D., & Gasperini, P., 2020b. Horus - homogenized instrumental seismic catalog – <http://horus.bo.ingv.it/>.
- Molina, I., Velsquez, J. S., Rubinstein, J. L., Garcia-Aristizabal, A., & Dionicio, V., 2020. Seismicity induced by massive wastewater injection near Puerto Gaitn, Colombia, *Geophysical Journal International*, **223**(2), 777–791.
- Mulargia, F. & Bizzarri, A., 2014. Anthropogenic triggering of large earthquakes, *Scientific Reports*, **4**, 6100–6107.
- Ogata, Y., 1988. Statistical models for earthquake occurrences and residual analysis for point processes, *Journal of the American Statistical Association*, **83**(401), 9–27.
- Ogata, Y., 1998. Space-time point-process models for earthquake occurrences, *Annals of the Institute of Statistical Mathematics*, **50**(401), 379–402.
- Orlecka-Sikora, B., Lasocki, S., Kocot, J., Szeplieniec, T., Grasso, J. R., Garcia-Aristizabal, A., Schaming, M., Urban, P., Jones, G., Stimpson, I., Dineva, S., Sałek, P., Leptokaropoulos, K., Lizurek, G., Olszewska, D., Schmittbuhl, J., Kwiatek, G., Blanke, A., Saccorotti, G., Chodzińska, K., Rudziński, Ł., Dobrzycka, I., Mutke, G., Barański, A., Pierzyna, A., Kozlovskaya, E., Nevalainen, J., Kinscher, J., Sileny, J., Sterzel, M., Cielesta, S., & Fischer, T., 2020. An open data infrastructure for the study of anthropogenic hazards linked to georesource exploitation, *Scientific Data*, **7**(1), 89.
- Pagliuca, N. M., Battelli, P., Berardi, M., Modica, G., Castellano, C., Melorio, C., Nardi, A., Pirro, M., Arcoraci, L., & Baccheschi, P., 2020. Bollettino Sismico Italiano (BSI), III quadrimestre 2019.
- Parsons, T. & Velasco, A. A., 2009. On near-source earthquake triggering, *Journal of Geophysical Research: Solid Earth*, **114**(B10).
- Pintori, F., Serpelloni, E., Longuevergne, L., Garcia, A., Faenza, L., Dalberto, L., Gualandi, A., & Elina, B. M., 2021. Mechanical response of shallow crust to groundwater storage variations: Inferences from deformation and seismic observations in the eastern southern alps, Italy, *Journal of Geophysical Research: Solid Earth*, **126**.
- Schoenball, M., Davatzes, N. C., & Glen, J. M. G., 2015. Differentiating induced and natural seismicity using space-time-magnitude statistics applied to the coso geothermal field, *Geophysical Research Letters*, **42**(15), 62216228.
- Shapiro, S. A., Dinske, C., & Kummerow, J., 2007. Probability of a given-magnitude earthquake induced by a fluid injection, *Geophysical Research Letters*, **34**(22).
- Shapiro, S. A., Dinske, C., Langenbruch, C., & Wenzel, F., 2010. Seismogenic index and magnitude probability of earthquakes induced during reservoir fluid stimulations, *The Leading Edge*, **29**(3), 304309.
- Stabile, T. A., Giocoli, A., Lapenna, V., Perrone, A., Piscitelli, S., & Telesca, L., 2014a. Evidence of low-magnitude continued reservoir-induced seismicity associated with the pertusillo artificial lake (southern Italy), *Bulletin of the Seismological Society of America*, **104**(4), 1820–1828.
- Stabile, T. A., Giocoli, A., Perrone, A., Piscitelli, S., & Lapenna, V., 2014b. Fluid injection induced seismicity reveals a NE dipping fault in the southeastern sector of the high Agri valley (southern Italy), *Geophysical Research Letters*, **41**(16), 5847–5854.
- Tahir, M., Grasso, J.-R., & Amorse, D., 2012. The largest aftershock: How strong, how far away, how delayed?, *Geophysical Research Letters*, **39**(4).
- Toda, S., Stein, R. S., Richards-Dinger, K., & Bozkurt, S. B., 2005. Forecasting the evolution of seismicity in southern California: Animations built on earthquake stress transfer, *Journal of Geophysical Research: Solid Earth*, **110**(B5).
- UNMIG, 2020. Unmig databook 2020. ministero dello sviluppo economico, direzione generale per le infrastrutture e la sicurezza dei sistemi energetici e geominerari, ufficio nazionale minerario per gli idrocarburi e le georisorse. link: <https://unmig.mise.gov.it/images/stat/databook-2020.pdf> (last accessed: November 2020).
- Valoroso, L., Impropa, L., Chiaraluce, L., Di Stefano, R., Ferranti, L., Govoni, A., & Chiarabba, C., 2009. Active faults and induced seismicity in the Val d'Agri area (Southern Apennines, Italy), *Geophysical Journal International*, **178**(1), 488–502.
- Valoroso, L., Impropa, L., De Gori, P., & Chiarabba, C., 2011. Upper crustal structure, seismicity and pore pressure variations in an extensional seismic belt through 3-d and 4-d vp and vp/vs models: The example of the val d'agri area (southern Italy), *Journal of Geophysical Research: Solid Earth*, **116**(B7).
- van der Voort, N. & Vanclay, F., 2015. Social impacts of earthquakes caused by gas extraction in the province of Groningen, the Netherlands, *Environmental Impact Assessment Review*, **50**, 1–15.
- van Hinsbergen, D. J., Torsvik, T. H., Schmid, S. M., Maenco, L. C., Maffione, M., Vissers, R. L., Gürer, D., & Spakman, W., 2020. Orogenic architecture of the Mediterranean region and kinematic reconstruction of its tectonic evolution since the Triassic, *Gondwana Res.*, **81**, 79–229.
- van Stiphout, T., Zhuang, J., & Marsan, D., 2012. Seismicity declustering, community online resource for statistical seismicity analysis, available at <http://www.corssa.org>.
- Wells, D. L. & Coppersmith, K. J., 1994. New empirical relationships among magnitude, rupture length, rupture width, rupture area, and surface displacement, *Bulletin of the Seismological Society of America*, **84**(4), 974–1002.
- Wessel, P., Luis, J. F., Uieda, L., Scharroo, R., Wobbe, F., Smith, W. H. F., & Tian, D., 2019. The generic mapping tools version 6, *Geochemistry, Geophysics, Geosystems*, **20**(11), 5556–5564.
- Wiemer, S. & Wyss, M., 2000. Minimum magnitude of completeness in earthquake catalogs: Examples from Alaska, the western United States, and Japan, *Bulletin of the Seismological Society of America*, **90**(4), 859–869.
- Wonnacott, T. & Wonnacott, R. J., 1977. *Introductory Statistics*, Wiley.
- Zaliapin, I. & Ben-Zion, Y., 2013. Earthquake clusters in southern California I: Identification and stability, *Journal of Geophysical Research: Solid Earth*, **118**.
- Zaliapin, I. & Ben-Zion, Y., 2016. Discriminating Characteristics of Tectonic and Human-Induced Seismicity, *Bulletin of the Seismological Society of America*, **106**(3), 846–859.
- Zhang, H., Eaton, D. W., Li, G., Liu, Y., & Harrington, R. M., 2016. Discriminating induced seismicity from natural earthquakes using moment tensors and source spectra, *Journal of Geophysical Research: Solid Earth*, **121**(2), 972993.
- Zhuang, J., Ogata, Y., & Vere-Jones, D., 2002. Stochastic declustering of space-time earthquake occurrences, *Journal of the American Statistical Association*, **97**.