

EarthArxiv Preprint

5 The following unpublished, not yet peer-reviewed
manuscript was submitted to the journal
Geoscience Communication on March 22, 2021.

10

You can download the teaching tool detailed in this manuscript:

<https://doi.org/10.5281/zenodo.4597618>

15

Please feel free to contact the author with suggestions or tips:

Bryan Lougheed

Department of Earth Sciences, Uppsala University, Sweden

bryan.lougheed@geo.uu.se

20

bonne journée

Orbital, the Box - An interactive educational tool for in-depth understanding of astronomical climate forcing.

25 Bryan C. Lougheed

Department of Earth Sciences, Uppsala University, Uppsala, Sweden

bryan.lougheed@geo.uu.se

Abstract

30

“*Orbital, the Box*” provides an interactive tool with graphical user interface (GUI) for stimulating active, visual learning for understanding of astronomical climate forcing. This cross-platform tool can be run locally on a personal computer using a standard web browser environment with no need for plugins, thus maximising accessibility for students and teachers alike. The tool facilitates in the development of a holistic and quantitative understanding of astronomical climate forcing by allowing students to independently vary orbital parameters, after which they can instantaneously see the resulting effect upon the seasonal and latitudinal distribution of solar irradiance arriving at the top of the Earth’s atmosphere. Such an approach follows a classic controlled experimental design whereby one parameter can be changed while all others are kept constant. This experimental tool can be deployed as a virtual laboratory, including within a flipped classroom setting, to promote active learning of traditionally challenging concepts such as the roles of eccentricity and precession in astronomical climate forcing, and in particular their interaction with Kepler’s second law and the subsequent consequences for season length.

45 **1.0 Introduction**

Astronomical climate forcing is the theory that changes in a Earth’s orbital configuration can influence its climate (Herschel, 1832; Adhémar, 1842; Croll, 1864). A major breakthrough in the furthering of this theory was achieved by the exhaustive calculations of the Serbian civil engineer Milutin Milanković who, in the early 20th century, used known orbital parameters to calculate secular changes in the seasonal and latitudinal distribution of incoming irradiance at the top of the Earth’s atmosphere throughout the ages, and related these changes to the Quaternary ice ages. The decades of work is summed up in Milanković’s *Kanon der Erdebestrahlung und seine Anwendung auf das Eiszeitenproblem* (Milanković, 1941). Changes in the aforementioned seasonal and

55 latitudinal distribution are an important driver of long-term climate changes on planet Earth and understanding thereof is a foundational element of geoscience education. However, quantitative teaching of the separate contributions of the various orbital parameters upon the irradiance distribution can often be a challenging task.

60 The three main parameters that can change the aforementioned seasonal and latitudinal distribution are obliquity of the ecliptic, eccentricity of the orbital ellipse and general precession (the combined effect of axial and apsidal precession), hereafter referred to simply as obliquity, eccentricity and precession. As outlined by Wampler (2000a), introductory teaching of astronomical climate forcing within geosciences should include the following:

65

(1) A description of the changes in physical motion caused by the parameters and some explanation about the causes.

70

(2) A description of the influence of the parameters upon the seasonal and latitudinal distribution of incoming irradiance.

Regarding the first point, teaching of astronomical climate forcing within geosciences curricula does impart a general explanation of the physical motion of the three main parameters (eccentricity, obliquity and precession) that is sufficient enough for purposes of palaeoclimate teaching. The physical causes of the processes (i.e. gravitational interaction between the celestial bodies) are also touched upon.

The second point, a description of the influence of the parameters upon the seasonal and latitudinal distribution of incoming irradiance, is of great importance for understanding of palaeoclimate, and is generally more difficult to teach in the case of some parameters, especially within a lecture and/or textbook format. In the case of obliquity, the influence upon the irradiance distribution is fortunately relatively straightforward to teach once the understanding is established that seasons on Earth are caused by obliquity in combination with the Earth's orbit around the Sun. From that base understanding, one can subsequently convey that greater obliquity will lead to greater seasonality and vice-versa.

Teaching of the more nuanced changes caused by the combined influence of both eccentricity and precession can be challenging because it must necessarily invoke Kepler's second law, i.e. that the

Earth will orbit at a faster speed when closer to the Sun and vice-versa. In my own experience, the
90 influence of these phenomena upon the Earth's climate are difficult to convey in a lecture setting,
and exam questions relating to obliquity tend to be better answered than questions relating to
eccentricity and precession, indicating a failure to fully convey these concepts. Communication
with colleagues across multiple departments (Earth Sciences and Physics) confirmed that, in
particular, the concept of precession consistently represents a "muddiest point" in multiple
95 disciplines. Due to the challenges in teaching this particular concept, and possibly also due to the
fact that Keplerian concepts are difficult to convey graphically, textbooks traditionally touch on it
only briefly. This has the potential for students developing incomplete understanding regarding
astronomical climate forcing (Wampler, 2000b, 2000a). This incomplete understanding
subsequently has the potential to persist into the scientific literature in some cases, as noted by
100 Bol'shakov (2017). Misrepresentations of Milanković's work in the literature include the claim that
Milanković theory of astronomical forcing of glacial ablation/accumulation is based on high-
latitude irradiance received on the day of the summer solstice (often erroneously referred to as
"June 21" for all geological ages), whereas Milanković explicitly referred to not a single day, but
the caloric summer half of the year, the *sommerhalbjahr* (Milanković, 1941). This type of
105 misrepresentation, in essence an oversimplification of Milanković theory, likely stems from from
incomplete undergraduate teaching of the influence of the Keplerian orbit upon the Earth's season
lengths and subsequent solar irradiance profile. Relatively recent studies in the literature have
sought to comprehensively readdress such simplifications (Huybers, 2006, 2011; Berger et al.,
2010), but these studies are not aimed at an undergraduate audience. Other misrepresentations can
110 include, e.g., orbital tuning of climate data to single orbital parameters (such as eccentricity) and
citing Milanković as a justification, whereas Milanković theory pertains to the collective
contribution of all orbital parameters to the distribution of incoming irradiance, and specifically
with respect to Quaternary ice ages.

115 Interactive pedagogical method/tools can improve existing geoscience teaching methods, increasing
attainment of understanding and student motivation when implemented within, e.g. a "flipped
classroom" (Bykerk-Kauffman, 1995; Huguet et al., 2020). In the case of astronomical climate
forcing, it is obviously not possible to apply the traditional interactive methods used within
geosciences, e.g. a fieldwork or laboratory element. However, an interactive tool with graphical
120 user interface (GUI) can be developed to offer a virtual laboratory element (Kostadinov and Gilb,
2014). Here, *Orbital, the Box* is presented, which can be used to help imparting an understanding of
orbital parameters upon the distribution of irradiance across latitudes and seasons, with the aim of

developing both a conceptual (specifically, a sense of the physical mechanism) and quantitative understanding of processes. Quantitative understanding is particularly important and is often overlooked within the geosciences (Manduca et al., 2008). Such a holistic and quantitative understanding of the effect of orbital parameters upon the Earth's irradiance profile is vital before the student can proceed on to other related subjects, such as learning about the millennial-scale time frequencies of past changes in eccentricity, obliquity and precession, and how they may or may not be manifested in the palaeoclimate record across vast timescales.

130

2.0 Method

2.1 GUI interface

135

The user is presented with a control panel in the form of three slider controls: (1) eccentricity, (2) obliquity and (3) ω , the geocentric solar longitude (λ) at which perihelion occurs, changes of which are caused by precessional processes. The user can set 15 unique values for eccentricity (covering a range of typical Earth values from 0.001 to 0.058), 15 unique values for obliquity (covering a range of typical Earth values between 21.8° and 24.5°) and 13 unique values for ω (from 0° to 360°). Each time a slider is adjusted, the image to the right displaying the irradiance distribution and orbital information (distance from sun, orbital speed) is updated instantaneously, helping to facilitate rapid active learning. Each of the three controls can be set independently, meaning that there are a total of $15 \times 15 \times 13 = 2925$ possible unique combinations to be explored. The reason the results can be loaded instantaneously on a standard personal computer is because the images containing the irradiance distributions are not computed live, instead, the respective images for all 2925 unique combinations have been pre-computed and can thus be instantly called upon when required. The control panel system used to call the images is written in HTML, CSS and Javascript, all of which are included with the major web browsers on all major computing platforms. As such, the tool can be run on desktop computers, laptops or tablets. Publicly available tutorials were used to design the CSS and Javascript elements (w3schools.com, 2020).

145

150

2.2 Computation of solar irradiance

155 The 2925 unique irradiance scenarios were computed in Matlab 2019a using scripts based on established methodologies. These scripts were specifically written for the development of this

educational tool, but could be used in other settings (e.g. research), and have therefore been made publicly available on Github (http://www.github.com/bryanlougheed/orbital_the_box/).

160 The image of annual irradiance distribution is calculated as follows. For each unique combination of eccentricity, obliquity and ω , a 180° latitude by 365.2 day grid of irradiance is produced, using a respective resolution of 0.5° and 0.2 days. For the centre of each grid cell, 24 hr mean irradiance (daily Q_{mean}) is calculated in Wm^{-2} following Berger (1978), using a solar constant of 1361 Wm^{-2} . The aforementioned procedure requires λ as input, and I have calculated the λ associated with each
165 0.2 day increment following the standard Keplerian methods outlined by Meeus (1998) involving a binary search solution for the Kepler equation developed by Sinnott (1985). To the best of my knowledge, this computational procedure for the Kepler equation was first highlighted within the geosciences by Kostadinov and Gilb (2014). The Berger (1978) irradiance procedure furthermore calls for geocentric latitude as input, but I have substituted it here by geographic latitude to account
170 for the ellipsoidal shape of the Earth (using the WGS84 reference ellipsoid). I validated this substitution approach against the incoming irradiance angle correction approach of Van Hemelrijck (1983). In order to better facilitate visual interpretation of the irradiance grid showing daily Q_{mean} , it is contoured at 25 Wm^{-2} intervals from 0 to 650 Wm^{-2} . The annual Q_{mean} for each latitude is calculated by taking the mean of all 24 hr Q_{mean} values calculated for each 0.2 day increment. Earth's
175 distance from the Sun for each λ increment is calculated following the standard Keplerian methods outlined in Meeus (1998), assuming a semi-major axis of 1 AU. The corresponding orbital speed is calculated following the law of orbital energy invariance.

3.0 Overview of the GUI tool

180 3.1 Graphical representation

Within palaeoclimate teaching, current representations of the effect of orbital parameters upon the distribution of irradiance are sometimes not not described correctly (Wampler, 2000b, 2000a), possibly due to the limitations of the text format. Here, I have created a graphical representation of that captures many complex concepts in one glance, shown in Figure 1 for the orbital configuration
185 of the 21st century (eccentricity: 0.0167, obliquity: 23.4° , geocentric longitude of perihelion: 283°). This static graphical representation already allows a student to develop an understanding of the effects of eccentricity and precession upon the Earth's irradiance profile. For example, one sees that nowadays the Earth is closest to the Sun (i.e. at perihelion) approximately near the 280th day of the tropical year, approximately coinciding with southern hemisphere summer. This means that
190 southern hemisphere summer receives slightly higher peak irradiance than the northern hemisphere

summer does, as is visualised in the figure. However, one can also see how the Earth's orbital speed is slightly faster during southern hemisphere summer, which translates to southern hemisphere summer being slightly shorter than northern hemisphere summer (as indicated by the day durations indicated in the figure). The season length compensates for the difference in irradiance received, meaning that any latitude will receive the same mean irradiance throughout the course of the entire tropical year as its corresponding latitude in the opposite hemisphere (as can be seen in the right panel).




Figure 1. Graphical representation of the Earth's irradiance profile in the early 21st century (eccentricity: 0.0167, obliquity: 23.4°, geocentric longitude of perihelion: 283°). **Top panel:** The Earth's distance from the sun and the speed of its orbit throughout the tropical year, starting on the day of the spring equinox. Equinox and solstice days are indicated by dashed lines. **Bottom-left panel:** Distribution of irradiance (24-hour mean irradiance in Wm^{-2}) for the latitudes of Earth throughout the tropical year, also starting on the day of the spring equinox. Equinox and solstice days are indicated by dashed lines. **Bottom-right panel:** The mean irradiance received at each latitude throughout the course of the entire tropical year.

205

210

215




Figure 2. Graphical representation of the Earth's orbital configuration in the early 21st century (eccentricity: 0.0167, geocentric longitude of perihelion: 283°). The orbital shape has been drawn to scale. The size of the Earth and Sun have been increased for visual clarity.

220

In addition to the graphical representation of the irradiance distribution reaching the Earth, a scale drawing of the elliptical orbit of the Earth, with the Sun at the focal point, is also provided (Figure 2). This figure allows the student to visually analyse the aphelion and perihelion distances, and how the tropical year is oriented with regards to the orbit due to general precession.

3.2 Interactive representation

225

230

235

The graphical representations shown in Figure 1 and Figure 2 are a static representation of a single orbital configuration corresponding to the orbital parameters of the early 21st century. Here, this graphical representation is used to form the basis of an interactive tool. It has previously been demonstrated that such interactive tools with real-time updated results help students to more thoroughly build a conceptual understanding of a complex process, especially when the student is allowed to independently explore the possibilities provided by the tool (Wieman et al., 2008; National Research Council, 2015). In the case of “*Orbital, the Box*”, the student can experiment by using a control panel (Figure 3) to independently change orbital parameters and see how such changes influence the following: (1) the variation in Earth's orbital speed and distance from the Sun throughout the year; (2) the irradiance distribution across the Earth's latitudes and throughout the year; (3) the day length of the astronomical seasons; (4) the mean annual irradiance received at each latitude; (5) the shape and positioning of the Earth's orbit relative to the Sun. This setup constitutes a classic experimental environment with multiple variables (in this case three), whereby one variable is changed while the other two are kept constant.




Figure 3. The slider control panels used to control the graphical representations.

240

3.3 Digital textbook

Included in the interactive GUI is access to a very basic digital textbook explaining the concepts of obliquity, eccentricity and geocentric longitude of perihelion (which is governed by precession).

245 The hyperlinks above the slider controls in Figure 3 will take the students to pages with basic information about the orbital parameters and a tutorial containing example calculations and figures demonstrating basic principles. In the obliquity section, students learn how to calculate the declination latitude of the Sun, and the midday angle of the Sun for the solstices at their city, given a specific obliquity. In the eccentricity section, there is a tutorial on how to calculate aphelion and perihelion distances given a specific eccentricity, as well as information on how to calculate the speed of the Earth's orbit at aphelion and perihelion. Geocentric longitude of perihelion is explained and its cause (general precession) is explained using animations.

250

4.0 Conclusion and reuse potential

255 The GUI tool has great potential to be integrated into geosciences curricula, particularly within a classic experimental environment allowing for the isolated study of the individual effects of orbital parameters upon the seasonal and latitudinal distribution of irradiance. The tool provides an experimental environment that can be deployed within a flipped classroom setting with jigsaw activities, as has previously been successfully deployed in geosciences the case of, e.g. plate tectonics (Sawyer et al., 2005). An example lesson plan is to split students into three groups, with each group focussing on learning the effect of one (or all) of the three orbital parameters (obliquity, eccentricity and precession). Each group could investigate the effect (if any) of their parameter(s) upon orbital distance, speed, seasonal irradiance distribution, length of season, and mean annual irradiance received at each latitude. Afterwards, groups can rotate members and/or hold discussions to teach each the other groups about the particular orbital parameter that they studied.

265

It should be noted that the GUI tool developed here exclusively includes realistic values for the orbital parameters of planet Earth. A useful additional exercise/challenge is assigning groups unrealistic parameters for Earth (such as an obliquity value of 45°) and asking them to then sketch the irradiance profile that they would expect to be associated with such a parameter, as well as what latitude the polar circle and tropics would be. Such a task can promote metacognitive thinking, as it challenges the students to extrapolate an extreme hypothetical scenario, which can in turn help them better understand the functioning of more realistic scenarios.

Planet Earth's irradiance distribution and its relation to orbital parameters is a very challenging subject to teach. Geosciences has traditionally relied upon fieldwork and/or laboratory settings to stimulate active learning, but carrying out real-world experiments upon the Earth's orbit is beyond current technology and probably undesirable. A computer-based experimental environment can allow for similar interactive attainment of understanding as would be attained in a real-world experimental environment.

Software availability

The *Orbital, the Box* interactive teaching tool can be downloaded from Zenodo:

<https://doi.org/10.5281/zenodo.4597618>. Matlab scripts that can be used to recreate the irradiance

and orbital figures shown in the tool can be downloaded from Github:

https://github.com/bryanlougheed/orbital_the_box

Acknowledgements

The author acknowledges Swedish Research Council (*Vetenskapsrådet – VR*) Starting Grant number 2018-04992. Yunus Baykal, Jorijntje Henderiks, Boris-Theofanis Karatsolis, Patrik Thunström, Niklas Wahlström (all Uppsala University) and Brett Metcalfe (Wageningen UR & Vrije Universiteit Amsterdam) are thanked for discussions.

References

Adhémar, J. A.: Révolutions de la mer, déluges périodiques, Carilian-Goeury et V. Dalmont, Paris., 1842.

Berger, A., Loutre, M.-F. and Yin, Q.: Total irradiation during any time interval of the year using elliptic integrals, *Quaternary Science Reviews*, 29(17), 1968–1982, doi:10.1016/j.quascirev.2010.05.007, 2010.

Berger, A. L.: Long-Term Variations of Daily Insolation and Quaternary Climatic Changes, *J. Atmos. Sci.*, 35(12), 2362–2367, doi:10.1175/1520-0469(1978)035<2362:LTVODI>2.0.CO;2, 1978.

Bol'shakov, V. A.: On the ill-defined notion of the Milankovitch Theory and its influence on the development of the orbital theory of the paleoclimate, *Her. Russ. Acad. Sci.*, 87(4), 356–369, doi:10.1134/S1019331617040025, 2017.

Bykerk-Kauffman, A.: Using Cooperative Learning in College Geology Classes, *Journal of Geological Education*, 43(4), 309–316, doi:10.5408/0022-1368-43.4.309, 1995.

Croll, J.: XIII. On the physical cause of the change of climate during geological epochs, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 28(187), 121–137, 1864.

Herschel, J. F. W.: XVII.—On the Astronomical Causes which may influence Geological Phænomena., *Transactions of the Geological Society of London*, S2-3(2), 293–300, doi:10.1144/transgslb.3.2.293, 1832.

Huguet, C., Pearse, J., Noè, L. F., Valencia, D. M., Ruiz, N. C., Heredia, A. J. and Avedaño, M. A. P.: Improving the motivation of students in a large introductory geoscience course through active learning, *Journal of Geoscience Education*, 68(1), 20–32, doi:10.1080/10899995.2019.1588489, 2020.

Huybers, P.: Early Pleistocene Glacial Cycles and the Integrated Summer Insolation Forcing, *Science*, 313(5786), 508–511, doi:10.1126/science.1125249, 2006.

Huybers, P.: Combined obliquity and precession pacing of late Pleistocene deglaciations, *Nature*, 480(7376), 229–232, doi:10.1038/nature10626, 2011.

Kostadinov, T. S. and Gilb, R.: Earth Orbit v2.1: a 3-D visualization and analysis model of Earth's orbit, Milankovitch cycles and insolation, *Geosci. Model Dev.*, 7(3), 1051–1068, doi:10.5194/gmd-7-1051-2014, 2014.

Manduca, C. A., Baer, E., Hancock, G., Macdonald, R. H., Patterson, S., Savina, M. and Wenner, J.: Making Undergraduate Geoscience Quantitative, *Eos Trans. AGU*, 89(16), 149, doi:10.1029/2008EO160001, 2008.

Meeus, J.: *Astronomical Algorithms*, 2nd ed., Willmann-Bell, Inc., Richmond, Virginia., 1998.

Milanković, M.: *Kanon der Erdebestrahlung und seine Anwendung auf das Eiszeitenproblem*, Königlich Serbische Akademie., 1941.

National Research Council: *Reaching Students: What Research Says About Effective Instruction in Undergraduate Science and Engineering*, National Academies Press, Washington, D.C., 2015.

Sawyer, D. S., Henning, A. T., Shipp, S. and Dunbar, R. W.: A Data Rich Exercise for Discovering Plate Boundary Processes, *Journal of Geoscience Education*, 53(1), 65–74, doi:10.5408/1089-9995-53.1.65, 2005.

Sinnott, R. W.: A computer assault on Kepler's equation, *Sky & Telescope*, 70, 158–159, 1985.

Van Hemelrijck, E.: The oblateness effect on the extraterrestrial solar radiation, *Solar Energy*, 31(2), 223–228, doi:10.1016/0038-092X(83)90085-3, 1983.

Wampler, J. M.: Misconceptions – A Column about Errors in Geoscience Textbooks: Confusion about the Changing Eccentricity of the Earth's Orbit, *Journal of Geoscience Education*, 48(4), 541–543, doi:10.5408/1089-9995-48.4.541, 2000a.

Wampler, J. M.: Misconceptions - A Column about Errors in Geoscience Textbooks: Misconceptions of Changes in Obliquity of the Ecliptic, *Journal of Geoscience Education*, 48(5), 701–704, doi:10.5408/1089-9995-48.5.701, 2000b.

Wieman, C. E., Adams, W. K. and Perkins, K. K.: PhET: Simulations That Enhance Learning, *Science*, 322(5902), 682–683, doi:10.1126/science.1161948, 2008.

w3schools.com. How To - Range Sliders. Available at

https://www.w3schools.com/howto/howto_js_rangeslider.asp [last accessed 28 December 2020].