

JOURNAL OF GLACIOLOGY



CAMBRIDGE
UNIVERSITY PRESS

THIS MANUSCRIPT HAS BEEN SUBMITTED TO THE JOURNAL OF GLACIOLOGY AND HAS NOT BEEN PEER-REVIEWED.

Mass loss of the Greenland ice sheet until the year 3000 under a sustained late-21st-century climate

Journal:	<i>Journal of Glaciology</i>
Manuscript ID	JOG-21-0089
Manuscript Type:	Letter
Date Submitted by the Author:	05-Jul-2021
Complete List of Authors:	Greve, Ralf; Hokkaido University, Institute of Low Temperature Science Chambers, Christopher; Hokkaido University Institute of Low Temperature Science, Glaciology
Keywords:	Ice-sheet modelling, Arctic glaciology, Climate change, Ice and climate
Abstract:	We conduct extended versions of the ISMIP6 future climate experiments for the Greenland ice sheet until the year 3000 with the model SICOPOLIS. Beyond 2100, the climate forcing is kept fixed at late-21st-century conditions. For the unabated warming pathway RCP8.5/SSP5-8.5, the ice sheet suffers a severe mass loss, which amounts to ~ 1.8 m SLE (sea-level equivalent) for the twelve-experiment mean, and ~ 3.5 m SLE ($\sim 50\%$ of the entire mass) for the most sensitive experiment. For the reduced emissions pathway RCP2.6/SSP1-2.6, the mass loss is limited to a two-experiment mean of ~ 0.28 m SLE. Climate-change mitigation during the next decades will therefore be an efficient means for limiting the contribution of the Greenland ice sheet to sea-level rise in the long term.



SCHOLARONE™
Manuscripts

Mass loss of the Greenland ice sheet until the year 3000 under a sustained late-21st-century climate

Ralf GREVE,^{1,2} Christopher CHAMBERS¹

¹*Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan*

²*Arctic Research Center, Hokkaido University, Sapporo, Japan*

Correspondence: Ralf Greve <greve@lowtem.hokudai.ac.jp>

ABSTRACT. We conduct extended versions of the ISMIP6 future climate experiments for the Greenland ice sheet until the year 3000 with the model SICOPOLIS. Beyond 2100, the climate forcing is kept fixed at late-21st-century conditions. For the unabated warming pathway RCP8.5/SSP5-8.5, the ice sheet suffers a severe mass loss, which amounts to ~ 1.8 m SLE (sea-level equivalent) for the twelve-experiment mean, and ~ 3.5 m SLE ($\sim 50\%$ of the entire mass) for the most sensitive experiment. For the reduced emissions pathway RCP2.6/SSP1-2.6, the mass loss is limited to a two-experiment mean of ~ 0.28 m SLE. Climate-change mitigation during the next decades will therefore be an efficient means for limiting the contribution of the Greenland ice sheet to sea-level rise in the long term.

1 INTRODUCTION

Global warming, also referred to as climate change, is the observed rise in the average temperature of the Earth's climate system and its related effects. It is established scientific consensus that the climate system is warming, and that human influence has been the dominant cause of the observed warming since the mid-20th century (e.g., IPCC, 2013). A major consequence of global warming is sea-level rise, currently occurring at a global mean rate of 3.6 ± 0.5 millimetres per year (for the period 2006–2015; Oppenheimer and others, 2019). The main sources are melting/discharge of ice sheets, ice caps and glaciers, and thermal expansion of ocean water. In the long term, the two ice sheets of Antarctica (AIS) and Greenland (GIS)

26 are the largest potential contributors to global sea-level rise because of their enormous volumes, together
27 amounting to ~ 65 mSLE (sea-level equivalent) (Morlighem and others, 2017, 2020). The ice sheets have
28 therefore been the focus of intensive observational as well as modelling efforts.

29 The Coupled Model Intercomparison Project Phase 6 (CMIP6) is a major international climate mod-
30 elling initiative (Eyring and others, 2016). As a part of this project, the Ice Sheet Model Intercomparison
31 Project for CMIP6 (ISMIP6) brought together a consortium of ice-sheet modellers to explore the sea-level-
32 rise contribution from the GIS and AIS (Nowicki and others, 2016, 2020). ISMIP6 focussed on the CMIP6
33 period from 2015 until the end of 2100. The main findings for the GIS, when forced by output from CMIP5
34 global climate models (GCMs), were contributions of 90 ± 50 and 32 ± 17 mm SLE for the unabated warm-
35 ing pathway RCP8.5 [RCP: Representative Concentration Pathway] and the reduced emissions pathway
36 RCP2.6, respectively (Goelzer and others, 2020). CMIP6 GCMs generally feature a warmer atmosphere,
37 which results in higher mass loss due to increased surface melt (Payne and others, 2021). For the AIS and
38 CMIP5 climate forcings, ISMIP6 found a mass loss in the range of -7.8 to 30.0 cm SLE under RCP8.5
39 (Seroussi and others, 2020). The limited number of results for RCP2.6 fall within this range, and so do
40 the results obtained with CMIP6 climate forcings (Payne and others, 2021). This rather unclear picture
41 for the AIS is a consequence of the counteracting effects of mass loss due to ocean warming and mass gain
42 from increased snowfall.

43 The full suite of ISMIP6 experiments with both CMIP5 and CMIP6 forcings was carried out with
44 the ice-sheet model SICOPOLIS (“SIMulation COde for POLythermal Ice Sheets”, www.sicopolis.net), as
45 documented in detail by Greve and others (2020a,b). Chambers and others (2021) extend the ISMIP6 sim-
46 ulations for the AIS with SICOPOLIS until the year 3000, assuming a sustained late-21st-century climate
47 beyond 2100. Compared to the uncertain response projected over the ISMIP6 period, a radically different
48 picture emerges, demonstrating that the consequences of the high-emissions scenario RCP8.5/SSP5-8.5
49 [SSP: Shared Socioeconomic Pathway] are much greater in the long term even if no further climate trend
50 is applied beyond 2100.

51 The response of the GIS to longer-term climate change has also been investigated. Vizcaino and others
52 (2015) carried out simulations until 2300 with a coupled ECHAM5.2/MPI-OM/SICOPOLIS model for the
53 pathways RCP2.6, RCP4.5, and a modified RCP8.5 with a $4 \times \text{CO}_2$ limit. More recently, Aschwanden
54 and others (2019) used projections from four CMIP5 GCMs until 2300 for RCP2.6, RCP4.5 and RCP8.5,
55 extrapolated until 3000, to force the ice-sheet model PISM. In this study, we transfer the approach by

56 Chambers and others (2021) to the GIS. The objective is to assess its long-term response to late-21st-
57 century climatic conditions for the full ensemble of ISMIP6 climate forcings, which consists of fourteen
58 scenarios from ten different CMIP5 and CMIP6 GCMs.

59 **2 METHODS**

60 The main tool used for this study is the ice-sheet model SICOPOLIS. We apply it to the GIS with
61 hybrid shallow-ice–shelfy-stream dynamics (Bernales and others, 2017), a Weertman-Budd-type sliding
62 law tuned separately for 20 different regions (Greve and others, 2020b), and ice thermodynamics treated
63 by the one-layer melting-CTS enthalpy scheme (CTS: cold-temperate transition surface; Blatter and Greve,
64 2015; Greve and Blatter, 2016). The horizontal resolution is 5 km. In the vertical, we use terrain-following
65 coordinates (sigma transformation) with 81 layers in the ice domain and 41 layers in the thermal lithosphere
66 layer below. The detailed set-up is described by Greve and others (2020b) and shall not be repeated here.

67 Following the ISMIP6 protocol, climate forcing from 2015 until the end of 2100 has an atmospheric
68 and an oceanic component. The atmospheric forcing consists of the 1960–1989 reference climatology, plus
69 space-time-dependent anomalies for the surface mass balance (SMB), the surface temperature (ST) and
70 their vertical gradients. The oceanic forcing translates ocean thermal forcing (ambient water temperature
71 relative to the freezing point) at the margin of the GIS to prescribed maps for average retreat rates,
72 specified for seven ice–ocean sectors around Greenland. For further details, see Goelzer and others (2020),
73 Nowicki and others (2020), Payne and others (2021), and references therein.

74 For the period from 2101 until the end of 3000, we extend the simulations in a similar way than
75 Chambers and others (2021) do for the AIS. For every year of this extended period, the atmospheric
76 forcing (SMB, ST, vertical gradients) for the 10-year interval 2091–2100 is randomly sampled such that
77 no further trend is applied, but some inter-annual fluctuations remain (similar to Calov and others, 2018).
78 The oceanic forcing (prescribed retreat maps) does not show any notable year-to-year fluctuations, so we
79 simply keep it fixed at 2100 conditions.

80 An overview of our extended ISMIP6 experiments is given in Table 1. Twelve experiments are for
81 the 21st-century unabated warming pathway RCP8.5 (CMIP5) / SSP5-8.5 (CMIP6), and two are for the
82 reduced emissions pathway RCP2.6 (CMIP5) / SSP1-2.6 (CMIP6) that is largely in line with the commit-
83 ments of the Paris Agreement (maintaining the global mean temperature well below a 2°C increase above
84 pre-industrial levels). In two experiments, the impact of different sensitivities of the retreat parameteriza-

85 tion due to oceanic forcing (“high” and “low” vs. the normal, “medium” sensitivity) is tested. In addition,
86 a control simulation (“ctrl_proj”) employs constant climate conditions based on a 1960–1989 climatology
87 and no explicit oceanic forcing.

88 3 RESULTS

89 The simulated mass change of the GIS, expressed as a sea-level contribution, and ice area are shown in
90 Figure 1. The historical run (“hist”) bridges the gap between our initialization year 1990 and the start
91 date of the projections in January 2015 by employing MIROC5/RCP8.5 SMB and ST forcing (Greve and
92 others, 2020b). Like the regular projections, the projection control run (“ctrl_proj”) starts from January
93 2015 and runs until the end of 3000. In ctrl_proj, the ice sheet remains nearly stable, showing a slight
94 mass gain of 6.4 mm SLE and area loss of $4.7 \times 10^3 \text{ km}^2$ during the 986 years model time, which is of the
95 order of permilles of the present-day values.

96 For all future projections, the ice sheet keeps losing both mass and extent over the entire period. The
97 largest rate of change occurs typically around the year 2100, beyond which it slows down to some extent;
98 however, without reaching or coming close to a new steady state. This demonstrates that the committed
99 mass loss due to 21st-century climate change extends way beyond the 21st century and impacts the ice
100 sheet on a much longer time scale. Corroborating the findings for the 21st century (Goelzer and others,
101 2020; Greve and others, 2020b), the GIS responds much more strongly to the ensemble of RCP8.5/SSP5-8.5
102 simulations than to the two RCP2.6/SSP1-2.6 simulations. By the year 3000, the mass loss amounts to
103 $1.78 \pm 0.80 \text{ m SLE}$ (mean \pm 1-sigma range) for RCP8.5/SSP5-8.5, while it is limited to $0.28 \pm 0.12 \text{ m SLE}$
104 for RCP2.6/SSP1-2.6. In relative terms, the area loss is similar to that of the mass loss.

105 The influence of the ice retreat due to oceanic forcing is explored by Exps. 5, 9, 10 (MIROC5/RCP8.5
106 with “medium”, “high” and “low” sensitivity, respectively). The results are shown by the green lines
107 and green-shaded regions in Figure 1. By 3000, the simulated mass loss is $1.63_{-0.031}^{+0.039} \text{ m SLE}$. Thus, the
108 uncertainty due to these three calibrations is very small in the long range. Relative to the uncertainty due
109 to the different climate forcings, it is more pronounced for the 21st century (Greve and others, 2020b).
110 This is because the continued retreat of the ice sheet decreases its contact with the ocean, so that the
111 oceanic forcing plays a smaller role in the longer term.

112 As reported by Greve and others (2020b) and Payne and others (2021), for both the 21st-century
113 RCP8.5/SSP5-8.5 and RCP2.6/SSP1-2.6 pathways, the CMIP6 climate models produce a larger response

114 of the ice sheet than the CMIP5 ones. While the significance of this statement is limited in the case of
115 RCP2.6/SSP1-2.6 (only one experiment each), it is more robust for RCP8.5/SSP5-8.5, where the ensemble
116 contains eight and four experiments forced by CMIP5 and CMIP6 models, respectively. By 3000, the mean
117 mass loss for the four CMIP6 SSP5-8.5 experiments is 2.72 m SLE, and the maximum value from Exp. B4
118 (CESM2/SSP5-8.5) is as large as 3.54 m SLE, almost 50% of the entire present-day ice mass.

119 We now discuss in more detail the results of Exp. 5 (MIROC5/RCP8.5), a typical representative for
120 which the mass loss is close to the ensemble average. Figure 2 shows the components of the global mass
121 balance: surface mass balance (SMB), basal mass balance (BMB), calving and ice volume change (dV/dt)
122 [all counted as positive for mass gain]. On a mean-annual basis, the residual, $\text{Res} = |\text{SMB} + \text{BMB} +$
123 $\text{Calving} - dV/dt|$, is always less than $10^6 \text{ m}^3 \text{ a}^{-1}$. This is five to six orders of magnitude smaller than the
124 typical range of values in the figure, so that the model conserves mass very well. As already discussed
125 above, the ice sheet keeps losing volume (\propto mass) over the entire period, with maximal rates of change
126 occurring around the year 2100. The SMB is initially positive, but changes its sign in the second half of
127 the 21st century and stays negative beyond that. Calving into the surrounding ocean peaks during 2080–
128 2085, when it contributes approximately the same amount to ice volume loss than negative SMB. After
129 that, calving decreases continuously due to ice-sheet retreat from the coast and becomes almost negligible
130 towards the end of the 3rd millennium. BMB is small over the entire model time. The noise of the volume
131 change is due to the noise of the SMB, which reflects the inter-annual variability of the atmospheric forcing.

132 Figure 3 shows snapshots of the ice thickness and surface velocity for Exp. 5 for the initial year 2015
133 and the final year 3000. Comparing the thickness distributions demonstrates nicely that the ice sheet
134 retreats from the coast almost all around its perimeter, and contact to the ocean is very limited by the
135 end of the simulation, which entails the low calving rates mentioned above. By contrast, the ice sheet
136 does not suffer much change in its interior parts north of $\sim 68^\circ\text{N}$. The large-scale pattern of the ice flow
137 and the organization of the ice sheet into major drainage basins remain largely intact. However, on the
138 regional scale, more pronounced changes occur. The fast-flowing outlet glaciers in south-western Greenland
139 disappear entirely due to the extreme retreat in this area. The north-western outlet glaciers, including
140 Petermann Glacier, also slow down substantially. The central-western Jakobshavn Ice Stream loses its
141 clear delimitation to the surrounding glaciers, but remains an area of fast-flowing ice. The major features
142 in East Greenland, e.g., the North-East Greenland Ice Stream, Kangerdlugssuaq and Helheim glaciers, are
143 less affected and remain well identifiable.

144 4 DISCUSSION AND CONCLUSION

145 The future climate simulations carried out in this study for the GIS over the 3rd millennium confirm and
146 continue the trends that were reported by ISMIP6-Greenland for the 21st century (Goelzer and others,
147 2020; Greve and others, 2020b; Payne and others, 2021). The response of the ice sheet is mainly governed
148 by a negative SMB due to increased surface melting near the ice margin. Marine-terminating glacier
149 retreat, triggered by increasing oceanic thermal forcing, constitutes a further negative contribution to the
150 total mass balance, but becomes less important in the longer term. Under the unabated warming pathway
151 RCP8.5/SSP5-8.5, this leads to a severe mass loss during the 3rd millennium, while the loss is much smaller
152 under the reduced emissions pathway RCP2.6/SSP1-2.6. Results obtained with forcings from the newer
153 CMIP6 climate models consistently produce larger mass losses than those obtained with the older CMIP5
154 models, for SSP5-8.5 in the range of a $\sim 25\text{--}50\%$ loss of the present-day ice mass (and area) by 3000. For
155 comparison, Aschwanden and others (2019) reported a mass loss of $\sim 75\text{--}100\%$ by 3000 for their ensemble
156 of RCP8.5 simulations, for which a warming trend is assumed to continue until 2500. Efficient climate
157 change mitigation during the next decades is therefore crucial for limiting the contribution of the GIS to
158 long-term sea-level rise.

159 Our study is limited to investigating the impact of a sustained late-21st-century climate (without
160 imposing a further trend beyond 2100) on the GIS. However, in reality, climate change will continue beyond
161 2100 (e.g., Bakker and others, 2016; Lyon and others, 2020). Further, the unidirectional coupling approach
162 (climate model \rightarrow ice-sheet model) employed by ISMIP6, and thus here, lacks a detailed accounting of
163 feedbacks of the changing ice sheet on the climate. As we explained in Sect. 2, the climate forcing for
164 Greenland includes vertical gradients of the surface mass balance and surface temperature. Therefore, the
165 changing ice-sheet geometry acts back on these climatic forcing fields. However, the linearized approach was
166 derived for small perturbations of the present-day state only, and it cannot be validated for large changes of
167 the ice sheet. This shortcoming becomes more severe in our simulations over almost a millennium compared
168 to the 86-year scope of ISMIP6, adding to the uncertainty of the results.

169 Therefore, future work in the direction of long-term simulations of ice-sheet response to climate change
170 should aim at employing GCM projections beyond 2100 and improving the representation of feedback
171 processes. The ultimate solution would be to carry out such simulations in a fully coupled way, with the
172 ice-sheet model integrated in the global climate model. This approach has been pursued (e.g., Vizcaino and

173 others, 2015; Gregory and others, 2020); however, fully coupled simulations are demanding and computationally expensive, which makes it difficult to run large ensembles, involving many different climate and ice-sheet models, over long time scales and at adequate resolution. Intermediary, more manageable solutions may consist of involving snapshots of climate-model results combined with more refined parameterizations for the climatic forcing, similar to the approach by Abe-Ouchi and others (2013) for the paleo-glaciation of the Northern Hemisphere.

179 ACKNOWLEDGEMENTS

180 We thank Jorge Bernales (MARUM Bremen), Reinhard Calov (PIK Potsdam), Takashi Obase (University of Tokyo) and Fuyuki Saito (JAMSTEC Yokohama) for their recent contributions to the development of the SICOPOLIS model, and Ayako Abe-Ouchi for fruitful discussions about ice-sheet and climate modelling. We thank the Climate and Cryosphere (CliC) effort, which provided support for ISMIP6 through sponsoring of workshops, hosting the ISMIP6 website and wiki, and promoting ISMIP6. We acknowledge the World Climate Research Programme, which, through its Working Group on Coupled Modelling, coordinated and promoted CMIP5 and CMIP6. We thank the climate modelling groups for producing their model output and making it available; the Earth System Grid Federation (ESGF) for archiving the CMIP data and providing access to it; the University at Buffalo for ISMIP6 data distribution and upload; and the multiple funding agencies who support CMIP5, CMIP6, and ESGF. We thank the ISMIP6 steering committee, the ISMIP6 model selection group and ISMIP6 dataset preparation group for their continuous engagement in defining ISMIP6. This is ISMIP6 contribution No. xxx.

192 Ralf Greve and Christopher Chambers were supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grant No. JP17H06323, and by a Leadership Research Grant (Category 2) of Hokkaido University's Institute of Low Temperature Science. Ralf Greve was supported by JSPS KAKENHI Grant No. JP17H06104, and by the Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) through the Arctic Challenge for Sustainability project ArCS II (program grant number JP-MXD1420318865).

198 REFERENCES

199 Abe-Ouchi A, Saito F, Kawamura K, Raymo ME, Okuno J, Takahashi K and Blatter H (2013) Insolation-driven 200 100,000-year glacial cycles and hysteresis of ice-sheet volume. *Nature*, **500**(7461), 190–193 (doi: 10.1038/na-

- 201 ture12374)
- 202 Aschwanden A, Fahnestock MA, Truffer M, Brinkerhoff DJ, Hock R, Khroulev C, Mottram R and Khan SA (2019)
- 203 Contribution of the Greenland Ice Sheet to sea level over the next millennium. *Science Advances*, **5**(6), eaav9396
- 204 (doi: 10.1126/sciadv.aav9396)
- 205 Bakker P, Schmittner A, Lenaerts JTM, Abe-Ouchi A, Bi D, van den Broeke MR, Chan WL, Hu A, Beadling RL,
- 206 Marsland SJ, Mernild SH, Saenko OA, Swingedouw D, Sullivan A and Yin J (2016) Fate of the Atlantic Meridional
- 207 Overturning Circulation: Strong decline under continued warming and Greenland melting. *Geophysical Research*
- 208 *Letters*, **43**, 12252–12260 (doi: 10.1002/2016GL070457)
- 209 Bernales J, Rogozhina I, Greve R and Thomas M (2017) Comparison of hybrid schemes for the combination of
- 210 shallow approximations in numerical simulations of the Antarctic Ice Sheet. *The Cryosphere*, **11**(1), 247–265 (doi:
- 211 10.5194/tc-11-247-2017)
- 212 Blatter H and Greve R (2015) Comparison and verification of enthalpy schemes for polythermal glaciers and ice
- 213 sheets with a one-dimensional model. *Polar Science*, **9**(2), 196–207 (doi: 10.1016/j.polar.2015.04.001)
- 214 Calov R, Beyer S, Greve R, Beckmann J, Willeit M, Kleiner T, Rückamp M, Humbert A and Ganopolski A (2018)
- 215 Simulation of the future sea level contribution of Greenland with a new glacial system model. *The Cryosphere*,
- 216 **12**(10), 3097–3121 (doi: 10.5194/tc-12-3097-2018)
- 217 Chambers C, Greve R, Obase T, Saito F and Abe-Ouchi A (2021) Extended ISMIP6 projections for the Antarc-
- 218 tic ice sheet with the model SICOPOLIS. *Journal of Glaciology*, submitted (preprint at EarthArXiv, doi:
- 219 10.31223/X5CP7C)
- 220 Eyring V, Bony S, Meehl GA, Senior CA, Stevens B, Stouffer RJ and Taylor KE (2016) Overview of the Coupled Model
- 221 Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*,
- 222 **9**(5), 1937–1958 (doi: 10.5194/gmd-9-1937-2016)
- 223 Goelzer H, Nowicki S, Payne A, Larour E, Seroussi H, Lipscomb WH, Gregory J, Abe-Ouchi A, Shepherd A, Simon E,
- 224 Agosta C, Alexander P, Aschwanden A, Barthel A, Calov R, Chambers C, Choi Y, Cuzzzone J, Dumas C, Edwards
- 225 T, Felikson D, Fettweis X, Golledge NR, Greve R, Humbert A, Huybrechts P, Le clec'h S, Lee V, Leguy G, Little
- 226 C, Lowry DP, Morlighem M, Nias I, Quiquet A, Rückamp M, Schlegel NJ, Slater D, Smith R, Straneo F, Tarasov
- 227 L, van de Wal R and van den Broeke M (2020) The future sea-level contribution of the Greenland ice sheet: a
- 228 multi-model ensemble study of ISMIP6. *The Cryosphere*, **14**(9), 3071–3096 (doi: 10.5194/tc-14-3071-2020)
- 229 Gregory JM, George SE and Smith RS (2020) Large and irreversible future decline of the Greenland ice sheet. *The*
- 230 *Cryosphere*, **14**(12), 4299–4322 (doi: 10.5194/tc-14-4299-2020)

- 231 Greve R and Blatter H (2016) Comparison of thermodynamics solvers in the polythermal ice sheet model SICOPOLIS.
232 *Polar Science*, **10**(1), 11–23 (doi: 10.1016/j.polar.2015.12.004)
- 233 Greve R, Calov R, Obase T, Saito F, Tsutaki S and Abe-Ouchi A (2020a) ISMIP6 future projections for the Antarctic
234 ice sheet with the model SICOPOLIS. Technical report, Zenodo (doi: 10.5281/zenodo.3971232)
- 235 Greve R, Chambers C and Calov R (2020b) ISMIP6 future projections for the Greenland ice sheet with the model
236 SICOPOLIS. Technical report, Zenodo (doi: 10.5281/zenodo.3971251)
- 237 IPCC (2013) Summary for policymakers. In TF Stocker, D Qin, GK Plattner, M Tignor, SK Allen, J Boschung,
238 A Nauels, Y Xia, V Bex and PM Midgley (eds.), *Climate Change 2013: The Physical Science Basis. Contribution*
239 *of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, 3–29,
240 Cambridge University Press, Cambridge, UK and New York, NY, USA
- 241 Lyon C, Saupe E, Smith C, Hill D, Beckerman A, Stringer L, Marchant R, McKay J, Burke A, O’Higgins P, Dunhill A,
242 Allen B, Riel-Salvatore J and Aze T (2020) Climate change research and action must look beyond 2100. *EarthArXiv*
243 (doi: 10.31223/x5qg7d), preprint
- 244 Morlighem M, Williams CN, Rignot E, An L, Arndt JE, Bamber JL, Catania G, Chauché N, Dowdeswell JA, Dorschel
245 B, Fenty I, Hogan K, Howat I, Hubbard A, Jakobsson M, Jordan TM, Kjeldsen KK, Millan R, Mayer L, Mouginot
246 J, Noël BPY, O’Cofaigh C, Palmer S, Rysgaard S, Seroussi H, Siegert MJ, Slabon P, Straneo F, van den Broeke
247 MR, Weinrebe W, Wood M and Zinglensen KB (2017) BedMachine v3: Complete bed topography and ocean
248 bathymetry mapping of Greenland from multibeam echo sounding combined with mass conservation. *Geophysical*
249 *Research Letters*, **44**(21), 11051–11061 (doi: 10.1002/2017GL074954)
- 250 Morlighem M, Rignot E, Binder T, Blankenship D, Drews G, Rand Eagles, Eisen O, Ferraccioli F, Forsberg R, Fretwell
251 P, Goel V, Greenbaum JS, Gudmundsson H, Guo J, Helm V, Hofstede C, Howat I, Humbert A, Jokat W, Karlsson
252 NB, Lee WS, Matsuoka K, Millan R, Mouginot J, Paden J, Pattyn F, Roberts J, Rosier S, Ruppel A, Seroussi H,
253 Smith EC, Steinhage D, Sun B, van den Broeke MR, van Ommen TD, van Wessem M and Young DA (2020) Deep
254 glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet. *Nature Geoscience*,
255 **13**(2), 132–137 (doi: 10.1038/s41561-019-0510-8)
- 256 Nowicki S, Goelzer H, Seroussi H, Payne AJ, Lipscomb WH, Abe-Ouchi A, Agosta C, Alexander P, Asay-Davis
257 XS, Barthel A, Bracegirdle TJ, Cullather R, Felikson D, Fettweis X, Gregory JM, Hattermann T, Jourdain NC,
258 Kuipers Munneke P, Larour E, Little CM, Morlighem M, Nias I, Shepherd A, Simon E, Slater D, Smith RS, Straneo
259 F, Trusel LD, van den Broeke MR and van de Wal R (2020) Experimental protocol for sea level projections from
260 ISMIP6 stand-alone ice sheet models. *The Cryosphere*, **14**(7), 2331–2368 (doi: 10.5194/tc-14-2331-2020)

- 261 Nowicki SMJ, Payne A, Larour E, Seroussi H, Goelzer H, Lipscomb W, Gregory J, Abe-Ouchi A and Shepherd A
262 (2016) Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6. *Geoscientific Model Develop-*
263 *ment*, **9**(12), 4521–4545 (doi: 10.5194/gmd-9-4521-2016)
- 264 Oppenheimer M, Glavovic BC, Hinkel J, van de Wal R, Magnan AK, Abd-Elgawad A, Cai R, Cifuentes-Jara M,
265 DeConto RM, Ghosh T, Hay J, Isla F, Marzeion B, Meysignac B and Sebesvari Z (2019) Sea level rise and
266 implications for low-lying islands, coasts and communities. In HO Pörtner, DC Roberts, V Masson-Delmotte,
267 P Zhai, M Tignor, E Poloczanska, K Mintenbeck, A Alegría, M Nicolai, A Okem, J Petzold, B Rama and NM Weyer
268 (eds.), *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, 321–445, Intergovernmental
269 Panel on Climate Change, Geneva, Switzerland
- 270 Payne AJ, Nowicki S, Abe-Ouchi A, Agosta C, Alexander P, Albrecht T, Asay-Davis X, Aschwanden A, Barthel
271 A, Bracegirdle TJ, Calov R, Chambers C, Choi Y, Cullather R, Cuzzone J, Dumas C, Edwards TL, Felikson
272 D, Fettweis X, Galton-Fenzi BK, Goelzer H, Gladstone R, Golledge NR, Gregory JM, Greve R, Hattermann T,
273 Hoffman MJ, Humbert A, Huybrechts P, Jourdain NC, Kleiner T, Munneke PK, Larour E, Le clec'h S, Lee V,
274 Leguy G, Lipscomb WH, Little CM, Lowry DP, Morlighem M, Nias I, Pattyn F, Pelle T, Price SF, Quiquet A,
275 Reese R, Rückamp M, Schlegel NJ, Seroussi H, Shepherd A, Simon E, Slater D, Smith RS, Straneo F, Sun S,
276 Tarasov L, Trusel LD, Van Breedam J, van de Wal R, van den Broeke M, Winkelmann R, Zhao C, Zhang T and
277 Zwinger T (2021) Future sea level change under CMIP5 and CMIP6 scenarios from the Greenland and Antarctic
278 ice sheets. *Geophysical Research Letters* (doi: 10.1029/2020GL091741), in press
- 279 Seroussi H, Nowicki S, Payne AJ, Goelzer H, Lipscomb WH, Abe-Ouchi A, Agosta C, Albrecht T, Asay-Davis X,
280 Barthel A, Calov R, Cullather R, Dumas C, Galton-Fenzi BK, Gladstone R, Golledge N, Gregory JM, Greve R,
281 Hatterman T, Hoffman MJ, Humbert A, Huybrechts P, Jourdain NC, Kleiner T, Larour E, Leguy GR, Lowry DP,
282 Little CM, Morlighem M, Pattyn F, Pelle T, Price SF, Quiquet A, Reese R, Schlegel NJ, Shepherd A, Simon E,
283 Smith RS, Straneo F, Sun S, Trusel LD, Van Breedam J, van de Wal RSW, Winkelmann R, Zhao C, Zhang T and
284 Zwinger T (2020) ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st
285 century. *The Cryosphere*, **14**(9), 3033–3070 (doi: 10.5194/tc-14-3033-2020)
- 286 Vizcaino M, Mikolajewicz U, Ziemen F, Rodehacke CB, Greve R and van den Broeke MR (2015) Coupled simulations
287 of Greenland Ice Sheet and climate change up to A.D. 2300. *Geophysical Research Letters*, **42**(10), 3927–3935 (doi:
288 10.1002/2014GL061142)

#	exp_id	Scenario	GCM	Ocean forcing	
5	exp05	RCP8.5	MIROC5	Medium	Core experiments (Tier 1)
6	exp06	RCP8.5	NorESM1-M	Medium	
7	exp07	RCP2.6	MIROC5	Medium	
8	exp08	RCP8.5	HadGEM2-ES	Medium	
9	exp09	RCP8.5	MIROC5	High	
10	exp10	RCP8.5	MIROC5	Low	
A1	expa01	RCP8.5	IPSL-CM5A-MR	Medium	Extended
A2	expa02	RCP8.5	CSIRO-Mk3.6.0	Medium	ensemble
A3	expa03	RCP8.5	ACCESS1.3	Medium	(Tier 2)
B1	expb01	SSP5-8.5	CNRM-CM6-1	Medium	CMIP6 extension (Tier 2)
B2	expb02	SSP1-2.6	CNRM-CM6-1	Medium	
B3	expb03	SSP5-8.5	UKESM1-0-LL	Medium	
B4	expb04	SSP5-8.5	CESM2	Medium	
B5	expb05	SSP5-8.5	CNRM-ESM2-1	Medium	

Table 1. Extended ISMIP6-Greenland Tier-1 and 2 future climate experiments discussed in this study. See Nowicki and others (2020) for references for the GCMs.




Fig. 1. Extended ISMIP6-Greenland historical run (hist), projection control run (ctrl_proj) and Tier-1 and 2 future climate experiments: (a) Simulated ice mass change (counted positively for loss and expressed as sea-level contribution), (b) ice area. The red and blue boxes to the right show the mean ± 1 -sigma ranges for RCP8.5/SSP5.8.5 and RCP2.6/SSP1.2.6, respectively; the whiskers show the corresponding full ranges.




Fig. 2. Main components of the global mass balance for Exp. 5 (MIROC5/RCP8.5): Surface mass balance (SMB, red), basal mass balance (BMB, blue), calving (brown) and ice volume change (dV/dt , green). Note the shifted, right axis for the latter. The black and green dashed lines indicate the zero levels for the left and right axis, respectively.




Fig. 3. Ice thickness (panels a, b) and surface velocity (panels c, d) for the initial time (2015; panels a, c) and final time (3001; panels b, d) of Exp. 5 (MIROC5/RCP8.5).