

JOURNAL OF GLACIOLOGY



CAMBRIDGE
UNIVERSITY PRESS

THIS MANUSCRIPT HAS BEEN SUBMITTED TO THE JOURNAL OF GLACIOLOGY AND HAS NOT BEEN PEER-REVIEWED.

Mass loss of the Greenland ice sheet until the year 3000 under a sustained late-21st-century climate

Journal:	<i>Journal of Glaciology</i>
Manuscript ID	JOG-21-0089.R3
Manuscript Type:	Letter
Date Submitted by the Author:	23-Jan-2022
Complete List of Authors:	Greve, Ralf; Hokkaido University, Institute of Low Temperature Science Chambers, Christopher; Hokkaido University Institute of Low Temperature Science, Glaciology
Keywords:	Ice-sheet modelling, Arctic glaciology, Climate change, Ice and climate
Abstract:	We conduct extended versions of the ISMIP6 future climate experiments for the Greenland ice sheet until the year 3000 with the model SICOPOLIS. Beyond 2100, the climate forcing is kept fixed at late-21st-century conditions. For the unabated warming pathway RCP8.5/SSP5-8.5, the ice sheet suffers a severe mass loss, which amounts to ~ 1.8 m SLE (sea-level equivalent) for the twelve-experiment mean, and ~ 3.5 m SLE ($\sim 50\%$ of the entire mass) for the most sensitive experiment. For the reduced emissions pathway RCP2.6/SSP1-2.6, the mass loss is limited to a two-experiment mean of ~ 0.28 m SLE. Climate-change mitigation during the next decades will therefore be an efficient means for limiting the contribution of the Greenland ice sheet to sea-level rise in the long term.



SCHOLARONE™
Manuscripts

Mass loss of the Greenland ice sheet until the year 3000 under a sustained late-21st-century climate

Ralf GREVE,^{1,2} Christopher CHAMBERS¹

¹*Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan*

²*Arctic Research Center, Hokkaido University, Sapporo, Japan*

Correspondence: Ralf Greve <greve@lowtem.hokudai.ac.jp>

ABSTRACT. We conduct extended versions of the ISMIP6 future climate experiments for the Greenland ice sheet until the year 3000 with the model SICOPOLIS. Beyond 2100, the climate forcing is kept fixed at late-21st-century conditions. For the unabated warming pathway RCP8.5/SSP5-8.5, the ice sheet suffers a severe mass loss, which amounts to ~ 1.8 m SLE (sea-level equivalent) for the twelve-experiment mean, and ~ 3.5 m SLE ($\sim 50\%$ of the entire mass) for the most sensitive experiment. For the reduced emissions pathway RCP2.6/SSP1-2.6, the mass loss is limited to a two-experiment mean of ~ 0.28 m SLE. Climate-change mitigation during the next decades will therefore be an efficient means for limiting the contribution of the Greenland ice sheet to sea-level rise in the long term.

1 INTRODUCTION

It is established scientific consensus that the Earth's climate system is warming, and that human influence has been the dominant cause of the observed warming since the mid-20th century (e.g., IPCC, 2021). A major consequence of global warming is sea-level rise, currently (for the period 2006–2018) occurring at a global mean rate of 3.69 ± 0.48 millimetres per year. The main sources are melting/discharge of ice sheets, ice caps and glaciers ($\sim 45\%$), thermal expansion of ocean water ($\sim 38\%$), and changes in land water storage ($\sim 17\%$) (Fox-Kemper and others, 2021). In the long term, the two ice sheets of Antarctica (AIS) and Greenland (GrIS) are the largest potential contributors to global sea-level rise because of their

26 enormous volumes, together amounting to ~ 65 m SLE (sea-level equivalent) (Morlighem and others, 2017,
27 2020). The ice sheets have therefore been the focus of intensive observational as well as modelling efforts.

28 The Coupled Model Intercomparison Project Phase 6 (CMIP6) is a major international climate mod-
29 elling initiative (Eyring and others, 2016). As a part of this project, the Ice Sheet Model Intercomparison
30 Project for CMIP6 (ISMIP6) brought together a consortium of ice-sheet modellers to explore the sea-level-
31 rise contribution from the GrIS and AIS (Nowicki and others, 2016, 2020). ISMIP6 focussed on the CMIP6
32 period from 2015 until the end of 2100. The main findings for the GrIS, when forced by output from CMIP5
33 global climate models (GCMs), were contributions of 90 ± 50 and 32 ± 17 mm SLE for the unabated warm-
34 ing pathway RCP8.5 [RCP: Representative Concentration Pathway] and the reduced emissions pathway
35 RCP2.6, respectively (Goelzer and others, 2020). The CMIP6 GCMs tend to feature a warmer atmosphere,
36 which results in higher mass loss due to increased surface melt (Payne and others, 2021). For the AIS and
37 CMIP5 climate forcings, ISMIP6 found a mass loss in the range of -7.8 to 30.0 cm SLE under RCP8.5
38 (Seroussi and others, 2020). The limited number of results for RCP2.6 fall within this range, and so do
39 the results obtained with CMIP6 climate forcings (Payne and others, 2021). This rather unclear picture
40 for the AIS is a consequence of the counteracting effects of mass loss due to ocean warming and mass gain
41 from increased snowfall.

42 The full suite of ISMIP6 experiments with both CMIP5 and CMIP6 forcings was carried out with
43 the ice-sheet model SICOPOLIS (“SIMulation COde for POLythermal Ice Sheets”, www.sicopolis.net), as
44 documented in detail by Greve and others (2020a,b). Chambers and others (2021) extend the ISMIP6 sim-
45 ulations for the AIS with SICOPOLIS until the year 3000, assuming a sustained late-21st-century climate
46 beyond 2100. Compared to the uncertain response projected over the ISMIP6 period, a radically different
47 picture emerges, demonstrating that the consequences of the high-emissions scenario RCP8.5/SSP5-8.5
48 [SSP: Shared Socioeconomic Pathway] are much greater in the long term even if no further climate trend
49 is applied beyond 2100.

50 The response of the GrIS to longer-term climate change has also been investigated. In addition to
51 an ensemble of projections for the 21st century with their higher-order ice-sheet model, the study by
52 Fürst and others (2015) also conducted projections until 2300 for the two low-emission scenarios RCP2.6
53 and RCP4.5, forced by selected CMIP5 GCMs. Vizcaino and others (2015) carried out simulations until
54 2300 with a coupled ECHAM5.2/MPI-OM/SICOPOLIS model for the pathways RCP2.6, RCP4.5, and a
55 modified RCP8.5 with a $4 \times \text{CO}_2$ limit. Calov and others (2018) drove an extended version of SICOPOLIS

56 (“IGLOO”) with RCP4.5 and RCP8.5 surface temperature and surface mass balance anomalies created by
57 the regional climate model MAR with boundary conditions from simulations with three CMIP5 GCMs.
58 Similar to our approach, prolongation of the climatic forcing beyond 2100 was done by assuming no
59 further warming trend. Aschwanden and others (2019) used projections based on four CMIP5 GCMs
60 until 2300 for RCP2.6, RCP4.5 and RCP8.5, extrapolated until 3000, to force the ice-sheet model PISM.
61 Their climatic forcing was processed by the regional climate model HIRHAM5, delivering the surface-
62 temperature anomaly as the main driver, from which precipitation changes were parameterized, and runoff
63 was computed by a positive-degree-day (PDD) method. Van Breedam and others (2020) projected the
64 response of the GrIS and AIS 10,000 years into the future with the Earth system model of intermediate
65 complexity LOVECLIMv1.3 (including the ice sheet model AGISM), forced by the extended concentration
66 pathways ECP2.6, 4.5, 6.0 and 8.5 until 2300 and zero emissions thereafter. In the present study, we
67 transfer the approach by Chambers and others (2021) to the GrIS. The objective is to assess its long-term
68 response to late-21st-century climatic conditions for the full ensemble of ISMIP6 climate forcings, which
69 consists of fourteen scenarios from ten different CMIP5 and CMIP6 GCMs.

70 2 METHODS

71 The main tool used for this study is the ice-sheet model SICOPOLIS. We apply it to the GrIS with
72 hybrid shallow-ice-shelfy-stream dynamics (Bernales and others, 2017), a Weertman-Budd-type sliding
73 law tuned separately for 20 different regions (Greve and others, 2020b), and ice thermodynamics treated
74 by the one-layer melting-CTS enthalpy scheme (CTS: cold-temperate transition surface; Blatter and Greve,
75 2015; Greve and Blatter, 2016). The horizontal resolution is 5 km. In the vertical, we use terrain-following
76 coordinates (sigma transformation) with 81 layers in the ice domain and 41 layers in the thermal lithosphere
77 layer below. For details on the set-up, the initialization procedure by a paleoclimatic spin-up, comparisons
78 between the simulated and observed ice thickness and surface velocity for our initialization year 1990, as
79 well as the historical run (“hist”) that bridges the gap between 1990 and the start date of the projections in
80 January 2015 by employing MIROC5/RCP8.5 surface mass balance (SMB) and surface temperature (ST)
81 forcing, we refer to Greve and others (2020b).

82 Following the ISMIP6 protocol, climate forcing from 2015 until the end of 2100 has an atmospheric and
83 an oceanic component. The atmospheric forcing consists of a 1960–1989 reference climatology, plus space-
84 time-dependent anomalies for SMB and ST. These were derived from a systematic sampling of CMIP5

85 GCMs that reflects their spread in future projections (Barthel and others, 2020), while CMIP6 GCMs were
86 added on the basis of availability only (Payne and others, 2021). All GCM forcings were downscaled to
87 the GrIS surface with the regional climate model MAR v3.9.6 (Fettweis and others, 2017; Delhasse and
88 others, 2020). Although MAR uses a static GrIS, it also provides vertical gradients of SMB and ST, thus
89 allowing to include SMB–height and ST–height feedbacks in the ice-sheet simulations (Franco and others,
90 2012; Nowicki and others, 2020). The oceanic forcing is based on a retreat parameterization for tidewater
91 glaciers, forced by MAR runoff and ocean temperature changes specified for seven ice–ocean sectors around
92 Greenland (Slater and others, 2019, 2020).

93 For the period from 2101 until the end of 3000, we extend the simulations in a similar way than
94 Chambers and others (2021) do for the AIS. For every year of this extended period, the atmospheric
95 forcing (SMB, ST, vertical gradients) for the 10-year interval 2091–2100 is randomly sampled such that
96 no further trend is applied, but some inter-annual fluctuations remain (similar to Calov and others, 2018).
97 The oceanic forcing (prescribed retreat maps) does not show any notable year-to-year fluctuations, so we
98 simply keep it fixed at 2100 conditions.

99 An overview of our extended ISMIP6 experiments is given in Table 1, and the magnitude of the
100 atmospheric forcing is shown in supplementary Table S1. Twelve experiments are for the 21st-century
101 unabated warming pathway RCP8.5 (CMIP5) / SSP5-8.5 (CMIP6), and two are for the reduced emissions
102 pathway RCP2.6 (CMIP5) / SSP1-2.6 (CMIP6) that is largely in line with the commitments of the Paris
103 Agreement (maintaining the global mean temperature well below a 2°C increase above pre-industrial levels).
104 In two experiments, the impact of different sensitivities of the retreat parameterization due to oceanic
105 forcing (“high” and “low” vs. the normal, “medium” sensitivity, thereby exploring the uncertainty of the
106 parameterization; Slater and others, 2019, 2020) is tested. In addition, a projection control run for the
107 period 2015–3000 (“ctrl_proj”) employs constant climate conditions based on a 1960–1989 climatology and
108 no explicit oceanic forcing.

109 **3 RESULTS**

110 The simulated mass change of the GrIS, expressed as a sea-level contribution, and ice area are shown
111 in Figure 1. For the control run ctrl_proj, the ice sheet remains nearly stable, showing a slight mass
112 gain of 6.4 mm SLE and area loss of $4.7 \times 10^3 \text{ km}^2$ during the 986 years model time, which is of the order
113 of permilles of the present-day values. For all future projections, the ice sheet keeps losing both mass

114 and extent over the entire period. (Values of the resulting sea-level contribution by 2100, 2300 and 3000
115 relative to `ctrl_proj` are provided in the supplementary Table S2.) The largest rate of change occurs
116 typically around the year 2100, beyond which it slows down to some extent; however, without reaching or
117 coming close to a new steady state. This demonstrates that the committed mass loss due to 21st-century
118 climate change extends way beyond the 21st century and impacts the ice sheet on a much longer time
119 scale. Corroborating the findings for the 21st century (Goelzer and others, 2020; Greve and others, 2020b),
120 the GrIS responds much more strongly to the ensemble of RCP8.5/SSP5-8.5 simulations than to the two
121 RCP2.6/SSP1-2.6 simulations. By the year 3000, the mass loss amounts to 1.79 ± 0.80 m SLE (mean \pm
122 1-sigma range) for RCP8.5/SSP5-8.5, while it is limited to 0.28 ± 0.12 m SLE for RCP2.6/SSP1-2.6.

123 The percentage area loss (Fig. 1b) is similar to the mass loss. However, it is striking that the variability
124 in area varies significantly between the different simulations. This results from the different variability of
125 the SMB forcing and affects mainly the thin, near-margin parts of the ice sheet, which do not contribute
126 much to the total ice mass. Therefore, the variability is not paralleled in Fig. 1a.

127 The influence of the ice retreat due to oceanic forcing is explored by Exps. 5, 9, 10 (MIROC5/RCP8.5
128 with “medium”, “high” and “low” sensitivity, respectively). The results are shown by the olive lines
129 and olive-shaded regions in Figure 1. By 3000, the simulated mass loss is $1.62^{+0.051}_{-0.031}$ m SLE. Thus, the
130 uncertainty due to these three calibrations is very small in the long range. Relative to the uncertainty due
131 to the different climate forcings, it is more pronounced for the 21st century (Greve and others, 2020b).
132 This is because the continued retreat of the ice sheet decreases its contact with the ocean, so that the
133 oceanic forcing plays a smaller role in the longer term.

134 As reported by Greve and others (2020b) and Payne and others (2021), for both the 21st-century
135 RCP8.5/SSP5-8.5 and RCP2.6/SSP1-2.6 pathways, the CMIP6 climate models produce a larger response
136 of the ice sheet than the CMIP5 ones. While the significance of this statement is limited in the case of
137 RCP2.6/SSP1-2.6 (only one experiment each), it is more robust for RCP8.5/SSP5-8.5, where the ensemble
138 contains eight and four experiments forced by CMIP5 and CMIP6 models, respectively. By 3000, the mean
139 mass loss for the four CMIP6 SSP5-8.5 experiments is 2.73 m SLE, and the maximum value from Exp. B4
140 (CESM2/SSP5-8.5) is as large as 3.54 m SLE, almost 50% of the entire present-day ice mass.

141 We now discuss in more detail the results of Exp. 5 (MIROC5/RCP8.5), which was already focused on
142 in the original ISMIP6-Greenland study (Goelzer and others, 2020). Mainly due to its large SMB forcing,
143 it produces, along with Exp. A1 (IPSL-CM5A-MR/RCP8.5), the strongest mass loss among the CMIP5

144 forcings, while the mass loss is about average for our combined CMIP5/CMIP6 ensemble. Figure 2 shows
145 the components of the global mass balance (integrated over the ice sheet, all counted as positive for mass
146 gain): surface mass balance (SMB), basal mass balance (BMB), calving and ice volume change (dV/dt). On
147 a mean-annual basis, the residual, $\text{Res} = |\text{SMB} + \text{BMB} + \text{Calving} - dV/dt|$, is always less than $10^6 \text{ m}^3 \text{ a}^{-1}$.
148 This is five to six orders of magnitude smaller than the typical range of values in the figure, so that the
149 model conserves mass very well (see also Calov and others, 2018).

150 As already stated above, the ice sheet keeps losing volume (\propto mass) over the entire period, with
151 maximal rates of change occurring shortly before the year 2100. The SMB is initially positive, but changes
152 its sign in the second half of the 21st century and stays negative beyond that. Calving into the surrounding
153 ocean peaks during 2080–2085, when it contributes approximately the same amount to ice volume loss than
154 negative SMB. After that, calving decreases continuously due to ice-sheet retreat from the coast (loss of
155 ocean contact) and becomes almost negligible towards the end of the 3rd millennium. The decrease is
156 likely accelerated by a limitation of the oceanic forcing approach: the fixed retreat mask beyond 2100
157 does not follow the retreating ice margin further, making it ineffective as the ice sheet recedes beyond its
158 reach. The negative SMB shows a slightly decreasing trend after 2100 because, as the ice sheet shrinks,
159 less area is available for further melting. Both effects together cause the magnitude of dV/dt (loss rate of
160 ice volume) to decrease gradually, which results in the slight flattening of the curves in Figure 1 (the same
161 mechanisms are effective for the other experiments). BMB is relatively small over the entire model time.
162 The inter-annual variability of the volume change is due to that of the SMB, which reflects the variability
163 of the atmospheric forcing.

164 Figure 3 shows snapshots of the ice thickness and surface velocity for Exp. 5 for the initial year 2015
165 and the final year 3000. Comparing the thickness distributions demonstrates nicely that the ice sheet
166 retreats from the coast almost all around its perimeter, and contact to the ocean is very limited by the
167 end of the simulation, which entails the low calving rates mentioned above. By contrast, the ice sheet
168 does not suffer much change in its interior parts north of $\sim 68^\circ\text{N}$. The large-scale pattern of the ice flow
169 and the organization of the ice sheet into major drainage basins remain largely intact. However, on the
170 regional scale, more pronounced changes occur. The fast-flowing outlet glaciers in south-western Greenland
171 disappear entirely due to the extreme retreat in this area. The north-western outlet glaciers, including
172 Petermann Glacier, also slow down substantially. The central-western Jakobshavn Ice Stream loses its
173 clear delimitation to the surrounding glaciers, but remains an area of fast-flowing ice. The major features

174 in East Greenland, e.g., the North-East Greenland Ice Stream, Kangerdlugssuaq and Helheim glaciers, are
175 less affected and remain well identifiable.

176 4 DISCUSSION AND CONCLUSION

177 The future climate simulations carried out in this study for the GrIS over the 3rd millennium confirm and
178 continue the trends that were reported by ISMIP6-Greenland for the 21st century (Goelzer and others,
179 2020; Greve and others, 2020b; Payne and others, 2021). The response of the ice sheet is mainly governed
180 by a negative SMB due to increased surface melting near the ice margin. Marine-terminating glacier
181 retreat, caused by increased oceanic thermal forcing and increased meltwater runoff, constitutes a further
182 negative contribution to the total mass balance, but becomes less important in the longer term. Under the
183 unabated warming pathway RCP8.5/SSP5-8.5, this leads to a severe mass loss during the 3rd millennium,
184 while the loss is much smaller under the reduced emissions pathway RCP2.6/SSP1-2.6. Results obtained
185 with forcings from the newer CMIP6 climate models consistently produce larger mass losses than those
186 obtained with the older CMIP5 models, for SSP5-8.5 in the range of a $\sim 25\text{--}50\%$ loss of the present-day
187 ice mass (and area) by 3000. For comparison, Aschwanden and others (2019) reported a mass loss of
188 $\sim 75\text{--}100\%$ by 3000 for their ensemble of RCP8.5 simulations, for which a warming trend is assumed to
189 continue until 2500. Efficient climate change mitigation during the next decades is therefore crucial for
190 limiting the contribution of the GrIS to long-term sea-level rise.

191 As for interpreting the stronger response of the GrIS to CMIP6-derived forcings compared to CMIP5-
192 derived ones, the different strategies of sampling the respective GCM ensembles for ISMIP6-Greenland must
193 be considered. As we mentioned in Sect. 2, the six CMIP5 GCMs were chosen systematically, whereas the
194 four CMIP6 GCMs were the only ones available for downscaling at the time. Subsequent analysis of the
195 entire CMIP6 model ensemble revealed that the results cluster around two groups of climate sensitivities
196 (global mean temperature response to doubled atmospheric CO_2), and the four models that were available
197 for ISMIP6-Greenland all fall in the high-sensitivity group (Meehl and others, 2020; Payne and others,
198 2021). Future work should therefore aim at conducting additional simulations with a more representative
199 sampling of the CMIP6 GCMs.

200 Our study is limited to investigating the impact of a sustained late-21st-century climate (without
201 imposing a further trend beyond 2100) on the GrIS. However, climate change is projected to continue beyond
202 2100 (e.g., Bakker and others, 2016; Lyon and others, 2022), with potentially even more devastating effects

203 on the GrIS than reported here, plus a significant decay of the AIS in the long term (Van Breedam and
204 others, 2020). Further, the unidirectional coupling approach (climate model → ice-sheet model) employed
205 by ISMIP6, and thus here, lacks a detailed accounting of feedbacks of the changing ice sheet on the climate.
206 As we explained in Sect. 2, the climate forcing for Greenland includes vertical gradients of the surface mass
207 balance and surface temperature. Therefore, the changing ice-sheet geometry acts back on these climatic
208 forcing fields. However, the linearized approach was derived for small perturbations of the present-day
209 state only, and it cannot be validated for large changes of the ice sheet. This shortcoming becomes more
210 severe in our simulations over almost a millennium compared to the 86-year scope of ISMIP6, and it is
211 not possible to judge a priori whether it rather leads to an over- or underestimation of the simulated mass
212 loss. Using the PDD method for modelling runoff, like in the study by Aschwanden and others (2019),
213 allows in principle to handle arbitrarily large changes of the ice sheet. PDD is also easy to implement in
214 an ice-sheet model, which adds to its appeal. However, it is a grossly simplifying parameterization of the
215 complex energy balance at the surface of an ice sheet and difficult to calibrate for climatic conditions other
216 than present-day ones (e.g., Bauer and Ganopolski, 2017), so that we do not consider it superior to the
217 ISMIP6-type approach of our study.

218 Future work in the direction of long-term simulations of ice-sheet response to climate change should aim
219 at employing GCM projections beyond 2100 and improving the representation of feedback processes. The
220 ultimate solution would be to carry out such simulations in a fully coupled way, with the ice-sheet model
221 integrated in the GCM. This approach has been pursued (e.g., Vizcaino and others, 2015; Sellevold and
222 others, 2019; Gregory and others, 2020; Muntjewerf and others, 2020a,b); however, fully coupled simulations
223 are demanding and computationally expensive, which makes it difficult to run large ensembles, involving
224 many different climate and ice-sheet models, over long time scales and at adequate resolution. Intermediary,
225 more manageable solutions may consist of involving snapshots of climate-model results combined with more
226 refined parameterizations for the climatic forcing, similar to the approach by Abe-Ouchi and others (2013)
227 for the paleo-glaciation of the Northern Hemisphere.

228 SUPPLEMENTARY MATERIAL

229 The supplementary material for this article can be found at <https://doi.org/10.1017/jog.xxxx.xxxx>.

230 **CODE AND DATA AVAILABILITY**

231 SICOPOLIS is free and open-source software, available through a persistent Git repository hosted by the
232 Alfred Wegener Institute for Polar and Marine Research (AWI) in Bremerhaven, Germany (Greve and
233 SICOPOLIS Developer Team, 2021). Detailed instructions for obtaining and compiling the code are at
234 <http://www.sicopolis.net> (last access: 2 November 2021). The output data produced for this study are
235 available at Zenodo, <https://doi.org/10.5281/zenodo.5880518>.

236 **ACKNOWLEDGEMENTS**

237 We thank the two anonymous reviewers, the Scientific Editor Frank Pattyn and the Chief Editor Hester
238 Jiskoot for constructive remarks and suggestions that helped to improve the manuscript. We thank Jorge
239 Bernales (MARUM Bremen), Reinhard Calov (PIK Potsdam), Takashi Obase (University of Tokyo) and
240 Fuyuki Saito (JAMSTEC Yokohama) for their recent contributions to the development of the SICOPOLIS
241 model, and Ayako Abe-Ouchi for fruitful discussions about ice-sheet and climate modelling. Some colour
242 schemes of our figures were taken from Paul Tol's (SRON Netherlands Institute for Space Research) online
243 resource at <https://personal.sron.nl/~pault>. We thank the Climate and Cryosphere (CliC) effort, which
244 provided support for ISMIP6 through sponsoring of workshops, hosting the ISMIP6 website and wiki,
245 and promoting ISMIP6. We acknowledge the World Climate Research Programme, which, through its
246 Working Group on Coupled Modelling, coordinated and promoted CMIP5 and CMIP6. We thank the
247 climate modelling groups for producing their model output and making it available; the Earth System Grid
248 Federation (ESGF) for archiving the CMIP data and providing access to it; the University at Buffalo for
249 ISMIP6 data distribution and upload; and the multiple funding agencies who support CMIP5, CMIP6, and
250 ESGF. We thank the ISMIP6 steering committee, the ISMIP6 model selection group and ISMIP6 dataset
251 preparation group for their continuous engagement in defining ISMIP6. This is ISMIP6 contribution No. 26.

252 Ralf Greve and Christopher Chambers were supported by Japan Society for the Promotion of Science
253 (JSPS) KAKENHI Grant No. JP17H06323, and by a Leadership Research Grant (Category 2) of Hokkaido
254 University's Institute of Low Temperature Science. Ralf Greve was supported by JSPS KAKENHI Grant
255 No. JP17H06104, and by the Japanese Ministry of Education, Culture, Sports, Science and Technology
256 (MEXT) through the Arctic Challenge for Sustainability project ArCS II (program grant number JP-
257 MXD1420318865).

258 **REFERENCES**

- 259 Abe-Ouchi A, Saito F, Kawamura K, Raymo ME, Okuno J, Takahashi K and Blatter H (2013) Insolation-driven
260 100,000-year glacial cycles and hysteresis of ice-sheet volume. *Nature*, **500**(7461), 190–193 (doi: 10.1038/nature12374)
261 (doi: 10.1038/nature12374)
- 262 Aschwanden A, Fahnestock MA, Truffer M, Brinkerhoff DJ, Hock R, Khroulev C, Mottram R and Khan SA (2019)
263 Contribution of the Greenland Ice Sheet to sea level over the next millennium. *Science Advances*, **5**(6), eaav9396
264 (doi: 10.1126/sciadv.aav9396)
- 265 Bakker P, Schmittner A, Lenaerts JTM, Abe-Ouchi A, Bi D, van den Broeke MR, Chan WL, Hu A, Beadling RL,
266 Marsland SJ, Mernild SH, Saenko OA, Swingedouw D, Sullivan A and Yin J (2016) Fate of the Atlantic Meridional
267 Overturning Circulation: Strong decline under continued warming and Greenland melting. *Geophysical Research
268 Letters*, **43**, 12252–12260 (doi: 10.1002/2016GL070457)
- 269 Barthel A, Agosta C, Little CM, Hattermann T, Jourdain NC, Goelzer H, Nowicki S, Seroussi H, Straneo F and
270 Bracegirdle TJ (2020) CMIP5 model selection for ISMIP6 ice sheet model forcing: Greenland and Antarctica. *The
271 Cryosphere*, **14**(3), 855–879 (doi: 10.5194/tc-14-855-2020)
- 272 Bauer E and Ganopolski A (2017) Comparison of surface mass balance of ice sheets simulated by positive-degree-day
273 method and energy balance approach. *Climate of the Past*, **13**(7), 819–832 (doi: 10.5194/cp-13-819-2017)
- 274 Bernales J, Rogozhina I, Greve R and Thomas M (2017) Comparison of hybrid schemes for the combination of
275 shallow approximations in numerical simulations of the Antarctic Ice Sheet. *The Cryosphere*, **11**(1), 247–265 (doi:
276 10.5194/tc-11-247-2017)
- 277 Blatter H and Greve R (2015) Comparison and verification of enthalpy schemes for polythermal glaciers and ice
278 sheets with a one-dimensional model. *Polar Science*, **9**(2), 196–207 (doi: 10.1016/j.polar.2015.04.001)
- 279 Calov R, Beyer S, Greve R, Beckmann J, Willeit M, Kleiner T, Rückamp M, Humbert A and Ganopolski A (2018)
280 Simulation of the future sea level contribution of Greenland with a new glacial system model. *The Cryosphere*,
281 **12**(10), 3097–3121 (doi: 10.5194/tc-12-3097-2018)
- 282 Chambers C, Greve R, Obase T, Saito F and Abe-Ouchi A (2021) Mass loss of the Antarctic ice sheet until the year
283 3000 under a sustained late-21st-century climate. *Journal of Glaciology* (doi: 10.1017/jog.2021.124)
- 284 Delhasse A, Kittel C, Amory C, Hofer S, van As D, S Fausto R and Fettweis X (2020) Brief communication:
285 Evaluation of the near-surface climate in ERA5 over the Greenland Ice Sheet. *The Cryosphere*, **14**(3), 957–965
286 (doi: 10.5194/tc-14-957-2020)

- 287 Eyring V, Bony S, Meehl GA, Senior CA, Stevens B, Stouffer RJ and Taylor KE (2016) Overview of the Coupled Model
288 Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*,
289 **9**(5), 1937–1958 (doi: 10.5194/gmd-9-1937-2016)
- 290 Fettweis X, Box JE, Agosta C, Amory C, Kittel C, Lang C, van As D, Machguth H and Gallée H (2017) Recon-
291 structions of the 1900–2015 Greenland ice sheet surface mass balance using the regional climate MAR model. *The*
292 *Cryosphere*, **11**(2), 1015–1033 (doi: 10.5194/tc-11-1015-2017)
- 293 Fox-Kemper B, Hewitt HT, Xiao C, Aðalgeirsdóttir G, Drijfhout SS, Edwards TL, Golledge NR, Hemer M, Kopp
294 RE, Krinner G, Mix A, Notz D, Nowicki S, Nurhati IS, Ruiz L, Sallée JB, Slangen ABA and Yu Y (2021) Ocean,
295 cryosphere and sea level change. In V Masson-Delmotte, P Zhai, A Pirani, SL Connors, C Péan, S Berger, N Caud,
296 Y Chen, L Goldfarb, MI Gomis, M Huang, K Leitzell, E Lonnoy, JBR Matthews, TK Maycock, T Waterfield,
297 O Yelekçi, R Yu and B Zhou (eds.), *Climate Change 2021: The Physical Science Basis. Contribution of Working*
298 *Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University
299 Press, in press
- 300 Franco B, Fettweis X, Lang C and Erpicum M (2012) Impact of spatial resolution on the modelling of the Greenland
301 ice sheet surface mass balance between 1990–2010, using the regional climate model MAR. *The Cryosphere*, **6**(3),
302 695–711 (doi: 10.5194/tc-6-695-2012)
- 303 Fürst JJ, Goelzer H and Huybrechts P (2015) Ice-dynamic projections of the Greenland ice sheet in response to
304 atmospheric and oceanic warming. *The Cryosphere*, **9**(3), 1039–1062 (doi: 10.5194/tc-9-1039-2015)
- 305 Goelzer H, Nowicki S, Payne A, Larour E, Seroussi H, Lipscomb WH, Gregory J, Abe-Ouchi A, Shepherd A, Simon E,
306 Agosta C, Alexander P, Aschwanden A, Barthel A, Calov R, Chambers C, Choi Y, Cuzzzone J, Dumas C, Edwards
307 T, Felikson D, Fettweis X, Golledge NR, Greve R, Humbert A, Huybrechts P, Le clec'h S, Lee V, Leguy G, Little
308 C, Lowry DP, Morlighem M, Nias I, Quiquet A, Rückamp M, Schlegel NJ, Slater D, Smith R, Straneo F, Tarasov
309 L, van de Wal R and van den Broeke M (2020) The future sea-level contribution of the Greenland ice sheet: a
310 multi-model ensemble study of ISMIP6. *The Cryosphere*, **14**(9), 3071–3096 (doi: 10.5194/tc-14-3071-2020)
- 311 Gregory JM, George SE and Smith RS (2020) Large and irreversible future decline of the Greenland ice sheet. *The*
312 *Cryosphere*, **14**(12), 4299–4322 (doi: 10.5194/tc-14-4299-2020)
- 313 Greve R and Blatter H (2016) Comparison of thermodynamics solvers in the polythermal ice sheet model SICOPOLIS.
314 *Polar Science*, **10**(1), 11–23 (doi: 10.1016/j.polar.2015.12.004)
- 315 Greve R and SICOPOLIS Developer Team (2021) SICOPOLIS. GitLab, Alfred Wegener Institute for Polar and
316 Marine Research, Bremerhaven, Germany, URL <https://gitlab.awi.de/sicopolis/sicopolis>

- 317 Greve R, Calov R, Obase T, Saito F, Tsutaki S and Abe-Ouchi A (2020a) ISMIP6 future projections for the Antarctic
318 ice sheet with the model SICOPOLIS. Technical report, Zenodo (doi: 10.5281/zenodo.3971232)
- 319 Greve R, Chambers C and Calov R (2020b) ISMIP6 future projections for the Greenland ice sheet with the model
320 SICOPOLIS. Technical report, Zenodo (doi: 10.5281/zenodo.3971251)
- 321 IPCC (2021) Summary for policymakers. In V Masson-Delmotte, P Zhai, A Pirani, SL Connors, C Péan, S Berger,
322 N Caud, Y Chen, L Goldfarb, MI Gomis, M Huang, K Leitzell, E Lonnoy, JBR Matthews, TK Maycock, T Wa-
323 terfield, O Yelekçi, R Yu and B Zhou (eds.), *Climate Change 2021: The Physical Science Basis. Contribution of*
324 *Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge
325 University Press, in press
- 326 Lyon C, Saupé EE, Smith CJ, Hill DJ, Beckerman AP, Stringer LC, Marchant R, McKay J, Burke A, O'Higgins P,
327 Dunhill AM, Allen BJ, Riel-Salvatore J and Aze T (2022) Climate change research and action must look beyond
328 2100. *Global Change Biology*, **28**(2), 349–361 (doi: 10.1111/gcb.15871)
- 329 Meehl GA, Senior CA, Eyring V, Flato G, Lamarque JF, Stouffer RJ, Taylor KE and Schlund M (2020) Context for
330 interpreting equilibrium climate sensitivity and transient climate response from the CMIP6 earth system models.
331 *Science Advances*, **6**(26), eaba1981 (doi: 10.1126/sciadv.aba1981)
- 332 Morlighem M, Williams CN, Rignot E, An L, Arndt JE, Bamber JL, Catania G, Chauché N, Dowdeswell JA, Dorschel
333 B, Fenty I, Hogan K, Howat I, Hubbard A, Jakobsson M, Jordan TM, Kjeldsen KK, Millan R, Mayer L, Mouginot
334 J, Noël BPY, O'Cofaigh C, Palmer S, Rysgaard S, Seroussi H, Siegert MJ, Slabon P, Straneo F, van den Broeke
335 MR, Weinrebe W, Wood M and Zinglarsen KB (2017) BedMachine v3: Complete bed topography and ocean
336 bathymetry mapping of Greenland from multibeam echo sounding combined with mass conservation. *Geophysical*
337 *Research Letters*, **44**(21), 11051–11061 (doi: 10.1002/2017GL074954)
- 338 Morlighem M, Rignot E, Binder T, Blankenship D, Drews G, Rand Eagles, Eisen O, Ferraccioli F, Forsberg R, Fretwell
339 P, Goel V, Greenbaum JS, Gudmundsson H, Guo J, Helm V, Hofstede C, Howat I, Humbert A, Jokat W, Karlsson
340 NB, Lee WS, Matsuoka K, Millan R, Mouginot J, Paden J, Pattyn F, Roberts J, Rosier S, Ruppel A, Seroussi H,
341 Smith EC, Steinhage D, Sun B, van den Broeke MR, van Ommen TD, van Wessem M and Young DA (2020) Deep
342 glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet. *Nature Geoscience*,
343 **13**(2), 132–137 (doi: 10.1038/s41561-019-0510-8)
- 344 Muntjewerf L, Petrini M, Vizcaino M, Ernani da Silva C, Sellevold R, Scherrenberg MDW, Thayer-Calder K, Bradley
345 SL, Lenaerts JTM, Lipscomb WH and Lofverstrom M (2020a) Greenland Ice Sheet contribution to 21st century
346 sea level rise as simulated by the coupled CESM2.1-CISM2.1. *Geophysical Research Letters*, **47**(9), e2019GL086836
347 (doi: 10.1029/2019GL086836)

- 348 Muntjewerf L, Sellevold R, Vizcaino M, Ernani da Silva C, Petrini M, Thayer-Calder K, Scherrenberg MDW, Bradley
349 SL, Katsman CA, Fyke J, Lipscomb WH, Lofverstrom M and Sacks WJ (2020b) Accelerated Greenland Ice Sheet
350 mass loss under high greenhouse gas forcing as simulated by the coupled CESM2.1-CISM2.1. *Journal of Advances
351 in Modeling Earth Systems*, **12**(10), e2019MS002031 (doi: 10.1029/2019MS002031)
- 352 Nowicki S, Goelzer H, Seroussi H, Payne AJ, Lipscomb WH, Abe-Ouchi A, Agosta C, Alexander P, Asay-Davis
353 XS, Barthel A, Bracegirdle TJ, Cullather R, Felikson D, Fettweis X, Gregory JM, Hattermann T, Jourdain NC,
354 Kuipers Munneke P, Larour E, Little CM, Morlighem M, Nias I, Shepherd A, Simon E, Slater D, Smith RS, Straneo
355 F, Trusel LD, van den Broeke MR and van de Wal R (2020) Experimental protocol for sea level projections from
356 ISMIP6 stand-alone ice sheet models. *The Cryosphere*, **14**(7), 2331–2368 (doi: 10.5194/tc-14-2331-2020)
- 357 Nowicki SMJ, Payne A, Larour E, Seroussi H, Goelzer H, Lipscomb W, Gregory J, Abe-Ouchi A and Shepherd A
358 (2016) Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6. *Geoscientific Model Develop-
359 ment*, **9**(12), 4521–4545 (doi: 10.5194/gmd-9-4521-2016)
- 360 Payne AJ, Nowicki S, Abe-Ouchi A, Agosta C, Alexander P, Albrecht T, Asay-Davis X, Aschwanden A, Barthel
361 A, Bracegirdle TJ, Calov R, Chambers C, Choi Y, Cullather R, Cuzzone J, Dumas C, Edwards TL, Felikson
362 D, Fettweis X, Galton-Fenzi BK, Goelzer H, Gladstone R, Golledge NR, Gregory JM, Greve R, Hattermann T,
363 Hoffman MJ, Humbert A, Huybrechts P, Jourdain NC, Kleiner T, Kuipers Munneke P, Larour E, Le clec'h S, Lee
364 V, Leguy G, Lipscomb WH, Little CM, Lowry DP, Morlighem M, Nias I, Pattyn F, Pelle T, Price SF, Quiquet
365 A, Reese R, Rückamp M, Schlegel NJ, Seroussi H, Shepherd A, Simon E, Slater D, Smith RS, Straneo F, Sun
366 S, Tarasov L, Trusel LD, Van Breedam J, van de Wal R, van den Broeke M, Winkelmann R, Zhao C, Zhang T
367 and Zwinger T (2021) Future sea level change under Coupled Model Intercomparison Project Phase 5 and Phase
368 6 scenarios from the Greenland and Antarctic ice sheets. *Geophysical Research Letters*, **48**(16), e2020GL091741
369 (doi: 10.1029/2020GL091741)
- 370 Sellevold R, van Kampenhout L, Lenaerts JTM, Noël B, Lipscomb WH and Vizcaino M (2019) Surface mass balance
371 downscaling through elevation classes in an Earth system model: application to the Greenland ice sheet. *The
372 Cryosphere*, **13**(12), 3193–3208 (doi: 10.5194/tc-13-3193-2019)
- 373 Seroussi H, Nowicki S, Payne AJ, Goelzer H, Lipscomb WH, Abe-Ouchi A, Agosta C, Albrecht T, Asay-Davis X,
374 Barthel A, Calov R, Cullather R, Dumas C, Galton-Fenzi BK, Gladstone R, Golledge N, Gregory JM, Greve R,
375 Hatterman T, Hoffman MJ, Humbert A, Huybrechts P, Jourdain NC, Kleiner T, Larour E, Leguy GR, Lowry DP,
376 Little CM, Morlighem M, Pattyn F, Pelle T, Price SF, Quiquet A, Reese R, Schlegel NJ, Shepherd A, Simon E,
377 Smith RS, Straneo F, Sun S, Trusel LD, Van Breedam J, van de Wal RSW, Winkelmann R, Zhao C, Zhang T and
378 Zwinger T (2020) ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st
379 century. *The Cryosphere*, **14**(9), 3033–3070 (doi: 10.5194/tc-14-3033-2020)

- 380 Slater DA, Straneo F, Felikson D, Little CM, Goelzer H, Fettweis X and Holte J (2019) Estimating Greenland
381 tidewater glacier retreat driven by submarine melting. *The Cryosphere*, **13**(9), 2489–2509 (doi: 10.5194/tc-13-
382 2489-2019)
- 383 Slater DA, Felikson D, Straneo F, Goelzer H, Little CM, Morlighem M, Fettweis X and Nowicki S (2020) Twenty-first
384 century ocean forcing of the Greenland ice sheet for modelling of sea level contribution. *The Cryosphere*, **14**(3),
385 985–1008 (doi: 10.5194/tc-14-985-2020)
- 386 Van Breedam J, Goelzer H and Huybrechts P (2020) Semi-equilibrated global sea-level change projections for the
387 next 10 000 years. *Earth System Dynamics*, **11**(4), 953–976 (doi: 10.5194/esd-11-953-2020)
- 388 Vizcaino M, Mikolajewicz U, Ziemen F, Rodehacke CB, Greve R and van den Broeke MR (2015) Coupled simulations
389 of Greenland Ice Sheet and climate change up to A.D. 2300. *Geophysical Research Letters*, **42**(10), 3927–3935 (doi:
390 10.1002/2014GL061142)

#	exp_id	Scenario	GCM	Ocean forcing	
5	exp05	RCP8.5	MIROC5	Medium	Core experiments (Tier 1)
6	exp06	RCP8.5	NorESM1-M	Medium	
7	exp07	RCP2.6	MIROC5	Medium	
8	exp08	RCP8.5	HadGEM2-ES	Medium	
9	exp09	RCP8.5	MIROC5	High	
10	exp10	RCP8.5	MIROC5	Low	
A1	expa01	RCP8.5	IPSL-CM5A-MR	Medium	Extended
A2	expa02	RCP8.5	CSIRO-Mk3.6.0	Medium	ensemble
A3	expa03	RCP8.5	ACCESS1.3	Medium	(Tier 2)
B1	expb01	SSP5-8.5	CNRM-CM6-1	Medium	CMIP6 extension (Tier 2)
B2	expb02	SSP1-2.6	CNRM-CM6-1	Medium	
B3	expb03	SSP5-8.5	UKESM1-0-LL	Medium	
B4	expb04	SSP5-8.5	CESM2	Medium	
B5	expb05	SSP5-8.5	CNRM-ESM2-1	Medium	

Table 1. Extended ISMIP6-Greenland Tier-1 and 2 future climate experiments discussed in this study. See Nowicki and others (2020) for references for the GCMs.




Fig. 1. Extended ISMIP6-Greenland historical run (hist, 1990–2015), projection control run (ctrl_proj) and Tier-1 and 2 future climate experiments: (a) Simulated ice mass change (counted positively for loss and expressed as sea-level contribution), (b) ice area. The red and blue boxes to the right show the mean ± 1 -sigma ranges for RCP8.5/SSP5-8.5 and RCP2.6/SSP1-2.6, respectively; the whiskers show the corresponding full ranges.




Fig. 2. Main components of the global mass balance for Exp. 5 (MIROC5/RCP8.5): Surface mass balance (SMB, purple), basal mass balance (BMB, blue), calving (yellow) and ice volume change (dV/dt , green). Note the shifted, right axis for the latter. The black and green dashed lines indicate the zero levels for the left and right axis, respectively.




Fig. 3. Ice thickness (panels a, b) and surface velocity (panels c, d) for the initial time (2015; panels a, c) and final time (3001; panels b, d) of Exp. 5 (MIROC5/RCP8.5).

Supplement of
**Mass loss of the Greenland ice sheet until
the year 3000 under a sustained
late-21st-century climate**

Ralf Greve^{1,2}, Christopher Chambers¹

¹Institute of Low Temperature Science, Hokkaido University,
Sapporo, Japan

²Arctic Research Center, Hokkaido University, Sapporo, Japan

Author for correspondence:

Ralf Greve, E-mail: greve@lowtem.hokudai.ac.jp

Tables

#	Scenario	GCM	aST 2091–2100 (°C)	caSMB 2015–2100 (m ice equiv.)
5	RCP8.5	MIROC5	6.2000	–27.385
6	RCP8.5	NorESM1-M	5.0206	–18.387
7	RCP2.6	MIROC5	1.4073	–10.391
8	RCP8.5	HadGEM2-ES	8.3467	–23.132
9	RCP8.5	MIROC5	(same as Exp. #5)	
10	RCP8.5	MIROC5	(same as Exp. #5)	
A1	RCP8.5	IPSL-CM5A-MR	6.3040	–27.454
A2	RCP8.5	CSIRO-Mk3.6.0	5.0795	–12.083
A3	RCP8.5	ACCESS1.3	5.9924	–18.392
B1	SSP5-8.5	CNRM-CM6-1	7.6037	–31.615
B2	SSP1-2.6	CNRM-CM6-1	1.8968	–12.941
B3	SSP5-8.5	UKESM1-0-LL	10.0087	–49.421
B4	SSP5-8.5	CESM2	6.7026	–48.311
B5	SSP5-8.5	CNRM-ESM2-1	7.2940	–30.458

Table S1: Surface temperature anomaly (aST, 2091–2100 mean) and cumulative SMB anomaly (caSMB, 2015–2100), spatially averaged over the present-day Greenland ice sheet, for all future climate experiments of this study. The anomalies are relative to the 1960–1989 means of the reference climatology. See also Table 1 of the main paper.

Note (2022-05-17):

In the original version of this document, the values for “aST 2091–2100” in Table S1 were wrong. This has been corrected here.

#	Scenario	GCM	SLC 2100 (m)	SLC 2300 (m)	SLC 3000 (m)
5	RCP8.5	MIROC5	0.1212	0.5486	1.6305
6	RCP8.5	NorESM1-M	0.0892	0.3697	1.0492
7	RCP2.6	MIROC5	0.0406	0.0803	0.1705
8	RCP8.5	HadGEM2-ES	0.1010	0.4534	1.2424
9	RCP8.5	MIROC5	0.1407	0.5872	1.6810
10	RCP8.5	MIROC5	0.1086	0.5229	1.5992
A1	RCP8.5	IPSL-CM5A-MR	0.1207	0.5292	1.5874
A2	RCP8.5	CSIRO-Mk3.6.0	0.0743	0.2749	0.7207
A3	RCP8.5	ACCESS1.3	0.0934	0.3743	1.0500
B1	SSP5-8.5	CNRM-CM6-1	0.1428	0.7535	2.2911
B2	SSP1-2.6	CNRM-CM6-1	0.0525	0.1663	0.4068
B3	SSP5-8.5	UKESM1-0-LL	0.2086	1.0614	3.0782
B4	SSP5-8.5	CESM2	0.1963	1.1498	3.5473
B5	SSP5-8.5	CNRM-ESM2-1	0.1381	0.6721	2.0060

Table S2: Sea-level contribution (SLC) by 2100, 2300 and 3000 relative to the projection control run (ctrl_proj) for all future climate experiments of this study. See also Table 1 of the main paper.