

## The ‘europium anomaly’ in plants: facts and fiction

Olivier Pourret<sup>1</sup>, Antony van der Ent<sup>2</sup>, Andrew Hursthouse<sup>3</sup>, Dasapta Erwin Irawan<sup>4</sup>, Haiyan Liu<sup>5</sup>, Oliver Wiche<sup>6</sup>

<sup>1</sup>UniLaSalle, AGHYLE, Beauvais, France.

<sup>2</sup>Centre for Mined Land Rehabilitation, Sustainable Minerals Institute, The University of Queensland, Queensland, Australia.

<sup>3</sup>School of Computing, Engineering & Physical Sciences, University of the West of Scotland, Paisley, Scotland PA1 2BE, UK.

<sup>4</sup>Applied Geology Research Group, Faculty of Earth Sciences and Technology, Institut Teknologi Bandung, West Java, Indonesia.

<sup>5</sup>School of Water Resources and Environmental Engineering, East China University of Technology, Nanchang 330013, PR China.

<sup>6</sup>Institute for Biosciences, Biology/Ecology Group, TU Bergakademie Freiberg, Germany.

\*corresponding author: [olivier.pourret@unilasalle.fr](mailto:olivier.pourret@unilasalle.fr)

### **Abstract**

Aims

Rare earth elements (REEs) and normalized REE patterns determined in plant and soil samples represent powerful tools to trace biogeochemical processes during weathering, soil genesis and processes in the rhizosphere, and thus publications reporting rare earth elements and normalized REE patterns in soil systems and plants are rapidly increasing.

## Methods

A normalized REE pattern allows for the recognition of an anomalous concentrations of an individual REE. In the literature anomalies are predominantly reported/focused for/on the redox-sensitive elements cerium (Ce) and europium (Eu) that can shift their oxidation state during interactions with organic and inorganic soil phases and the biological processes affecting their mobility in soil and uptake by plants. Thus positive Eu anomalies in plants are often interpreted as a consequence of reduction of  $\text{Eu}^{3+}$  to  $\text{Eu}^{2+}$  in the rhizosphere followed by a preferential uptake of  $\text{Eu}^{2+}$ .

## Results


Due to an analytical artefact in ICP-MS analysis, a false Eu anomaly may be reported. This can be avoided by using a barium (Ba) interference correction. We draw attention to the possibility of this problem and to being aware of its potential occurrence when Eu anomalies are reported

## Conclusions

We recommend (i) including information on how this potential problem was dealt with in the Materials and Methods section of articles and (ii) how to implement Findable Accessible Interoperable and Reusability (FAIR) guiding principles in that section (including data availability in an open repository).

## Keywords

Artifact; Isobaric Correction; Europium anomaly; Interference; REEs; FAIR guiding principles.



## Introduction

The rare earth elements (REEs) comprise a group of 16 elements from the lanthanide series including lanthanum (La), yttrium (Y) and scandium (Sc) that occur in most all soils at concentrations similar to most plant essential micronutrients, such as copper and zinc, and are therefore of increasing interest in plant and soil research (Tyler 2005; Liang et al. 2008; Liu et al. 2018). For the sake of convenience, the REE distribution in natural samples is usually expressed by normalized REE patterns where concentrations in a target material are referred to concentrations in a reference material (see Davranche et al. 2017). A distinct change in this pattern typically underlies slight differences in the chemical properties of the REEs caused by the decrease in the ionic radius with increasing atomic number, leading to slight changes in the stability constants with soil organic and inorganic soil phases and differences in solution complexation between light to heavy REEs. As a consequence, these differences result in a systematic REE fractionation pattern called the 'lanthanide contraction'. Additionally, a normalized REE pattern allows for the recognition of an anomalous concentrations of an individual REE as either a positive or negative anomaly in an otherwise smooth pattern. This type of anomaly can occur in response to the redox

properties of Eu which can exist in the trivalent or divalent state (Bau 1991). The anomalous behavior of Eu is quantified *via* calculation of Eu anomaly  $(Eu/Eu^*) = 2Eu_N / (Sm_N + Gd_N)$  where  $N$  equals the normalized abundance to a reference material (Figure 1) (Bau 1991; Davranche et al. 2017). REE concentrations in plants, and especially in shoot biomass are reported in an increasing number of publications (Ichihashi et al. 1992; Liang et al. 2008; Brioschi et al. 2013; Censi et al. 2014; Liu et al. 2018; Martinez et al. 2018). However, in some of publications, the analytical techniques employed are not fully documented, which may lead to questions about the validity of the results. In particular, the risk of reporting false positive anomalies in plants caused by barium (Ba) interference with europium (Eu). This Letter highlights this issue, and raises a caution for researchers to avoid this analytical error in their work, recommending improved laboratory documentation to offer a route to reduce the impact of this problem.

### **Analytical techniques**


Environmental samples typically contain concentrations of REEs at trace and ultra-trace levels (Davranche et al. 2017). Particularly in plants, concentrations of single REEs rarely exceed 1 mg/kg for the light REEs (La, Nd, Ce) and are frequently substantially lower than 1 µg/kg, especially in case of the very low abundance heavy REEs (e.g. Dy, Ho, Lu) following the Oddo-Harkins rule (Tyler 2005). Since the 1990's, progress in the determination of REE in environmental samples has moved hand in hand with the development of advanced instrumental analytical techniques. Among the most commonly utilized approaches are neutron activation analysis (NAA), X-ray fluorescence (XRF) spectrometry, Inductively Coupled Plasma Optical Emission Spectrometry (ICP-AES) and Inductively Coupled Plasma Mass Spectrometry (ICP-MS). Due to its high selectivity and sensitivity, ICP-MS is currently the most frequently used method, characterized by its high sensitivity and the selectivity needed for the direct determination of all REEs in complex sample matrices with supreme limits of detection (routinely parts per billion *i.e.*, µg/kg). Apart from hyphenated methods,

coupling ICP-MS with laser ablation (LA), standard ICP-MS devices typically require aqueous samples, and thus, solid environmental samples require a sample pretreatment where REEs are transferred into solution. Where total concentrations of solids are required, sample pretreatment/digestion methods for the dissolution of REE-bearing geogenic or biogenic silicates are achieved by fusion with borates or carbonates at high temperatures followed by a dissolution in acid or through microwave-assisted heating of powerful acid mixtures (perchloric acid, hydrochloric acid, nitric acid, hydrofluoric acid), with hydrofluoric (HF) acid used for the dissolution of sparingly soluble silicates. If HF is required, the amount of HF should be adjusted to the amount of Si in the samples because an excess of F in the resulting solutions may precipitate REEs together with Ca. To overcome this problem, boric acid or digestion with  $\text{HBF}_4$  can improve REE recovery by preventing the precipitation of Ca and REE fluorides (Krachler et al. 2002). All other procedures such as aqua regia or single acid microwave digestion without hydrofluoric acid the REE concentrations in environmental or geochemical materials, including plant material, will be significantly underestimated and just give pseudo-totals of the desired elements. Additionally, it is worth noting that closed vessel digestions should be used to avoid losses of volatile element species.

The resulting solutions are not typically problematic for analysis by ICP-MS. The solution stability of REEs is good with  $\text{HNO}_3$  or HCl, but fluoride, phosphate and sulfate salts and carbonate are particularly insoluble, and a neutral/alkaline pH is also an issue. The typical approach for ICP-MS is to separate from the matrix using ion exchange resins and sequential elution (Yan et al. 2018). It adds complexity but improves detection and has been used for precise isotope ratio determinations (Zhao et al. 2019). Other methods for the quantification of mobile and potentially plant available fractions of REEs in soils deploy single step or sequential extractions (Wiche and Heilmeyer 2016; Wiche et al. 2017). For ICP-MS, however, the determination of REEs requires solutions with low concentrations of REEs (of the order of  $<10$  mg/L) in order to measure all 16 naturally occurring REEs simultaneously without any separation from the matrix. Ions produced in high temperature plasma are identified based on their mass to charge ratio,  $m/z$ , which is characteristic for any given

isotope. Each REE has at least one isotope that is free from isobaric overlaps and their sensitivity is relatively uniform from  $^{139}\text{La}$  to  $^{175}\text{Lu}$ .

In addition to the time consuming and sometimes challenging sample pretreatment, a second more latent problem may hamper the reliable quantification of REEs by ICP-MS due to the occurrence of spectral interferences in the mass spectrum. In particular, the most important challenges with interferences encountered in ICP-MS analysis are caused by the formation oxides in the plasma (Longerich et al. 1987; Jarvis et al. 1992). The REEs form a continuous group from 139 to 175 m/z, and the formation of light REEs and barium oxide can produce significant middle REE interferences. Refractory oxide ions are influenced by the plasma operating parameters. The oxide production level is close to 1% to 2% (Davranche et al. 2004; Davranche et al. 2017). Mathematical corrections are required to suppress these specific interferences and these corrections are calculated as a function of the oxide level. Previous studies have shown the importance of these corrections for accurate REE quantification (Dulski 1994; Aries et al. 2000; Raut et al. 2005a; Raut et al. 2005b). In the case of Eu, Ba which forms the  $\text{BaO}^+$  species and the isotopes  $^{135}\text{Ba}$  ( $135+16=151$ ) and  $^{137}\text{Ba}$  ( $137+16=153$ ) (Jarvis et al. 1989; Smirnova et al. 2006; Zhao et al. 2019) and  $\text{BaOH}$  species and the isotope  $^{134}\text{Ba}$  ( $134+17=151$ ) and  $^{136}\text{Ba}$  ( $136+17=153$ ) (Möller et al. 1992) isobarically interferes with  $^{151}\text{Eu}$  and  $^{153}\text{Eu}$ . Eventually, progress has been made with introduction of collision cell techniques that allow to remove interfering ions through ion/neutral reactions (Yip and Sham 2007) and higher resolution multiple quadrupole systems which can successfully remove the influence of oxide interferences (Zhu et al. 2021). To increase the precision of REE measurements, a sample can be enriched with a pre-concentrated multi-spike REE solution, and the matrix can be removed by using ion chromatography (Baker et al. 2002; Rousseau et al. 2013). In addition, a suitable reference material should be systematically analysed in order to assess the effect of isobaric interferences on the different REE masses.



**Figure 1** Example of Upper continental crust (UCC)-normalized REE patterns in *Anisopappus chinensis* leaves sampled at Mikembo (Democratic Republic of Congo), (a) without Ba interference correction (modified after (Pourret et al. 2014)) and (b) with Ba interference correction (modified after (Pourret et al. 2019)). The UCC values are from McLennan (2001).

### Specific behavior of europium during absorption by plants

At low to very low Ca concentrations in the soil, REE concentrations have been described to decrease simultaneously with the appearance of a strong Eu anomaly in the biomass of plants growing on such soils (Stille et al. 2006). The shoot biomass is then characterized by positive Eu anomalies, as has been reported by Ding et al. (2006) for soil-grown wheat and Brioschi et al. (2013) for plants growing on granite-derived soils. Ding et al. (2006) proposed that such Eu anomalies were caused by precipitation of Eu-enriched phosphate particles in the different plant organs. Indeed, the possible deposition of authigenic phosphates in plant roots may occur. Due to the strong REE affinity towards phosphates, this process is an opportunity for REE fractionation at the plant-soil interface and can modify the features of normalised REE patterns in plants (Romero-Freire et al. 2021). Subsequently Stille et al. (2006) and Brioschi et al. (2013) hypothesized that this anomaly was a linked to the strong depletion of Ca in the soils where plants may suffer from Ca-deficiency (Brioschi et al. 2013). Gao et al. (2003) have shown that Eu can be absorbed and bound to the inner membranes

of different plant cell organelles. Even if transport proteins that can mediate REE uptake by plant cells have not yet been fully characterized (Wehrmann et al. 2019; Cheng et al. 2021) and REE may inhibit cation channels (White 2000), most studies to date demonstrated that ion channels, especially  $\text{Ca}^{2+}$ ,  $\text{K}^+$  and  $\text{Na}^+$  mediate REE diffusion and uptake in plant root cells (Han et al. 2005) which also explains the preferential accumulation of light REEs (e.g., La to Eu + Sc) (LREE) in plants due to their similar ionic radii and the physiochemical properties of LREEs and  $\text{Ca}^{2+}$  (Brioschi et al. 2013; Censi et al. 2014; Censi et al. 2017; Martinez et al. 2018). Additionally, there is evidence that during root–shoot translocation heavy REEs are preferably transported to the shoots which has been attributed to the higher stability of heavy REE complexes formed with organic substances, especially amino and organic acids in the xylem (Yuan et al. 2017; Yuan et al. 2018). Overall, this suggests that most plants discriminate between cations differing by their ionic radius and charge during uptake and transport within plant tissues. The effects of these processes can be usually observed in fractionation pattern of concentrations of LREEs relative to HREEs depending on plant species, growing conditions and plant tissue (Yuan et al. 2017; Yuan et al. 2018). Calcium and Eu have almost identical ionic radii (Shannon 1976), which means that Eu is likely to competitively replace Ca ions in biological systems (Evans 1983). Europium may be transported apoplastically across the root to the xylem like Ca (White 2001). Although, a biological role for lanthanides is not known, previous studies have shown that Eu can substitute for Ca in plants and/or replace Ca in cation binding sites due to similar ionic radii, in particular in soils with elevated Eu/Ca ratios (Zeng et al. 2003; Shtangeeva and Ayrault 2007). Moreover, positive Eu anomalies in plant shoots suggest that  $\text{Eu}^{3+}$  can form stable organic complexes thereby replacing  $\text{Ca}^{2+}$  in some biological processes, such as in xylem fluid during normal nutrient fluxes (Zeng et al. 2003). The possibility that Eu mobility in these fluids can be enhanced by its reductive speciation to  $\text{Eu}^{2+}$  cannot be ruled out. In rice, a positive Eu anomaly has been found in the aerial plant parts, whereas HREE enrichment (e.g., Gd to Lu + Y) has been observed in grains. A positive Eu anomaly has also been found in rice roots suggesting Eu accumulation in plant nutrition fluxes (Liang et al. 2008).

However, these results were not found in hydroponic studies performed by Martinez et al. (2018) on REE accumulation in rice plants. Whether or not Eu may be a Ca physiological substitute remains unknown.

Studies using biological enrichment indices have shown that the different enrichment patterns of REE may occur in different plant species and because of different growth environments. Fractionation of REEs with both variable valences (*i.e.*, Ce and Eu) and stable valences (*i.e.*, La and Lu) can be easily observed in plants owing to the combined effects of physical, chemical, and biological factors. REEs have a dual effect on the physiological and biochemical reactions in the plant development, Zeng et al. (2003) showed that  $\text{Eu}^{3+}$  could replace  $\text{Ca}^{2+}$  in plants (e.g. *Amaranthus caudatus*) and promote Ca transport across plasma membrane. Therefore, possibly some Ca, which plays a key role in plant cellular walls and membranes, might have been replaced by  $\text{Eu}^{3+}$ . The concentration of free and active  $\text{Ca}^{2+}$  in the cytoplasm of plant cell must remain low and may be controlled by the precipitation of oxalate crystals which allows the storage of biologically inactive Ca and probably also of Eu in older plant organs. It must be noted that not all plants precipitate Ca-oxalate. Cytosolic  $\text{Ca}^{2+}$  concentration is also regulated by  $\text{Ca}^{2+}$  fluxes across the plasma membrane and across the membranes of organelles, particularly the vacuole, where Ca is stored (White 2015).

### **A false europium positive anomaly?**

The presence of a major positive Eu anomaly, as shown in Figure 1a, is highly suspicious and is caused by an analytical artifact due to insufficient correction for the BaO and BaOH interferences (that may be avoided if the analysis uses collision cell technology). The analytical quality of the literature that reports positive Eu anomalies in plant material samples should be scrutinized, especially if information on any REE pre-concentration method; previously shown to have a significant positive impact on isobaric interferences (Zhu and Itoh 2021); is not provided in the Materials and Methods. It raises the issue that we cannot rule out Ba oxide interference. As we have emphasized, Eu analysis in low-REE samples suffers

from important interferences of BaO and BaOH on the Eu signal, often leading to false positives in normalized patterns. Barium is ubiquitous in soils and all plants contain a small quantity of it usually in concentrations around 4 to 50 µg/kg dry weight of plants (Chaudhry et al. 1977). If Ba concentrations are not reported, it is impossible to know whether this interference may have occurred. The authors should make sure that the Eu-signal is real (rule of thumb: Ba/Eu<1000), and that the reported Eu values are not a false positive (Shabani et al. 1992). The Inductively Coupled Plasma Atomic Emission Spectrometry (ICP-AES) method has gained in sensitivity over recent years (routinely low parts per million *i.e.*, µg/g), and does not suffer from the same problematic Eu interference of ICPMS, although other spectral interferences may still occur, it offers the opportunity to cross validate detection issues (Amaral et al. 2017; He et al. 2017). Such detail procedural descriptions are not usually included in the methods section of many reports.

### **Concluding remarks**

In summary, the Eu anomaly may be reported because of an analytical artifact in ICP-MS analysis. This error can be avoided by using isobaric corrections considering Ba and alternative detection strategies are increasingly available. Not all studies using ICP-MS for REE analysis of soil and plant material samples provide comprehensive information on how this error was avoided or dealt with. We draw attention to the possibility of this problem and recommend being cautious, recognizing its potential occurrence and to take appropriate measures during analysis. The use of material reference certified for REE (Gills 1995; Samczyński et al. 2012) may be part of the solution and making analytical data available considering Findable Accessible Interoperable and Reusability (FAIR) principles (Pourret et al. 2020) another part. Further, we also propose that FAIR principles should be used to document laboratory methods. Detailed procedures, including the failures in experiment, usually are not described sufficiently in common journal articles, due to the editorial guidelines. The usage of electronic laboratory notebook could be an option and data

availability statements are now commonplace in publication guidelines. The successes and failures would contribute to the development of a knowledge base for REE tests in plants, including those related to Eu.

## Declarations

**Funding** This research did not receive any specific grant from funding agencies in the public, commercial or non-profit sectors.

**Conflicts of interest/Competing interests** No potential conflict of interest was reported by the authors.

**Availability of data and material** Not applicable.

**Code availability** Not applicable.

**Authors' contributions** all the authors contributed to writing the article.

## References

- Amaral CDB, Barros JAVA, Virgilio A, Nóbrega JA, Nogueira ARA, Schiavo D, Machado RC (2017) Determination of rare earth elements in geological and agricultural samples by ICP-OES. *Spectroscopy (Santa Monica)* 32: 32-36.
- Aries S, Valladon M, Polvé M, Dupré B (2000) A routine method for oxide and hydroxide interference corrections in ICP-MS chemical analysis of environmental and geological samples. *Geostandard Newslet* 24: 19-31. <https://doi.org/10.1111/j.1751-908X.2000.tb00583.x>
- Baker J, Waight T, Ulfbeck D (2002) Rapid and highly reproducible analysis of rare earth elements by multiple collector inductively coupled plasma mass spectrometry. *Geochim Cosmochim Acta* 66: 3635-3646. [https://doi.org/10.1016/S0016-7037\(02\)00921-3](https://doi.org/10.1016/S0016-7037(02)00921-3)
- Bau M (1991) Rare-earth element mobility during hydrothermal and metamorphic fluid-rock interaction and the significance of the oxidation state of europium. *Chem Geol* 93: 219-230. [https://doi.org/10.1016/0009-2541\(91\)90115-8](https://doi.org/10.1016/0009-2541(91)90115-8)
- Brioschi L, Steinmann M, Lucot E, Pierret MC, Stille P, Prunier J, Badot PM (2013) Transfer of rare earth elements (REE) from natural soil to plant systems: Implications for the environmental availability of anthropogenic REE. *Plant Soil* 366: 143-163. <https://doi.org/10.1007/s11104-012-1407-0>
- Censi P, Cibella F, Falcone EE, Cuttitta G, Saiano F, Inguaggiato C, Latteo V (2017) Rare earths and trace elements contents in leaves: A new indicator of the composition of

- atmospheric dust. *Chemosphere* 169: 342-350.  
<https://doi.org/10.1016/j.chemosphere.2016.11.085>
- Censi P, Saiano F, Pisciotta A, Tuzzolino N (2014) Geochemical behaviour of rare earths in *Vitis vinifera* grafted onto different rootstocks and growing on several soils. *Sci Total Environ* 473–474: 597-608. <http://dx.doi.org/10.1016/j.scitotenv.2013.12.073>
- Chaudhry FM, Wallace A, Mueller RT (1977) Barium toxicity in plants. *Commun Soil Sci Plan* 8: 795-797. <https://doi.org/10.1080/00103627709366776>
- Cheng M, Wang L, Zhou Q, Chao D, Nagawa S, He D, Zhang J, Li H, Tan L, Gu Z, et al. (2021) Lanthanum(III) triggers AtrbohD- and jasmonic acid-dependent systemic endocytosis in plants. *Nat Commun* 12: 4327. <https://doi.org/10.1038/s41467-021-24379-z>
- Davranche M, Gruau G, Dia A, Le Coz-Bouhnik M, Marsac R, Pédrot M, Pourret O (2017) Chapter 7. Rare Earth Elements in Wetlands. In: Rinklebe J., Knox AS, Paller M eds. *Trace Elements in Waterlogged Soils and Sediments*: Taylor & Francis Group/CRC Press, 135-162. <https://doi.org/10.1201/9781315372952-9>
- Davranche M, Pourret O, Gruau G, Dia A (2004) Impact of humate complexation on the adsorption of REE onto Fe oxyhydroxide. *J Colloid Interface Sci* 277: 271-279. <https://doi.org/10.1016/j.jcis.2004.04.007>
- Ding S, Liang T, Zhang C, Huang Z, Xie Y, Chen T (2006) Fractionation mechanisms of rare earth elements (REEs) in hydroponic wheat: An application for metal accumulation by plants. *Environ Sci Technol* 40: 2686-2691. <https://doi.org/10.1021/es052091b>
- Dulski P (1994) Interferences of oxide, hydroxide and chloride analyte species in the determination of rare earth elements in geological samples by inductively coupled plasma-mass spectrometry. *Fresen J Anal Chem* 350: 194-203. <https://doi.org/10.1007/bf00322470>
- Evans JR (1983) Nitrogen and Photosynthesis in the Flag Leaf of Wheat (*Triticum aestivum* L.). *Plant Physiol* 72: 297-302. <https://doi.org/10.1104/pp.72.2.297>
- Gao Y, Zeng F, Yi A, Ping S, Jing L (2003) Research of the entry of rare earth elements Eu<sup>3+</sup> and La<sup>3+</sup> into plant cell. *Biol Trace Elem Res* 91: 253-265. <https://doi.org/10.1385/BTER:91:3:253>
- Gills TE (1995) Standard reference material 1537a-tomatoe leaves-certificate of analysis: 5.
- Han F, Shan X-Q, Zhang J, Xie Y-N, Pei Z-G, Zhang S-Z, Zhu Y-G, Wen B (2005) Organic acids promote the uptake of lanthanum by barley roots. *New Phytol* 165: 481-492. <https://doi.org/10.1111/j.1469-8137.2004.01256.x>
- He M, Hu B, Chen B, Jiang Z (2017) Inductively Coupled Plasma Optical Emission Spectrometry for Rare Earth Elements Analysis. *Physical Sciences Reviews* 2. <https://doi.org/10.1515/psr-2016-0059>
- Ichihashi H, Morita H, Tatsukawa R (1992) Rare earth elements (REEs) in naturally grown plants in relation to their variation in soils. *Environ Pollut* 76: 157-162. [http://dx.doi.org/10.1016/0269-7491\(92\)90103-H](http://dx.doi.org/10.1016/0269-7491(92)90103-H)
- Jarvis KE, Gray AL, Houk RS (1992) *Handbook of Inductively Coupled Plasma Mass Spectrometry*. Springer.
- Jarvis KE, Gray AL, McCurdy E (1989) Avoidance of spectral interference on europium in inductively coupled plasma mass spectrometry by sensitive measurement of the doubly charged ion. *J Anal At Spectrom* 4: 743-747. <https://doi.org/10.1039/JA9890400743>
- Krachler M, Mohl C, Emons H, Shotyk W (2002) Analytical procedures for the determination of selected trace elements in peat and plant samples by inductively coupled plasma mass spectrometry. *Spectrochim Acta B* 57: 1277-1289. [https://doi.org/10.1016/S0584-8547\(02\)00068-X](https://doi.org/10.1016/S0584-8547(02)00068-X)
- Liang T, Ding S, Song W, Chong Z, Zhang C, Li H (2008) A review of fractionations of rare earth elements in plants. *J Rare Earth* 26: 7-15. [https://doi.org/10.1016/S1002-0721\(08\)60027-7](https://doi.org/10.1016/S1002-0721(08)60027-7)
- Liu C, Yuan M, Liu W-S, Guo M-N, Huot H, Tang Y-T, Laubie B, Simonnot M-O, Morel JL, Qiu R-L (2018) Element Case Studies: Rare Earth Elements. In: Van der Ent A, Echevarria G, Baker AJM, Morel JL eds. *Agromining: Farming for Metals: Extracting*

- Unconventional Resources Using Plants*. Cham: Springer International Publishing, 297-308. [https://doi.org/10.1007/978-3-319-61899-9\\_19](https://doi.org/10.1007/978-3-319-61899-9_19)
- Longerich HP, Fryer BJ, Strong DF, Kantipuly CJ (1987) Effects of operating conditions on the determination of the rare earth elements by inductively coupled plasma-mass spectrometry (ICP-MS). *Spectrochim Acta B* 42: 75-92. [https://doi.org/10.1016/0584-8547\(87\)80051-4](https://doi.org/10.1016/0584-8547(87)80051-4)
- Martinez RE, Pourret O, Faucon MP, Dian C (2018) Effect of rare earth elements on rice plant growth. *Chem Geol* 489: 28-37. <https://doi.org/10.1016/j.chemgeo.2018.05.012>
- McLennan SM (2001) Relationships between the trace element composition of sedimentary rocks and upper continental crust. *Geochem Geophys Geosy* 2: 109. <https://doi.org/10.1029/2000gc000109>
- Möller P, Dulski P, Luck J (1992) Determination of rare earth elements in seawater by inductively coupled plasma-mass spectrometry. *Spectrochim Acta B* 47: 1379-1387. [https://doi.org/10.1016/0584-8547\(92\)80128-4](https://doi.org/10.1016/0584-8547(92)80128-4)
- Pourret O, Bollinger J-C, van Hullebusch ED (2020) On the difficulties of being rigorous in environmental geochemistry studies: some recommendations for designing an impactful paper. *Environ Sci Pollut Res* 27: 1267-1275. <https://doi.org/10.1007/s11356-019-06835-y>
- Pourret O, Lange B, Jitaru P, Mahy G, Faucon MP (2014) Transfer of rare earth elements from natural metalliferous (copper and cobalt rich) soils into plant shoot biomass of metallophytes from Katanga (Democratic Republic of Congo). *Geophys Res Abstr* 16: EGU2014-6272.
- Pourret O, Lange B, Martinez RE, Wiche O, Faucon M-P (2019) Relationships between soil chemical properties and rare earth element concentrations in the aboveground biomass of a tropical herbaceous plant. Preprint EarthArXiv. <https://doi.org/10.31223/osf.io/w4hqs>
- Raut NM, Huang L-S, Aggarwal SK, Lin K-C (2005a) Mathematical Correction for Polyatomic Isobaric Spectral Interferences in Determination of Lanthanides by Inductively Coupled Plasma Mass Spectrometry. *J Chin Chem Soc* 52: 589-597.
- Raut NM, Huang L-S, Lin K-C, Aggarwal SK (2005b) Uncertainty propagation through correction methodology for the determination of rare earth elements by quadrupole based inductively coupled plasma mass spectrometry. *Anal Chim Acta* 530: 91-103. <https://doi.org/10.1016/j.aca.2004.08.067>
- Romero-Freire A, González V, Groenenberg JE, Qiu H, Auffan M, Cotelle S, Giamberini L (2021) Cytotoxicity and genotoxicity of lanthanides for *Vicia faba* L. are mediated by their chemical speciation in different exposure media. *Sci Total Environ* 790: 148223. <https://doi.org/10.1016/j.scitotenv.2021.148223>
- Rousseau TCC, Sonke JE, Chmeleff J, Candaudap F, Lacan F, Boaventura G, Seyler P, Jeandel C (2013) Rare earth element analysis in natural waters by multiple isotope dilution - sector field ICP-MS. *J Anal At Spectrom* 28: 573-584. <http://doi.org/10.1039/C3JA30332B>
- Samczyński Z, Dybczyński RS, Polkowska-Motrenko H, Chajduk E, Pyszynska M, Danko B, Czerska E, Kulisa K, Doner K, Kalbarczyk P (2012) Two New Reference Materials Based on Tobacco Leaves: Certification for over a Dozen of Toxic and Essential Elements. *The Scientific World Journal* 2012: 216380. <https://doi.org/10.1100/2012/216380>
- Shabani MB, Akagi T, Masuda A (1992) Preconcentration of trace rare-earth elements in seawater by complexation with bis(2-ethylhexyl) hydrogen phosphate and 2-ethylhexyl dihydrogen phosphate adsorbed on a C18 cartridge and determination by inductively coupled plasma mass spectrometry. *Anal Chem* 64: 737-743. <https://doi.org/10.1021/ac00031a008>
- Shannon RD (1976) Revised effective ionic radii and systematic studies of interatomic distances in halides and chalcogenides. *Acta Crystallogr A* 32: 751-767. <https://doi.org/10.1107/S0567739476001551>

- Shtangeeva I, Ayrault S (2007) Effects of Eu and Ca on yield and mineral nutrition of wheat (*Triticum aestivum*) seedlings. *Environ Exp Bot* 59: 49-58. <https://doi.org/10.1016/j.envexpbot.2005.10.011>
- Smirnova EV, Mysovskaya IN, Lozhkin VI, Sandimirova GP, Pakhomova NN, Smagunova AA (2006) Spectral interferences from polyatomic barium ions in inductively coupled plasma mass spectrometry. *J Appl Spectrosc* 73: 911-917. <https://doi.org/10.1007/s10812-006-0175-0>
- Stille P, Steinmann M, Pierret MC, Gauthier-Lafaye F, Chabaux F, Viville D, Pourcelot L, Matera V, Aouad G, Aubert D (2006) The impact of vegetation on REE fractionation in stream waters of a small forested catchment (the Strengbach case). *Geochim Cosmochim Acta* 70: 3217-3230. <https://doi.org/10.1016/j.gca.2006.04.028>
- Tyler G (2005) Rare earth elements in soil and plant systems - A review. *Plant Soil* 267: 191-206. <https://doi.org/10.1007/s11104-005-4888-2>
- Wehrmann M, Berthelot C, Billard P, Klebensberger J (2019) Rare Earth Element (REE)-Dependent Growth of *Pseudomonas putida* KT2440 Relies on the ABC-Transporter PedA1A2BC and Is Influenced by Iron Availability. *Front Microbiol* 10. <https://doi.org/10.3389/fmicb.2019.02494>
- White PJ (2000) Calcium channels in higher plants. *BBA-Biomembranes* 1465: 171-189. [https://doi.org/10.1016/S0005-2736\(00\)00137-1](https://doi.org/10.1016/S0005-2736(00)00137-1)
- White PJ (2001) The pathways of calcium movement to the xylem. *J Exp Bot* 52: 891-899. <https://doi.org/10.1093/jexbot/52.358.891>
- White PJ (2015) Calcium. *Handbook of Plant Nutrition*: CRC Press. <https://doi.org/10.1201/b18458-8>
- Wiche O, Heilmeyer H (2016) Germanium (Ge) and rare earth element (REE) accumulation in selected energy crops cultivated on two different soils. *Miner Eng* 92: 208-215. <http://dx.doi.org/10.1016/j.mineng.2016.03.023>
- Wiche O, Zertani V, Hentschel W, Achtziger R, Midula P (2017) Germanium and rare earth elements in topsoil and soil-grown plants on different land use types in the mining area of Freiberg (Germany). *J Geochem Explor* 175: 120-129. <http://dx.doi.org/10.1016/j.gexplo.2017.01.008>
- Yan X, Dai S, Graham IT, He X, Shan K, Liu X (2018) Determination of Eu concentrations in coal, fly ash and sedimentary rocks using a cation exchange resin and inductively coupled plasma mass spectrometry (ICP-MS). *Int J Coal Geol* 191: 152-156. <https://doi.org/10.1016/j.coal.2018.03.009>
- Yip Y-c, Sham W-c (2007) Applications of collision/reaction-cell technology in isotope dilution mass spectrometry. *TrAC Trends in Analytical Chemistry* 26: 727-743. <https://doi.org/10.1016/j.trac.2007.03.007>
- Yuan M, Guo M-N, Liu W-S, Liu C, van der Ent A, Morel JL, Huot H, Zhao W-Y, Wei X-G, Qiu R-L, et al. (2017) The accumulation and fractionation of Rare Earth Elements in hydroponically grown *Phytolacca americana* L. *Plant Soil* 421: 67-82. <https://doi.org/10.1007/s11104-017-3426-3>
- Yuan M, Liu C, Liu W-S, Guo M-N, Morel JL, Huot H, Yu H-J, Tang Y-T, Qiu R-L (2018) Accumulation and fractionation of rare earth elements (REEs) in the naturally grown *Phytolacca americana* L. in southern China. *Int J Phytoremediat* 20: 415-423. <https://doi.org/10.1080/15226514.2017.1365336>
- Zeng F, Tian H, Wang Z, An Y, Gao F, Zhang L, Li F, Shan L (2003) Effect of rare earth element europium on amaranthin synthesis in *Amaranthus caudatus* seedlings. *Biol Trace Elem Res* 93: 271-282. <https://doi.org/10.1385/bter:93:1-3:271>
- Zhao W, Zong K, Liu Y, Hu Z, Chen H, Li M (2019) An Effective Oxide Interference Correction on Sc and REE for Routine Analyses of Geological Samples by Inductively Coupled Plasma-Mass Spectrometry. *J Earth Sci* 30: 1302-1310. <https://doi.org/10.1007/s12583-019-0898-5>
- Zhu Y, Itoh A (2021) Pseudo isotope dilution (PID) as an approach for correcting barium-related spectral interferences on the measurement of europium by inductively coupled

- plasma mass spectrometry (ICP-MS). *Anal Chim Acta* 1180: 338854. <https://doi.org/10.1016/j.aca.2021.338854>
- Zhu Y, Nakano K, Shikamori Y, Itoh A (2021) Direct determination of rare earth elements in natural water samples by inductively coupled plasma tandem quadrupole mass spectrometry with oxygen as the reaction gas for separating spectral interferences. *Spectrochim Acta B* 179: 106100. <https://doi.org/10.1016/j.sab.2021.106100>