
Disclaimer

This manuscript has been submitted to NATURE COMMUNICATIONS EARTH & ENVIRONMENT, and is not peer-reviewed.

Please feel free to contact any of the authors with feedback and suggestions for improvements.

Document history

Date	Action
03/Aug/2022	MS sent to co-authors for final draft acceptance
	Supplementary materials uploaded to Zenodo
	MS Submitted to EarthArXiv
	MS Submitted to Communications Earth and Environment

THE INFLUENCE OF REEF ISOSTASY, DYNAMIC TOPOGRAPHY, AND GLACIAL ISOSTATIC ADJUSTMENT ON THE LAST INTERGLACIAL SEA-LEVEL RECORD OF NORTHEASTERN AUSTRALIA.

PREPRINT, COMPILED AUGUST 3, 2022

Alessio Rovere^{1,2*}, Tamara Pico³, Frederick Richards⁴, Michael J. O’Leary⁵, Jerry X. Mitrovica⁶, Ian D. Goodwin^{7,8},
Jacqueline Austermann⁹, and Konstantin Latychev⁶

¹Department of Environmental Sciences, Informatics and Statistics, Ca’ Foscari University of Venice, IT

²MARUM - Center for Marine Environmental Sciences, University of Bremen, DE Germany

³Earth & Planetary Sciences Department, UC Santa Cruz, Santa Cruz, USA

⁴Department of Earth Science & Engineering, Imperial College London, London, UK

⁵School of Earth Sciences, University of Western Australia Oceans Institute, Perth, AU

⁶Department of Earth and Planetary Sciences, Harvard University, Boston, USA

⁷Climalab, New South Wales, AU

⁸Climate Change Research Centre and Australian Centre for Excellence in Antarctic Science, University of New South Wales, AU

⁹Department of Earth and Environmental Sciences & Lamont-Doherty Earth Observatory, Columbia University, New York, USA

ABSTRACT

NOTE FOR THE READER: this is a preprint of a manuscript submitted to Nature Communications Earth & Environment. This manuscript has not been peer-reviewed.

Understanding sea level during the warmest peak of the Last Interglacial (125,000 yrs ago; Marine Isotope Stage 5e) is important for assessing future ice-sheet dynamics in response to climate change, and relies on the measurement and interpretation of paleo sea-level indicators, corrected for post-depositional vertical land motions. The coasts and continental shelves of northeastern Australia (Queensland) preserve an extensive Last Interglacial record in the facies of coastal strandplains onland and fossil reefs offshore. However, there is a discrepancy (amounting to tens of meters) in the elevation of sea-level indicators between offshore and onshore sites. Here, we assess the influence of geophysical processes that may have changed the elevation of these sea-level indicators since the Last Interglacial. We modeled sea-level change due to: i) dynamic topography; ii) glacial isostatic adjustment, and iii) isostatic adjustment due to coral reef loading, which we term "reef isostasy". These processes caused relative sea-level changes on the order of, respectively, 10 m, 5 m, and 0.3 m since the Last Interglacial. Of these geophysical processes, the dynamic topography predictions most closely match the tilting observed between onshore and offshore sea-level markers. However, we found that these combined geophysical processes cannot explain the full amplitude of the observed discrepancy between onshore and offshore sea-level indicators.

Keywords Last Interglacial · Sea level changes · NE Australia · Great Barrier Reef

1 INTRODUCTION

2 Reef coring typically encountered LIG reefs between 5 and
3 20 m below the modern GBR reef flats. Strikingly, along the
4 Queensland and far northern New South Wales coastline, LIG
5 strandplains are identified at higher elevations with ridge/swale
6 heights (ranging from +3 to +9m) than offshore LIG reefs [60,
7 33]. These onshore markers are not as precisely dated as the
8 coral sea-level markers, however they were arguably formed
9 during the LIG. The higher elevations of these coastal strandplains
10 are roughly consistent with estimates for peak LIG global mean
11 sea level (GMSL). Such estimates are consistently above modern
12 mean sea level (0 m), albeit they vary substantially depending
13 on study sites analyzed and corrections for vertical land motions
14 applied to the proxy record (from 6 to 9 m 43, 8 m 23, 2 m 78,
15 and 1-5 m 24).

16 The most obvious explanation of the discrepancy between on-
17 shore and offshore LIG sea-level indicators in Northeastern
18 Australia is that these two areas are subject to differential ver-
19 tical land motions. When reconstructing past global mean sea

level (GMSL) from geological sea-level proxies, it is essential to
disentangle the components causing globally averaged sea-level
changes from other regional processes that may have caused ver-
tical displacement of past sea-level indicators [65, 69]. Among
these, the most relevant are glacial isostatic adjustment (GIA)
[25], tectonic deformation processes [54] and mantle dynamic
topography (DT) [5].

Crustal loading due to local processes can also cause the vertical
displacement of observed sea-level indicators through isostatic
adjustment. For example, sediment loading can cause regional
sea level to depart significantly from the global mean along ma-
jor deltaic systems [18, 64, 27, 64, 71, 26, 88]. Karst erosion is
another mechanism that induces isostatic adjustment, through
mass unloading, causing a net crustal uplift. This process is rep-
resented in the Plio-Pleistocene shoreline complexes in Florida,
that were uplifted following isostatic response to the karstifica-
tion (leading to rock mass loss) of the landscape [16, 61, 1, 89].
To date, estimates of peak LIG GMSL from tropical areas have
not accounted for the isostatic response to coral reef loading

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38




Figure 1: Map (left panel) and elevation plot (right panel) of LIG paleo RSL obtained from fossil reefs (blue markers) and beach barriers (yellow markers) along the GBR and the Queensland Coasts. Error bars represent 1-sigma ranges.

39 over the last glacial cycle. This process stems from the fact that
 40 corals can grow into spatially extensive reefs, reaching thick-
 41 nesses of several tens of meters during interglacials. The effect
 42 of reef accretion and related loading on local sea-level histories
 43 remains largely unexplored.

44 In this work, we model the influence of geophysical processes
 45 that may have changed the elevation of geologic sea-level indi-
 46 cators along the Queensland coasts and offshore, on the GBR,
 47 since the LIG. We assess the extent to which the combined geo-
 48 physical processes of glacial isostatic adjustment and dynamic
 49 topography may have impacted the LIG sea-level record in this
 50 region. Importantly, in this study we also isolate the process of
 51 coral reef loading, and assess its importance in causing regional
 52 departures from GMSL. While we find that the combined geo-
 53 physical processes modeled in this study cannot fully explain
 54 the amplitude of the observed discrepancy between onshore
 55 and offshore sea-level markers in the study area, we identify

that dynamic topography may represent the key to solve this
 conundrum.

1 LIG SEA-LEVEL INDICATORS

The study of past sea-level changes relies on the measurement
 and dating of relative sea-level (RSL) indicators, i.e. geological
 proxies that formed in connection with former positions of the
 sea. Once a sea-level indicator is measured and dated, it is
 necessary to establish its indicative meaning [84, 76] to quantify
 the relationship between the elevation or depth of an indicator
 and the position of the former sea level, including associated
 uncertainties due to the environmental range of formation. Once
 the elevation of a sea-level indicator is corrected, taking into
 account its indicative meaning, it reflects paleo relative sea level
 (RSL), i.e., the paleo position of the sea including both barystatic
 (i.e., eustatic, 35) changes and elevation changes due to vertical
 land motions of different origin.




Figure 2: Digital Elevation Model [31] and topographic profile (a-b) of the Wyvuri Embayment, where [30] identified LIG coastal sediments in a core under a dune/beach barrier. The star indicates the approximate point where core JW4 of [30] was drilled. The blue dot indicates the inner part of the LIG barrier, that was used as a sea-level proxy in this study. The blue transparent overlay indicates the paleo RSL calculated using the elevation of the inner margin of the barrier and the indicative meaning calculator tool [52].

ogy for these strandplains is in progress [33], and shows that complete LIG strandplains are located inboard of the modern Holocene equivalents. In far north Queensland, Gagan et al. [30] describes a LIG dune/beach barrier located onshore with respect to the Holocene equivalent at Wyvuri Embayment (Figure 2). The top of the barrier, composed of aeolian sediments, is located at +6 m above modern sea-level, while the beach barrier sands were intercepted about 4 meters below the surface, in drill cores. This elevation roughly corresponds to a break in slope on the coastal plain (3.4 ± 1.5 m), which can be interpreted as a shoreline angle. Considering this a beach deposit, and using the indicative meaning calculator [52], we calculate that this strandplain indicates a LIG paleo RSL of 3.44 ± 2.26 m (Figure 2). At the nearby Cowley Beach strandplain, Brooke et al. [9] established that the strandplain beach ridge morphology tracked Holocene sea-level trends.

The surface expression of the Wyvuri Embayment LIG beach barrier can be found at other locations along the Queensland coast, with the shoreline angle located roughly at the same elevation as that of Wyvuri Embayment (yellow markers in Figure 1).

Starting from the description of Gagan et al. [30] and high-resolution (5m) Digital Elevation Models from [31], we identified other locations scattered along the Queensland coast where the LIG beach barrier is visible and where sea-level index points can be derived (see Supplementary Materials for detailed maps of each area and a spreadsheet containing sea-level interpretations, similar to those shown in Figure 2). The elevation of these barriers is consistent with those identified in northern New South Wales, which preserve a LIG sea-level trend from a highstand at $+6 \pm 0.5$ m at 129 ka BP to $+4$ m by 116 ka BP [33]. The SE Queensland and northern New South Wales studies revealed that regional coastal fault reactivation has occurred during the Late Quaternary that has influenced the accommodation space for strandplain deposition. Overall the Late Quaternary onshore strandplains extending from far North Queensland to far northern New South Wales records indicate that the coastline and relative sea-levels since MIS 7 are preserved in the $+3$ to $+6$ m elevation. This is in stark contrast to the offshore submerged record, suggesting a LIG paleo relative sea level below the modern one.

The fact that LIG reefs in the GBR are found below the typical elevation of reefs of the same age on passive continental margins was discussed by [53], who attributed it to a combination of long-term subsidence of the continental margin and erosion of the Pleistocene reef framework during glacial times. Differential Holocene reef growth rates seem to indicate that the Central GBR is subsiding with respect to the Northern and Southern GBR [20], and this subsidence may be related to the re-activation of NNW-SSE extensional faults along the eastern Queensland margin [73, and references therein].

2 REEF ISOSTASY

Coral reefs are created by the fixation of calcium carbonate mostly by hermatypic corals and calcareous algae [90], that respond to variations in sea-level by catching up, keeping up or giving up. From the geological perspective, this results in the creation of a mass of reef framework, which can exert a

On the GBR, corals of LIG age are presently preserved under a subsurface unconformity, which occurs between 3-5 to 20-25 meters below present sea level, depending on the site [39, 53, 72]. Murray-Wallace and Belperio [60] highlight that, while low-lying islands are scattered throughout the GBR, outcrops of Pleistocene reefs above modern sea level are absent. The only exception may be an exposed reef of supposed Pleistocene age at 1-4m above present sea level [39] at Digby Island [47, 48]. However, the age of this reef has never been confirmed with absolute dating, and it will not be discussed further. Retrieval of LIG reef sections on the GBR has been historically done by coring through the Holocene reef down to the Holocene/LIG unconformity. A full account of the best-preserved and best-dated Last Interglacial corals on the GBR, alongside their indicative meaning, is provided by Dechnik et al. [19]. These data were recently compiled into the standardized WALIS (World Atlas of Last Interglacial Shorelines) database by Chutcharavan and Dutton [15] (blue markers in Figure 1)

Murray-Wallace and Belperio [60] report the presence of scattered coastal deposits of LIG age along the continental coasts of Queensland. These deposits become ubiquitous along the SE Queensland Fraser Island Coast and far north New South Wales coasts [33]. In contrast to LIG reef sequences in the GBR, most of these strandplains are rarely assigned an age with absolute dating techniques. Their MIS 5e age has been inferred via chronostratigraphic correlation with lower younger (Holocene) units, and infinite radiocarbon ages. An expanding OSL chronol-




Figure 3: Illustration of reef isostasy caused by the buildup of the reef complex since the Last Interglacial. **A.** The LIG reef is built on top of an older reef (or the bedrock). The addition of this load leads to isostatic subsidence of the underlying bedrock. **B.** As GMSL falls (e.g., under glacial conditions), the reef is partially eroded and/or dissolved (e.g., by karst processes), resulting in isostatic rebound. **C.** As sea level rises a second time, the reef starts to build again on top of previous structures, causing additional subsidence. ΔRSL represents the relative sea-level change caused by reef isostasy. The colored dashed lines represent the elevation of the coral during the LIG (red) and its present-day elevation (violet). Note that the uplift and subsidence following reef loading and unloading are transient through glacial-interglacial times, and that in our study we do not model the uplift following reef erosion, which we consider to be balanced with Holocene re-growth.

156 significant load on the underlying crust. This loading causes an
 157 isostatic response that is non-negligible. Hereafter, we define
 158 the isostatic adjustment induced by coral reef building as “reef
 159 isostasy”.

160 An illustration of how reef isostasy impacts the elevation of
 161 a LIG reef measured today is shown in Figure 3. During the
 162 LIG, reef builds on top of an older reef surface (or the basement,
 163 Figure 3A). This loading induces isostatic adjustment, causing
 164 subsidence, or equivalently a relative sea-level rise. The sea-
 165 level change ΔRSL magnitude induced by reef isostasy depends
 166 on reef thickness as well as its geographic extent. Areas with
 167 widespread reef coverage (larger in areal extent than the effective
 168 lithospheric thickness) produce a longer wavelength isostatic sig-
 169 nal, and therefore a larger magnitude relative sea-level change
 170 associated with reef isostasy. In contrast, less extensive reef
 171 coverage, smaller in areal extent than the effective lithospheric
 172 thickness of 50–100 km in this region [6], produce a minimal iso-
 173 static response. During a subsequent glacial period of lower sea
 174 level, erosion and karstification may lead to unloading-induced
 175 uplift that partially compensates for the subsidence during reef-
 176 building Figure 3B)

177 An increase in local relative sea-level from crustal subsidence
 178 induced by reef isostasy results in lower elevation LIG coral
 179 sea-level markers today, compared to their original elevation at
 180 the LIG. Therefore LIG coral reef sea-level marker elevations
 181 must be corrected upwards to account for reef isostasy, poten-
 182 tially resulting in higher reconstructed LIG GMSL than prior
 183 estimates.

184 3 RESULTS & DISCUSSION

185 3.1 Reef isostasy: high vs coarse resolution

186 Figure 4 (right panels) shows the elevation change a LIG sea
 187 level marker would undergo from 122 ka to 0 ka due to reef
 188 isostasy (negative values signify that sea-level markers experi-
 189 enced subsidence since the LIG). Our high-resolution simulation
 190 of reef isostasy in the Great Barrier Reef predicts a maximum

relative sea level change of 0.34 m since the Last Interglacial
 (Figure 4B). These maximum values are reached in Northeast-
 ern Queensland and along the coastline of the southern GBR.
 Our predictions for relative sea level change due to reef isostasy
 suggest this process is negligible compared to other uncertain-
 ties on the paleoelevation of LIG coral reefs (for example coral
 growth depths, tides etc.). In contrast, the coarse resolution
 reef isostasy calculations (using a 1D GIA model set up and a
 loading scenario that does not account for reef coverage area)
 predict a maximum relative sea level change of 1.45 m since the
 Last Interglacial (Figure 4D). The discrepancy between high vs.
 coarse resolution models is due to the fact that the high res-
 olution calculation involves a more localized loading geometry
 (and thus reduced crustal deflection) due to elastic compensation
 within the lithosphere.

Because high-resolution modeling using the 3D sea-level model
 is computationally expensive, we also tested whether a 1D sea-
 level model could accurately capture the pattern and magnitude
 of relative sea level change due to reef isostasy. We used the
 high resolution coral reef loading scenario (paired with the 3D
 sea-level model) and first multiplied the loading grid by the
 fractional area of reef coverage on a 1 km scale. We then inter-
 polated this loading scenario onto a grid with ~ 34 km resolution
 to create a coarse grid that accounts for fractional area of reef
 coverage (Figure 4E). We ran a 1D sea-level model with this
 loading scenario using the same Earth model as in the other 1D
 calculation. This simulation resulted in a similar magnitude of
 reef isostasy as in the 3D sea-level high-resolution model, with
 a maximum value of 0.4 m of RSL change since the LIG (Figure
 4F). However, the spatial pattern does not reproduce the signal
 along the southern Great Barrier Reef coastline shown in the 3D
 sea-level high resolution simulations. This difference is likely
 due to the higher resolution associated with the 3D sea-level
 simulation rather than 3D earth structure, as the coarse resolu-
 tion 1D calculation does not capture the reef loading regions
 along the central and southern Great Barrier Reef coastline.

To assess the sensitivity of our results to Earth structure pa-
 rameters, we also performed 1D sea-level simulations using an

191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228




Figure 4: **A.** High resolution coral reef thickness (122-0 ka) for the reef isostasy loading scenario. **B.** Predicted marker elevation change since LIG due to reef isostasy in response to loading in frame A. **C-D.** As in A-B, except for the coarse resolution modeling. **E-F.** As in C-D, except for the coarse resolution treatment of reef thickness (122-0 ka) accounting for reef area coverage. Yellow and blue dots in each map represent the sites shown in Figure 1

229 alternate Earth model, VM2 [63]. We found that changing the
 230 Earth model had a negligible effect, perturbing the predicted RSL
 231 change by a maximum of 3% at the Queensland/GBR sea-level
 232 indicator sites.

233 3.2 Contribution of other geodynamic processes

234 We predicted the elevation change due to reef isostasy (Figure
 235 5A), dynamic topography (Figure 5B), and glacial isostatic ad-
 236 justment (Figure 5C) from 127 ka to present day. These values
 237 represent the elevation change a LIG sea-level indicator would
 238 undergo from 127 ka to 0 ka (negative values signify that sea-
 239 level indicators experienced subsidence, positive values signify
 240 that sea-level indicators experienced uplift since the LIG). The
 241 total predicted influence on Last Interglacial sea-level indicator
 242 elevation from these geodynamic processes is shown in Figure
 243 5D.

244 Our dynamic topography predictions show an elevation change
 245 of -10 to 10 m from 127 ka to present day. This means that dy-
 246 namic topography would have uplifted the Australian continent
 247 by up to 10 m, while offshore regions on the continental shelf
 248 would have subsided up to 5 to 10 m since the LIG. Variations in
 249 input density and viscosity structure lead to $\sim \pm 1$ m uncertainty
 250 in post-LIG dynamic topography change (based on standard
 251 deviation of 15 model predictions), and the spatial pattern is
 252 remarkably consistent amongst the 15 models investigated here.
 253 These results suggest that our predictions of convectively driven
 254 onshore-offshore tilting are robust. This inference is corroborated
 255 by ~ 100 m Myr⁻¹ uplift rates inferred from river profile
 256 modelling [17] and patterns of Late Cenozoic age-independent
 257 magmatism [7], both features that have been attributed to the
 258 presence of an active small-scale convection cell beneath the
 259 Queensland margin. Although the dynamic topography maxima
 260 and minima are offset with respect to the observed relative sea
 261 level maxima and minima, the highest horizontal resolution for
 262 the dynamic topography predictions is ~ 200 km, and therefore
 263 it may not be possible to precisely match the observed tilting at
 264 this resolution.

265 Similarly, glacial isostatic adjustment would have produced
 266 uplift on the continent and subsidence offshore. Our predictions
 267 show that the continent may have uplifted 6 m and offshore
 268 regions subsided 2 m since the Last Interglacial. The spatial
 269 variability in elevation change due to glacial isostatic adjustment
 270 is caused by the process known as continental levering, where
 271 uplift occurs along continental margins as sea-level rise causes
 272 subsidence in ocean basins due to additional loading [57].

273 In this study, we did not model some other potential mechanisms
 274 that may cause departure from eustasy in the study area. For
 275 example, crustal deformation due to re-activation of older faults
 276 has been inferred to affect Holocene reefs [see 73, and references
 277 therein]. While such mechanism might have relevant local effect,
 278 any fault system causing crustal motions would have to be active
 279 (with roughly the same deformation rates) over nearly 2000 km
 280 of coast to reconcile the observed onshore-offshore tilting trend.
 281 This seems an unlikely pattern in an intraplate margin setting
 282 such as the Queensland-GBR area. Another process we did not
 283 model is the isostatic response to siliciclastic sediment loading.
 284 While this process may be relevant at some locations (e.g., on
 285 the shelf in front of large rivers, with relevant sediment inputs on
 286 the shelf), studies on the Central GBR shelf suggested that the

thickness of Holocene sediments is rather limited [<2.5 m 44],
 hence siliciclastic sediment isostasy is unlikely to explain the
 difference between onshore and offshore LIG sea-level proxies,
 recorded over such a large latitudinal gradient.

4 CONCLUSIONS

The Queensland - GBR area is characterized by an enigmatic
 difference in the elevation of LIG sea-level indicators between
 offshore (GBR) and onshore (Queensland coast) sites. This
 offset motivated our study's modeling of local post-depositional
 vertical land motion. We modelled sea-level change due to reef
 isostasy, dynamic topography, and glacial isostatic adjustment
 since the LIG in this area, which is located on a passive margin
 spanning a latitudinal range of almost 2000 km. Our models
 explored whether reef isostasy, which is considered here for
 the first time, may play a role in the vertical displacement of
 LIG fossil reefs, which are among the most frequently used
 geological sea-level proxies [82, 21, 62].

In our study area, the contribution of reef isostasy to vertical
 land motions is negligible, reaching maximum values of 0.34m.
 In terms of GMSL, this is roughly equivalent to half the contribu-
 tion of mountain glaciers melting and thermal expansion during
 the LIG (estimated as up to 1m; 22). Reef isostasy therefore
 produces a small change in RSL since the LIG at the GBR, and
 is insufficient in magnitude to explain discrepancies between
 observed LIG RSL markers offshore and onshore. However,
 we highlight that this mechanism may represent a potentially
 important contribution to vertical land motions in areas with
 dense and widespread coral reef coverage. Therefore, it should
 be always considered as a potential bias towards higher GMSL
 in areas with widespread reef coverage.

To realistically represent coral reef loading since the LIG in
 a given area, it is important to gather direct measurements of
 reef thickness, extent, density and porosity, together with esti-
 mates of mass loss since the LIG (e.g., due to erosion or karst
 processes, which we do not model here) and, in the case of
 wide lagoons, carbonate sediment production from the reef. Our
 results also underscore the importance of high resolution mod-
 eling, especially in accounting for the areal coverage of coral
 reefs, to accurately reproduce relative sea level change due to
 reef isostasy. Although 1D sea-level models are more computa-
 tionally efficient, for small-scale loading patterns such as coral
 reefs, it may be important to use grid refinement in 3D modeling
 (or high resolution, accounting for reef coverage) in order to
 accurately capture relative sea level response to reef loading.

Comparing the modeled relative contributions of reef isostasy,
 dynamic topography, and glacial isostatic adjustment, we sur-
 mise that only the predicted changes due to dynamic topography
 across sites has a similar magnitude to the differences in sea-
 level indicators elevation between onshore and offshore. Our
 dynamic topography simulations, in contrast to reef isostasy
 and GIA modeling, predict trends of uplift on the continent and
 subsidence offshore of a magnitude similar to the observed in
 relative sea-level proxies. This result strengthens the idea that
 dynamic topography may play a major role in the vertical dis-
 placement of LIG sea-level indicators [5]. Therefore we suggest
 that, along the GBR, dynamic topography driven by mantle con-
 vention movements may capture the majority of the "long-term"




Figure 5: A-C. Predicted elevation change to sea-level indicators from 127 to 0 ka due to: A. reef isostasy B. dynamic topography C. glacial isostatic adjustment. Colored circles represent LIG sea-level indicators as shown in Figure 1. D. Total predicted elevation change to sea-level indicators from 127 to 0 ka. E-F. Gray represents observed elevation range and black line represents mean values for transect $\alpha-\alpha'$ (left) and $\beta-\beta'$ (right). G-H. Light blue line and envelope represents the observed range in reef thicknesses in coral reef loading scenario from LIG to present. Dark blue line and envelope represents the predicted elevation change to sea-level markers due to reef isostasy (as in Figure 5A). Lines represent mean values based on spatial uncertainty of 100 km on either side of transect and intermodel variation uncertainty; envelopes represent the 2 sigma combined uncertainty. I-J. GBR LIG sea-level data points projected onto transects $\alpha-\alpha'$ (left) and $\beta-\beta'$ as a function of distance between the data point and the closest point on the transect. Colored circles/triangles represent LIG sea-level indicator ages. Predicted elevation change projected onto transect A (left) and B (right) for reef isostasy (blue), dynamic topography (red), glacial isostatic adjustment (green), and total (pink). Lines and envelope calculated as in G-H

344 subsidence that was noted by previous studies [53] attempting to
 345 explain the conundrum of lower-than-present LIG reefs on the
 346 Great Barrier Reef. This is thus an important avenue for future
 347 work as improved models of mantle heterogeneity beneath the
 348 area become available.

349 5 METHODS

350 5.0.1 Constructing the coral reef loading scenario

351 As a baseline dataset for the presence/absence of coral reefs, we
 352 used the 500x500m raster dataset [11, 12, 42] of the warm-water
 353 reefs map compiled by UNEP-WCMC, WorldFish Centre, WRI,
 354 TNC [83, 40, 41, 79]. We created a coral reef loading scenario
 355 since the Last Interglacial (122-0 ka) using two methods, with
 356 different resolutions. For the "coarse resolution grid", we used
 357 standard approach for sea-level model calculations and placed
 358 our coral loading scenario onto a ~34 km resolution grid. For
 359 the "high resolution grid", we placed our coral loading scenario
 360 onto a 1 km resolution grid, and accounted for the areal fraction
 361 of coral reef coverage within each 1 km x 1 km grid cell.

362 Because the GBR reef is characterized by narrow, sometimes
 363 isolated, strips of coral reef, we were concerned that the stan-
 364 dard grid resolution (~34 km) used in sea-level models may
 365 unrealistically smooth out the reef loading signal. Thus, for the
 366 "high resolution grid" we interpolated a high-resolution Digital
 367 Elevation Model for bathymetry in the Great Barrier Reef area
 368 onto a 1 km resolution grid [8]. We then assessed the fractional
 369 area of reef coverage within each 1 km x 1 km grid cell using
 370 the "Fishnet" tool of ArcGIS. Of grid cells with non-zero reef
 371 coverage, 44% had full reef coverage (Figure 6). We then multi-
 372 plied the coral reef thickness in our 1 km x 1 km grids by the
 373 areal fraction of reef coverage to produce our "high resolution
 374 grid" coral reef loading scenario.




Figure 6: Fractional area of present-day reef coverage. Yellow and blue dots represent the sites shown in Figure 1.

375 We also used a standard approach for constructing a loading
 376 scenario by interpolating a high-resolution bathymetric Digital

Elevation Model of the GBR area onto a Gauss Legendre grid
 with ~34 km resolution (maximum spherical harmonic degree
 512) commonly used in sea-level calculations. This approach
 does not account for coral reef coverage since the coral reef
 thickness is smoothed over a wide area relative to the lateral
 extent of coral reefs. We term this coral reef loading scenario
 the "coarse resolution grid" (Figure 4C).

In both scenarios, we assumed that regions with any reef cover-
 age (fractional area of reef coverage > 0; Figure 6A) had coral
 reefs that had grown since the Last Interglacial. We assigned the
 total coral reef thickness deposited since the Last Interglacial
 as the modern basement depth (as in, we assumed the coral
 reef surface grew to modern sea level) in regions with basement
 depths shallower than 55 m. Below this bathymetry, we con-
 sidered that no reef was present in the LIG. To partition coral
 reef loading across 122 to 0 ka, we made the assumption that
 the Last Interglacial reef thickness would represent 1.5 times
 the thickness of Holocene coral reef growth, given the longer
 time available for LIG reefs to grow with respect to Holocene
 ones. In our models, we assumed a reef porosity of 40% (that is,
 the porosity of reefs in sand flats/lagoons in the GBR reported
 by 38) and a coral reef density of 1600 kg/m³ (equivalent to the
 average coral colony density as reported by 10 in 38).

For the "high resolution grid" coral loading scenario, we mul-
 tiplied our map of reef thickness by the fractional area of reef
 coverage (Figure 6A). This assumes that the coverage hasn't
 changed since 120 ka. To isolate the impact of reef loading, we
 did not include ice sheet loading changes in our modeling. Our
 reef loading scenario introduced the LIG coral thickness at 120
 ka and the Holocene coral thickness at 8 ka. Although coral
 reefs built over a longer time span, we simplified our calculation
 by introducing the load at a single timestep, assuming that the
 timing of the load will have a negligible impact at present-day
 after several thousand years of isostatic adjustment. To con-
 serve mass, we uniformly removed a layer of sediment from the
 continents with a mass equivalent to the total reef load globally.

Although reef loading prior to the LIG would have induced an
 ongoing isostatic response at the LIG, our analysis is limited to
 estimating sea-level change since the LIG due to reef loading
 over only the last glacial cycle. Thus, we limited our modeling
 to the period from 122 to 0 ka to assess the magnitude of sea
 level change due to reef loading since 122 ka.

5.0.2 Modeling Isostatic Adjustment: Reef isostasy

1D calculation (coarse resolution). To calculate relative sea-
 level change (Δ RSL) in response to reef loading over the last ice
 age, we used a gravitationally self-consistent sea-level model.
 We used the coarse resolution coral reef loading scenario as in-
 put to a 1D sea-level model, which assumes radially symmetric
 Earth structure. Our calculations are based on the theory and
 pseudo-spectral algorithm described by Kendall et al. [46] with
 a spherical harmonic truncation at degree and order 512 (spatial
 resolution of ~34 km). These calculations include the impact of
 load-induced Earth rotation changes on sea level [55, 59], evol-
 ving shorelines and the migration of grounded, marine-based ice
 [45, 56, 50, 46]. Our predictions require models for Earth's vis-
 coelastic structure. We adopted an earth model characterized by
 a lithospheric thickness of 96 km, and upper and lower mantle
 viscosities of 5×10^{20} and 5×10^{21} Pa s, respectively.

435 **3D calculation (high resolution).** The predicted magnitude of
 436 relative sea level change is sensitive to the spatial scale of the
 437 load, in addition to the load thickness. To assess whether the
 438 coarser resolution accurately captures the crustal deformation
 439 (and thus relative sea level) response to reef loading, we next
 440 performed calculations using a 3D sea-level model, and the
 441 "high resolution grid" coral reef loading scenario with a regional
 442 spatial resolution of 1 km that accounts for the fractional area of
 443 reef coverage in each grid cell.

444 To solve for relative sea level change in response to coral reef
 445 loading on a higher resolution of 1 km, we used a global 3D
 446 finite volume sea level and Earth deformation model [51]. The
 447 numerical approach incorporates lateral variations in Earth structure
 448 and calculates the resulting gravitationally self-consistent
 449 sea level change [58]. Previous studies have adopted this computational
 450 model in order to account for 3D earth structure (e.g.,
 451 4, 32, 49). The 3D glacial isostatic adjustment model is capable
 452 of km-scale resolution, which is achieved through regional grid
 453 refinement for computational efficiency [32]. The importance
 454 of high resolution GIA modeling has been demonstrated for the
 455 solid Earth response to marine grounding line migration in
 456 Antarctica [87]. Grid refinement is achieved by incrementally
 457 bisecting grid edges in the selected region to achieve the desired
 458 1 km x 1 km resolution, and a final smoothing operation along
 459 the region boundary to ensure a well-behaved transition.

460 Our simulation uses a 3D viscoelastic earth model. Here, we
 461 apply the hybrid model described in Austermann et al. [6], which
 462 infers mantle viscosity from seismic tomography using anelastic
 463 scaling relationships and additional information on the thermal
 464 and rheological state of the upper mantle. In the upper 400 km,
 465 a calibrated parameterisation of anelastic behaviour at seismic
 466 frequencies is used to self-consistently determine lithospheric
 467 thickness (assumed here to be equivalent to 1175°C isotherm
 468 depth) and viscosity variations from the shear-wave velocity (V_S)
 469 structure of the tomographic model, SL2013sv [66, 75]. Below
 470 400km, viscosities are derived from the shear wave tomography
 471 model SEMUCB-WM1 [29]. Austermann et al. (2021) provides
 472 details on the V_S to viscosity conversion.

473 In our 3D GIA calculations, viscosity variations are shifted at
 474 each depth to average to 5×10^{20} Pa s in the upper mantle
 475 viscosity 5×10^{21} Pa s in the lower mantle viscosity [65], identical
 476 to the earth model used in the 1D GIA calculations. The effective
 477 lithospheric thickness in this region varies from 50–100
 478 km (Figure SX). We paired this model with the high resolution
 479 coral reef loading scenario (Figure 4A) which accounts for reef
 480 coverage area at 1 km resolution (Figure 6A).

481 5.1 Modeling Glacial Isostatic Adjustment: Ice loading

482 We modeled relative sea level change in response to ice sheet
 483 and ocean loading changes since the LIG using the 1D pseudo-
 484 spectral approach described in Kendall et al. [46]. We used
 485 the same model and earth structure described in the 1D reef
 486 loading sea-level calculations (an Earth model characterized by a
 487 lithospheric thickness of 96 km, and ρ_{55} upper and lower mantle
 488 viscosities (5×10^{20} Pa s and 5×10^{21} Pa s, respectively).

489 We used an ice history characterized by the GMSL history in
 490 Waelbroeck et al. [85] over the last glacial cycle. The ice history
 491 was constructed using the ICE-6G deglacial ice geometry history

and has no excess melt across the LIG (as in 6). The GMSL
 history was adjusted at the LIG since the Waelbroeck GMSL
 history assumes a value of -75 m at 128 ka, which is at odds with
 coral evidence from the many locations that indicate sea level
 must have been close to present at that time. To account for this
 discrepancy, the timing of the GMSL curve is shifted prior to the
 LIG back by 3.5 ka. This shift allows for a longer interglacial
 time period without changing the deglaciation pattern of the
 original curve and places the MIS 6 sea-level low stand at 135.5
 ka (as in 24).

5.2 Dynamic Topography

Observational estimates indicate that mantle flow-driven vertical
 motions can reach rates of ~ 0.1 -1 m kyr⁻¹ in certain locations,
 suggesting a significant fraction of relative sea-level change along
 the Great Barrier Reef from the LIG to present day could result
 from evolving mantle dynamic topography [36, 86, 5, 81]. To investigate
 this possibility, we simulate rates of global dynamic topography
 change using the mantle convection code ASPECT and an ensemble
 of Earth models based on 5 seismic tomographic inversions of deep
 Earth structure (LLNL-G3D-JPS, 77; S40RTS, 68; SAVANI, 3; SEMUCB-
 WM1, 29; TX2011, 34) and 3 radial viscosity profiles (S10, 80; F10V1,
 28; F10V2, 28).

Above 300 km, input temperature and density fields are derived
 from a modified version of the RHGW20 model of Richards et al.
 [66], which accounts for anelasticity at seismic frequencies and
 has been demonstrated to yield acceptable fits to present-day
 short-wavelength dynamic topography. Unlike RHGW20, which is
 based exclusively on the SL2013sv global surface wave tomographic
 model [75], the upper mantle model we adopt here is augmented
 with regional high-resolution tomographic studies in North America
 (SL2013NA; 74), Africa (AF2019; 14), and South America and the
 South Atlantic Ocean (SA2019; 13; see 37 and 67 for further details).
 Below 400 km, a thermodynamic modelling approach is used to
 obtain thermochemical buoyancy structures for each combination
 of seismic tomographic and rheological input that are compatible
 with present-day geophysical observables, including geoid anomalies,
 dynamic topography, and CMB excess ellipticity, and comprise
 thermochemical anomalies within the base of LLVPs (67; see
 Supplementary Material for further details). Between 300 and 400
 km, temperatures and densities derived from these two independent
 parameterisations are smoothly merged by taking their weighted
 average as a function of depth.

The time-dependent geodynamic simulations derived from these
 Earth models assume free-slip conditions at the surface and core-
 mantle boundary, account for lithospheric cooling by including
 shallow mantle buoyancy variations and representative thermal
 conductivity, and incorporate temperature- and composition-
 dependent viscosity variations (see Supplementary Material for
 further details). Following [5], we run our models forward in
 time and, to avoid the potential for transient numerical artefacts
 in early time steps to affect our results, we assume the average
 rate of dynamic topography change between 0.5 and 1.5 Ma is
 representative of that experienced between the LIG and the
 present day. Change in dynamic topography at specific sea-level
 sites is calculated by combining perturbations due to the
 evolving mantle flow pattern with those caused by rigid plate

550 motion across the convective planform. This is accomplished by
 551 translating the dynamic topography field calculated for the LIG
 552 into its present-day coordinates using plate velocities taken from
 553 MORVEL [2], before calculating the difference between this
 554 rotated LIG field and the predicted present-day field, yielding
 555 a total of 15 individual model predictions. Note that the maxi-
 556 mum horizontal resolution of the tomographically derived Earth
 557 models is ~200 km, placing an important limit on the minimum
 558 wavelength of predicted dynamic topography variations.

559 6 DATA AVAILABILITY

560 Supplementary figures and the datasets used in this study are
 561 available open-access as Rovere et al. [70].

562 7 AUTHOR CONTRIBUTIONS

563 The manuscript was written jointly by A.R. and T.P. The initial
 564 concept of this work was developed by A.R., M.J.O, I.D.G. and
 565 J.X.M. Models of reef isostasy were developed by T.P. Models
 566 of dynamic topography and glacial isostatic adjustment were
 567 developed by F.R., J.A. and K.L. The parts of the manuscript
 568 related to field observations was written by A.R., M.J.O. and
 569 I.D.G. The parts of the manuscript related to modelled vertical
 570 land motions were written by T.P. and F.R. with inputs from
 571 J.X.M., J.A. and K.L.

572 8 ACKNOWLEDGMENTS

573 This work was funded by the European Research Council (ERC)
 574 under the European Union’s Horizon 2020 research and inno-
 575 vation programme (grant agreement n. 802414 to A.R.). T.P.
 576 acknowledges funding from the NSF EAR Postdoctoral Fellow-
 577 ship, the University of California President’s Postdoctoral Fel-
 578 lowship, and NSF OCE – 2054757. F.D.R. acknowledges fund-
 579 ing from the Imperial College Research Fellowship Scheme. J.A.
 580 acknowledges funding from NSF grant OCE-1841888. Fund-
 581 ing is also acknowledged from Harvard University (J.X.M. and
 582 K.L.)

583 REFERENCES

584 [1] P. N. Adams, N. D. Opdyke, and J. M. Jaeger. Isostatic up-
 585 lift driven by karstification and sea-level oscillation: Mod-
 586 eling landscape evolution in north florida. *Geology*, 38(6):
 587 531–534, 2010.

588 [2] D. F. Argus, R. G. Gordon, and C. DeMets. Geologically
 589 current motion of 56 plates relative to the no-net-rotation
 590 reference frame. *Geochemistry, Geophysics, Geosystems*,
 591 12(11), 2011.

592 [3] L. Auer, L. Boschi, T. Becker, T. Nissen-Meyer, and D. Gi-
 593 ardini. Savani: A variable resolution whole-mantle model
 594 of anisotropic shear velocity variations based on multiple
 595 data sets. *Journal of Geophysical Research: Solid Earth*,
 596 119(4):3006–3034, 2014.

597 [4] J. Austermann, J. X. Mitrovica, K. Latychev, and G. A.
 598 Milne. Barbados-based estimate of ice volume at last
 599 glacial maximum affected by subducted plate. *Nature*
 600 *Geoscience*, 6(7):553–557, 2013.

[5] J. Austermann, J. X. Mitrovica, P. Huybers, and A. Ro-
 vere. Detection of a dynamic topography signal in last
 interglacial sea-level records. *Science Advances*, 3(7):
 e1700457, 2017.

[6] J. Austermann, M. J. Hoggard, K. Latychev, F. D. Richards,
 and J. X. Mitrovica. The effect of lateral variations in earth
 structure on last interglacial sea level. *Geophysical Journal*
International, 227(3):1938–1960, 2021.

[7] P. Ball, K. Czarnota, N. White, M. Klöcking, and D. Davies.
 Thermal structure of eastern australia’s upper mantle and
 its relationship to cenozoic volcanic activity and dynamic
 topography. *Geochemistry, Geophysics, Geosystems*, 22
 (8):e2021GC009717, 2021.

[8] R. Beaman. High-resolution depth model for the great
 barrier reef and coral sea - 100 m, 2020.

[9] B. P. Brooke, Z. Huang, W. A. Nicholas, T. S. Oliver,
 T. Tamura, C. D. Woodroffe, and S. L. Nichol. Rela-
 tive sea-level records preserved in holocene beach-ridge
 strandplains—an example from tropical northeastern aus-
 tralia. *Marine Geology*, 411:107–118, 2019.

[10] R. W. Buddemeier, J. E. Maragos, and D. W. Knutson.
 Radiographic studies of reef coral exoskeletons: rates and
 patterns of coral growth. *Journal of Experimental Marine*
Biology and Ecology, 14(2):179–199, 1974.

[11] L. Burke, K. Reytar, M. Spalding, and A. Perry. Reefs at
 risk revisited: technical notes on modeling threats to the
 world’s coral reefs. *Washington, DC: World Resources*
Institute, 2011.

[12] L. Burke, K. Reytar, M. Spalding, and A. Perry. *Reefs at*
risk revisited. World Resources Institute, 2011.

[13] N. L. Celli, S. Lebedev, A. J. Schaeffer, and C. Gaina.
 African cratonic lithosphere carved by mantle plumes. *Nature*
Communications, 11(92):1–10, 2020. ISSN 20411723.
 doi: 10.1038/s41467-019-13871-2.

[14] N. L. Celli, S. Lebedev, A. J. Schaeffer, M. Ravenna, and
 C. Gaina. The upper mantle beneath the South Atlantic
 Ocean, South America and Africa from waveform tomog-
 raphy with massive data sets. *Geophysical Journal In-*
ternational, 221:178–204, 2020. ISSN 0956-540X. doi:
 10.1093/gji/ggz574.

[15] P. M. Chutcharavan and A. Dutton. A global com-
 pilation of u-series-dated fossil coral sea-level indica-
 tors for the last interglacial period (marine isotope stage
 5e). *Earth System Science Data*, 13(7):3155–3178, 2021.
 doi: 10.5194/essd-13-3155-2021. URL <https://essd.copernicus.org/articles/13/3155/2021/>.

[16] J. R. Creveling, J. Austermann, and A. Dutton. Uplift of
 trail ridge, florida, by karst dissolution, glacial isostatic ad-
 justment, and dynamic topography. *Journal of Geophysical*
Research: Solid Earth, 2019.

[17] K. Czarnota, G. Roberts, N. White, and S. Fishwick. Spa-
 tial and temporal patterns of australian dynamic topogra-
 phy from river profile modeling. *Journal of Geophysical*
Research: Solid Earth, 119(2):1384–1424, 2014.

[18] A. Dalca, K. Ferrier, J. X. Mitrovica, J. Perron, G. Milne,
 and J. Creveling. On postglacial sea level—iii. incorporat-

- ing sediment redistribution. *Geophysical Journal International*, 194(1):45–60, 2013.
- [19] B. Dechnik, J. M. Webster, G. E. Webb, L. Nothdurft, A. Dutton, J. C. Braga, J.-x. Zhao, S. Duce, and J. Sadler. The evolution of the great barrier reef during the last interglacial period. *Global and Planetary Change*, 149:53–71, 2017.
- [20] B. Dechnik, J. M. Webster, G. E. Webb, L. Nothdurft, and J.-x. Zhao. Successive phases of holocene reef flat development: Evidence from the mid-to outer great barrier reef. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 466:221–230, 2017. ISSN 0031-0182. doi: <https://doi.org/10.1016/j.palaeo.2016.11.030>. URL <https://www.sciencedirect.com/science/article/pii/S0031018216304412>.
- [21] A. Dutton and K. Lambeck. Ice volume and sea level during the last interglacial. *science*, 337(6091):216–219, 2012.
- [22] A. Dutton, A. E. Carlson, A. J. Long, G. A. Milne, P. U. Clark, R. DeConto, B. P. Horton, S. Rahmstorf, and M. E. Raymo. Sea-level rise due to polar ice-sheet mass loss during past warm periods. *science*, 349(6244):aaa4019, 2015.
- [23] A. Dutton, J. M. Webster, D. Zwart, K. Lambeck, and B. Wohlfarth. Tropical tales of polar ice: evidence of last interglacial polar ice sheet retreat recorded by fossil reefs of the granitic seychelles islands. *Quaternary Science Reviews*, 107:182–196, 2015.
- [24] B. Dyer, J. Austermann, W. J. D’Andrea, R. C. Creel, M. R. Sandstrom, M. Cashman, A. Rovere, and M. E. Raymo. Sea-level trends across the Bahamas constrain peak last interglacial ice melt. *Proceedings of the National Academy of Sciences of the United States of America*, 118(33):1–11, 2021. ISSN 10916490. doi: 10.1073/pnas.2026839118.
- [25] W. Farrell and J. A. Clark. On postglacial sea level. *Geophysical Journal International*, 46(3):647–667, 1976.
- [26] K. L. Ferrier, J. X. Mitrovica, L. Giosan, and P. D. Clift. Sea-level responses to erosion and deposition of sediment in the indus river basin and the arabian sea. *Earth and Planetary Science Letters*, 416:12–20, 2015.
- [27] K. L. Ferrier, J. Austermann, J. X. Mitrovica, and T. Pico. Incorporating sediment compaction into a gravitationally self-consistent model for ice age sea-level change. *Geophysical Journal International*, 211(1):663–672, 2017.
- [28] A. M. Forte, S. Quéré, R. Moucha, N. A. Simmons, S. P. Grand, J. X. Mitrovica, and D. B. Rowley. Joint seismic-geodynamic-mineral physical modelling of African geodynamics: A reconciliation of deep-mantle convection with surface geophysical constraints. *Earth and Planetary Science Letters*, 295(3-4):329–341, 2010.
- [29] S. W. French and B. Romanowicz. Broad plumes rooted at the base of the Earth’s mantle beneath major hotspots. *Nature*, 525(7567):95–99, 2015. ISSN 0028-0836. doi: 10.1038/nature14876.
- [30] M. K. Gagan, D. P. Johnson, and G. M. Crowley. Sea level control of stacked late quaternary coastal sequences, central great barrier reef. *Sedimentology*, 41(2):329–351, 1994.
- [31] Geoscience Australia. *Digital Elevation Model (DEM) of Australia derived from LiDAR 5 Metre Grid*. Australian Government, Canberra, 2015. doi: <https://doi.org/10.26186/89644>.
- [32] N. Gomez, K. Latychev, and D. Pollard. A coupled ice sheet–sea level model incorporating 3d earth structure: Variations in antarctica during the last deglacial retreat. *Journal of Climate*, 31(10):4041–4054, 2018.
- [33] I. D. Goodwin. Last interglacial sea-level and wave climate change in the subtropical south-west pacific. in prep.
- [34] S. P. Grand. Mantle shear–wave tomography and the fate of subducted slabs. *Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences*, 360(1800):2475–2491, 2002.
- [35] J. M. Gregory, S. M. Griffies, C. W. Hughes, J. A. Lowe, J. A. Church, I. Fukimori, N. Gomez, R. E. Kopp, F. Landerer, G. L. Cozannet, R. Ponte, D. Stammer, M. E. Tamisiea, and R. S. W. van de Wal. Concepts and terminology for sea level: Mean, variability and change, both local and global. *Surveys in Geophysics*, 40(6):1251–1289, 2019.
- [36] R. A. Hartley, G. G. Roberts, N. White, and C. Richardson. Transient convective uplift of an ancient buried landscape. *Nature Geoscience*, 4(8):562–565, 2011.
- [37] M. J. Hoggard, K. Czarnota, F. D. Richards, D. L. Huston, A. L. Jaques, and S. Ghelichkhan. Global distribution of sediment-hosted metals controlled by craton edge stability. *Nature Geoscience*, 13:504–510, 2020.
- [38] D. Hopley. *Density and Porosity Density and Porosity: Influence on Reef Accretion Rates*, pages 303–304. Springer Netherlands, Dordrecht, 2011. ISBN 978-90-481-2639-2. doi: 10.1007/978-90-481-2639-2_275. URL https://doi.org/10.1007/978-90-481-2639-2_275.
- [39] D. Hopley, S. G. Smithers, and K. Parnell. *The geomorphology of the Great Barrier Reef: development, diversity and change*. Cambridge University Press, 2007.
- [40] IMaRS-USF. Millennium coral reef mapping project. unvalidated maps, 2005.
- [41] IMaRS-USF, IRD. Millennium coral reef mapping project. validated maps, 2005.
- [42] Institute for Marine Remote Sensing, University of South Florida (IMaRS/USF) Institut de Recherche pour le Développement (IRD), UNEP-WCMC, The WorldFish Center, and WRI,. Global Coral Reefs composite dataset compiled from multiple sources for use in the Reefs at Risk Revisited project incorporating products from the Millennium Coral Reef Mapping Project prepared by IMaRS/USF and IRD, 2011.
- [43] IPCC. *Ipcc special report on the ocean and cryosphere in a changing climate*. 2019.
- [44] D. Johnson and D. Searle. Post-glacial seismic stratigraphy, central great barrier reef, australia. *Sedimentology*, 31(3):335–352, 1984.
- [45] P. Johnston. The effect of spatially non-uniform water loads on prediction of sea-level change. *Geophysical Journal International*, 114(3):615–634, 1993.

- 771 [46] R. A. Kendall, J. X. Mitrovica, and G. A. Milne. On post-
772 glacial sea level—ii. numerical formulation and comparative
773 results on spherically symmetric models. *Geophysical*
774 *Journal International*, 161(3):679–706, 2005.
- 775 [47] J. A. Kleypas. *Geological development of fringing reefs in*
776 *the Southern Great Barrier Reef, Australia*. James Cook
777 University of North Queensland, 1991.
- 778 [48] J. A. Kleypas and D. Hopley. Reef development across
779 a broad continental shelf, southern great barrier reef, aus-
780 tralia. In *Proc*, volume 7, pages 1129–1141, 1992.
- 781 [49] J. Kuchar, G. Milne, and K. Latychev. The importance of
782 lateral earth structure for north american glacial isostatic
783 adjustment. *Earth and Planetary Science Letters*, 512:
784 236–245, 2019.
- 785 [50] K. Lambeck, A. Purcell, P. Johnston, M. Nakada, and
786 Y. Yokoyama. Water-load definition in the glacio-hydro-
787 isostatic sea-level equation. *Quaternary Science Reviews*,
788 22(2-4):309–318, 2003.
- 789 [51] K. Latychev, J. X. Mitrovica, J. Tromp, M. E. Tamisiea,
790 D. Komatitsch, and C. C. Christara. Glacial isostatic ad-
791 justment on 3-d earth models: a finite-volume formulation.
792 *Geophysical Journal International*, 161(2):421–444, 2005.
- 793 [52] T. Lorscheid and A. Rovere. The indicative meaning
794 calculator—quantification of paleo sea-level relationships
795 by using global wave and tide datasets. *Open Geospatial*
796 *Data, Software and Standards*, 4(1):1–8, 2019.
- 797 [53] J. F. Marshall and P. J. Davies. Last interglacial reef growth
798 beneath modern reefs in the southern great barrier reef.
799 *Nature*, 307(5946):44–46, 1984.
- 800 [54] M. T. McCulloch and T. Esat. The coral record of last inter-
801 glacial sea levels and sea surface temperatures. *Chemical*
802 *Geology*, 169(1-2):107–129, 2000.
- 803 [55] G. A. Milne and J. X. Mitrovica. Postglacial sea-level
804 change on a rotating earth: first results from a gravita-
805 tionally self-consistent sea-level equation. *Geophysical*
806 *Journal International*, 126(3):F13–F20, 1996.
- 807 [56] G. A. Milne, J. X. Mitrovica, and J. L. Davis. Near-field
808 hydro-isostasy: the implementation of a revised sea-level
809 equation. *Geophysical Journal International*, 139(2):464–
810 482, 1999.
- 811 [57] J. X. Mitrovica and G. Milne. On the origin of late
812 holocene sea-level highstands within equatorial ocean
813 basins. *Quaternary Science Reviews*, 21(20-22):2179–
814 2190, 2002.
- 815 [58] J. X. Mitrovica and G. A. Milne. On post-glacial sea level:
816 I. general theory. *Geophysical Journal International*, 154
817 (2):253–267, 2003.
- 818 [59] J. X. Mitrovica, J. Wahr, I. Matsuyama, and A. Paulson.
819 The rotational stability of an ice-age earth. *Geophysical*
820 *Journal International*, 161(2):491–506, 2005.
- 821 [60] C. Murray-Wallace and A. Belperio. The last interglacial
822 shoreline in australia—a review. *Quaternary Science Re-*
823 *views*, 10(5):441–461, 1991.
- 824 [61] N. Opdyke, D. Spangler, D. Smith, D. Jones, and
825 R. Lindquist. Origin of the epeirogenic uplift of pliocene-
826 pleistocene beach ridges in florida and development of the
827 florida karst. *Geology*, 12(4):226–228, 1984.
- [62] M. J. O’Leary, P. J. Hearty, W. G. Thompson, M. E. Raymo,
J. X. Mitrovica, and J. M. Webster. Ice sheet collapse
following a prolonged period of stable sea level during the
last interglacial. *Nature Geoscience*, 6(9):796–800, 2013.
- [63] W. Peltier and R. G. Fairbanks. Global glacial ice volume
and last glacial maximum duration from an extended bar-
bados sea level record. *Quaternary Science Reviews*, 25
(23-24):3322–3337, 2006.
- [64] T. Pico, J. X. Mitrovica, K. L. Ferrier, and J. Braun. Global
ice volume during mis 3 inferred from a sea-level analy-
sis of sedimentary core records in the yellow river delta.
Quaternary Science Reviews, 152:72–79, 2016.
- [65] M. E. Raymo, J. X. Mitrovica, M. J. O’Leary, R. M.
DeConto, and P. J. Hearty. Departures from eustasy in
pliocene sea-level records. *Nature Geoscience*, 4(5):328–
332, 2011.
- [66] F. D. Richards, M. J. Hoggard, N. White, and S. Ghe-
lichkhan. Quantifying the relationship between short-
wavelength dynamic topography and thermomechanical
structure of the upper mantle using calibrated parame-
terization of anelasticity. *Journal of Geophysical Re-*
search: Solid Earth, 125:e2019JB019062, 2020. doi:
10.1029/2019JB019062.
- [67] F. D. Richards, M. J. Hoggard, S. Ghelichkhan, P. Koele-
meijer, and H. Lau. Geodynamic, geodetic, and seismic
constraints favour deflated and dense-cored llvps. *Earth-*
ArXiv, 2021.
- [68] J. Ritsema, A. Deuss, H. J. Van Heijst, and J. H. Wood-
house. S40RTS: A degree-40 shear-velocity model for
the mantle from new Rayleigh wave dispersion, teleseis-
mic traveltimes and normal-mode splitting function mea-
surements. *Geophysical Journal International*, 184:1223–
1236, 2011. ISSN 0956540X. doi: 10.1111/j.1365-246X.
2010.04884.x.
- [69] A. Rovere, P. Stocchi, and M. Vacchi. Eustatic and relative
sea level changes. *Current Climate Change Reports*, 2(4):
221–231, 2016.
- [70] A. Rovere, T. Pico, F. Richards, M. J. O’Leary, J. X. Mitro-
vica, I. D. Goodwin, J. Austermann, and K. Latychev.
Supplementary data for: "The influence of reef isostasy,
dynamic topography, and glacial isostatic adjustment on
the Last Interglacial sea- level record of Northeastern Aus-
tralia", Aug. 2022. URL [https://doi.org/10.5281/
zenodo.6957644](https://doi.org/10.5281/zenodo.6957644).
- [71] G. A. Ruetenik, K. L. Ferrier, J. R. Creveling, and M. Fox.
Sea-level responses to rapid sediment erosion and deposi-
tion in taiwan. *Earth and Planetary Science Letters*, 538:
116198, 2020.
- [72] E. Ryan, S. Smithers, S. Lewis, T. Clark, J.-X. Zhao, and
Q. Hua. Fringing reef growth over a shallow last inter-
glacial reef foundation at a mid-shelf high island: Hol-
bourne island, central great barrier reef. *Marine Geology*,
398:137–150, 2018.
- [73] A. Sansoleimani, G. E. Webb, D. L. Harris, S. R. Phinn,
and C. M. Roelfsema. Antecedent topography and active
tectonic controls on holocene reef geomorphology in the
great barrier reef. *Geomorphology*, 413:108354, 2022.

- 885 ISSN 0169-555X. doi: [https://doi.org/10.1016/j.geomorph.](https://doi.org/10.1016/j.geomorph.2022.108354) 940
 886 2022.108354. URL [https://www.sciencedirect.](https://www.sciencedirect.com/science/article/pii/S0169555X22002471) 941
 887 [com/science/article/pii/S0169555X22002471](https://www.sciencedirect.com/science/article/pii/S0169555X22002471). 942
- 888 [74] A. Schaeffer and S. Lebedev. Imaging the north american 943
 889 continent using waveform inversion of global and usarray 944
 890 data. *Earth and Planetary Science Letters*, 402:26–41, 945
 891 2014. 946
- 892 [75] A. J. Schaeffer and S. Lebedev. Global shear speed struc- 947
 893 ture of the upper mantle and transition zone. *Geophys- 948
 894 ical Journal International*, 194:417–449, 2013. ISSN 949
 895 0956540X. doi: 10.1093/gji/ggt095. 950
- 896 [76] I. Shennan, A. J. Long, and B. P. Horton. *Handbook of 951
 897 sea-level research*. John Wiley & Sons, 2015. 952
- 898 [77] N. Simmons, S. Myers, G. Johannesson, E. Matzel, and 953
 899 S. Grand. Evidence for long-lived subduction of an an- 954
 900 cient tectonic plate beneath the southern Indian Ocean. 955
 901 *Geophysical Research Letters*, 42(21):9270–9278, 2015. 956
- 902 [78] A. Skrivanek, J. Li, and A. Dutton. Relative sea- 957
 903 level change during the last interglacial as recorded 958
 904 in bahamian fossil reefs. *Quaternary Science 959
 905 Reviews*, 200:160 – 177, 2018. ISSN 0277- 960
 906 3791. doi: [https://doi.org/10.1016/j.quascirev.2018.](https://doi.org/10.1016/j.quascirev.2018.09.033) 961
 907 09.033. URL [http://www.sciencedirect.com/](http://www.sciencedirect.com/science/article/pii/S0277379118304384) 962
 908 [science/article/pii/S0277379118304384](http://www.sciencedirect.com/science/article/pii/S0277379118304384). 963
- 909 [79] M. Spalding, M. D. Spalding, C. Ravilious, and E. P. Green. 964
 910 *World atlas of coral reefs*. Univ of California Press, 2001. 965
- 911 [80] B. Steinberger, S. C. Werner, and T. H. Torsvik. Deep 966
 912 versus shallow origin of gravity anomalies, topography 967
 913 and volcanism on Earth, Venus and Mars. *Icarus*, 207(2): 968
 914 564–577, 2010. 969
- 915 [81] S. N. Stephenson, N. J. White, T. Li, and L. F. Robinson. 970
 916 Disentangling interglacial sea level and global dynamic 971
 917 topography: Analysis of madagascar. *Earth and Planetary 972
 918 Science Letters*, 519:61–69, 2019. 973
- 919 [82] C. Stirling, T. Esat, K. Lambeck, and M. McCulloch. Tim- 974
 920 ing and duration of the last interglacial: evidence for a 975
 921 restricted interval of widespread coral reef growth. *Earth 976
 922 and Planetary Science Letters*, 160(3-4):745–762, 1998. 977
- 923 [83] UNEP-WCMC, WorldFish Centre, WRI, TNC. Global 978
 924 distribution of warm-water coral reefs, compiled from mul- 979
 925 tiple sources including the millennium coral reef mapping 980
 926 project. version 4.0, 2010. [http://data.unep-wcmc.](http://data.unep-wcmc.org/datasets/1) 981
 927 [org/datasets/1](http://data.unep-wcmc.org/datasets/1). 982
- 928 [84] O. Van de Plassche. *Sea-level research: A manual for the 983
 929 collection and evaluation of data: Norwich. UK, Geobooks,* 984
 930 1986. 985
- 931 [85] C. Waelbroeck, L. Labeyrie, E. Michel, J.-C. Duplessy, J. F. 986
 932 Mcmanus, K. Lambeck, E. Balbon, and M. Labracherie. 987
 933 Sea-level and deep water temperature changes derived 988
 934 from benthic foraminifera isotopic records. *Quaternary 989
 935 science reviews*, 21(1-3):295–305, 2002. 990
- 936 [86] R. Walker, M. Telfer, R. Kahle, M. Dee, B. Kahle, J.-L. 991
 937 Schwenninger, R. Sloan, and A. Watts. Rapid mantle- 992
 938 driven uplift along the angolan margin in the late quater- 993
 939 nary. *Nature Geoscience*, 9(12):909–914, 2016. 994
- [87] J. X. W. Wan, N. Gomez, K. Latychev, and H. K. Han. 940
 Resolving glacial isostatic adjustment (gia) in response to 941
 modern and future ice loss at marine grounding lines in 942
 west antarctica. *The Cryosphere*, 16(6):2203–2223, 2022. 943
- [88] M. Wolstencroft, Z. Shen, T. E. Törnqvist, G. A. Milne, 944
 and M. Kulp. Understanding subsidence in the mississippi 945
 delta region due to sediment, ice, and ocean loading: In- 946
 sights from geophysical modeling. *Journal of Geophysical 947
 Research: Solid Earth*, 119(4):3838–3856, 2014. 948
- [89] H. B. Woo, M. P. Panning, P. N. Adams, and A. Dut- 949
 ton. Karst-driven flexural isostasy in north-central florida. 950
Geochemistry, Geophysics, Geosystems, 18(9):3327–3339, 951
 2017. 952
- [90] R. Wood. *General Evolution of Carbonate Reefs*, 953
 pages 452–469. Springer Netherlands, Dordrecht, 954
 2011. ISBN 978-90-481-2639-2. doi: 10.1007/ 955
 978-90-481-2639-2_16. URL [https://doi.org/10.](https://doi.org/10.1007/978-90-481-2639-2_16) 956
[1007/978-90-481-2639-2_16](https://doi.org/10.1007/978-90-481-2639-2_16). 957