

Cover Letter for "*Volcanic Lightning and Prebiotic Chemistry on the Early Earth*" by Jeffrey L. Bada, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA 92024, jbada@ucsd.edu

This is a Matters Arising submission to *Nature* which was received on 12 September, 2022. It is presently under consideration. The manuscript is related to the recent paper by Pan, Z., Mao, F., Rosenfeld, D, et. al., "Coarse sea spray inhibits lightning." *Nature Communications* 13.1, 1-7 (2022).

Abstract: Based on the paper by Pan et al., on the early Earth >4.2 Ga with limited exposed land areas, coupled with an ice covered ocean, lightning could have been rare. This presents a conundrum because lightning is considered to be an important energy source needed for the synthesis of prebiotic compounds required for the origin of life. Lightning occurrence during eruptions on wide spread volcanic islands coupled with their associated release of reduced gases provides a solution.

The manuscript has 1030 words including references (#13) and has one Figure. It has not been submitted or presented elsewhere.

Matters Arising Contribution 9-12-2022

Volcanic Lightning and Prebiotic Chemistry on the Early Earth

Jeffrey L. Bada

Scripps Institution of Oceanography

University of California, San Diego

La Jolla, CA 92024

Abstract: Based on the paper by Pan et al., on the early Earth >4.2 Ga with limited exposed land areas, coupled with an ice covered ocean, lightning could have been rare. This presents a conundrum because lightning is considered to be an important energy source needed for the synthesis of prebiotic compounds required for the origin of life. Lightning occurrence during eruptions on wide spread volcanic islands coupled with their associated release of reduced gases provides a solution.

Pan et. al.¹ recently reported that lightning is rare over the oceans because of sea salt spray interactions. The sea salt caused depression of lightning is reflected in the average global lightning² of 44 ± 5 flashes/sec (fl s^{-1}), with the land/ocean occurrence being 10:1. However, on erupting volcanic islands, lightning can be much more intense. For example, the huge eruption on January 15, 2022 of Hunga Tonga–Hunga Ha‘apai, a submarine volcano 65 km (40 mi) north of Tongatapu on Tonga’s main island, produced enormous volcanic lightning³ with a stunning >400,000 flashes in total, with 200,000 flashes (56 fl s^{-1}) during the most active hour!

Continental, island and coastal regions constitute 88% of the total global lightning on the modern Earth⁴. However, in the Hadean eon (>4.2 Ga) there was likely only a limited amount (~20%) of exposed sub-arial continental area⁵. The Hadean was thus truly a ‘water world’. If the same sea salt caused interference existed then, global lightning could have been significantly less than today. Also, because of reduced solar luminosity, the early oceans could have been ice covered⁶. Lightning over the Arctic and Antarctica today is infrequent⁷. Thus, it’s possible there was a significantly reduced level of overall global lightning on the early Earth.

This presents a conundrum: Lightning is considered to have played a role in the synthesis of organic compounds considered essential for the origin of life⁸. If lightning on the early Earth was relatively rare, this could in turn have limited the production of prebiotic organic compounds. However, there were likely sub-aerial volcanic islands present where lightning would have been common⁹. As shown above, the Hunga Tonga-Hunga Ha'apai eruption had a peak fl^{-2} somewhat greater than that over the entire Earth during its most vigorous hour of the eruption. The occurrence of volcanic island lightning scales with the island size⁹, with the maximum being in the 10^2 to 10^3 km^2 range, perhaps similar to those on the early Earth.

The island volcanoes also would have expelled water and reduced gases such as carbon monoxide, hydrogen sulfide and hydrogen^{10,11}. With the ash, water, reduced gases and lightning, important prebiotic reagents such as hydrogen cyanide, phosphide, aldehydes and ketones as well as some complex compounds could have been generated^{8,12}. The discharged water would have produced lakes, pools and ponds on volcano flanks¹². These prebiotic reagents would have been rinsed off by precipitation and accumulated in these water reservoirs. There, with periodic evaporative concentration, prebiotic syntheses would have taken place, producing important compounds such as amino acids and nucleobases^{8,12}.

Volcanic islands today tend to be short-lived.¹³ Whether this was the case on the early Earth is unknown. Nevertheless, eventually the accumulated prebiotic products would have rinsed off into the oceans during island subsidence. There, further prebiotic molecular evolution could have taken place. Where and when the emergence of life took place in this scenario requires further refinement of the underlying mechanisms and processes. But one thing is certain: "Thunder is good, thunder is impressive; but it is lightning that does the work. *Mark Twain*".

1. Pan, Z., Mao, F., Rosenfeld, D, et. al., "Coarse sea spray inhibits lightning." *Nature Communications* 13.1, 1-7 (2022).
2. Christian, H. J., Blakeslee, R. J., Boccippio, D. J., et. al., Global frequency and distribution of lightning as observed from space by the Optical Transient Detector. *J. Geophys, Res.: Atm.*, 108(D1), ACL-4 (2003).
3. Yuen, D. A., Scruggs, M. A., Spera, F. J., et. al., "Under the surface: Pressure-induced planetary-scale waves, volcanic lightning, and gaseous clouds caused by the submarine eruption of Hunga Tonga-Hunga Ha'apai volcano." *Earthquake Res. Adv.*, 100134 (2022).
4. Williams, E., Chan, T., Boccippio, D., Islands as miniature continents: Another look at the land-ocean lightning contrast. *Journal of Geophysical Research: Atmospheres*, 109(D16) (2004).
5. Korenaga, J., "Was There Land on the Early Earth?" *Life* <https://doi.org/10.3390/life11111142> (2021).
6. Bada, J. L., Bigham, C., Miller, S. L. "Impact melting of frozen oceans on the early Earth: implications for the origin of life." *Proc. Nat. Acad. Sci.*, 91(4), 1248-1250 (1994).
7. Holzworth, R. H., Brundell J. B. et. al. "Lightning in the Arctic." *Geophys. Res. Lett.* 48, no. 7: e2020GL091366 (2021) and Yang Chen per, comm.).
8. Bada, J. L. "New insights into prebiotic chemistry from Stanley Miller's spark discharge experiments" *Chem. Soc. Rev.*, 42(5), 2186-2196 (2013).
9. Yuen. D., Scruggs, M. A., et. al., Under the surface: Pressure-induced planetary-scale waves, volcanic lightning, and gaseous clouds caused by the submarine eruption of Hung-a Tonga-Hunga Ha'apai volcano, *Earthquake Res. Adv.*, 2(3), (2022),
10. Bada, J. L., Korean, J. "Exposed areas above sea level on Earth >3.5 Gyr ago: Implications for prebiotic and primitive biotic chemistry." *Life* 8.4, 55-64 (2018).
11. Hess, B. L., Piazzolo, S., & Harvey, J. (2021). Lightning strikes as a major facilitator of prebiotic phosphorus reduction on early Earth. *Nature Comm.*, 12(1), 1-8.
12. Prosdocimi, F., Torres de Farias, S., José, M. J., "Prebiotic chemical refuge: multifaceted scenario for the formation of biomolecules in primitive Earth." *Theory in Biosciences* (2022): 1-9.

13. Kimberly L., J., "Hotspot swells and the lifespan of volcanic ocean islands." *Sci. Adv.* 6.1: eaaw6906. (2020).

Acknowledgments: I thank Steve McNutt, Melissa Scruggs, and Robert Holzworth for helpful comments and suggestions. Canon E. J. Vossen for pointing out the Mark Tain quote.



Figure: A small eruption of Mount Rinjani, with volcanic lightning on the island of Lombok in Indonesia. Photo credit: Oliver Spalt, Taken from Wikimedia Commons.