

Geodetic Evidence for a Buoyant Mantle Plume Beneath the Eifel Volcanic Area, NW Europe

Corné Kreemer*

Nevada Bureau of Mines and Geology and Nevada Seismological Laboratory

University of Nevada, Reno

1664 N. Virginia Street, Reno, NV 89557-0178, USA

kreemer@unr.edu

ORCID: 0000-0001-6882-9809

Geoffrey Blewitt

Nevada Bureau of Mines and Geology

University of Nevada, Reno

1664 N. Virginia Street, Reno, NV 89557-0178, USA

gblewitt@unr.edu

ORCID: 0000-0002-7490-5983

Paul M. Davis

Department of Earth, Planetary, and Space Sciences

University of California Los Angeles

595 Charles Young Drive East, Box 951567, Los Angeles, CA 90095-1567, USA

pdavisucla@gmail.com

Summary

The volcanism of the Eifel volcanic field (EVF), in west-central Germany, is often considered an example of hotspot volcanism given its geochemical signature and the putative mantle plume imaged underneath. EVF's setting in a stable continental area provides a rare natural laboratory to image surface deformation and test the hypothesis of there being a thermally buoyant plume. Here we use Global Positioning System (GPS) data to robustly image vertical land motion (VLM) and horizontal strain rates over most of intraplate Europe. We find a spatially-coherent positive VLM anomaly over an area much larger than the EVF and with a maximum uplift of $\sim 1 \text{ mm yr}^{-1}$ at the EVF (when corrected for glacial isostatic adjustment). This rate is considerably higher than averaged over the Late-Quaternary. Over the same area that uplifts, we find significant horizontal extension surrounded by a radial pattern of shortening, a superposition that strongly suggests a common dynamic cause. Besides the Eifel, no other area in NW Europe shows significant positive VLM coupled with extensional strain rates, except for the much broader region of glacial isostatic adjustment. We refer to this 3D deformation anomaly as the Eifel Anomaly. We also find an extensional strain rate anomaly near the Massif Central volcanic field surrounded by radial shortening, but we do not detect a significant positive VLM signal there. The fact that the Eifel Anomaly is located above the Eifel plume suggests the plume causes the anomaly. Indeed, we show that buoyancy forces induced by the plume at the bottom of the lithosphere can explain this remarkable surface deformation. Plume-induced deformation can also explain the relatively high rate of regional seismicity, particularly along the Lower Rhine Embayment.

Keywords: Satellite geodesy – Europe – Continental tectonics: extensional – Intra-plate processes – Dynamics: convection currents, and mantle plumes

1. Introduction

Intraplate volcanic activity in west-central Europe has long been associated with mantle upwellings (Granet *et al.* 1995, Hoernle *et al.* 1995). Most effort has focused on the Eifel Volcanic Field (EVF) where a period of late Quaternary volcanism (Fig. 1) continued until ~11 ka and included the explosive eruption of Laacher See in the eastern EVF at 12.9 ka (Nowell *et al.* 2006, Schmincke 2007). Although the volcanism does not exhibit a clear space-time progression indicative of a hotspot track, geochemical analyses (Griesshaber *et al.* 1992, Hoernle *et al.* 1995, Aeschbach-Hertig *et al.* 1996, Wedepohl & Baumann 1999, Buikin *et al.* 2005, Bräuer *et al.* 2013, Caracausi *et al.* 2016) have shown that EVF (and some other central European) volcanic rocks and gases have the characteristics of a mantle source, while seismological studies have shown evidence for a mantle plume underneath the Eifel (Ritter 2007). Specifically, there exists a low seismic velocity anomaly up to ~410 km depth (Ritter *et al.* 2001, Keyser *et al.* 2002, Pilidou *et al.* 2005, Budweg *et al.* 2006, Montelli *et al.* 2006, Zhu *et al.* 2012). Some studies have imaged a low seismic velocity anomaly underneath this location in the lower mantle (Goes *et al.* 1999, Grunewald *et al.* 2001, Zhao 2007), but the 660 km discontinuity seems unaffected (Budweg *et al.* 2006), which suggests that there is no physical connection between the lower and upper mantle anomalies. There is evidence for a broader low velocity zone at ~50 km depth that could be interpreted as the plume head (Budweg *et al.* 2006, Mathar *et al.* 2006), which is consistent with the lithosphere-asthenosphere boundary (LAB) underneath the Eifel being relatively shallow at ~40-50 km depth (Plomerová & Babuška 2010, Seiberlich *et al.* 2013). For the Massif Central (France), the other major Quaternary volcanic field in western Europe (with the most recent eruption at ~7 ka of Lac Pavin (Juvigné & Gilot 1986, Nowell *et al.* 2006, Chapron *et al.* 2010)), some earlier tomographic studies (Granet & Trampert 1989, Granet *et al.* 1995, Sobolev *et al.* 1997) suggested an underlying plume, but only few studies have convincingly reproduced those findings (Spakman & Wortel 2004, Barth *et al.* 2007, Koulakov *et al.* 2009). Limited geochemical analyses have revealed a mantle signature there (Matthews *et al.* 1987, Aeschbach-Hertig *et al.* 1999, Zangana *et al.* 1999).

Despite the above indications suggesting EVF volcanism being the result of decompression melting of a buoyant mantle upwelling, the fact that volcanism seems to flare up at the end of glacial periods (also at Massif Central) has been interpreted as volcanism instead being caused (or modulated) by glacial unloading (Nowell *et al.* 2006). Alternatively, the location of the EVF near the major European Cenozoic Rift System has been used to argue that volcanism is the product of fluid/gas pathways caused by Alpine-collision induced dilatancy along shear bands in the upper mantle and lower crust (Regenauer-Lieb 1998) and/or passive partial melting of the asthenospheric mantle induced by lithospheric stretching (Wilson & Downes 1992, Lustrino & Carminati 2007).

If a thermally buoyant plume is present, it is predicted to cause significant domal uplift at the surface, although the amount of uplift and spatial extent thereof depend on the stage of upwelling, temperature contrast, viscosity, width of the plume (head) (Griffiths & Campbell 1991, Hill 1991), the plume's inherent composition/density (Dannberg & Sobolev 2015), and lateral strength variations in the lithosphere induced by thermal weakening (Garcia-Castellanos *et al.* 2000). Vertical strength stratification within the lithosphere may even yield topographic undulations rather than a singular dome (Burov & Guillou-Frottier 2005, Burov & Gerya 2014), and one study argued this to be the case for the Eifel (Guillou-Frottier *et al.* 2007). By measuring the rate and spatial extent of present-day surface motion, the specifics and mere existence of a buoyant Eifel or Massif Central plume can be assessed. More generally, northwestern Europe provides a unique natural laboratory to contrast current surface

deformation with plume model predictions, because other active hotspots are either underneath oceans or in areas that are tectonically and/or volcanically too active to assess the secular deformation (i.e., Yellowstone and Afar).

Hints of significant present-day uplift of the central Rhenish Massif (i.e., the area surrounding the EVF, Fig. 1) have existed for decades (Mälzer *et al.* 1983), but inconsistencies between different geodetic studies in the surrounding area have put into question whether measured VLM even reflected tectonic movement (Demoulin & Collignon 2000, Camelbeeck *et al.* 2002, J. Campbell *et al.* 2002, Demoulin & Collignon 2002, Francis *et al.* 2004). Long-running absolute gravity measurements revealed significant regional uplift at a number of stations in the Belgium part of the Rhenish Massif (i.e., Ardennes) (Van Camp *et al.* 2011), but that study lacked the density and broad spatial extent of measurements, nor the horizontal sensitivity, that our study provides.

The results of the 3D surface motion in NW Europe presented here are part of a larger study encompassing most of intraplate Europe (Fig 2.). By investigating both vertical and horizontal deformation, by using robust estimations techniques (of the model and their uncertainties), and by doing this over most of intraplate Europe, our study is better suited to detect significant regional anomalies than previous studies, which had limited spatial reach and/or focused on only vertical (Kontny & Bogusz 2012, Serpelloni *et al.* 2013, Husson *et al.* 2018, Bogusz *et al.* 2019) or horizontal deformation (Ward 1998, Marotta *et al.* 2004, Nocquet *et al.* 2005, Tesauro *et al.* 2006, Bogusz *et al.* 2014, Neres *et al.* 2018, Masson *et al.* 2019). This paper focusses mostly on the only significant 3D deformation anomaly observed in our model (i.e., the ‘Eifel Anomaly’), but results for the Massif Central will be presented as well.

2. GPS Data Analysis

2.1 Processing Details

We processed all the available geodetic-quality GPS data from continuously-operating stations in our study area (red polygon inset Fig.1). Our study benefited tremendously from the data made available to us from many regional commercial and state networks and that made this type of study possible for the first time for all of NW Europe (see Acknowledgements). The GPS data were retrieved from archives in the form of daily station-specific RINEX files, which contain raw, dual-frequency carrier phase and pseudorange data, typically for every 15- or 30-second epoch. We then reduced the data for each station to a time-series of daily precise point positions (Zumberge *et al.* 1997) using the GipsyX version 1.0 software (released January 2019) licensed by the Jet Propulsion Laboratory (JPL), together with JPL's high-precision GPS orbit and clock products. Data were processed for each station independently, and thus the results are insulated from potential data problems at other individual stations. Metadata required for correct processing options were provided by the RINEX header information, corrected by custom-software and alias tables for commonly known errors such as field misplacement, typos, and non-standard receiver-type and antenna-type fields. One exception is that metadata from the International GNSS Service (IGS) was used for antenna types if available, which is the case for approximately 4% of the data. Another exception is that we used our own metadata for approximate coordinates, which should be accurate to <10 meters to ensure the validity of linearised observation equations. After editing the data using the TurboEdit algorithm (Blewitt 1990) available with GipsyX, the GPS data were reduced to 300-second epochs using decimation for carrier phase, and using carrier-smoothing for pseudorange.

In addition to the RINEX files, we used essential input data produced by JPL, including precise GPS orbits, Earth orientation, eclipse shadow events, clock offsets, and WLPB biases (Bertiger *et al.* 2010) for carrier phase ambiguity resolution. The satellite orbits were minimally constrained, which requires that we transform the estimated station coordinates every day by a global, 7-parameter transformation into the IGS14 reference frame, using daily values provided by JPL. For modelling ocean tidal loading, we used coefficients of the FES2004 model (Lyard *et al.* 2006) computed by Chalmers University, Sweden, produced by the email interface to the server described at <http://holt.oso.chalmers.se/loading/>. For modelling neutral atmospheric delay (commonly known as “tropospheric delay”, but includes stratospheric delay), we used VMF1GRID gridded map products from University of Vienna, Austria (Boehm *et al.* 2006), which is based on the ECMWF numeric weather model based on global meteorological data. VMF1GRID allowed us to calibrate for nominal so-called dry and wet delays, and epoch-dependent mapping function parameters for each station. For first-order ionospheric calibration, GipsyX implements the so-called ionosphere-free linear combination of observables. For higher-order ionospheric calibration, we used JPL’s IONEX gridded map product available in daily files, together with the NOAA’s 12th generational magnetic field model IGRF12. For antennas, we used calibrations made available by the IGS (Schmid *et al.* 2007). Following IGS standards, if a calibration was not available for a particular antenna, we did not use the data. If a calibration was not available for particular radome, we used the no-radome calibration for an antenna. Internally, the GipsyX software implements JPL’s planetary ephemeris to compute tidal effects.

The estimation strategy was set by default according to the distributed version of GipsyX Version 1.0, which we now describe. Satellite positions and clocks are held to their nominal values, whereas station clocks are estimated freely as white noise every 300-s epoch. Carrier phase biases are initially estimated freely as constants in between detected cycle-slips (integer-wavelength discontinuities), then double-difference carrier phase biases to IGS stations (using the WLPB file produced daily by JPL) are constrained to integer-wavelength values on both frequencies (which are linearly combined for the ionosphere-free bias). Parameters of the neutral atmosphere are estimated as random walk processes at every 300-s epoch. These parameters include wet zenith delay (with constraints at 5 cm per square-root hour), and two gradients parameters (with constraints at 5 mm per square-root hr). The three station Cartesian coordinates were estimated as a constant over each 24 hour period. Data are processed with several iteration cycles to clean up the data using a series of post-fit residual outlier tests set by default.

2.2 Time-series analysis

The output of each station-day process includes a 24-hour estimate of a constant station position in the IGS14 frame which is obtained by using daily 7-parameter transformations produced by JPL. The IGS14 reference frame is computed by the International GNSS Service (IGS) as a GPS-compatible realization of the International Terrestrial Reference Frame ITRF2014 (Altamimi *et al.* 2016). For this study we consider 3D position time-series between 1 January, 2000, and 5 October, 2019, but use only stations for which the time-series span at least 2.5 year and for which there are at least 304 data points (which translates into a theoretical minimum 33% and 4.2% completeness for the shortest and longest time-series considered). We estimate velocities for 2383 stations (Fig. 2), of which 1414 are in the area presented here (blue polygon inset Fig.1).

The GPS velocities are estimated in a multi-step process. Firstly, we fit to the position time-series a station motion model that includes an intercept, trend, annual and semi-annual periodic signal, and

offsets due to known equipment changes or other unknown causes. We then iteratively identify position outliers in the residual time-series and remove them from the position time-series. Outliers are being defined as a position that exceeds 3σ from the median, and σ is defined as $1.4826 \times MAD$, where MAD (‘median absolute deviation’) is the absolute deviation around the median, i.e., $\text{median}(|res(t) - \text{median}(res(t))|)$ (Gauss 1816, Hampel 1974). With this definition we follow the general suggestion by Leys *et al.* (2013) and a GPS time-series specific practice advocated by Klos *et al.* (2016). The 1.4826 factor is there so that σ equals the traditional standard deviation describing the variance in $res(t)$ in case $res(t)$ is normally distributed (Huber 1981).

Next, the cleaned residual time-series are used to construct time-series of regional common-mode components (CMC), which are removed from the original time-series. CMC removal is important because otherwise strain rates or VLM anomalies can arise from combining velocities inferred from different time-spans for which the time-series can be biased by long-period CMC signals (Márquez-Azúa & DeMets 2003, Serpelloni *et al.* 2013) (Supplemental Fig. S3 and S8). The CMC estimation procedure used here is described in (Kreemer 2020); it uses cross-correlation-based weights, is explicitly robust against outlier data, and takes advantage of all data (including stations with short time-series). The key features are: 1) the CMC of stations with longer time-series are used to correct the shorter time-series of other stations in a hierarchical scheme, 2) as a result, all stations can potentially be ‘filter stations’ (although we identify outliers) and the precision and the spatio-temporal appropriateness of the CMC is dictated by the spatio-temporal density of the data, and 3) each step in the procedure is based on robust median statistics.

After we fit the station model once more to the ‘filtered’ time-series, and we correct the offsets, we estimate the final trend and uncertainty using the robust MIDAS algorithm (Blewitt *et al.* 2016). MIDAS-derived velocity uncertainties are typically similar to, and tend to be more conservative than those derived from stochastic models (Santamaría-Gómez *et al.* 2017, Simon *et al.* 2018, Murray *et al.* 2019).

To help constrain the far-field glacial isostatic adjustment (GIA) signal, we add 80 published velocities in southern Scandinavia (Kierulf *et al.* 2013, 2014, Lahtinen *et al.* 2019). We also exclude three stations located above abandoned coal mines in the Kempen (Belgium) and Limburg (The Netherlands) areas and which experience anomalously fast uplift due to underground groundwater flooding (Caro Cuenca *et al.* 2013, Vervoort 2016).

3. Methodology

3.1 VLM Imaging

We interpolate the GPS station vertical velocities to obtain a robust VLM map using a method that is based on the GPS Imaging algorithm of Hammond *et al.* (2016), but differs in some aspects. The major difference is that we do not use a spatial structure function to determine how the weight the rates of stations as function of their distance from a station/point-of-interest. Such function implies a similar spatial coherence in vertical rates across the entire study area. This is an inappropriate assumption here, because our study area includes both potential long-wavelength (e.g., GIA (e.g., Nocquet *et al.* 2005, Husson *et al.* 2018)) and short wavelength (e.g., subsidence above the Groningen gas field (Ketelaar 2009)) signals. Instead, we aim to consider a set of stations that are as local as possible. We do this by applying a Delaunay triangulation of the station locations (Renka 1997). To reduce the influence of

stations that are relatively far away, we estimate the median distance from the station/point-of-interest to all neighbouring stations (in a Delaunay sense) and include the rates of additional stations in the median estimation when the distance to those stations is less than the median distance. In case the rate of a relatively far away station is anomalous, the addition of these extra stations would effectively down-weight the outlier (even more so than a typical median approach of the neighbouring stations would already do). Moreover, the addition of other nearby stations would make the median as local as possible and constrained by the density of stations and spatial variation therein.

In practice, the addition of extra station rates is not effective when station density is rather uniform, and in our case it matters only for places where the station density changes laterally, such as along coast lines. It is also important to note that in the algorithm of Hammond *et al.* (2016) one needs to subjectively choose the weight of the station itself, while here the station itself has objectively the same weight as the other nearby stations (i.e., in case the standard deviations in the observations are the same, see below). Another important difference between our algorithm and that of Hammond *et al.* (2016) is how we estimate uncertainties and how we estimate the weighted median (see below)

In short, our alternative GPS Imaging approach (which we call “Robust Network Imaging” (RNI) since it could be applied to any type of geospatial network data) includes the following steps. First, the vertical rate of each station i (x_i) is replaced by a weighted median (or ‘despeckled’) value (\hat{x}_i) derived from the rates of the N most local stations. These most local stations include the station itself, those connected to it in a Delaunay scheme, and any other station that is closer to the station itself than the median distance between the station itself and those connected to it. The weights are given by $1/\sigma_i$, where σ_i is the standard deviation in the observed rate. The weighted median algorithm is based on that most recently presented by Bowden *et al.* (2016), and also implemented by Kreemer (2020). For the standard deviation in \hat{x}_i we use the MAD, similarly as was defined for the outlier detection.

In a second step we use these despeckled rates and standard deviation therein ($\hat{x}_i, \hat{\sigma}_i$) to estimate the weighted median vertical rate at a gridded set of M evaluation points j within the convex hull of the station locations. A new Delaunay triangulation is calculated on a set of points that includes the evaluation point together with all station locations. The weighted median \hat{x}_j is then estimated from \hat{x}_i and $\hat{\sigma}_i$ of all local stations defined by the same procedure as in the first step. The standard deviation for the median at the evaluation point $\hat{\sigma}_j$ is estimated similarly as above. For this we use the absolute deviation from typically the original observed rates (i.e., x_i) which would yield conservative uncertainties, but occasionally (Supplemental Fig. S2b and the dark shaded blue in Fig. 6) from the despeckled values (\hat{x}_i), which would typically yield a smaller uncertainty that reflects the spatial variation in the imaged VLM rather than in the underlying data. Either way, these robust estimates of standard deviation differ from those in the method of Hammond *et al.* (2016), which were either defined as the root-mean-square residual scatter in x_i , or as the weighted mean of x_i .

In practice, the Delaunay triangulation does not work if two stations are exactly collocated. We therefore allow for station coordinates and rates to be averaged when stations are within 1 m distance from each other. This is done before RNI starts. Also, when we add stations that are within a median distance, we add 100 m to that distance so that, when the median distance is in fact the distance to a station that is nominally collocated with others, the rates of those stations are considered as well.

3.2 Strain Rate Imaging

The horizontal strain rates are estimated with a modification of the ‘MELD’ (Median Estimation of Local Deformation) imaging algorithm of Kreemer *et al.* (2018). As was shown by Kreemer *et al.* (2018), MELD is very robust against outlier velocities, and gives realistic uncertainties. MELD-derived strain rates are derived at an evaluation point from the multivariate median of a set of strain rates from a number of station-based local triangles (see Kreemer *et al.* (2018)) and requires two parameters. The first one is the minimum number of triangles to use for estimating the strain rate (N_{\min}). We use here $N_{\min}=56$, which can theoretically be reached with as few as 6 stations. The second MELD input parameter is based on the theoretical standard deviation in strain rate in a triangle of stations based on the triangle’s geometry and size. The maximum allowed value (σ_{\max}) is used to exclude triangles that are too small or skinny, which would make the strain rate estimate uncertain. We set $\sigma_{\max}=11.1364$, which was chosen such that noise of the amplitude of the median 1-D standard deviation in the GPS velocities ($\sim 0.11 \text{ mm yr}^{-1}$) does not result in strain rates $> 1 \times 10^{-9} \text{ yr}^{-1}$. That is, no strain rates from a triangle of stations are considered if the standard deviation in strain rate in any one component exceeds σ_{\max} . In general, $\sigma_{\max} \approx 1.225/\sigma_{\text{GPS}}$. An equilateral triangle that yields a standard deviation in strain rate of $11.1364 \times 10^{-9} \text{ yr}^{-1}$ has an area with equivalent circle radius of 47.2 km, which is thus our minimum theoretical spatial resolution. The general relationship between σ_{\max} and this minimum radius (R_{\min}) is $R_{\min} \approx 525.2/\sigma_{\max}$.

We make one change to the MELD algorithm as presented by Kreemer *et al.* (2018). This change is applied after the strain rate at each evaluation point (typically part of a grid of points) is estimated. Because the strain rates are based on triangles with areas equivalent to circles with radii typically larger than the distance between evaluation points, we replace the results at the evaluation points with a weighted spatial average based on all evaluation points within a distance R_{median} from the evaluation point considered (including the point itself). Here R_{median} is the corresponding radius for the median area of all triangles considered at a point. R_{median} is by definition $\geq R_{\min}$. The spatial smoothing not only better aligns the spatial resolution of the model results to a corresponding spatial smoothness, the smoothed field also minimizes significant model differences between two neighbouring evaluation points which could exist if each of those points are inside a different station triangle which could have yielded a significantly different set of stations to be considered. Appendix A details the estimation of standard deviation in each spatial average given that model parameters between neighbouring grid points are derived from many common velocity data yielding a large degree of correlation.

4. Results

4.1 Vertical land motion

Observed VLM (Fig. 3a) for individual GPS stations shows some regional patterns, but also large variability. In a first step, we pre-process the VLM by applying our RNI algorithm. Based on this despeckled VLM field (Fig. 3b), we then use RGI again to estimate the VLM at a grid of evaluation points (Fig. 3c). Offshore areas are clipped in the figures, because checkerboard tests (Fig 4. and Fig. S1) reveal an expected lack of resolution there. The same tests show that at the Eifel area $200 \times 200 \text{ km}$ and $100 \times 100 \text{ km}$ VLM patches (i.e., Fig 4. and Fig. S1, resp.) can be significantly recovered at the 2σ , but that at the Massif Central area only the larger patches can be resolved only at the 1σ level, because of there being fewer stations there compared to the Eifel area. Because the results presented here are

part of a larger study that covers most of intraplate western Europe (Fig. 2), they do not suffer from boundary effects.

We observe an area of positive VLM over most of the Rhenish Massif, including the EVF, and it is centred slightly northerly thereof (Fig. 3c). This VLM signal is anomalous given that most of intraplate Europe south of Scandinavia is subsiding, which likely reflects forebulge collapse related to GIA. Given that GIA models can differ significantly, we choose to subtract from our results the VLM predicted by the GIA model that is most consistent with our regional observations of the rate of forebulge collapse (Husson *et al.* 2018). When we present our result relative to this GIA model, we find that the EVF uplift anomaly is the only coherent significant signal (at the 2σ level) in NW Europe (Fig. 3d). The area of anomalous uplift covers a roughly circular/oval area and includes most of the west-central Rhenish Massif as well as southeastern Netherlands (i.e., Limburg). The highest uplift is slightly above 1 mm/yr relative to the regional GIA-associated subsidence. While VLM at the Massif Central is higher than in surrounding areas in France, the VLM signal is insignificant (that is, at the 2σ level, but even so at the 1σ level, which suggest there either does not exist a significant VLM anomaly or the anomaly is much less than 200×200 km in scale for which our model lacks resolution (Fig. S1)).

Note that our study also contains a couple areas of anomalous subsidence (i.e., >1 mm yr⁻¹). Examples are the western Paris Basin, the western part of The Netherlands, and the northernmost part of The Netherlands (Groningen). Subsidence in the western part of The Netherlands has previously been observed with InSAR and is explained by peat decomposition (Caro Cuenca & Hanssen 2008). Subsidence in Groningen has previously also been detected by InSAR (Ketelaar 2009) and is due to gas extraction (van Thienen-Visser & Breunese 2015, Jagt *et al.* 2017). This anthropogenically-induced subsidence is the only significant signal (at the 2σ level) in our model, with imaged subsidence up to ~ 6 mm yr⁻¹ (not corrected for GIA).

4.2 Horizontal deformation

In an independent analysis from the VLM, we use the velocities derived from the horizontal time-series of the same set of GPS stations used in the VLM analysis to infer the horizontal strain rate field (a subset of the velocities is shown in Fig. 5a). For all models the results are estimated on a 0.1° grid of evaluation points. For about all areas in our preferred model the strain rates represent deformation for an area with ~ 50 - 60 km radius (Supplemental Fig. S4a) and this is inherently controlled by the station spacing and, more importantly, our choice to exclude strain rate estimates based on stations that are close together. The uncertainty in the dilatational strain rates is shown in Supplemental Fig. S4b.

We visualize the results in terms of style (Fig. 5b) and the dilatational amplitude of the strain rate field (Fig. 5c-d). We show the dilatational strain rates, because in most places the strain rate tensor in the study area is dominated by either extension or contraction. A large part of the area is dominated by contractional strain rates, which can be explained by contraction of the GIA forebulge, as evidenced by the systematic rotation of the contraction direction in the regions directly surrounding the former ice-sheet. In the southeastern part of our model area, i.e., southern Germany, we see enhanced contraction that could be associated with Alpine shortening protruding into intraplate Europe. The pattern of the wide-scale contraction appears to be interrupted by a dilatational strain rate anomaly centred NNW of the EVF and defined by significant \sim N-S oriented extension and a maximum dilatation rate of

$\sim 3.6 \pm 0.9 \times 10^{-9} \text{ yr}^{-1}$. Furthermore, the orientations of the contractional strain rates in most of the areas directly surrounding the extension anomaly exhibit a radial pattern oriented towards that anomaly. For the Massif Central we find a significant (at the 2σ level) extensional strain rate anomaly just west of the area of the most recent volcanic activity, with extension being bi-axial and a magnitude of $\sim 1-2 \times 10^{-9} \text{ yr}^{-1}$. The pattern of contractional strain rate around this extensional anomaly also shows a radial pattern, as seen for the Eifel, but it is less convincing.

To illustrate the robustness of our results, we also show the results for models with minimum triangle sizes at 50% (i.e. minimum $\sim 23 \text{ km}$ radius, Supplemental Fig. S5) and 150% (i.e., $\sim 71 \text{ km}$, Supplemental Fig. S5) compared to our preferred model. When the spatial scale is reduced, other strain rate features appear but the only significant feature is a high-dilatation feature centred at the same place as the extensional anomaly NNW of the Eifel in our preferred model. When the spatial scale is increased, the same anomaly persists but at a lower rate than in our preferred model. We do observe extensional strain rates $> 2 \times 10^{-9} \text{ yr}^{-1}$ just west of recent volcanism in the Massif Central when we reduce the spatial scale (Fig. S5b), but for that model (as well as for the one with increased spatial scale) the dilatational strain rate anomaly is not significant at the 2σ level (Fig. S5c and S6c).

To illustrate that our preferred MELD-derived model is robust against outlier velocities, we also present a model based on a set of velocities minus outliers and that looks very similar (Supplemental Fig. S7) to the one presented in Fig. 5. Fig. S8 shows a model that is based on velocities derived from time-series that do not have CMC removed. That model shows a significant contractional area of $2-3 \times 10^{-9} \text{ yr}^{-1}$ (with contraction in NS orientation) in northern France, which is also present in the model with the reduced spatial resolution (Fig. S5b), and this anomaly was also recently observed by Masson *et al.* (2019). We now show this anomaly to be mostly insignificant. Masson *et al.*'s observation may either be an artefact of CMC in the time-series possibly not entirely removed by their first-order filtering and/or the fact that they assumed a spatial resolution that is too small (given the data noise) and an underestimation of their uncertainties (given that spatial scale). Finally, in Supplemental Fig. S9 we show, for comparison, a result based on the original MELD algorithm that does not include the added spatial averaging and which inherently more scattered, but still reveals the major anomalies.

5. The Eifel Anomaly

To further illustrate the significance of the VLM result across the Eifel, we plot the results across the profile defined in Fig. 3d (Fig. 6). The profile highlights the broad nature and significance of positive VLM across the broader Eifel/Rhenish Massif area. We define the 'Eifel Anomaly' by the area where we see superimposed significant uplift and significant extensional strain rate, which is the same area where there is enhanced seismicity, the EVF, and which sits above the imaged mantle plume (Fig. 7a). The area undergoing the most significant extension is a bit offset NNW from the area of largest uplift. The centre of the uplift anomaly is, in turn, slightly offset to the north from the projected mantle plume and coincides with the EVF. Across the anomaly, the maximum horizontal separation rate is $\sim 0.33 \text{ mm yr}^{-1}$ (in a roughly NS direction), compared to the maximum uplift of $\sim 1 \text{ mm/yr}$.

6. Plume modelling

We use a simple model to test if a buoyant plume can explain the long-wavelength 3D surface deformation. A rising plume will exert dynamic and buoyancy forces on the elastic lithosphere and we focus here on the buoyancy forces. In order to mimic those forces coming from a mantle plume head, we model them as a bi-modal Gaussian areal distribution of half-space vertical forces (Mindlin 1936,

Anderson 1937) exerted on a plane at depth. The relationship between surface deformation is schematically given as: $[u,v,w]=A(x,y:\sigma_1,\sigma_2,\theta,c)\times F(u,v,w:x,y,c)$. The 3D surface deformation is represented here as displacement rates (u,v,w , being east, north and vertical velocities, respectively), but we also invert for strain rates. The gain function, $A(x,y)$, is a normalized bi-modal Gaussian amplitude distribution with standard deviations (σ_1,σ_2), centred on x_0,y_0 , with a rotation relative the East axis (θ), and a fixed depth of $c=50$ km, and F is the vertical force. This 6-parameter model is fit to the measured displacement and strain rates, with regional trends removed. Removing the regional trends involves linear fits that added another 9 parameters, for a total of 15.

For our best-fitting model, we find the half-widths of the bi-modal Gaussian distribution to be 174 and 98 km in the roughly EW and NS directions, respectively, and the centre is found at 50.5°N 6.4°E. Table 1 lists all the model parameters, and Fig. 8 graphically shows function A . Our simple model adequately fits the long-wavelength measured displacement and strain rates (Fig. 7b). We constrain the plume head to be at 50 km depth (consistent with imaged depth of LAB and plume head (Budweg *et al.* 2006, Mathar *et al.* 2006)), because of trade-offs between the Gaussian widths and the depth of the model. Supplementary Fig. S10 shows the predicted surface deformation for the best fitting model that forces the depth to be at 27 km and 100 km.

7. Discussion and Conclusions

The remarkable superimposition of significant uplift, horizontal extension, and volcanism in the Eifel area strongly suggests a causal relationship with the putative underlying mantle plume. The circular VLM pattern is consistent with the Quaternary uplift but at odds with studies, some specific to the Eifel area, that predict an undulating pattern (Burov & Guillou-Frottier 2005, Guillou-Frottier *et al.* 2007, Burov & Gerya 2014). To first order, most model studies would predict to find uplift (I. H. Campbell 2005, Dannberg & Sobolev 2015) and extension (Burov *et al.* 2007, Cloetingh *et al.* 2013) above a buoyant mantle plume. Indeed, we obtain a good regional fit to the long-wavelength aspects of the surface deformation by applying buoyancy forces related to the plume head at the bottom of the lithosphere. This is the simplest model consistent with seismic evidence, but it should be noted that an actual inversion for the depth is very poorly constrained. For example, placing the force distribution at crustal level could fit our observations too (Supplementary Figure S10). There is, however, no evidence of a regional magmatic sill, although seismic velocities in the lowermost crust underneath most of the Rhenish Massif are found to be significantly reduced (Prodehl *et al.* 2006). We also note that the surface deformation contains some details (such as some sharp edges in the uplift anomaly) which our simple model does not fit. The fit may improve when considering lateral variations in the strength of the lithosphere as caused by plume-induced thermal weakening, which would also require a shallower source (Garcia-Castellanos *et al.* 2000). Furthermore, the asymmetry of the deformation pattern and its offset from the imaged plume could be a consequence of the plume being tilted and/or the interaction between a rising plume and a moving plate (Wüllner *et al.* 2006).

Based on river terrace elevation data, the Rhenish Massif is known to have uplifted since ~700-800 ka (i.e., the same time as Quaternary Eifel volcanism commenced), with maximum uplift between ~140 m (Westaway 2001, Demoulin & Hallot 2009) and 250 m (Van Balen *et al.* 2000, Meyer & Stets 2002) centred on the Eifel area, where we see the highest uplift. These data imply ~0.1–0.3 mm yr⁻¹ of average uplift since ~700–800 ka. Such rate is considerably lower than we find here and could either be evidence that uplift has increased since the onset of the volcanism in the late Quaternary or that the loading/unloading effect related to glacial periods causes the net VLM to vary considerably over time.

From our spatially integrated force and the first-order assumption that the plume has effectively been buoyant since between 250 ka (to explain 250 m Quaternary uplift) and 800 ka (at today's rate) ago, we estimate that a 360 km (i.e., 410 minus 50 km) high plume requires a $\sim 57\text{--}184\text{ kg m}^{-3}$ density reduction (i.e., $\sim 0.7\text{--}5.6\%$ of a 3300 kg m^{-3} dense reference mantle), which is consistent with observed seismic velocity reductions (Ritter 2007).

Although the EVF is found in what is typically referred to as intraplate Europe, the area near the EVF is seismically unusually active (Fig. 1). Much of that seismicity is attributed to faulting within the Lower Rhine Embayment (LRE) (Hinzen & Reamer 2007), and those faults appear to have increased their slip rate since the same time as the onset of late Quaternary uplift and volcanism ($\sim 700\text{--}800$ ka) (Gold *et al.* 2017). In this area of highest seismic activity we also find the highest extension rates. We particularly note that the roughly N–S oriented extension we find above the LRE is favourably orientated to generate extensional earthquakes along the normal faults within the LRE, as evidenced by the orientation of the mostly extensional earthquakes there (Plenefisch & Bonjer 1997, Hinzen 2003, Camelbeeck *et al.* 2007) (Fig. 7a). Our findings suggest that the surface deformation imposed by the Eifel plume explains why the LRE is so much more seismically active than many of the faults associated with other failed rifts in Europe (such as the Upper Rhine Graben (URG) between the Eifel and Alps). In fact, we see no deformation anomaly (horizontal or vertical) along the URG that could be interpreted as localized extension; a finding that is consistent with some prior studies (Rózsa *et al.* 2005, Tesauro *et al.* 2005), but inconsistent with others (J. Campbell *et al.* 2002, Fuhrmann *et al.* 2013, 2015).

Recently, it was also found that low-frequency seismic swarms occur in the lower crust underneath the Laacher See (Hensch *et al.* 2019). This activity was interpreted as the vertical migration of magma or magmatic fluids. Those findings, when combined with ours (as well as observations of continual gas emissions (Griesshaber *et al.* 1992, Aeschbach-Hertig *et al.* 1996, Buikin *et al.* 2005, Bräuer *et al.* 2013, Caracausi *et al.* 2016) and degassing events being correlated with seismicity (Berberich *et al.* 2019)), strongly suggest that the EVF is an active dynamic system.

Acknowledgements

We are extremely grateful to the many agencies, companies, and networks that have made GPS RINEX data available. In many cases the data were provided specifically to make our study possible. We specifically thank the following networks and institutions for raw RINEX data: EUREF (Bruyninx *et al.* 2019), IGS and the many agencies working under its umbrella (Johnston *et al.* 2017), CDDIS (Noll 2010), Technical University Delft (The Netherlands), NETPOS (The Netherlands), 06-GPS (The Netherlands), LNR Globalcom (The Netherlands), NAM/Shell (The Netherlands), ESTEC (The Netherlands), Royal Observatory of Belgium (Bruyninx & Defraigne 2018), FLEPOS (Belgium), WALCORS (Belgium), Institut Géographique National (France), Réseau GNSS Permanent (France), ORPHEON (France), RENAG (France) (RESIF 2017), SONEL (France) (Gravelle *et al.* 2019), SPSLux (Luxembourg), Natural Environment Research Council's British Isles continuous GNSS Facility (United Kingdom), Leica Smartnet (Poland), ASG-EUPOS (Poland), Pecny Observatory (Czech Republic), GEONAS (Czech Republic) (Schenk *et al.* 2010), VESOG (Czech Republic) (Kostecký & Kostecký 2005), CZEPOS (Czech Republic) (Kostecký & Kostecký 2005), FreDNet (Italy), Regione Autonoma Friuli Venezia Giulia (Italy), STPOS Bolzano (Italy), Politecnico di Torino (Italy), ARPA Piemonte (Italy), Rete GPS Veneto (Italy), Rete Dinamica Nazionale (Italy),

Instituto Nazionale di Geofisica e Vulcanologia (Italy), ITALPOS (Italy), Geodetski Institute (Slovenia), NIEP (Romania), ROMPOS (Romania), ESTPOS (Metsar *et al.* 2018) (Estonia), LATPOS (Latvia), LITPOS (Lithuania), Danish GPS Center Aalborg (Denmark), Danish Geodata Agency (Denmark), Instituto Geográfico Nacional (Spain), Topcon (Spain), CATNET Catalonia (Spain), GNSS Activa del Principado de Asturias (Spain), Red Activa de Estaciones GNSS de Cantabria (Spain), Red de estaciones GNSS de Castilla y León (Spain), Red Extremeña de Posicionamiento (Spain), Carto Galicia (Spain), REGAM Murcia (Spain), Meristenum Murcia (Spain), Government of La Rioja (Spain), Universidad de Oviedo (Spain), Ayuntamiento de Leganes (Spain), ETSI Topografía Geodesia y Cartografía (Spain), Leica Smartnet (Poland), ASG-EUPOS (Poland), ReNEP - Instituto Geografico Portugues (Portugal), SERVIR (Portugal), Norwegian Mapping Authority (Norway), Norwegian Metrology Service (Norway), SWEPOS (Sweden), Finnish Geodetic Institute (Finland), LIPOS (Austria), Bundesamt für Kartographie und Geodäsie (Germany), GFZ German Research Centre for Geosciences (Germany) (Uhlemann *et al.* 2016), Bundesanstalt für Gewässerkunde (Germany), Deutsches Geodätisches Forschungsinstitut (Germany) (Seitz *et al.* 2014), ASCOS (Germany), and SAPOS networks operated by various German States (i.e., State Office for Spatial Information and Land Development Baden-Württemberg, Hessian State Office for Land Management and Geoinformation, Landesamt für innere Verwaltung Mecklenburg-Vorpommern, Landesvermessung und Geobasisinformation Brandenburg, Staatsbetrieb Geobasisinformation und Vermessung Sachsen, Landesamt für Vermessung und Geoinformation Thüringen, Landesamt für Vermessung und Geobasisinformation Rheinland-Pfalz, Bezirksregierung Köln, Geoinformation und Landentwicklung Saarland, and Landesamt für Digitalisierung, Breitband und Vermessung Bayern). ORPHEON GNSS data were provided to the authors for scientific use in the framework of the GEODATA-INSU-CNRS convention. The services of the UK Natural Environment Research Council (NERC) British Isles continuous GNSS Facility (BIGF), www.bigf.ac.uk, in providing archived GNSS data for this station to this project, are gratefully acknowledged. A special thanks to H. van der Marel for helping to make the Kadaster/06-GPS/NAM data in the Netherlands available, A. Araszkiwicz for helping to make the Polish Smartnet data available, and T. van Dam for helping to make the SPSLux data in Luxembourg available. The GPS velocities can be found here: <https://doi.org/10.7910/DVN/ONATFP>, and GPS time-series at <http://geodesy.unr.edu> (Blewitt *et al.* 2018). Model results are given in the Supplementary Information. We thank J. Harvey for the locations of the volcanic centres, and I. Zaliapin for his help with the material presented in Appendix A. All figures were made with the Generic Mapping Tools (Wessel *et al.* 2013). This work got started through financial support to CK from the Royal Dutch Academy of Sciences. Additional support for CK was provided by USGS NEHRP award G18AP00019 and for CK and GB by NASA ROSES grants NNX16AK89G and 80NSSC19K1044.

References

- Aeschbach-Hertig, W., Hofer, M., Kipfer, R., Imboden, D.M. & Wieler, R. (1999) Accumulation of mantle gases in a permanently stratified volcanic lake (Lac Pavin, France). *Geochimica et Cosmochimica Acta*, **63**, 3357–3372. doi:10.1016/S0016-7037(99)00257-4
- Aeschbach-Hertig, W., Kipfer, R., Hofer, M., Imboden, D.M., Wieler, R. & Signer, P. (1996) Quantification of gas fluxes from the subcontinental mantle: The example of Laacher See, a maar lake in Germany. *Geochimica et Cosmochimica Acta*, **60**, 31–41. doi:10.1016/0016-7037(95)00370-3
- Altamimi, Z., Rebischung, P., Métivier, L. & Collilieux, X. (2016) ITRF2014: A new release of the International Terrestrial Reference Frame modeling nonlinear station motions. *Journal of Geophysical Research: Solid Earth*, **121**, 6109–6131. doi:10.1002/2016JB013098

- Anderson, E.M. (1937) The dynamics of the formation of cone-sheets, ring-dykes, and caldron-subsidences. *Proceedings of the Royal Society of Edinburgh*, **56**, 128–157. doi:10.1017/S0370164600014954
- Barth, A., Jordan, M. & Ritter, J.R.R. (2007) Crustal and upper mantle structure of the French Massif Central plume. in *Mantle Plumes: A Multidisciplinary Approach* eds. Ritter, J.R.R. & Christensen, U.R., pp. 159–184, Berlin, Heidelberg: Springer Berlin Heidelberg. doi:10.1007/978-3-540-68046-8_5
- Basili, R., Kastelic, V., Demircioglu, M.B., Garcia Moreno, D., Nemser, E.S., Petricca, P., Sboras, S.P., *et al.* (2013) The European Database of Seismogenic Faults (EDSF) Compiled in the framework of the Project SHARE. doi:10.6092/INGV.IT-SHARE-EDSF
- Berberich, G.M., Berberich, M.B., Ellison, A.M. & Wöhler, C. (2019) First identification of periodic degassing rhythms in three mineral springs of the East Eifel Volcanic Field (EEVF, Germany). *Geosciences*, **9**, 189. doi:10.3390/geosciences9040189
- Bertiger, W., Desai, S.D., Haines, B., Harvey, N., Moore, A.W., Owen, S. & Weiss, J.P. (2010) Single receiver phase ambiguity resolution with GPS data. *J Geod*, **84**, 327–337. doi:10.1007/s00190-010-0371-9
- Blewitt, G. (1990) An automatic editing algorithm for GPS data. *Geophys. Res. Lett.*, **17**, 199–202. doi:10.1029/GL017i003p00199
- Blewitt, G., Hammond, W.C. & Kreemer, C. (2018) Harnessing the GPS data explosion for interdisciplinary science. *Eos*, **99**. doi:10.1029/2018EO104623
- Blewitt, G., Kreemer, C., Hammond, W.C. & Gazeaux, J. (2016) MIDAS robust trend estimator for accurate GPS station velocities without step detection. *J. Geophys. Res. Solid Earth*, **121**, 2054–2068. doi:10.1002/2015JB012552
- Boehm, J., Werl, B. & Schuh, H. (2006) Troposphere mapping functions for GPS and very long baseline interferometry from European Centre for Medium-Range Weather Forecasts operational analysis data. *Journal of Geophysical Research: Solid Earth*, **111**. doi:10.1029/2005JB003629
- Bogusz, J., Kłos, A., Grzempowski, P. & Kontny, B. (2014) Modelling the velocity field in a regular grid in the area of Poland on the basis of the velocities of European permanent stations. *Pure Appl. Geophys.*, **171**, 809–833. doi:10.1007/s00024-013-0645-2
- Bogusz, J., Kłos, A. & Pokonieczny, K. (2019) Optimal strategy of a GPS position time series analysis for post-glacial rebound investigation in Europe. *Remote Sensing*, **11**, 1209. doi:10.3390/rs11101209
- Bowden, J., Davey Smith, G., Haycock, P.C. & Burgess, S. (2016) Consistent estimation in Mendelian Randomization with some invalid instruments using a weighted median estimator. *Genet Epidemiol*, **40**, 304–314. doi:10.1002/gepi.21965
- Bräuer, K., Kämpf, H., Niedermann, S. & Strauch, G. (2013) Indications for the existence of different magmatic reservoirs beneath the Eifel area (Germany): A multi-isotope (C, N, He, Ne, Ar) approach. *Chemical Geology*, **356**, 193–208. doi:10.1016/j.chemgeo.2013.08.013
- Bruyninx, C. & Defraigne, P. (2018) ROB GNSS Network Data. doi:10.24414/fst8-p256
- Bruyninx, C., Legrand, J., Fabian, A. & Pottiaux, E. (2019) GNSS metadata and data validation in the EUREF Permanent Network. *GPS Solut*, **23**, 106. doi:10.1007/s10291-019-0880-9
- Budweg, M., Bock, G. & Weber, M. (2006) The Eifel Plume—imaged with converted seismic waves. *Geophys J Int*, **166**, 579–589. doi:10.1111/j.1365-246X.2005.02778.x
- Buikin, A., Trieloff, M., Hopp, J., Althaus, T., Korochantseva, E., Schwarz, W.H. & Altherr, R. (2005) Noble gas isotopes suggest deep mantle plume source of late Cenozoic mafic alkaline

- volcanism in Europe. *Earth and Planetary Science Letters*, **230**, 143–162. doi:10.1016/j.epsl.2004.11.001
- Burov, E. & Gerya, T. (2014) Asymmetric three-dimensional topography over mantle plumes. *Nature*, **513**, 85–89. doi:10.1038/nature13703
- Burov, E. & Guillou-Frottier, L. (2005) The plume head–continental lithosphere interaction using a tectonically realistic formulation for the lithosphere. *Geophys J Int*, **161**, 469–490. doi:10.1111/j.1365-246X.2005.02588.x
- Burov, E., Guillou-Frottier, L., Acremont, E. d', Le Pourhiet, L. & Cloetingh, S. (2007) Plume head–lithosphere interactions near intra-continental plate boundaries. *Tectonophysics*, **434**, 15–38. doi:10.1016/j.tecto.2007.01.002
- Camelbeeck, T., Camp, M.V., Jongmans, D., Francis, O. & Dam, T. van. (2002) Comment on “Nature of the recent vertical ground movements inferred from high-precision leveling data in an intraplate setting: NE Ardenne, Belgium” by A. Demoulin and A. Collignon. *Journal of Geophysical Research: Solid Earth*, **107**, ETG 6-1-ETG 6-6. doi:10.1029/2001JB000397
- Camelbeeck, T., Vanneste, K., Alexandre, P., Verbeeck, K., Petermans, T., Rosset, P., Everaerts, M., *et al.* (2007) Relevance of active faulting and seismicity studies to assessments of long-term earthquake activity and maximum magnitude in intraplate northwest Europe, between the Lower Rhine Embayment and the North Sea. in *Special Paper 425: Continental Intraplate Earthquakes: Science, Hazard, and Policy Issues*, Vol. 425, pp. 193–224, Geological Society of America. doi:10.1130/2007.2425(14)
- Campbell, I.H. (2005) Large igneous provinces and the mantle plume hypothesis. *Elements*, **1**, 265–269. doi:10.2113/gselements.1.5.265
- Campbell, J., Kümpel, H.-J., Fabian, M., Fischer, D., Görres, B., Keyzers, C.J. & Lehmann, K. (2002) Recent movement pattern of the Lower Rhine Embayment from tilt, gravity and GPS data. *Netherlands Journal of Geosciences*, **81**, 223–230. doi:10.1017/S0016774600022472
- Caracausi, A., Avice, G., Burnard, P.G., Füre, E. & Marty, B. (2016) Chondritic xenon in the Earth's mantle. *Nature*, **533**, 82–85. doi:10.1038/nature17434
- Caro Cuenca, M. & Hanssen, R. (2008) Subsidence due to peat decomposition in The Netherlands, kinematic observations from radar interferometry. *Proc. of FRINGE 2007 Workshop*, 7.
- Caro Cuenca, M., Hooper, A.J. & Hanssen, R.F. (2013) Surface deformation induced by water influx in the abandoned coal mines in Limburg, The Netherlands observed by satellite radar interferometry. *Journal of Applied Geophysics*, **88**, 1–11. doi:10.1016/j.jappgeo.2012.10.003
- Chapron, E., Albéric, P., Jézéquel, D., Versteeg, W., Bourdier, J.-L. & Sitbon, J. (2010) Multidisciplinary characterisation of sedimentary processes in a recent maar lake (Lake Pavin, French Massif Central) and implication for natural hazards. *Natural Hazards and Earth System Sciences*, **10**, 1815–1827. doi:http://dx.doi.org/10.5194/nhess-10-1815-2010
- Cloetingh, S., Burov, E. & Francois, T. (2013) Thermo-mechanical controls on intra-plate deformation and the role of plume-folding interactions in continental topography. *Gondwana Research*, **24**, 815–837. doi:10.1016/j.gr.2012.11.012
- Dannberg, J. & Sobolev, S.V. (2015) Low-buoyancy thermochemical plumes resolve controversy of classical mantle plume concept. *Nature Communications*, **6**, 6960. doi:10.1038/ncomms7960
- Demoulin, A. & Collignon, A. (2000) Nature of the recent vertical ground movements inferred from high-precision leveling data in an intraplate setting: NE Ardenne, Belgium. *Journal of Geophysical Research: Solid Earth*, **105**, 693–705. doi:10.1029/1999JB900345
- Demoulin, A. & Collignon, A. (2002) Reply to comment by T. Camelbeeck *et al.* on “Nature of the recent vertical ground movements inferred from high-precision leveling data in an intraplate

- setting: NE Ardenne, Belgium”: COMMENTARY. *Journal of Geophysical Research: Solid Earth*, **107**, ETG 7-1-ETG 7-5. doi:10.1029/2002JB001837
- Demoulin, A. & Hallot, E. (2009) Shape and amount of the Quaternary uplift of the western Rhenish shield and the Ardennes (western Europe). *Tectonophysics*, **474**, 696–708. doi:10.1016/j.tecto.2009.05.015
- Francis, O., Van Camp, M., Dam, T. van, Warnant, R. & Hendrickx, M. (2004) Indication of the uplift of the Ardenne in long-term gravity variations in Membach (Belgium). *Geophys J Int*, **158**, 346–352. doi:10.1111/j.1365-246X.2004.02310.x
- Fuhrmann, T., Caro Cuenca, M., Knöpfler, A., Leijen, F.J. van, Mayer, M., Westerhaus, M., Hanssen, R.F., *et al.* (2015) Estimation of small surface displacements in the Upper Rhine Graben area from a combined analysis of PS-InSAR, levelling and GNSS data. *Geophysical Journal International*, **203**, 614–631. doi:10.1093/gji/ggv328
- Fuhrmann, T., Heck, B., Knöpfler, A., Masson, F., Mayer, M., Ulrich, P., Westerhaus, M., *et al.* (2013) Recent surface displacements in the Upper Rhine Graben — Preliminary results from geodetic networks. *Tectonophysics TOPO-EUROPE III*, **602**, 300–315. doi:10.1016/j.tecto.2012.10.012
- Garcia-Castellanos, D., Cloetingh, S. & Van Balen, R. (2000) Modelling the Middle Pleistocene uplift in the Ardennes–Rhenish Massif: Thermo-mechanical weakening under the Eifel? *Global and Planetary Change Environmental Tectonics and Climate: The Netherlands Environmental Earth System Dynamics Initiative*, **27**, 39–52. doi:10.1016/S0921-8181(01)00058-3
- Gauss, C.F. (1816) Bestimmung der genauigkeit der beobachtungen. *Zeitschrift für Astronomie und verwandte Wissenschaften*, **1**, 187–197.
- Goes, S., Spakman, W. & Bijwaard, H. (1999) A lower mantle source for central European volcanism. *Science*, **286**, 1928–1931.
- Gold, R.D., Friedrich, A., Kübler, S. & Salamon, M. (2017) Apparent Late Quaternary fault-slip rate increase in the southern Lower Rhine Graben, Central Europe. *Bulletin of the Seismological Society of America*, **107**, 563–580. doi:10.1785/0120160197
- Granet, M. & Trampert, J. (1989) Large-scale P-velocity structures in the Euro-Mediterranean area. *Geophys J Int*, **99**, 583–594. doi:10.1111/j.1365-246X.1989.tb02043.x
- Granet, M., Wilson, M. & Achauer, U. (1995) Imaging a mantle plume beneath the French Massif Central. *Earth and Planetary Science Letters*, **136**, 281–296. doi:10.1016/0012-821X(95)00174-B
- Gravelle, M., Guichard, M., Poirier, E., Prouteau, E., Testut, L. & Woppelmann, G. (2019) Status of SONEL: the GNSS at Tide Gauge Data Assembly Center for the Global Sea Level Observing System (GLOSS), Presented at the AGU Fall Meeting 2019, AGU.
- Griesshaber, E., O’Nions, R.K. & Oxburgh, E.R. (1992) Helium and carbon isotope systematics in crustal fluids from the Eifel, the Rhine Graben and Black Forest, F.R.G. *Chemical Geology*, **99**, 213–235. doi:10.1016/0009-2541(92)90178-8
- Griffiths, R.W. & Campbell, I.H. (1991) Interaction of mantle plume heads with the Earth’s surface and onset of small-scale convection. *Journal of Geophysical Research*, **96**, 18295. doi:10.1029/91JB01897
- Grunewald, S., Weber, M. & Kind, R. (2001) The upper mantle under Central Europe: indications for the Eifel plume. *Geophysical Journal International*, **147**, 590–601. doi:10.1046/j.1365-246x.2001.01553.x
- Grünthal, G., Wahlström, R. & Stromeyer, D. (2013) The SHARE European Earthquake Catalogue (SHEEC) for the time period 1900–2006 and its comparison to the European-Mediterranean Earthquake Catalogue (EMEC). *J Seismol*, **17**, 1339–1344. doi:10.1007/s10950-013-9379-y

- Guillou-Frottier, L., Burov, E., Nehlig, P. & Wyns, R. (2007) Deciphering plume–lithosphere interactions beneath Europe from topographic signatures. *Global and Planetary Change TOPO-EUROPE: the Geoscience of Coupled Deep Earth-Surface Processes*, **58**, 119–140. doi:10.1016/j.gloplacha.2006.10.003
- Hammond, W.C., Blewitt, G. & Kreemer, C. (2016) GPS Imaging of vertical land motion in California and Nevada: Implications for Sierra Nevada uplift. *Journal of Geophysical Research: Solid Earth*, **121**, 7681–7703. doi:10.1002/2016JB013458
- Hampel, F.R. (1974) The influence curve and its role in robust estimation. *Journal of the American Statistical Association*, **69**, 383–393. doi:10.2307/2285666
- Hensch, M., Dahm, T., Ritter, J., Heimann, S., Schmidt, B., Stange, S. & Lehmann, K. (2019) Deep low-frequency earthquakes reveal ongoing magmatic recharge beneath Laacher See Volcano (Eifel, Germany). *Geophys J Int*, **216**, 2025–2036. doi:10.1093/gji/ggy532
- Hill, R.I. (1991) Starting plumes and continental break-up. *Earth and Planetary Science Letters*, **104**, 398–416. doi:10.1016/0012-821X(91)90218-7
- Hinzen, K.-G. (2003) Stress field in the Northern Rhine area, Central Europe, from earthquake fault plane solutions. *Tectonophysics*, **377**, 325–356. doi:10.1016/j.tecto.2003.10.004
- Hinzen, K.-G. & Reamer, S.K. (2007) Seismicity, seismotectonics, and seismic hazard in the northern Rhine area. *Geological Society of America Special Papers*, **425**, 225–242. doi:10.1130/2007.2425(15)
- Hoernle, K., Zhang, Y.-S. & Graham, D. (1995) Seismic and geochemical evidence for large-scale mantle upwelling beneath the eastern Atlantic and western and central Europe. *Nature*, **374**, 34–39. doi:10.1038/374034a0
- Huber, P.J. (1981) *Robust Statistics*, New York: Wiley.
- Husson, L., Bodin, T., Spada, G., Choblet, G. & Kreemer, C. (2018) Bayesian surface reconstruction of geodetic uplift rates: Mapping the global fingerprint of Glacial Isostatic Adjustment. *Journal of Geodynamics*, **122**, 25–40. doi:10.1016/j.jog.2018.10.002
- Jagt, L., Ruigrok, E. & Paulssen, H. (2017) Relocation of clustered earthquakes in the Groningen gas field. *Netherlands Journal of Geosciences*, **96**, s163–s173. doi:10.1017/njg.2017.12
- Johnston, G., Riddell, A. & Hausler, G. (2017) The International GNSS Service. in *Springer Handbook of Global Navigation Satellite Systems* Springer Handbooks eds. Teunissen, P.J.G. & Montenbruck, O., pp. 967–982, Cham: Springer International Publishing. doi:10.1007/978-3-319-42928-1_33
- Juvigné, E. & Gilot, E. (1986) Ages et zones de dispersion de tephra émises par les volcans du Montcineyre et du lac Pavon (Massif Central, France). *Zeitschrift der Deutschen Geologischen Gesellschaft*, **137**, 613–623.
- Ketelaar, V.B.H. (Ed.). (2009) Subsidence Due to Hydrocarbon Production in the Netherlands. in *Satellite Radar Interferometry: Subsidence Monitoring Techniques Remote Sensing and Digital Image Processing*, pp. 7–26, Dordrecht: Springer Netherlands. doi:10.1007/978-1-4020-9428-6_2
- Keyser, M., Ritter, J.R.R. & Jordan, M. (2002) 3D shear-wave velocity structure of the Eifel plume, Germany. *Earth and Planetary Science Letters*, **203**, 59–82. doi:10.1016/S0012-821X(02)00861-0
- Kierulf, H.P., Ouassou, M., Simpson, M.J.R. & Vestøl, O. (2013) A continuous velocity field for Norway. *J Geod*, **87**, 337–349. doi:10.1007/s00190-012-0603-2
- Kierulf, H.P., Steffen, H., Simpson, M.J.R., Lidberg, M., Wu, P. & Wang, H. (2014) A GPS velocity field for Fennoscandia and a consistent comparison to glacial isostatic adjustment models. *J. Geophys. Res. Solid Earth*, 2013JB010889. doi:10.1002/2013JB010889

- Klos, A., Bogusz, J., Figurski, M. & Kosek, W. (2016) On the handling of outliers in the GNSS time series by means of the noise and probability analysis. *IAG 150 Years International Association of Geodesy Symposia* eds. Rizos, C. & Willis, P., pp. 657–664, Springer International Publishing.
- Kontny, B. & Bogusz, J. (2012) Models of vertical movements of the earth crust surface in the area of Poland derived from leveling and GNSS data. *Acta Geodynamica Et Geomaterialia*, **9**, 331–337.
- Kostelecký, J. & Kostelecký, J. (2005) New permanent GNSS networks in the Czech Republic and their use in geosciences. *Reports on Geodesy*, **2**, 151–157.
- Koulakov, I., Kaban, M.K., Tesauro, M. & Cloetingh, S. (2009) P- and S-velocity anomalies in the upper mantle beneath Europe from tomographic inversion of ISC data. *Geophys J Int*, **179**, 345–366. doi:10.1111/j.1365-246X.2009.04279.x
- Kreemer, C. (2020) Robust estimation of spatio-temporal common mode components in GPS time-series. *Journal of Geodesy*, **in preparation**.
- Kreemer, C., Hammond, W.C. & Blewitt, G. (2018) A robust estimation of the 3-D intraplate deformation of the North American plate from GPS. *Journal of Geophysical Research-Solid Earth*, **123**, 4388–4412. doi:10.1029/2017JB015257
- Lahtinen, S., Jivall, L., Häkli, P., Kall, T., Kollo, K., Kosenko, K., Galinauskas, K., *et al.* (2019) Densification of the ITRF2014 position and velocity solution in the Nordic and Baltic countries. *GPS Solut*, **23**, 95. doi:10.1007/s10291-019-0886-3
- Leys, C., Ley, C., Klein, O., Bernard, P. & Licata, L. (2013) Detecting outliers: Do not use standard deviation around the mean, use absolute deviation around the median. *Journal of Experimental Social Psychology*, **49**, 764–766. doi:10.1016/j.jesp.2013.03.013
- Lustrino, M. & Carminati, E. (2007) Phantom plumes in Europe and the circum-Mediterranean region. in *Special Paper 430: Plates, Plumes and Planetary Processes*, Vol. 430, pp. 723–745, Geological Society of America. doi:10.1130/2007.2430(33)
- Lyard, F., Lefevre, F., Letellier, T. & Francis, O. (2006) Modelling the global ocean tides: modern insights from FES2004. *Ocean Dynamics*, **56**, 394–415. doi:10.1007/s10236-006-0086-x
- Mälzer, H., Hein, G. & Zippelt, K. (1983) Height changes in the Rhenish Massif: Determination and analysis. *Plateau Uplift* eds. Fuchs, K., Gehlen, K. von, Mälzer, H., Murawski, H. & Semmel, A., pp. 164–176, Springer Berlin Heidelberg.
- Marotta, A.M., Mitrovica, J.X., Sabadini, R. & Milne, G. (2004) Combined effects of tectonics and glacial isostatic adjustment on intraplate deformation in central and northern Europe: Applications to geodetic baseline analyses. *J. Geophys. Res.*, **109**, B01413, doi:10.1029/2002JB002337.
- Márquez-Azúa, B. & DeMets, C. (2003) Crustal velocity field of Mexico from continuous GPS measurements, 1993 to June 2001: Implications for the neotectonics of Mexico. *J. Geophys. Res.*, **108**, 2450. doi:10.1029/2002JB002241
- Masson, C., Mazzotti, S., Vernant, P. & Doerflinger, E. (2019) Extracting small deformation beyond individual station precision from dense Global Navigation Satellite System (GNSS) networks in France and western Europe. *Solid Earth*, **10**, 1905–1920. doi:https://doi.org/10.5194/se-10-1905-2019
- Mathar, J.P., Ritter, J.R.R. & Friederich, W. (2006) Surface waves image the top of the Eifel plume. *Geophys J Int*, **164**, 377–382. doi:10.1111/j.1365-246X.2006.02835.x
- Matthews, A., Fouillac, C., Hill, R., O’Nions, R.K. & Oxburgh, E.R. (1987) Mantle-derived volatiles in continental crust: the Massif Central of France. *Earth and Planetary Science Letters*, **85**, 117–128. doi:10.1016/0012-821X(87)90026-4

- Metsar, J., Kollo, K. & Ellmann, A. (2018) Modernization of the Estonian national GNSS reference station network. *Geodesy and Cartography*, **44**, 55–62. doi:10.3846/gac.2018.2023
- Meyer, W. & Stets, J. (2002) Pleistocene to Recent tectonics in the Rhenish Massif (Germany). *Netherlands Journal of Geosciences*, **81**, 217–221. doi:10.1017/S0016774600022460
- Mindlin, R.D. (1936) Force at a point in the interior of a semi-infinite solid. *Physics*, **7**, 195. doi:10.1063/1.1745385
- Montelli, R., Nolet, G., Dahlen, F.A. & Masters, G. (2006) A catalogue of deep mantle plumes: New results from finite-frequency tomography. *Geochem. Geophys. Geosyst.*, **7**, Q11007. doi:10.1029/2006GC001248
- Murray, K.D., Murray, M.H. & Sheehan, A.F. (2019) Active deformation near the Rio Grande Rift and Colorado Plateau as inferred from continuous Global Positioning System measurements. *Journal of Geophysical Research: Solid Earth*, **124**, 2166–2183. doi:10.1029/2018JB016626
- Neres, M., Neves, M.C., Custódio, S., Palano, M., Fernandes, R., Matias, L., Carafa, M., *et al.* (2018) Gravitational potential energy in Iberia: A driver of active deformation in high-topography regions. *Journal of Geophysical Research: Solid Earth*, **123**, 10,277–10,296. doi:10.1029/2017JB015002
- Nocquet, J.-M., Calais, E. & Parsons, B. (2005) Geodetic constraints on glacial isostatic adjustment in Europe. *Geophys. Res. Lett.*, **32**, L06308, doi:10.1029/2004GL022174.
- Noll, C.E. (2010) The crustal dynamics data information system: A resource to support scientific analysis using space geodesy. *Advances in Space Research*, **45**, 1421–1440. doi:10.1016/j.asr.2010.01.018
- Nowell, D.A.G., Jones, M.C. & Pyle, D.M. (2006) Episodic Quaternary volcanism in France and Germany. *Journal of Quaternary Science*, **21**, 645–675. doi:10.1002/jqs.1005
- Pilidou, S., Priestley, K., Debayle, E. & Gudmundsson, O. (2005) Rayleigh wave tomography in the North Atlantic: high resolution images of the Iceland, Azores and Eifel mantle plumes. *Lithos*, **79**, 453–474. doi:10.1016/j.lithos.2004.09.012
- Plenefisch, T. & Bonjer, K.-P. (1997) The stress field in the Rhine Graben area inferred from earthquake focal mechanisms and estimation of frictional parameters. *Tectonophysics*, **275**, 71–97. doi:10.1016/S0040-1951(97)00016-4
- Plomerová, J. & Babuška, V. (2010) Long memory of mantle lithosphere fabric — European LAB constrained from seismic anisotropy. *Lithos*, **120**, 131–143. doi:10.1016/j.lithos.2010.01.008
- Prodehl, C., Mueller, St. & Haak, V. (2006) The European Cenozoic rift system. in *Developments in Geotectonics Continental rifts: evolution, structure, tectonics* ed. Olsen, K.H., Vol. 25, pp. 133–212, Elsevier. doi:10.1016/S0419-0254(06)80012-1
- Regenauer-Lieb, K. (1998) Dilatant plasticity applied to Alpine collision: ductile void growth in the intraplate area beneath the Eifel volcanic field. *Journal of Geodynamics*, **27**, 1–21. doi:10.1016/S0264-3707(97)00024-0
- Renka, R.J. (1997) Algorithm 772: STRIPACK: Delaunay triangulation and Voronoi diagram on the surface of a sphere. *ACM Trans. Math. Softw.*, **23**, 416–434. doi:10.1145/275323.275329
- RESIF. (2017) RESIF-RENAG French national Geodetic Network. RESIF - Réseau Sismologique et géodésique Français. doi:10.15778/resif.rg
- Ritter, J.R.R. (2007) The seismic signature of the Eifel plume. in *Mantle Plumes: A Multidisciplinary Approach* eds. Ritter, J.R.R. & Christensen, U.R., pp. 379–404, Berlin, Heidelberg: Springer Berlin Heidelberg. doi:10.1007/978-3-540-68046-8_12
- Ritter, J.R.R., Jordan, M., Christensen, U.R. & Achauer, U. (2001) A mantle plume below the Eifel volcanic fields, Germany. *Earth and Planetary Science Letters*, **186**, 7–14. doi:10.1016/S0012-821X(01)00226-6

- Rózsa, Sz., Heck, B., Mayer, M., Seitz, K., Westerhaus, M. & Zippelt, K. (2005) Determination of displacements in the upper Rhine graben Area from GPS and leveling data. *Int J Earth Sci (Geol Rundsch)*, **94**, 538–549. doi:10.1007/s00531-005-0478-z
- Santamaría-Gómez, A., Gravelle, M., Dangendorf, S., Marcos, M., Spada, G. & Wöppelmann, G. (2017) Uncertainty of the 20th century sea-level rise due to vertical land motion errors. *Earth and Planetary Science Letters*, **473**, 24–32. doi:10.1016/j.epsl.2017.05.038
- Schenk, V., Schenková, Z., Cajthamlová, M. & Fučík, Z. (2010) GEONAS - Geodynamic network of permanent GNSS stations within the Czech Republic. *Acta Geodynamica et Geomaterialia*, **7**, 99–111.
- Schmid, R., Steigenberger, P., Gendt, G., Ge, M. & Rothacher, M. (2007) Generation of a consistent absolute phase-center correction model for GPS receiver and satellite antennas. *J. Geod.*, **81**, 781–798. doi:10.1007/s00190-007-0148-y
- Schmincke, H.-U. (2007) The Quaternary volcanic fields of the East and West Eifel (Germany). in *Mantle Plumes: A Multidisciplinary Approach* eds. Ritter, J.R.R. & Christensen, U.R., pp. 241–322, Berlin, Heidelberg: Springer Berlin Heidelberg. doi:10.1007/978-3-540-68046-8_8
- Seiberlich, C.K.A., Ritter, J.R.R. & Wawerzinek, B. (2013) Topography of the lithosphere–asthenosphere boundary below the Upper Rhine Graben Rift and the volcanic Eifel region, Central Europe. *Tectonophysics*, **603**, 222–236. doi:10.1016/j.tecto.2013.05.034
- Seitz, F., Arenz, H. & Leismüller, F. (2014, July 21) Position measurements at five permanent GPS stations in the Bavarian Alps as part of the Geodetic Alpine Integrated Network (GAIN) of the ALPS-GPSQUAKENET project. *Deutsches Geodätisches Forschungsinstitut der Technischen Universität München*. doi:https://doi.org/10.1594/PANGAEA.834193
- Serpelloni, E., Faccenna, C., Spada, G., Dong, D. & Williams, S.D.P. (2013) Vertical GPS ground motion rates in the Euro-Mediterranean region: New evidence of velocity gradients at different spatial scales along the Nubia-Eurasia plate boundary. *J. Geophys. Res. Solid Earth*, **118**, 6003–6024. doi:10.1002/2013JB010102
- Simon, K.M., Riva, R.E.M., Kleinherenbrink, M. & Frederikse, T. (2018) The glacial isostatic adjustment signal at present day in northern Europe and the British Isles estimated from geodetic observations and geophysical models. *Solid Earth*, **9**, 777–795. doi:https://doi.org/10.5194/se-9-777-2018
- Sobolev, S.V., Zeyen, H., Granet, M., Achauer, U., Bauer, C., Werling, F., Altherr, R., *et al.* (1997) Upper mantle temperatures and lithosphere-asthenosphere system beneath the French Massif Central constrained by seismic, gravity, petrologic and thermal observations. *Tectonophysics*, **275**, 143–164. doi:10.1016/S0040-1951(97)00019-X
- Spakman, W. & Wortel, M.J.R. (2004) Tomographic View on Western Mediterranean Geodynamics. in *The TRANSMED Atlas, The Mediterranean Region from Crust to Mantle* eds. Cavazza, W., Roure, F., Spakman, W., Stampfli, G.M. & Ziegler, P., pp. 31–52, New York, NY: Springer-Verlag.
- Stucchi, M., Rovida, A., Gomez Capera, A.A., Alexandre, P., Camelbeeck, T., Demircioglu, M.B., Gasperini, P., *et al.* (2013) The SHARE European Earthquake Catalogue (SHEEC) 1000–1899. *J Seismol*, **17**, 523–544. doi:10.1007/s10950-012-9335-2
- Tesauro, M., Hollenstein, C., Egli, R., Geiger, A. & Kahle, H.-G. (2005) Continuous GPS and broad-scale deformation across the Rhine Graben and the Alps. *Int J Earth Sci (Geol Rundsch)*, **94**, 525–537. doi:10.1007/s00531-004-0453-0
- Tesauro, M., Hollenstein, C., Egli, R., Geiger, A. & Kahle, H.-G. (2006) Analysis of central western Europe deformation using GPS and seismic data. *J. Geodyn.*, **42**, 194–209. doi:10.1016/j.jog.2006.08.001

- Thienen-Visser, K. van & Breunese, J. (2015) Induced seismicity of the Groningen gas field: History and recent developments. *The Leading Edge*, **34**, 664–671. doi:10.1190/tle34060664.1
- Uhlemann, M., Gendt, G., Ramatschi, M. & Deng, Z. (2016) GFZ global multi-GNSS network and data processing results. *IAG 150 Years International Association of Geodesy Symposia* eds. Rizos, C. & Willis, P., pp. 673–679, Springer International Publishing.
- Van Balen, R.T., Houtgast, R.F., Van der Wateren, F.M., Vandenberghe, J. & Bogaart, P.W. (2000) Sediment budget and tectonic evolution of the Meuse catchment in the Ardennes and the Roer Valley Rift System. *Global and Planetary Change*, **27**, 113–129. doi:10.1016/S0921-8181(01)00062-5
- Van Camp, M.V., Viron, O. de, Scherneck, H.-G., Hinzen, K.-G., Williams, S.D.P., Lecocq, T., Quinif, Y., *et al.* (2011) Repeated absolute gravity measurements for monitoring slow intraplate vertical deformation in western Europe. *Journal of Geophysical Research: Solid Earth*, **116**, B08402. doi:10.1029/2010JB008174
- Vervoort, A. (2016) Surface movement above an underground coal longwall mine after closure. *Natural Hazards and Earth System Sciences*, **16**, 2107–2121. doi:https://doi.org/10.5194/nhess-16-2107-2016
- Ward, S.N. (1998) On the consistency of earthquake moment release and space geodetic strain rates: Europe. *Geophysical Journal International*, **135**, 1011–1018.
- Wedepohl, K.H. & Baumann, A. (1999) Central European Cenozoic plume volcanism with OIB characteristics and indications of a lower mantle source. *Contrib Mineral Petrol*, **136**, 225–239. doi:10.1007/s004100050534
- Wessel, P., Smith, W.H.F., Scharroo, R., Luis, J. & Wobbe, F. (2013) Generic Mapping Tools: Improved Version Released. *Eos, Transactions American Geophysical Union*, **94**, 409–410. doi:10.1002/2013EO450001
- Westaway, R. (2001) Flow in the lower continental crust as a mechanism for the Quaternary uplift of the Rhenish Massif, north-west Europe. in *River Basin Sediment Systems - Archives of Environmental Change* eds. Maddy, D., Macklin, M.G. & Woodward, J.C., pp. 87–167, London: CRC Press. doi:10.1201/9781439824672-7
- Wilson, M. & Downes, H. (1992) Mafic alkaline magmatism associated with the European Cenozoic rift system. *Tectonophysics*, **208**, 173–182. doi:10.1016/0040-1951(92)90343-5
- Wüllner, U., Christensen, U.R. & Jordan, M. (2006) Joint geodynamical and seismic modelling of the Eifel plume. *Geophys. J. Int.*, **165**, 357–372. doi:10.1111/j.1365-246X.2006.02906.x
- Zangana, N.A., Downes, H., Thirlwall, M.F., Marriner, G.F. & Bea, F. (1999) Geochemical variation in peridotite xenoliths and their constituent clinopyroxenes from Ray Pic (French Massif Central): implications for the composition of the shallow lithospheric mantle. *Chemical Geology*, **153**, 11–35. doi:10.1016/S0009-2541(98)00150-8
- Zhao, D. (2007) Seismic images under 60 hotspots: Search for mantle plumes. *Gondwana Research*, **12**, 335–355. doi:10.1016/j.gr.2007.03.001
- Zhu, H., Bozdağ, E., Peter, D. & Tromp, J. (2012) Structure of the European upper mantle revealed by adjoint tomography. *Nature Geoscience*, **5**, 493–498. doi:10.1038/ngeo1501
- Zumberge, J.F., Heflin, M.B., Jefferson, D.C., Watkins, M.M. & Webb, F.H. (1997) Precise point positioning for the efficient and robust analysis of GPS data from large networks. *J. Geophys. Res.*, **102**, 5005–5017. doi:10.1029/96JB03860

Table 1. Plume Model Parameters (see text for explanations)

| Parameter | value | Standard deviation |
|---------------------------------|--------------|---------------------------|
| Vertical Force F (1e11 N) | 15.8117 | 0.1564 |
| σ_1 (km) | 173.5312 | 1.7743 |
| σ_2 (km) | 98.0999 | 1.4502 |
| Lon_0 ($^{\circ}$ E) | 6.4000 | 0.0105 |
| Lat_0 ($^{\circ}$ N) | 50.5000 | 0.0055 |
| θ (radians) | 0.2831 | 0.0103 |
| u_0 (regional offset, mm) | 0.3544 | 0.0064 |
| v_0 (regional offset, mm) | -0.0112 | 0.0001 |
| w_0 (regional offset, mm) | -0.0050 | 0.0002 |
| du/dE (regional slope, mm/km) | 0.0962 | 0.0067 |
| du/dN (regional slope, mm/km) | -0.7008 | 0.0131 |
| dv/dE (regional slope, mm/km) | 0.0000 | 0.0002 |
| dv/dN (regional slope, mm/km) | 0.0058 | 0.0002 |
| dw/dE (regional slope, mm/km) | -0.0099 | 0.0002 |
| dw/dN (regional slope, mm/km) | -0.0099 | 0.0002 |




Figure 1. Black dots are epicentres of earthquakes between 1000-2006 in the SHEEC-SHARE database (Grünthal *et al.* 2013, Stucchi *et al.* 2013). Colours are epicentre density for circular areas with radius (R) of 30 km (with each event in a circle weighted by $(1-D/R)$, where D is distance from event to centre of circle). Blue dots are centres of Quaternary EVF activity, and blue triangle is location of latest activity in Massif Central. Light blue lines are seismogenic faults (Basili *et al.* 2013), and the Rhenish Massif is outlined in green. B = Belgium, NL = The Netherlands, LRE=Lower Rhine Embayment, URG=Upper Rhine Graben. Inset) red solid/dashed polygon is extent of data and model, resp., blue polygon is extent of results presented here, blue dashed polygon is area of Fig. 7.




Figure 2. Locations of GPS stations used, coloured by the length of the time-series. Black triangles are locations of velocities added from the literature (Kierulf *et al.* 2013, 2014, Lahtinen *et al.* 2019).




Figure 3. a) Observed VLM, b) despeckled VLM, c) imaged VLM (outline of Rhenish Massif (green polygon), centre of Eifel plume (Ritter *et al.* 2001) (open green triangle) and Massif Central (filled triangle) are shown for reference), d) imaged VLM corrected for GIA and shown only where corrected $VLM > 2\sigma$.




Figure 4. a) Input chequerboard with alternating $+1$ and -1 mm yr^{-1} VLM in 200×200 km cells. Open/closed triangle is location of Eifel and Massif Central, b) Imaged value at GPS locations using, c) Imaged VLM at 0.1° grid, d) Imaged VLM where absolute value is $\geq 1\sigma$ of imaged VLM, e) Imaged VLM where absolute value is $\geq 2\sigma$ of imaged VLM.




Figure 5. a) Observed horizontal velocities relative to the extension anomaly that encompasses the Eifel (Fig. 5b). For clarity, velocities which differ $>0.25 \text{ mm yr}^{-1}$ from the local velocity gradient are not shown. b) Colours and vectors show style and (normalized) principal axes of strain rate tensor (averaged over non-overlapping equal areas), respectively. Outline of Rhenish Massif and centre of Eifel and Massif Central are shown (triangles). Also shown for reference are green dashed lines, which are ellipses around Scandinavia and to which contractional strain rates in the northern part of the model are oriented orthogonally. Orange dashed circles indicates deviation from that pattern and highlights that contractional axes around the Eifel and Massif Central are oriented radially to their respective extension anomalies, c) contours are rate of dilatation (red is extension, blue is contraction) and model velocities (based on inferred strain and rotation rates) are in same reference frame and with same scale as in Fig. 5a, d) Same contours as in Fig. 5c, but only where dilatation rate is $>2\sigma$.




Figure 6. GIA-corrected VLM for profile shown in Fig 3d. Open circles are original VLM (inside profile box), orange squares are despeckled VLM (errors bars are 1σ), red line is imaged VLM along profile, dark and light blue outline is 1σ in imaged VLM defined, respectively, by deviation from original VLM and despeckled VLM.




Figure 7. a) Colours are imaged VLM corrected for GIA, vectors are principal axes of horizontal strain rate tensor (averaged over non-overlapping equal areas), purple dashed line outlines area of significant dilatation rate (at 2σ level). Focal mechanisms of regional studies (Hinzen 2003, Camelbeeck *et al.* 2007), colour coded for the implied extensional (red), contractional (blue), or strike-slip (green) deformation. Blue dots are centres of Quaternary EVF activity, and black lines are seismogenic faults (Basili *et al.* 2013). Green outline is Rhenish Massif and triangle is the projected centre of Eifel plume (Ritter *et al.* 2001). b) VLM and strain rate predicted from our best fitting plume model. Green star is centre of gain function applied to vertical force.




Figure 8. Graphical representation of gain function $A(x,y)$. See text for explanation.

Appendix A: Weighted Average and Its Variance

For each evaluation point we wish to obtain the weighted average (W) of the N strain rate estimates m_i , $i=1, \dots, N$, consisting of the estimate at the evaluation point itself and those at other evaluation points within a certain distance. The weights (w_i) are defined as σ_{ii}^{-1} , normalized by $\sum_{i=1}^N w_i$, and $\sigma_{ii} = \sigma_i^2$ with σ_i being the standard deviation in m_i . The expected value of W is given by:

$$E[W] = \sum_{i=1}^N w_i m_i \quad (1)$$

and the variance in W by:

$$Var[W] = \sum_{i=1}^N w_i^2 \sigma_{ii} + 2 \sum_{i \neq j} w_i w_j \sigma_{ij} \quad (2)$$

The question is what is σ_{ij} , i.e., the covariance between m_i and m_j ? In our case, each estimate of m_i and m_j can be thought of as average of strain rates (although it is really part of a multivariate median) for T_i and T_j number of triangles, respectively, and those are based on GPS velocities at P_i and P_j points, respectively. Unlike the triangles, the sets of used GPS points provide a straightforward measure of interdependency between m_i and m_j . We therefore propose the following definition of the covariance between two estimates (E_1 and E_2) based on P_1 and P_2 observations, respectively.

$$Cov[E_1, E_2] \approx \frac{p_0}{2} \left(\frac{P_1 Var(E_1) + P_2 Var(E_2)}{P_1 P_2} \right) \quad (3)$$

where p_0 is the number of common GPS stations that were used in both estimations.

The proposed covariance (3) has the following natural properties:

- 1) If the estimations do not use common stations, they are uncorrelated: i.e., $p_0=0$ and thus $Corr(E_1, E_2)=0$.
- 2) If the estimations use the same stations, they are fully correlated: i.e., $p_0=P_1=P_2$. and thus $Corr(E_1, E_2)=1$.
- 3) If the two estimations have very different variances (and the station number is approximately the same), the examined covariance is mostly affected by the largest variance.

The approximation in (3) is based on the following general and conceptual case. Consider a set of independent random variables $X_1, \dots, X_{p_0+p_1+p_2}$. Assume that the estimation E_1 at the first location uses $P_1=p_0+p_1$ observations, and the estimation E_2 at the first location uses $P_2=p_0+p_2$ observations in such a way that p_0 observations (and only those ones) are used at both locations. Specifically, we may assume that both locations use the common observations

$$\{X_1, \dots, X_{p_0}\}$$

In addition, the first location used the unique observations

$$\{X_{p_0+1}, \dots, X_{p_0+p_1}\}$$

and the second location uses the unique observations

$$\{X_{p_0+p_1+1}, \dots, X_{p_0+p_1+p_2}\}$$

For convenience, we denote the observations used at the first locations $Y_i, i = 1, \dots, P_1$, and the observations used at the second location $Z_i, i = 1, \dots, P_2$, in such a way that $Y_i = Z_i = X_i$ for all $i = 1, \dots, p_0$. Let us assume that the examined estimations are the sample averages:

$$E_1 \equiv \bar{X}_1 = \frac{1}{P_1} \sum_{i=1}^{P_1} Y_i \quad E_2 \equiv \bar{X}_2 = \frac{1}{P_2} \sum_{i=1}^{P_2} Z_i$$

The covariance between \bar{X}_1 and \bar{X}_2 is then given by:

$$\begin{aligned} \text{Cov}(\bar{X}_1, \bar{X}_2) &= \frac{1}{P_1 P_2} \text{Cov} \left(\sum_{i=1}^{P_1} Y_i, \sum_{i=1}^{P_2} Z_i \right) \\ &= \frac{1}{P_1 P_2} \text{Cov} \left(\sum_{i=1}^{n_0} Y_i + \sum_{i=n_0+1}^{P_1} Y_i, \sum_{i=1}^{n_0} Z_i + \sum_{i=n_0+1}^{P_2} Z_i \right) \\ &= \frac{1}{P_1 P_2} \text{Cov} \left(\sum_{i=1}^{n_0} X_i + \sum_{i=n_0+1}^{P_1} Y_i, \sum_{i=1}^{n_0} X_i + \sum_{i=n_0+1}^{P_2} Z_i \right) \\ &= \frac{1}{P_1 P_2} \text{Cov} \left(\sum_{i=1}^{n_0} X_i, \sum_{i=1}^{n_0} X_i \right) \\ &= \frac{1}{P_1 P_2} \text{Var} \left(\sum_{i=1}^{n_0} X_i \right) = \sigma^2 \frac{p_0}{P_1 P_2} \end{aligned}$$

Observe that

$$\sigma^2 = \text{Var}(\bar{X}_1) P_1 = \text{Var}(\bar{X}_2) P_2$$

and accordingly


$$\text{Cov}(\bar{X}_1, \bar{X}_2) = \text{Var}(\bar{X}_1) \frac{P_0}{P_2} = \text{Var}(\bar{X}_2) \frac{P_0}{P_1}$$

However, the assumption of equal variance σ^2 might be violated. We hence suggest, as a simple heuristic device, to estimate the covariance as the following average:


$$\text{Cov}[\bar{X}_1, \bar{X}_2] \approx \frac{1}{2} \left(\text{Var}(\bar{X}_1) \frac{P_0}{P_2} + \text{Var}(\bar{X}_2) \frac{P_0}{P_1} \right)$$

which is equivalent to (3).


Supplementary Figures




Supplementary Figure S1. a) Input checkerboard with alternating +1 and -1 mm yr⁻¹ VLM in 100x100 km cells. Open/closed triangle is location of Eifel and Massif Central, b) Imaged value at GPS locations using, c) Imaged VLM at 0.1° grid, d) Imaged VLM where absolute value is $\geq 1\sigma$ of imaged VLM, e) Imaged VLM where absolute value is $\geq 2\sigma$ of imaged VLM.




Supplementary Figure S2. a) Standard deviation defined relative to observed station VLM, b) Standard deviation defined relative to despeckled VLM.




Supplementary Figure S3. a-d) Same as Figure 3 and e) same as Figure 6 but all results based on VLM derived from time-series not filtered for common-mode errors




Supplementary Figure S4. a) Spatial scale of strain rate estimate expressed as the radius of a circle whose area is the median area of all triangles considered in the strain rate estimation. b) Standard deviation in the dilatation rate.




Supplementary Figure S5. Same as Fig 5b-d but with a spatial scale that is 50% of preferred model.




Supplementary Figure S6. Same as Fig 5b-d but with a spatial scale that is 150% of preferred model.




Supplementary Figure S7. Same as Fig 5b-d but with outlier velocities ($>1.5 \text{ mm yr}^{-1}$) removed before modelling.



Supplementary Figure S8. Same as Fig 5b-d but based on velocities derived from time-series that had no common-mode errors removed.



Supplementary Figure S9. This result is based on the MELD algorithm as originally presented (Kremer *et al.* 2018).



Supplementary Figure S10. Same as Fig. 7b but for different depths; a) 27 km, b) 100 km