

---

This manuscript is a preprint that has been submitted to EarthArXiv. The authors intend to submit the manuscript for publication. Subsequent versions of this manuscript may have slightly different content. If accepted, the final version of this manuscript will be available via the 'Peer-reviewed Publication DOI' link on the right-hand side of this webpage. Please feel free to contact any of the authors; we welcome feedback.

February 4, 2023

---



11 Through its vastness, resilience and biogeochemical complexity, the ocean offers  
12 humanity some of the largest potential natural pathways for removing carbon dioxide from the  
13 atmosphere while avoiding new sources of anthropogenic emissions. In proposing a network of  
14 new marine protected areas in service of global ocean conservation, Sala et al.<sup>1</sup> described a  
15 potentially large climate benefit of such a strategy: a reduction or elimination in carbon  
16 emissions that can result from the resuspension and subsequent remineralization of organic  
17 matter in ocean sediments following benthic trawling and dredging. Sala et al.<sup>1</sup> estimated that as  
18 much as 1.47 Pg CO<sub>2</sub> yr<sup>-1</sup> — equivalent to 4% of all global CO<sub>2</sub> emissions in 2020<sup>2</sup> — could be  
19 produced globally in the first year disturbance associated with benthic trawling, with CO<sub>2</sub>  
20 “emissions” from trawling activity declining after roughly a decade to 0.58 Pg CO<sub>2</sub> yr<sup>-1</sup>.

21 However, using the published output from a series of experiments in an inverse ocean  
22 circulation model<sup>3</sup> to account explicitly for the timescale of atmospheric ventilation, we show  
23 that the Sala et al.<sup>1</sup> estimate of global CO<sub>2</sub> efflux in the first year after trawling must be  
24 discounted by at least 33%, yielding an adjusted flux of 0.99 Pg CO<sub>2</sub> yr<sup>-1</sup>, if one aims to  
25 ultimately estimate emissions to the atmosphere (Fig. 1; Extended Data Fig. 2; Extended Data  
26 Table 1). Similarly, the cumulative CO<sub>2</sub> emissions attributable under the Sala et al.<sup>1</sup> model to a  
27 continuous pattern of global benthic trawling activity over a 30-year timescale must be  
28 discounted by at least 12% (compare 19.6 and 17.2 Pg CO<sub>2</sub>; Fig. 1; Extended Data Table 1).

29 We do not disagree that benthic trawling represents a severe source of disturbance to  
30 many benthic ecosystems<sup>4,5</sup> and that, under certain conditions, some of the organic carbon stored  
31 in ocean sediments may be remineralized back into CO<sub>2</sub> when subjected to disturbances such as  
32 benthic trawling and deep-sea mining. In addition, much of the world’s trawling activity takes  
33 place in the relatively shallow waters of continental shelves (Extended Data Fig 1; Extended

34 Data Table 1), where one might assume the shallow bottom water depth would result in rapid  
35 communication of remineralized carbon to the atmosphere. However, current empirical evidence  
36 and modeling studies, including our analysis, do not support a universal connection between  
37 physical disturbance, remineralization and consequent transmission of any remineralized CO<sub>2</sub>  
38 back to the atmosphere.<sup>6</sup> A complete description of our methods and calculations, including a  
39 validation of the use of the Siegel et al. inverse ocean circulation model output using an  
40 independent dataset of global ocean mixed layer depths<sup>7</sup>, is presented in the Supporting  
41 Information.

42 While Sala et al.<sup>1</sup> acknowledged they did not know what fraction of the remineralized  
43 CO<sub>2</sub> was likely to reach the atmosphere, their choice of the term “emissions” to describe the  
44 transformation and flux and of carbon from sediment to bottom waters, and their presentation of  
45 restrictions on bottom trawling as a promising nature-based solution to climate change predicated  
46 on avoided atmospheric carbon emissions, continue to support misinterpretation and confusion  
47 over their results, making imperative the need for adjusted estimates of the sort we present here.  
48 The original, uncorrected estimates of sediment CO<sub>2</sub> efflux contained in Sala et al.<sup>1</sup> have already  
49 been advanced as a low-cost-per-ton climate mitigation solution that could be included today in  
50 nationally determined contributions (NDCs) under the 2015 Paris Agreement<sup>8</sup>, and media have  
51 directly compared these uncorrected sediment fluxes with the global atmospheric emissions of  
52 the airline industry.<sup>9</sup>

53 Two overarching patterns can be seen in the results of our accounting for ventilation  
54 time. First, globally and for all exclusive economic zones (EEZs) for which we produced  
55 adjusted estimates, the relative discount that must be applied to account for ventilation time  
56 (measured in terms of percent deviation) becomes smaller as one considers cumulative emissions

57 from continuous trawling activity over longer time horizons (Extended Data Fig. 2; Extended  
58 Data Table 1; Supplementary Table 1).

59         Second, because atmospheric ventilation time is a function of both water depth and ocean  
60 circulation patterns<sup>3</sup>, the effect of this accounting does not manifest uniformly across different  
61 ocean basins and EEZs (Fig. 2). For nations whose EEZs are uniformly shallow, or where the  
62 benthic trawling activity is concentrated in waters of relatively shallow bottom depth, this  
63 accounting for ventilation time requires a relatively minor adjustment. In contrast, consideration  
64 of this fundamental feature of the ocean results in a dramatic adjustment for nations in whose  
65 EEZs the benthic trawling activity is concentrated in waters of relatively deep bottom depth. The  
66 average bottom depth of the trawled EEZ area, weighted by mass according to the efflux  
67 estimated by Sala et al.<sup>1</sup> for each model grid square, explained almost three-quarters of the  
68 variation in the percent deviation between our adjusted emissions estimates and the original  
69 estimates of CO<sub>2</sub> efflux.

70         For example, the Sala et al. estimate of CO<sub>2</sub> efflux from sediments in the Chinese EEZ  
71 must be discounted by just 3.1% to obtain an estimate of cumulative emissions to the atmosphere  
72 over 30 years (Fig. 2; Extended Data Table 1). China is responsible for an average of 4.4 Mt yr<sup>-1</sup>  
73 in landings from benthic trawling<sup>10</sup> — far more than any other country — but the nation's  
74 trawling activity is concentrated in waters whose average bottom depth, weighted by mass  
75 according to the efflux estimated by Sala et al.<sup>1</sup>, is just 25 m (Extended Data Table 1). However,  
76 a similar 30-year estimate of cumulative CO<sub>2</sub> efflux from continuous benthic trawling in the EEZ  
77 of Morocco — responsible for the fifth-most global landings from benthic trawling, at 1.9 Mt  
78 biomass yr<sup>-1</sup> — must be discounted by more than 45% (Fig. 2; Extended Data Table 1). The

79 mass-weighted average bottom depth of the portion of the Moroccan EEZ that is subjected to  
80 benthic trawling is 473 m (Extended Data Table 1).

81 While we present adjusted estimates of cumulative emissions for continuous trawling  
82 over several additional timescales (Extended Data Table 2; Supplementary Table 1), we  
83 considered a 30-year timescale most relevant to the time horizon over which the  
84 Intergovernmental Panel on Climate Change has concluded we must act to achieve global net  
85 zero emissions by mid-century.<sup>2</sup> We believe there are additional grounds on which one might  
86 question the estimates presented in Sala et al.<sup>1</sup> — for example, the implicit assumption that much  
87 of the sediment organic carbon hypothesized to be remineralized through benthic trawling would  
88 not otherwise be remineralized by natural biogeochemical processes, the failure to consider  
89 potential feedbacks associated with the release of nutrients, and apparent gaps in the AIS data  
90 used to drive much of the spatial analysis. Concerning the latter deficiency, which has been  
91 addressed in a previous reply<sup>11</sup>, we found there were no efflux data in the Sala et al.<sup>1</sup> dataset for  
92 model grid squares in the EEZs of Myanmar, Pakistan, Bangladesh or Guyana, even though  
93 these countries are responsible for at least 1.0 Mt yr<sup>-1</sup> in landings from benthic trawling<sup>10</sup>  
94 (Supplementary Table 1). We note also that the Sala et al.<sup>1</sup> analysis did not extend to areas of the  
95 emerging Arctic, where sea ice retreat and climate-driven migration of fish stocks could  
96 incentivize the trawling of large sections of seabed that have not been previously subjected to  
97 anthropogenic disturbance.<sup>12</sup>

98 Our results — based on an accounting for just one of many potential sources of  
99 uncertainty associated with the estimation of carbon emissions from anthropogenic disturbance  
100 of ocean sediments — suggest that any natural climate solution based on restricting or limiting  
101 the footprint of benthic trawling must be carefully scrutinized for scientific integrity, and that

102 ocean scientists with an understanding of both biogeochemistry and physical oceanography  
103 should be involved in the design and implementation of any such schemes. Perhaps most  
104 importantly, given the striking geographic heterogeneity we identified in the potential for  
105 atmospheric emissions from benthic trawling over timescales relevant for climate mitigation,  
106 there are serious implications for equity since not all coastal nations will likely be able to share  
107 equally in the benefits that would accrue from these restrictions or limitations.

108

### 109 **Data availability**

110 All input datasets used in this analysis are available at [https://github.com/jamesrco/global-](https://github.com/jamesrco/global-trawling-CO2)  
111 [trawling-CO2](https://github.com/jamesrco/global-trawling-CO2) or by direct download from the URLs provided in the text. The results of our  
112 analysis — including output data and adjusted emissions estimates by exclusive economic zone  
113 — are available at [https://github.com/jamesrco/global-trawling-](https://github.com/jamesrco/global-trawling-CO2/tree/main/data/global_trawling/derived/output)  
114 [CO2/tree/main/data/global\\_trawling/derived/output](https://github.com/jamesrco/global-trawling-CO2/tree/main/data/global_trawling/derived/output)

115

### 116 **Code availability**

117 The MATLAB and R code used in this analysis is available at  
118 <https://github.com/jamesrco/global-trawling-CO2>

119

### 120 **Acknowledgements**

121 This work was funded a gift from the Bezos Earth Fund to EDF. J.R.C. also acknowledges  
122 philanthropic support from Additional Ventures. We thank S. Doney for critical feedback on the  
123 design of our analysis, J. Mayorga for assistance with code used in the original Sala et al.<sup>1</sup>

124 analysis and D. Amrhein, G. Carroll, D. Cherian, S. Gaines, D. Gordon, M. Long, J. Mason, M.  
125 Moritsch, D. Rader, S. Saul, and E. Schwaab for discussions.

126

127 **Author contributions**

128 All authors contributed to development of the ideas presented in this Comment. J.R.C. performed  
129 the analysis, prepared the figures and wrote the first draft of the manuscript. All authors  
130 contributed to subsequent drafts of the manuscript and approved the final submitted version.

131

132 **Competing interests**

133 The authors declare no competing interests.

134

135 **Additional information**

136 **Supplementary Information** is available for this paper.


137 **Correspondence and requests for materials** should be addressed James R. Collins.



138 **References**


139

- 140 1. Sala, E. *et al.* Protecting the global ocean for biodiversity, food and climate. *Nature* **592**, 397–  
 141 402 (2021).
- 142 2. Riahi, K. *et al.* Mitigation pathways compatible with long-term goals. in *Climate Change*  
 143 *2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth*  
 144 *Assessment Report of the Intergovernmental Panel on Climate Change* (eds. Shukla, P. R. *et*  
 145 *al.*) (Cambridge University Press, 2022).
- 146 3. Siegel, D. A., DeVries, T., Doney, S. C. & Bell, T. Assessing the sequestration time scales of  
 147 some ocean-based carbon dioxide reduction strategies. *Environ. Res. Lett.* **16**, 104003 (2021).
- 148 4. Pitcher, C. R. *et al.* Trawl impacts on the relative status of biotic communities of seabed  
 149 sedimentary habitats in 24 regions worldwide. *Proc. Natl. Acad. Sci.* **119**, e2109449119  
 150 (2022).
- 151 5. Eigaard, O. R. *et al.* The footprint of bottom trawling in European waters: distribution,  
 152 intensity, and seabed integrity. *ICES J. Mar. Sci.* **74**, 847–865 (2017).
- 153 6. Epstein, G., Middelburg, J. J., Hawkins, J. P., Norris, C. R. & Roberts, C. M. The impact of  
 154 mobile demersal fishing on carbon storage in seabed sediments. *Glob. Change Biol.*  
 155 *gcb.16105* (2022) doi:10.1111/gcb.16105.
- 156 7. Holte, J., Talley, L. D., Gilson, J. & Roemmich, D. An Argo mixed layer climatology and  
 157 database. *Geophys. Res. Lett.* **44**, 5618–5626 (2017).
- 158 8. Claes, J., Hopman, D., Jaeger, G. & Rogers, M. *Blue carbon: The potential of coastal and*  
 159 *oceanic climate action*. 32 [https://www.mckinsey.com/business-functions/sustainability/our-](https://www.mckinsey.com/business-functions/sustainability/our-insights/Blue-carbon-The-potential-of-coastal-and-oceanic-climate-action)  
 160 [insights/Blue-carbon-The-potential-of-coastal-and-oceanic-climate-action](https://www.mckinsey.com/business-functions/sustainability/our-insights/Blue-carbon-The-potential-of-coastal-and-oceanic-climate-action) (2022).
- 161 9. McVeigh, K. Bottom trawling releases as much carbon as air travel, landmark study finds. *The*  
 162 *Guardian* (2021).
- 163 10. Pauly, D., Zeller, D. & Deng Palomares, M. L. Sea Around Us Concepts, Design and  
 164 Data. (2020).
- 165 11. Hilborn, R. & Kaiser, M. J. A path forward for analysing the impacts of marine protected  
 166 areas. *Nature* **607**, E1–E2 (2022).
- 167 12. Fauchald, P. *et al.* Poleward shifts in marine fisheries under Arctic warming. *Environ.*  
 168 *Res. Lett.* **16**, 074057 (2021).
- 169 13. Becker, J. J. *et al.* Global bathymetry and elevation data at 30 arc seconds resolution:  
 170 SRTM30\_PLUS. *Mar. Geod.* **32**, 355–371 (2009).
- 171 14. Flanders Marine Institute. Maritime Boundaries Geodatabase: Maritime Boundaries and  
 172 Exclusive Economic Zones (200NM), version 11. (2019).
- 173 15. Steadman, D. *et al.* *New perspectives on an old fishing practice: Scale, context and*  
 174 *impacts of bottom trawling*. 44 (2021).



175  
 176  
 177  
 178  
 179  
 180  
 181  
 182  
 183  
 184  
 185  
 186  
 187  
 188  
 189  
 190  
 191  
 192  
 193  
 194  
 195  
 196  
 197  
 198

**Fig. 1 | Cumulative carbon emissions from benthic trawling according to original model of Sala et al.<sup>1</sup> and an adjusted model (this study) that accounts for timescale of ventilation to the atmosphere.** To isolate the role of bottom depth and ocean circulation for purposes of our analysis, we assumed for our adjusted emissions estimates (solid lines) the same underlying model of sediment remineralization upon which Sala et al.<sup>1</sup> based their estimates (dashed lines). **a**, Cumulative carbon emissions from benthic trawling activity for all bottom depths within the Sala et al model domain. **b-d**, Estimates of emissions originating from trawling across three ranges of bottom depth: **(b)** Waters with bottom depths  $\leq 200$  m, **(c)** waters with bottom depths  $> 200$  m, and **(d)** waters with bottom depths  $> 400$  m. Trawling in waters with bottom depths  $\leq 200$  m accounted for 78% of the total C remineralization estimated by Sala et al.<sup>1</sup> for the first year of trawling, while trawling in waters with bottom depths  $> 400$  m accounted for 11% of total remineralization; see Extended Data Fig. 1 and Extended Data Table 1. **e-f**, Estimates of emissions originating from benthic trawling in two exclusive economic zones (EEZs) with divergent bottom depth profiles: **(e)** China, where the emissions-weighted mean bottom depth of the area subjected to benthic trawling is just 25 m, and **(f)** Morocco, where the weighted mean bottom depth of the area subjected to benthic trawling is 473 m. The mean bottom depth for each grid square in the Sala et al.<sup>1</sup> sediment CO<sub>2</sub> efflux dataset was retrieved from the database described in ref. <sup>13</sup>. EEZ emissions estimates were calculated by applying the EEZ boundaries described in ref. <sup>14</sup>; a link to the appropriate code is provided in the Code Availability statement. Estimates of cumulative emissions over a 200-year time scale for the same bottom depth ranges and EEZs are presented in Extended Data Fig. 2



199  
200


201 **Fig. 2 | Effect of mass-weighted EEZ bottom depth on carbon emissions to the atmosphere**  
 202 **from benthic trawling.** Values on the y-axis represent the percent deviation of our adjusted  
 203 estimate (accounting for the timescale of atmospheric ventilation) from the original emissions  
 204 estimate of Sala et al.<sup>1</sup> This percent deviation is plotted against the weighted mean bottom depth  
 205 of the EEZ area subjected to benthic trawling for the 30 countries with the greatest average  
 206 landings from benthic trawling from 2009–2018 (according to ref.<sup>10</sup>) and for the top 10 countries  
 207 ranked by percentage of total EEZ catch from bottom trawling (according to ref.<sup>15</sup>). Each circle  
 208 represents a different country; symbol size corresponds to the quantity of biomass landed within  
 209 that country from benthic trawling. The red line shows the result of a Model I linear regression  
 210 ( $R^2 = 0.74$ ;  $p < 0.0001$ ). The right-hand panel is an expansion of the shaded region shown on the  
 211 main panel. All supporting data is presented in Supplementary Table 1.

212 \*Percent deviation based on comparison of estimates of cumulative emissions after 30 years of  
 213 benthic trawling; percent deviations between our estimates and those of Sala et al. for other time  
 214 horizons are presented in Extended Data Table 1 and Supplementary Table 1.

215 †Geographic subsetting was performed using the exclusive economic zone boundaries described  
 216 in ref.<sup>14</sup>. We weighted the bottom depth of the area within each EEZ that was subjected to  
 217 benthic trawling according to the rates of remineralization estimated by Sala et al.<sup>1</sup> The mean  
 218 bottom depth for each grid square in the Sala et al. CO<sub>2</sub> efflux dataset was retrieved the database  
 219 described in ref.<sup>13</sup>; this depth was then weighted by the mass of CO<sub>2</sub> that Sala et al. estimated  
 220 was produced in that grid square in the first year of benthic trawling. A link to the appropriate  
 221 code is provided in the Code Availability statement.

222 ‡Catch data were retrieved from the Sea Around Us database<sup>10</sup> using the code to which a link is  
 223 provided in the Code Availability statement.

224  
225




226

227

228 **Extended Data Fig. 1 | Mass distribution of carbon remineralized as a result of benthic**  
229 **trawling according to bottom depth.** The figure was produced by segregating sediment CO<sub>2</sub>  
230 efflux estimates reported by Sala et al.<sup>1</sup> into 100 m bins according to bottom depth. The mean  
231 bottom depth for each grid square in the Sala et al. CO<sub>2</sub> efflux dataset was retrieved from the  
232 database described in ref. <sup>13</sup>. The fraction of the total mass present within selected depth bins is  
233 given in Extended Data Table 1; all supporting data are available via the link provided in the  
234 Code Availability statement.

235

236



237  
238

239 **Extended Data Fig. 2 | Cumulative carbon emissions from benthic trawling according to**  
 240 **original model of Sala et al.<sup>1</sup> and an adjusted model (this study) that accounts for timescale**  
 241 **of ventilation to the atmosphere.** This figure presents results for the same bottom depth ranges  
 242 and exclusive economic zones (EEZs) shown in Fig. 1, but over a 200-year time scale. We  
 243 assumed for our adjusted emissions estimates (solid lines) the same underlying model of  
 244 sediment remineralization upon which Sala et al.<sup>1</sup> based their estimates (dashed lines). **a,**  
 245 Cumulative carbon emissions from benthic trawling activity for all bottom depths within the Sala

246 et al model domain. **b-d**, Estimates of emissions originating from trawling across three ranges of  
247 bottom depth: **(b)** Waters with bottom depths  $\leq 200$  m, **(c)** waters with bottom depths  $> 200$  m,  
248 and **(d)** waters with bottom depths  $> 400$  m. **e-f**, Estimates of emissions originating from benthic  
249 trawling in two exclusive economic zones (EEZs) with divergent bottom depth profiles: **(e)**  
250 China, where the emissions-weighted mean bottom depth of the area subjected to benthic  
251 trawling is just 25 m, and **(f)** Morocco, where the weighted mean bottom depth of the area  
252 subjected to benthic trawling is 473 m.  
253

<b>Bottom depth* range</b>	<b>Fraction by mass of total est. sediment C remineralization in first year<sup>†</sup></b>
≤ 100 m	0.62
≤ 200 m	0.78
> 200 m	0.22
≤ 400 m	0.89
> 400 m	0.11

254

255 **Extended Data Table 1 | Fraction by mass of total estimated sediment C remineralization**  
 256 **from benthic trawling according to bottom depth.** The complete mass distribution, divided  
 257 into 100 m depth bins, is presented in Extended Data Fig. 1.

258 \*The mean bottom depth for each grid square in the Sala et al.<sup>1</sup> sediment CO<sub>2</sub> efflux dataset was  
 259 retrieved from the database described in ref. <sup>13</sup>.

260 <sup>†</sup>For the first year after start of benthic trawling activity, from the model of sediment  
 261 remineralization posited by Sala et al.<sup>1</sup>

Bottom depth range or exclusive economic zone (EEZ) area	Avg. landings from benthic trawling, 2009-2018*	Weighted mean bottom depth of area subjected to benthic trawling†	Fraction by mass of total est. C remineralization‡	Cumulative emissions after years of continuous trawling											
				Mt bio-mass	Rank by EEZ§	m	Rank by EEZ§	—	Rank by EEZ§	After 1 yr Pg CO <sub>2</sub>	Rank by EEZ§	After 5 yr Pg CO <sub>2</sub>	After 30 yr Pg CO <sub>2</sub>	Rank by EEZ§	After 200 yr Pg CO <sub>2</sub>
Global trawled ocean area, all bottom depths			1												
Global trawled ocean area, bottom depths <sup>  </sup> ≤ 200 m			0.78												
Global trawled ocean area, bottom depths <sup>  </sup> > 200 m			0.22												
Global trawled ocean area, bottom depths <sup>  </sup> > 400 m			0.11												



Bottom depth interval or exclusive economic zone (EEZ) area	Avg. landings from benthic trawling, 2009-2018*		Weighted mean bottom depth of area subjected to benthic trawling†		Fraction by mass of total est. C remineralization‡	Cumulative emissions after years of continuous trawling							
	Mt bio-mass	Rank by EEZ§	m	Rank by EEZ§		—	Rank by EEZ§	After 1 y Pg CO <sub>2</sub>	After 5 y Pg CO <sub>2</sub>	After 30 y Pg CO <sub>2</sub>	After 200 y Pg CO <sub>2</sub>	Rank by EEZ§	
China	4.4	1	25	27	0.52	1							
Original estimate (Sala et al.)							0.769	1	2.54	10.2	1	62.2	1
Adjusted estimate (this analysis)							0.605	1	2.22	9.92	1	61.8	1
Percent deviation							-21	25	-13	-3.1	26	-0.53	26
India	2.1	4	54	24	0.002	24							
Original estimate (Sala et al.)							$2.88 \times 10^{-3}$	24	$9.53 \times 10^{-3}$	$3.84 \times 10^{-3}$	24	0.233	24
Adjusted estimate (this analysis)							$2.02 \times 10^{-3}$	19	$8.06 \times 10^{-3}$	0.035	23	0.221	23
Percent deviation							-30	21	-15	-9	22	-5.4	16
Indonesia	2.4	3	39	25	$1.2 \times 10^{-8}$	31							
Original estimate (Sala et al.)							$1.71 \times 10^{-8}$	31	$5.64 \times 10^{-8}$	$2.27 \times 10^{-7}$	31	$1.38 \times 10^{-6}$	31
Adjusted estimate (this analysis)							$1.56 \times 10^{-8}$	31	$5.41 \times 10^{-8}$	$2.25 \times 10^{-7}$	31	$1.38 \times 10^{-6}$	31
Percent deviation							-8.4	27	-4	-0.9	27	-0.15	27
Morocco	1.9	5	473	2	0.0053	20							
Original estimate (Sala et al.)							$7.85 \times 10^{-3}$	20	$2.59 \times 10^{-3}$	0.105	20	0.635	20
Adjusted estimate (this analysis)							$1.06 \times 10^{-3}$	22	$8.42 \times 10^{-3}$	$5.75 \times 10^{-2}$	20	0.489	21
Percent deviation							-86	5	-68	-45	6	-23	6
Vietnam	2.5	2	29	26	0.00062	28							
Original estimate (Sala et al.)							$9.2 \times 10^{-4}$	28	$3.04 \times 10^{-3}$	$1.23 \times 10^{-2}$	28	$7.44 \times 10^{-2}$	28
Adjusted estimate (this analysis)							$5.66 \times 10^{-4}$	27	$2.12 \times 10^{-3}$	$1.11 \times 10^{-2}$	27	$7.31 \times 10^{-2}$	28
Percent deviation							-38	18	-30	-9.8	20	-1.7	24

264 **Extended Data Table 2 | Fisheries landings and cumulative CO<sub>2</sub> emissions estimates from benthic trawling for selected bottom**  
265 **depth ranges and exclusive economic zones, with rankings where appropriate.** Table presents cumulative emissions estimates  
266 from both the original Sala et al.<sup>1</sup> model and this study, which applies the methods described in the text to account for ventilation time  
267 to the atmosphere. Data and emissions estimates for additional time horizons and EEZs are presented in Supplementary Table 1.  
268 \*Catch data were retrieved from the Sea Around Us database<sup>10</sup> using the code to which a link is provided in the Code Availability  
269 statement.  
270 †We weighted the bottom depth of the area within each EEZ that was subjected to benthic trawling according to the rates of  
271 remineralization estimated by Sala et al.<sup>1</sup> The mean bottom depth for each grid square in the Sala et al.<sup>1</sup> CO<sub>2</sub> efflux dataset was  
272 retrieved from the database described in ref. <sup>13</sup>; this depth was then weighted by the mass of CO<sub>2</sub> that Sala et al.<sup>1</sup> estimated was  
273 produced in that grid square in the first year of benthic trawling.  
274 ‡For the first year after start of benthic trawling activity, from the model of sediment remineralization posited by Sala et al.<sup>1</sup>  
275 §Geographic subsetting was performed using the exclusive economic zone boundaries described in ref. <sup>14</sup>. A link to the appropriate  
276 code is provided in the Code Availability statement.  
277 ‖The mean bottom depth for each grid square in the Sala et al.<sup>1</sup> sediment CO<sub>2</sub> efflux dataset was retrieved from ref. <sup>13</sup>.  
278

Supplementary information

for

**Atmospheric carbon emissions from benthic trawling depend on water depth  
and ocean circulation**

James R. Collins<sup>1,2</sup>✉, Kristin M. Kleisner<sup>1</sup>, Rodney M. Fujita<sup>1</sup>, Robert E. Boenish<sup>1</sup>

<sup>1</sup> Environmental Defense Fund, New York, NY 10010, USA

<sup>2</sup> Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution,  
Woods Hole, MA 02543, USA

✉ Correspondence via e-mail: [jcollins@edf.org](mailto:jcollins@edf.org)

## Table of Contents

Description of methods .....	3
Supplementary Table 1 .....	6
Works referenced in Supplementary information file .....	12

## Methods

### Objective and assumptions

Sala et al.<sup>1</sup> combined a global dataset of sediment carbon concentrations<sup>2</sup>, fishing vessel track densities derived from automatic identification systems (AIS) data, prior findings concerning the sediment penetration depth and track width of various types of benthic trawling gear<sup>3,4</sup>, and a simple model of remineralization, to estimate the remineralization rate of sediment organic carbon due to benthic trawling and the consequent flux of CO<sub>2</sub> to overlying bottom waters. We sought to correct these flux estimates for ventilation time — the time it takes for an interior ocean water mass to come into contact with the atmosphere as a function of bottom depth and ocean circulation — to predict the fraction remineralized CO<sub>2</sub> one might expect to reach the atmosphere as a source of emissions. To focus our analysis on the importance of accounting for ventilation time, we used the same model of sediment remineralization upon which Sala et al. based their uncorrected estimates, including the authors' assumptions concerning the (lesser) fraction of carbon remineralized within a given model grid square with each successive year of continuous trawling activity. As we note in the main text, there are several potentially serious deficiencies in this underlying model; we did not formally address these sources of error in our analysis, leaving the development of a more realistic model of remineralization to future study.

### Adjustment of Sala et al.<sup>1</sup> efflux estimates for ventilation time

The primary input for our analysis was the Sala et al. data file containing estimates of sediment CO<sub>2</sub> efflux in Mg CO<sub>2</sub> km<sup>-2</sup> (1 km × 1 km resolution; file “co2\_efflux.tif,” available at <https://doi.org/10.25349/D9N89M>). Because these data were very sparse (flux values of zero were assigned to > 99% of grid squares), we performed our analysis on a subset of the data that contained only nonzero values while retaining the coordinates to facilitate spatial subsetting. The R and MATLAB codes we used to perform these and the other calculations described below are available at <https://github.com/jamesrco/global-trawling-CO2>. Calculations in R<sup>5</sup> were performed using the packages `sp`<sup>6,7</sup>, `rgdal`<sup>8</sup>, `sf`<sup>9</sup>, `raster`<sup>10</sup>, `data.table`<sup>11</sup>, `R.matlab`<sup>12</sup>, and `maptools`<sup>13</sup>.

To correct the Sala et al. flux estimates for ventilation time, we modified the published output from a series of experiments by Siegel et al.<sup>14</sup> in an inverse ocean circulation model. Siegel et al.<sup>14</sup> sought to estimate the fraction of injected CO<sub>2</sub> that would be retained at various locations in the ocean over different timescales; we sought to estimate the complementary quantity (i.e., the fraction of remineralized CO<sub>2</sub> that would be emitted to the atmosphere after a given time  $t$ ). We therefore used in our analysis the complement (i.e.,  $1 - f(t)$ ) of Siegel et al.'s *sequestration fractions*, which we termed *emissions fractions*. The Siegel et al. sequestration fractions (gridded data of 2° × 2° horizontal resolution and vertical resolution of 48 depths, proceeding from resolution of the underlying inverse circulation model) and accompanying MATLAB code were retrieved from <https://doi.org/10.6084/m9.figshare.15228690.v2>. A file containing the 48 depths in the model is available at [https://github.com/jamesrco/global-trawling-CO2/blob/main/data/global\\_trawling/derived/benthic\\_seqfractions/OCIM\\_modelDepths.csv](https://github.com/jamesrco/global-trawling-CO2/blob/main/data/global_trawling/derived/benthic_seqfractions/OCIM_modelDepths.csv). Assuming that trawling begins in some theoretical year 1, the emissions fraction for a given year  $n$  represents the fraction of CO<sub>2</sub> remineralized and discharged into bottom waters at a given location  $n$  years ago that will have reached the atmosphere by year  $1 + n$ .

For each year of interest (1 to 200 yr in one-year increments, and then 300 to 1000 yr in increments of 100 yr) we retrieved from the Siegel et al. model output the values closest in space (based on latitude and longitude) to the coordinates of the centroid of each grid square in the Sala et al. CO<sub>2</sub> efflux dataset. We then selected the depth in the Siegel et al. model output that was closest to the mean bottom depth of the Sala et al. 1 km × 1 km model grid square. We assigned bottom depths to the Sala et al. grid squares from version 8 of the SRTM30 digital elevation model, a 1 km × 1 km resolution global digital elevation model that includes ocean bathymetry.<sup>15</sup> Details concerning this DEM are available at <https://eatlas.org.au/data/uuid/80301676-97fb-4bdf-b06c-e961e5c0cb0b>.

We then adjusted the Sala et al. flux estimate for each grid square using the best-fit emissions fraction for each ventilation timescale. We calculated cumulative atmospheric emissions estimates for a variety of time horizons (presented in Fig. 1, Extended Data Fig. 2, Extended Data Table and Supplementary Table 1) by summing the adjusted flux estimates over the indicated timescale.

### First-order validation using mixed layer depth data

Because the inverse ocean circulation model used by Siegel et al. is not optimized for the relatively shallow, near coastal waters where most benthic trawling takes place (Extended Data Fig. 1; Extended Data Table 1), we validated our approach based on emissions fractions using an independent calculation based on the maximum annual mixed layer depth ( $MLD_{max}$ ). For purposes of this comparison, we assumed conservatively that any remineralized CO<sub>2</sub> discharged from sediments into waters whose bottom depth was  $\leq MLD_{max}$  at that location would reach the atmosphere within one year. By comparing an estimate of annual atmospheric emissions based on this criterion to the one we obtained for the first year of trawling using the Siegel et al. emissions fractions, we sought to make a crude, first-order validation of our approach.

From the April 2022 version of a 1° × 1° resolution global database of mixed layer depths<sup>16,17</sup> (available at <http://mixedlayer.ucsd.edu>) we first extracted the  $MLD_{max}$  for the coordinates nearest the centroid of each grid square in the Sala et al.<sup>1</sup> CO<sub>2</sub> efflux dataset. We then subsetted the Sala et al. efflux dataset to include only those grid squares whose bottom depth (assigned from ref.<sup>15</sup>, as described above) was  $\leq MLD_{max} \pm 10$  m. By summing the flux values within this subset of grid squares, we obtained an estimate of first-year atmospheric emissions of 0.92 Pg CO<sub>2</sub>. This estimate fell within 0.07 Pg (approx. 7%) of the atmospheric emissions estimate of 0.99 Pg CO<sub>2</sub> yr<sup>-1</sup> we produced for the first year using the emissions fractions derived from the Siegel et al.<sup>14</sup> model output (Extended Data Table 1). This favorable comparison suggested that the inverse ocean circulation model used in ref.<sup>14</sup> did not lead to a fatal bias, at least for purposes of our simple analysis. Further, almost one-fourth of the global potential CO<sub>2</sub> efflux (on a mass basis, as estimated by Sala et al.<sup>1</sup>) originates in sediments underlying waters of 200 m or greater bottom depth (Extended Data Table 1), suggesting that the consequences of benthic trawling are not confined only to shallow coastal waters. Substantial off-shelf sediment transport — in which sediment resuspended by benthic trawling was advected by ocean currents into deeper waters — was also documented independently in another recent study<sup>18</sup>, validating the use of an approach that attempts to account for the effects of ocean circulation.

### **Subsetting by exclusive economic zone (EEZ)**

We used the EEZ boundary shapefiles described in ref. <sup>19</sup> to subset our adjusted estimates (and those of Sala et al.) to waters of the 30 countries with the greatest average landings from benthic trawling from 2009-2018 (according to ref. <sup>20</sup>) and the top 10 countries ranked by percentage of total EEZ catch from bottom trawling (according to ref. <sup>21</sup>). Results are presented in Extended Data Table 1 and Supplementary Table 1.

### **Catch data**

Catch data for annual landings from benthic trawling (in Mt biomass yr<sup>-1</sup>) were extracted for these same countries from the Sea Around Us database<sup>20</sup> using the R code to which a link is provided in the Data availability statement. We extracted landings data for 2009-2018 and averaged these to obtain the values we report in Extended Data Table 1 and Supplementary Table 1.

### Supplementary Table 1 | Fisheries landings and cumulative CO<sub>2</sub> emissions estimates from benthic trawling for selected bottom depth ranges and EEZs, with rankings where appropriate

Bottom depth range or exclusive economic zone (EEZ) area	Avg. landings from benthic trawling, 2009-2018*		Weighted mean bottom depth of area subjected to benthic trawling <sup>†</sup>	Rank by EEZ <sup>§</sup>	Fraction by mass of total est. global sediment C remineralization <sup>‡</sup>	Cumulative emissions after years of continuous trawling											
	Mt biomass	Rank by EEZ <sup>§</sup>				Rank by EEZ <sup>§</sup>	Rank by EEZ <sup>§</sup>	1	5	10	25	30	50	75	100	200	Rank by EEZ <sup>§</sup>
			m	—		Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	
Global trawled ocean area, all bottom depths				1													
Original estimate (Sala et al.)						1.47	4.87	7.91	16.7	19.6	31.3	46	60.6	119			
Adjusted estimate (this analysis)						0.991	3.74	6.49	14.5	17.2	28.1	42	55.9	112			
Percent deviation						-33	-23	-18	-13	-12	-10	-8.7	-7.8	-5.9			
Global trawled ocean area, bottom depths <sup>  </sup> ≤ 200 m				0.78													
Original estimate (Sala et al.)						1.15	3.81	6.19	13.1	15.4	24.5	36	47.4	93.2			
Adjusted estimate (this analysis)						0.962	3.56	6.1	13.4	15.9	25.7	38	50.3	99.7			
Percent deviation						-16	-6.6	-1.5	2.8	3.4	4.8	5.7	6.2	7			
Global trawled ocean area, bottom depths <sup>  </sup> > 200 m				0.22													
Original estimate (Sala et al.)						0.324	1.07	1.74	3.68	4.32	6.9	10.1	13.3	26.2			
Adjusted estimate (this analysis)						0.0283	0.185	0.392	1.07	1.32	2.42	3.94	5.55	12.4			
Percent deviation						-91	-83	-77	-71	-69	-65	-61	-58	-53			
Global trawled ocean area, bottom depths <sup>  </sup> > 400 m				0.11													
Original estimate (Sala et al.)						0.163	0.54	0.877	1.85	2.18	3.47	5.1	6.72	13.2			
Adjusted estimate (this analysis)						0.0069	0.0481	0.11	0.328	0.411	0.784	1.33	1.92	4.61			
Percent deviation						-96	-91	-87	-82	-81	-77	-74	-71	-65			
Angola	0.29	22	527	1	0.011	8											
Original estimate (Sala et al.)						0.0166	8	0.0549	0.0893	0.188	0.221	8	0.354	0.519	0.684	1.34	8
Adjusted estimate (this analysis)						0.0008	24	0.0058	0.0124	0.0344	0.0428	21	0.0846	0.158	0.252	0.745	14
Percent deviation						-95	2	-89	-86	-82	-81	1	-76	-70	-63	-45	1
Argentina	0.65	11	85	21	0.002	23											
Original estimate (Sala et al.)						0.003	23	0.0098	0.016	0.0338	0.0397	23	0.0633	0.0929	0.122	0.241	23
Adjusted estimate (this analysis)						0.0024	17	0.0084	0.0141	0.0307	0.0363	22	0.0587	0.0871	0.116	0.231	22
Percent deviation						-21	26	-14	-12	-9.1	-8.6	23	-7.3	-6.2	-5.6	-4.1	18



Supplementary information for Collins et al. preprint; to be submitted for publication

Bottom depth interval or exclusive economic zone (EEZ) area	Avg. landings from benthic trawling, 2009-2018*		Weighted mean bottom depth of area subjected to benthic trawling†	Rank by EEZ <sup>s</sup>	Fraction by mass of total est. global sediment C remineralization‡	Rank by EEZ <sup>s</sup>	Cumulative emissions after years of continuous trawling										Rank by EEZ <sup>s</sup>	
	Mt biomass	Rank by EEZ <sup>s</sup>					1	5	10	25	30	50	75	100	200			
			Pg CO <sub>2</sub>	Rank by EEZ <sup>s</sup>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Rank by EEZ <sup>s</sup>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>			
Bangladesh <sup>§</sup>	0.15	28																
Original estimate (Sala et al.)																		
Adjusted estimate (this analysis)																		
Percent deviation																		
Brazil	0.16	27	77	22	0.0016	25												
Original estimate (Sala et al.)							0.0023	25	0.0076	0.0124	0.0262	0.0308	25	0.0491	0.0721	0.095	0.187	25
Adjusted estimate (this analysis)							0.0017	20	0.0066	0.0114	0.0249	0.0294	24	0.0475	0.0702	0.0929	0.184	25
Percent deviation							-28	22	-13	-8.2	-5	-4.5	25	-3.4	-2.6	-2.2	-1.4	25
Canada	0.28	23	301	7	0.0074	16												
Original estimate (Sala et al.)							0.0108	16	0.0358	0.0582	0.123	0.144	16	0.231	0.338	0.446	0.877	16
Adjusted estimate (this analysis)							0.0023	18	0.0141	0.0286	0.0729	0.0882	15	0.151	0.232	0.314	0.656	17
Percent deviation							-78	8	-61	-51	-41	-39	7	-35	-32	-30	-25	5
China	4.4	1	25	27	0.52	1												
Original estimate (Sala et al.)							0.769	1	2.54	4.13	8.71	10.2	1	16.3	24	31.6	62.2	1
Adjusted estimate (this analysis)							0.605	1	2.22	3.82	8.4	9.92	1	16	23.7	31.3	61.8	1
Percent deviation							-21	25	-13	-7.6	-3.6	-3.1	26	-2	-1.3	-1	-0.53	26
Côte d'Ivoire	0.1	31	104	20	0.0006	29												
Original estimate (Sala et al.)							0.0008	29	0.0028	0.0046	0.0096	0.0113	29	0.018	0.0265	0.0349	0.0686	29
Adjusted estimate (this analysis)							0.0002	28	0.0014	0.0031	0.008	0.0096	29	0.0162	0.0244	0.0327	0.066	29
Percent deviation							-78	9	-49	-31	-17	-15	14	-10	-7.9	-6.4	-3.8	19
Denmark	0.32	21	158	15	0.027	4												
Original estimate (Sala et al.)							0.0396	4	0.131	0.213	0.449	0.528	4	0.843	1.24	1.63	3.21	4
Adjusted estimate (this analysis)							0.0301	3	0.107	0.178	0.385	0.454	3	0.731	1.08	1.43	2.84	4
Percent deviation							-24	23	-18	-16	-14	-14	15	-13	-13	-12	-11	11
Germany	0.088	33	19	29	0.009	11												
Original estimate (Sala et al.)							0.0132	11	0.0436	0.0708	0.15	0.176	11	0.281	0.412	0.543	1.07	11
Adjusted estimate (this analysis)							0.0132	7	0.0436	0.0708	0.15	0.176	8	0.281	0.412	0.543	1.07	9
Percent deviation							-0.055	30	-0.009	-0.005	-0.002	-0.002	30	-0.001	-8 × 10 <sup>-4</sup>	-6 × 10 <sup>-4</sup>	-3 × 10 <sup>-4</sup>	30

Supplementary information for Collins et al. preprint; to be submitted for publication

Bottom depth interval or exclusive economic zone (EEZ) area	Avg. landings from benthic trawling, 2009-2018*		Weighted mean bottom depth of area subjected to benthic trawling†	Rank by EEZ <sup>s</sup>	Fraction by mass of total est. global sediment C remineralization†	Rank by EEZ <sup>s</sup>	Cumulative emissions after years of continuous trawling											
	Mt biomass	Rank by EEZ <sup>s</sup>					1	5	10	25	30	50	75	100	200	Rank by EEZ <sup>s</sup>		
	Pg CO <sub>2</sub>	Rank by EEZ <sup>s</sup>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Rank by EEZ <sup>s</sup>	
Guinea	0.49	17	211	13	0.0008	26												
Original estimate (Sala et al.)							0.0011	26	0.0037	0.006	0.0127	0.015	26	0.0239	0.0351	0.0463	0.0909	26
Adjusted estimate (this analysis)							0.0006	26	0.0024	0.004	0.0091	0.0109	28	0.0182	0.028	0.0382	0.0808	27
Percent deviation							-49	15	-37	-33	-29	-28	10	-24	-20	-17	-11	12
Guinea-Bissau	0.42	18	144	17	0.0051	21												
Original estimate (Sala et al.)							0.0075	21	0.0247	0.0401	0.0847	0.0996	21	0.159	0.233	0.307	0.604	21
Adjusted estimate (this analysis)							0.0048	12	0.0199	0.0339	0.0743	0.088	16	0.144	0.215	0.287	0.578	19
Percent deviation							-36	20	-19	-16	-12	-12	17	-9.6	-8	-6.8	-4.3	17
Guyana <sup>‡</sup>	0.036	35																
Original estimate (Sala et al.)																		
Adjusted estimate (this analysis)																		
Percent deviation																		
Iceland	0.25	24	278	8	0.0085	14												
Original estimate (Sala et al.)							0.0125	14	0.0413	0.067	0.142	0.166	14	0.266	0.39	0.514	1.01	14
Adjusted estimate (this analysis)							0.0042	13	0.0181	0.0316	0.0727	0.0867	17	0.144	0.217	0.291	0.606	18
Percent deviation							-67	11	-56	-53	-49	-48	5	-46	-44	-43	-40	2
India	2.1	4	54	24	0.002	24												
Original estimate (Sala et al.)							0.0029	24	0.0095	0.0155	0.0327	0.0384	24	0.0613	0.09	0.119	0.233	24
Adjusted estimate (this analysis)							0.002	19	0.0081	0.0136	0.0296	0.035	23	0.0565	0.0836	0.111	0.221	23
Percent deviation							-30	21	-15	-12	-9.5	-9	22	-7.8	-7	-6.6	-5.4	16
Indonesia	2.4	3	39	25	1.2 × 10 <sup>-8</sup>	31												
Original estimate (Sala et al.)							1.71 × 10 <sup>-8</sup>	31	5.64 × 10 <sup>-8</sup>	9.16 × 10 <sup>-8</sup>	1.93 × 10 <sup>-7</sup>	2.27 × 10 <sup>-7</sup>	31	3.63 × 10 <sup>-7</sup>	5.33 × 10 <sup>-7</sup>	7.02 × 10 <sup>-7</sup>	1.38 × 10 <sup>-6</sup>	31
Adjusted estimate (this analysis)							1.56 × 10 <sup>-8</sup>	31	5.41 × 10 <sup>-8</sup>	8.96 × 10 <sup>-8</sup>	1.91 × 10 <sup>-7</sup>	2.25 × 10 <sup>-7</sup>	31	3.61 × 10 <sup>-7</sup>	5.30 × 10 <sup>-7</sup>	7.00 × 10 <sup>-7</sup>	1.38 × 10 <sup>-6</sup>	31
Percent deviation							-8.4	27	-4	-2.2	-1.1	-0.9	27	-0.57	-0.39	-0.3	-0.15	27
Ireland	0.13	30	219	12	0.0095	10												
Original estimate (Sala et al.)							0.0141	10	0.0465	0.0755	0.159	0.187	10	0.299	0.439	0.579	1.14	10
Adjusted estimate (this analysis)							0.008	10	0.0342	0.0595	0.131	0.155	11	0.25	0.37	0.49	0.976	11
Percent deviation							-43	16	-26	-21	-18	-17	13	-16	-16	-15	-14	10
Japan	1.7	6	267	9	0.0098	9												
Original estimate (Sala et al.)							0.0144	9	0.0477	0.0775	0.164	0.192	9	0.307	0.45	0.594	1.17	9
Adjusted estimate (this analysis)							0.0016	21	0.0207	0.0518	0.139	0.167	9	0.282	0.425	0.568	1.14	8
Percent deviation							-89	4	-57	-33	-15	-13	16	-8.2	-5.7	-4.4	-2.3	21

Supplementary information for Collins et al. preprint; to be submitted for publication

Bottom depth interval or exclusive economic zone (EEZ) area	Avg. landings from benthic trawling, 2009-2018*		Weighted mean bottom depth of area subjected to benthic trawling†	Fraction by mass of total est. global sediment C remineralization‡	Cumulative emissions after years of continuous trawling													
	Mt biomass	Rank by EEZ <sup>§</sup>			m	Rank by EEZ <sup>§</sup>	—	Rank by EEZ <sup>§</sup>	1	5	10	25	30	50	75	100	200	Rank by EEZ <sup>§</sup>
							Pg CO <sub>2</sub>	Rank by EEZ <sup>§</sup>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Rank by EEZ <sup>§</sup>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Rank by EEZ <sup>§</sup>
Malaysia	0.65	10	10	30	0.0062	18												
Original estimate (Sala et al.)							0.0091	18	0.0301	0.0489	0.103	0.121	18	0.194	0.284	0.374	0.736	18
Adjusted estimate (this analysis)							0.0085	9	0.0296	0.0485	0.103	0.121	13	0.193	0.284	0.374	0.736	15
Percent deviation							-6.4	28	-1.6	-0.85	-0.39	-0.33	28	-0.21	-0.14	-0.11	-0.055	28
Mexico	0.64	13	146	16	$3.8 \times 10^{-5}$	30												
Original estimate (Sala et al.)							$5.59 \times 10^{-5}$	30	0.0002	0.0003	0.0006	0.0007	30	0.0012	0.0018	0.0023	0.0045	30
Adjusted estimate (this analysis)							$2.23 \times 10^{-5}$	30	$8.67 \times 10^{-5}$	0.0002	0.0004	0.0005	30	0.0008	0.0013	0.0019	0.004	30
Percent deviation							-60	13	-53	-49	-41	-38	8	-30	-24	-19	-11	13
Morocco	1.9	5	473	2	0.0053	20												
Original estimate (Sala et al.)							0.0079	20	0.0259	0.0422	0.089	0.105	20	0.167	0.245	0.323	0.635	20
Adjusted estimate (this analysis)							0.0011	22	0.0084	0.0174	0.0468	0.0575	20	0.103	0.163	0.226	0.489	21
Percent deviation							-86	5	-68	-59	-47	-45	6	-39	-34	-30	-23	6
Myanmar <sup>¶</sup>	0.64	12																
Original estimate (Sala et al.)																		
Adjusted estimate (this analysis)																		
Percent deviation																		
Namibia	0.37	19	403	3	0.0088	12												
Original estimate (Sala et al.)							0.013	12	0.0428	0.0696	0.147	0.173	12	0.276	0.404	0.533	1.05	12
Adjusted estimate (this analysis)							0.0007	25	0.0075	0.0175	0.0606	0.078	18	0.155	0.262	0.374	0.851	13
Percent deviation							-95	3	-83	-75	-59	-55	4	-44	-35	-30	-19	7
Netherlands	0.092	32	10	31	0.02	6												
Original estimate (Sala et al.)							0.0301	6	0.0993	0.161	0.341	0.401	6	0.64	0.938	1.24	2.43	6
Adjusted estimate (this analysis)							0.0301	4	0.0993	0.161	0.341	0.401	5	0.64	0.938	1.24	2.43	5
Percent deviation							-0.017	31	-0.002	-0.001	$-6.1 \times 10^{-4}$	$-5.2 \times 10^{-4}$	31	$-3.2 \times 10^{-4}$	$-2.2 \times 10^{-4}$	$-1.7 \times 10^{-4}$	$-8.5 \times 10^{-4}$	31
New Zealand	0.34	20	348	5	0.0054	19												
Original estimate (Sala et al.)							0.008	19	0.0263	0.0428	0.0903	0.106	19	0.169	0.249	0.328	0.644	19
Adjusted estimate (this analysis)							0.003	15	0.0133	0.0247	0.0596	0.0717	19	0.122	0.187	0.255	0.536	20
Percent deviation							-62	12	-50	-42	-34	-32	9	-28	-25	-22	-17	9



Supplementary information for Collins et al. preprint; to be submitted for publication

Bottom depth interval or exclusive economic zone (EEZ) area	Avg. landings from benthic trawling, 2009-2018*		Weighted mean bottom depth of area subjected to benthic trawling <sup>†</sup>	Fraction by mass of total est. global sediment C remineralization <sup>‡</sup>	Cumulative emissions after years of continuous trawling												
	Mt biomass	Rank by EEZ <sup>§</sup>			m	Rank by EEZ <sup>§</sup>	Rank by EEZ <sup>§</sup>	1	5	10	25	30	50	75	100	200	Rank by EEZ <sup>§</sup>
						Pg CO <sub>2</sub>	Rank by EEZ <sup>§</sup>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Rank by EEZ <sup>§</sup>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Pg CO <sub>2</sub>	Rank by EEZ <sup>§</sup>
Turkey	0.16	26	108	19	0.0085	13											
Original estimate (Sala et al.)						0.0125	13	0.0415	0.0674	0.142	0.167	13	0.267	0.392	0.516	1.01	13
Adjusted estimate (this analysis)						0.0079	11	0.0348	0.0597	0.132	0.156	10	0.253	0.375	0.498	0.993	10
Percent deviation						-37	19	-16	-11	-7.5	-6.8	24	-5.2	-4.1	-3.5	-2.1	22
United Kingdom	0.67	9	138	18	0.032	3											
Original estimate (Sala et al.)						0.0477	3	0.158	0.256	0.541	0.636	3	1.01	1.49	1.96	3.86	3
Adjusted estimate (this analysis)						0.0363	2	0.134	0.226	0.489	0.577	2	0.928	1.37	1.81	3.58	3
Percent deviation						-24	24	-15	-12	-9.6	-9.3	21	-8.6	-8.1	-7.9	-7.2	14
United States	1.2	7	170	14	0.021	5											
Original estimate (Sala et al.)						0.0307	5	0.101	0.165	0.348	0.409	5	0.653	0.959	1.26	2.48	5
Adjusted estimate (this analysis)						0.0182	5	0.0739	0.133	0.306	0.365	6	0.602	0.9	1.2	2.4	6
Percent deviation						-41	17	-27	-20	-12	-11	19	-7.9	-6.1	-5.1	-3.5	20
Vietnam	2.5	2	29	26	0.0006	28											
Original estimate (Sala et al.)						0.0009	28	0.003	0.0049	0.0104	0.0123	28	0.0196	0.0287	0.0378	0.0744	28
Adjusted estimate (this analysis)						0.0006	27	0.0021	0.0039	0.0092	0.0111	27	0.0183	0.0275	0.0366	0.0731	28
Percent deviation						-38	18	-30	-22	-11	-9.8	20	-6.2	-4.3	-3.3	-1.7	24

Supplementary Table 1 presents cumulative emissions estimates from both the original Sala et al.<sup>1</sup> model and this study, which applies the methods described in the text to account for ventilation time to the atmosphere. Data and emissions estimates are presented for the 30 countries with the greatest average landings from benthic trawling from 2009-2018 (according to ref. <sup>20</sup>; see note below) and for the top 10 countries ranked by percentage of total EEZ catch from bottom trawling (according to ref. <sup>21</sup>).

\*Catch data were retrieved from the Sea Around Us database<sup>20</sup> using the code to which a link is provided in the Code Availability statement.

<sup>†</sup>We weighted the bottom depth of the area within each EEZ that was subjected to benthic trawling according to the rates of remineralization estimated by Sala et al.<sup>1</sup> The mean bottom depth for each grid square in the Sala et al. CO<sub>2</sub> efflux dataset was retrieved from the database described in ref. <sup>15</sup>; this depth was then weighted by the mass of CO<sub>2</sub> that Sala et al.<sup>1</sup> estimated was produced in that grid square in the first year of benthic trawling.

<sup>‡</sup>For the first year after start of benthic trawling activity, from the model of sediment remineralization posited by Sala et al.<sup>1</sup>

<sup>§</sup>Geographic subsetting was performed using the exclusive economic zone boundaries contained in the database described in ref. <sup>19</sup>. A link to the appropriate code is provided in the Code Availability statement.

<sup>||</sup>The mean bottom depth for each grid square in the Sala et al.<sup>1</sup> sediment CO<sub>2</sub> efflux dataset was retrieved from the database described in ref. <sup>15</sup>.

<sup>¶</sup>Sala et al.<sup>1</sup> estimated no sediment CO<sub>2</sub> emissions for grid squares within this EEZ boundary.

## Supplementary References

1. Sala, E. *et al.* Protecting the global ocean for biodiversity, food and climate. *Nature* **592**, 397–402 (2021).
2. Atwood, T. B., Witt, A., Mayorga, J., Hammill, E. & Sala, E. Global patterns in marine sediment carbon stocks. *Front. Mar. Sci.* **7**, 165 (2020).
3. Hiddink, J. G. *et al.* Global analysis of depletion and recovery of seabed biota after bottom trawling disturbance. *Proc. Natl. Acad. Sci.* **114**, 8301–8306 (2017).
4. Eigaard, O. R. *et al.* Estimating seabed pressure from demersal trawls, seines, and dredges based on gear design and dimensions. *ICES J. Mar. Sci.* **73**, i27–i43 (2016).
5. R Core Team. R: a language and environment for statistical computing, version 4.2.1. (2022).
6. Pebesma, E. J. & Bivand, R. S. R package ‘sp’: Classes and methods for spatial data in R. (2005).
7. Bivand, R. S., Pebesma, E. J. & Gomez-Rubio, V. *Applied spatial data analysis with R.* (Springer, 2013).
8. Bivand, R., Keitt, T. & Rowlingson, B. R package ‘rgdal’: Bindings for the “Geospatial” Data Abstraction Library, ver. 1.5-32. (2022).
9. Pebesma, E. J. Simple features for R: standardized support for spatial vector data. *R J.* **10**, 439–446.
10. Hijmans, R. J. & van Etten, J. R package ‘raster’: Geographic analysis and modeling with raster data, ver. 3.5-29. (2022).
11. Dowle, M. & Srinivasan, A. R package ‘data.table,’ ver. 1.14.2. (2021).
12. Bengtsson, H. R package ‘R.matlab,’ ver. 3.7.0. (2022).
13. Bivand, R. & Lewin-Koh, N. R package ‘maptools’: Tools for handling spatial objects, ver. 1.1-4. (2022).
14. Siegel, D. A., DeVries, T., Doney, S. C. & Bell, T. Assessing the sequestration time scales of some ocean-based carbon dioxide reduction strategies. *Environ. Res. Lett.* **16**, 104003 (2021).
15. Becker, J. J. *et al.* Global bathymetry and elevation data at 30 arc seconds resolution: SRTM30\_PLUS. *Mar. Geod.* **32**, 355–371 (2009).
16. Holte, J. & Talley, L. A New algorithm for finding mixed layer depths with applications to Argo data and subantarctic mode water formation. *J. Atmospheric Ocean. Technol.* **26**, 1920–1939 (2009).
17. Holte, J., Talley, L. D., Gilson, J. & Roemmich, D. An Argo mixed layer climatology and database. *Geophys. Res. Lett.* **44**, 5618–5626 (2017).
18. Oberle, F. K. J., Storlazzi, C. D. & Hanebuth, T. J. J. What a drag: Quantifying the global impact of chronic bottom trawling on continental shelf sediment. *J. Mar. Syst.* **159**, 109–119 (2016).

19. Flanders Marine Institute. Maritime Boundaries Geodatabase: Maritime Boundaries and Exclusive Economic Zones (200NM), version 11. (2019).
20. Pauly, D., Zeller, D. & Deng Palomares, M. L. Sea Around Us Concepts, Design and Data. (2020).
21. Steadman, D. *et al.* *New perspectives on an old fishing practice: Scale, context and impacts of bottom trawling.* 44 (2021).