

Title: CDR modeled in MAGICC to return to preindustrial temperatures by 2100.

Author: Shannon A. Fiume, Autofracture, shannon@autofracture.com,

<https://twitter.com/safiume>, <https://github.com/safiume>

DOI: <https://doi.org/10.31223/X5K37C>

The enclosed paper, ‘CDR modeled in MAGICC to return to preindustrial temperatures by 2100’ is a non-peer reviewed preprint and submitted to EarthArXiv. It has not been previously submitted to any journal. All data is available at <https://github.com/hsbay/CDRMEx>, with the Creative Commons Attribution 4.0 International open source license. Referenced software is licensed by their respective licenses and listed as such.

CDR modeled in MAGICC to return to preindustrial temperatures by 2100.

S. A. Fiume

shannon@autofracture.com


Abstract

Experimental test of the theory proposed in an Alternative Method to Determine a Carbon Dioxide Removal Target. The multi-step experiment explores halting anthropogenic emissions from greenhouse gases and removing all historic cumulative anthropogenic carbon dioxide emissions in less than a hundred years to match the preindustrial temperature. The multi-step experiment was conducted on MAGICC 6.8, a reduced complexity model or model emulator, managed by pymagicc. The experiment compares the new experimental pathway 300x2050, marker SSP1 2.6, and SSP1 1.9 within the context of development under a green growth paradigm and explores large-scale linear carbon dioxide removal over the 80-year time frame. The multi-step experiment calibrated the experimental pathway to include recent historic emissions through 2020, and was subsequently tuned to model the recent average global temperatures and CO₂ concentration through 2020. Contrary to the prior proposed theory all anthropogenic emissions (from both fossil fuels and land-use change) needed to be removed to realize final temperatures close to 0.067°C relative to 1720-1800 mean. The experimental pathway evolved temperature to 0.13°C to 1850-1900 mean and realized a final CO₂ concentration of 278.82 ppm by 2550. The experimental pathway achieved nearly 0.069°C relative to 1720-1800 mean and 0.14°C over 1850-1900 of warming at 2100 after phasing out all greenhouse gases, excluding ammonia and removing all cumulative anthropogenic carbon dioxide.

Introduction

This letter explores climate modeling in MAGICC^{1,2,3} and outlines how to generate temperatures that roughly match preindustrial at 2100 through scaled Carbon Dioxide Removal. Although the large-scaled CDR experimental pathway of 300 x 2050 is highly implausible with the current technology, the state of the clean energy industry, and know-how, the modeling provides a likely lowest emissions bound and scaled CDR pathway generating temperatures far lower than the marker pathway of Shared Socioeconomic Pathway (SSP) 1 1.9^{5,6,7,8} within the SSP 1 storyline of ‘development under a green-growth paradigm’⁶. In order to constrain the total amount of carbon needed to be removed from the climate-carbon cycle, the pathway quickly reaches net-zero in the middle 2020s, followed by scaled fossil fuel free CDR.

Figure 1: Carbon Budget showing CDR from 1850-2100 for IMAGE SSP 1 2.6, IMAGE SSP 1 1.9, and 300 x 2050.



Results Summary

The novel high carbon dioxide removal experiment pathway named 300 x 2050 details how high rates of Carbon Dioxide Removal evolve Earth’s climate in the reduced complexity model¹: MAGICC 6.8^{2,3} over the time span of eighty years and continuing on for an additional 450 years. The results are in the jupyter python pymagicc (2.0)⁴ ONC CDRMEx workbook running MAGICC 6.8. Carbon Dioxide Removal (CDR) was accomplished by negative emissions of carbon dioxide and phase-outs of methane and carbon monoxide as well as all other industrial fossil fuel-based gases excluding ammonia ending by 2100. CO₂ removals in this text are specified as a negative quantity of emissions within the experiment parameters and not a technology or set of technologies to be agnostic to the type of implementation. The front-loaded CDR and phase-down of the other fossil fuel and agricultural greenhouse gases realize about 300 ppm. Even though the removals are agnostic to both technology and implementation, the dual conditions of the massive amounts of CDR and achieving 0°C by 2100 require a total and full phase-out of fossil fuels over the century.

The 300 x 2050 pathway removed roughly 237.85 GtC or 871.48 GtCO₂ for fossil fuel emissions by 2050, and 71.44 GtC or 261.76 GtCO₂ of Land-use change emissions by 2050. The pathway then continues on to remove all anthropogenic fossil fuel emissions totaling 481.90 GtC or 1765.7 GtCO₂ by 2100. It also removes all anthropogenic land-use change emissions totaling 162.16 GtC or 594.16 GtCO₂ by 2100.

As shown in Figure 2, the descent from peak emissions in the early 2020s is swiftly followed by net-zero in the mid-2020s (shown when the 300 x 2050 pathway crosses zero by late 2023), then steeper cuts of negative emissions are increased until 2030, when the yearly rate of negative emissions of Fossil Fuel emission added to AFOLU reach a total of 47 GtCO₂ yr-1 by





Figure 3 (a-d): Surface Temperature, Radiative Forcing, CO_{2-eq}, and CO₂ for the experimental 300 x 2050, SSP1 2.6 and SSP1 1.9 over the time 1850-2125, also shown in gray are the observational proxies of HadCRUT5 Analysis¹⁰ and the Keeling Curve, NOAA GML.¹¹

Table 1: CO₂ concentration, Radiative Forcing, and Temperature.

2030 continuing on until 2050, then lowered to total to 25.03 GtCO₂ yr-1 until 2100.

These types of removals, when run in the calibrated model, resulted in the following declines in global temperature, radiative forcing, CO_{2-eq} concentration, and CO₂ concentration. Listed for comparison are the marker SSPs for the SSP 1 storyline of high development of green-growth: IMAGE SSP1 2.6^{5,6} and IMAGE SSP 1 1.9^{5,6,7,8}; SSP data from the IIASA Explorer⁸.

Figure 2. Pathways SSP1 2.6^{5,6}, SSP1 1.9^{5,6,7,8}, and 300 x 2050 show emissions and emissions reductions starting in 2010 through 2125. Historical emissions through 2020 are included in the pathway 300 x 2050.



scenario	unit	variable	time 2020-01-01	2100-01-01	2550-01-01
SSP1-2.6	ppm	Atmospheric Concentrations CO2	419.1408	429.7285	374.5315
	W / m^2	Radiative Forcing	2.7057	2.8538	1.4417
	K	Surface Temperature	1.1880	1.8101	1.2480
SSP1-1.9	ppm	Atmospheric Concentrations CO2	415.8408	371.8397	344.8451
	W / m^2	Radiative Forcing	2.6656	2.0874	1.0784
	K	Surface Temperature	1.1856	1.3773	0.9568
300x2050	ppm	Atmospheric Concentrations CO2	421.3218	237.3759	278.8180
	W / m^2	Radiative Forcing	2.7738	-0.4094	0.0068
	K	Surface Temperature	1.2158	0.0692	0.0674

Methods

Experiment settings

The experiment was set up by calibrating and tuning MAGICC to better match present near-term temperature, concentration, and emissions data. The recent current anthropogenic carbon dioxide emissions data through 2020 from the Global Carbon Budget (2021)⁹ was used to calibrate and extend the carbon dioxide emissions data from 2009 to 2020 (estimated) and coupled to the removal pathway allowing for a closer near-term fit for peak CO₂ concentration and temperature. To emulate CMIP and present-day temperature evolution in MAGICC, the HadCRUT5 (2020)¹⁰ temperature data analysis was used to line-fit the average of the last five years. The carbon dioxide concentration from the Keeling Curve¹¹ was used to roughly line fit 2015 and 2020 and establish error, which under the tuned model yielding +8 ppm above the CO₂ concentration for 2020. These tunings were applied as MAGICC 6.8 was last harmonized with 2010 data.

Tuned Settings

The following settings were changed: climate sensitivity, ratio to land-ocean, heat exchange and amplification, north to south heat exchange, CO₂ fertilization and year start, land sink pools and fluxes, and soil feedback factors to allow the model to better line fit the HadCRUT5¹⁰ temperature analysis, GCB 2021⁹ emissions data, and Keeling Curve¹¹. Given the tunings to the land sink pools and fluxes, to simplify curve fitting and minimize overfitting only the soil feedback was needed and the other land sink feedbacks were disabled.

N₂O data caused a noticeable spike after after severely decreasing emissions; MAGICC 6.8, which was finalized in 2012, does seem to artificially hold this value high for a few years after a large emissions reduction resulting in the artifact around the year 2079 for the 300 x 2050 pathway or in 2100 for the standard scenarios visible in the N₂O, CO_{2-eq} and Surface Temperature graphs. In order to smooth the declining curves for 300 x 2050, all GHGs were declined to phase out by 2079, except ammonia and the negative carbon dioxide (both fossil fuel and land-use change) emissions. It is left to further study under models not subject to this same N₂O artifact if GHG phase-outs can happen closer to 2050 and would that lower the total number of removal necessary to reach 0°C.

See the Supplemental Data section of the Supplementary Information for a listing of all tuned MAGICC configuration settings.

Preindustrial Baseline, 1720-1800

A baseline of 1720-1800 was chosen per *Estimating Changes in Global Temperature since the Preindustrial Period, 2017*²¹ as mentioned in *Trajectories of the Earth System in the Anthropocene, 2018*²⁷ which resulted in temperatures 0.073°C warmer than the 1850-1900 baseline.

unit	variable	mean of 1720-1800	mean of 1850-1900	mean of 1861-1900	mean of 1880-1900	mean of 1951-1980	mean of 1961-1990
ppm	Atmospheric Concentrations CO ₂	2.7862e+02	289.8765	291.0738	293.9143	322.3647	333.5193
K	Surface Temperature	-3.0411e-18	-0.0730	-0.0805	-0.1440	0.1753	0.2177

Table 2: CO₂ concentration and temperature means for common various year spans.

The earlier baseline choice also avoids hysteresis seen if the model time series start is after 1765. A post-1765 start never reached a temperature of about 0°C by 2500. Additional details are in the Supplemental Data section.

Discussion

Model Temperature Calibration


When calibrating for the Northern and Southern regions for the HadCRUT5 analysis, the model evolved the northern hemisphere and southern hemisphere Ocean, and hemisphere respective land binning. The model was calibrated to achieve a delta of +0.0009°C for the world and a little over ±0.18°C for the northern hemisphere and the southern hemisphere. See Supplement Data Table 3 for temperature data in tabular form. The calibration to recent temperature forcing and recent emissions as of 2020, to match the final temperature and changed the amplitude of the temperature graphs over an untuned configuration. The calibration exercise additionally line fit global CO₂ concentration for 2020 in an attempt to reach 412.44 ppm and match CO₂ concentration per Dlugokencky, et. al (2021)¹¹. The experimental pathway 300 x 2050 for 2000-2020 reflects a closer match to the temperature and CO₂ evolution for near future dates, i.e., the 300 x 2050 pathway, 2021 was predicted to reach 1.25°C and a CO₂ concentration of 423.97±8 ppm. This is visually noticeable (see figure 2) by the divergence of 300 x 2050 from the harmonized

SSP1 2.6 and SSP1 1.9 starting about 2015. For an accounting of real-life emissions and temperature modeling, calibrating to the latest emissions, recent historical temperature, and CO₂ concentration data is highly recommended.

Calibration and Regionality

By combining the HadCRUT5 calibration for the world, the regions split between hemispheres and the CRUTEM5 land dataset¹² MAGICC predicts a temperature evolution lower than the mean of 2015-2020 in CRUTEM5 by +0.15°C. The northern hemisphere, with the most developed land, experiences the majority of the warming well beyond the other regions, then southern hemisphere land is followed by the northern hemisphere ocean, and finally the southern hemisphere ocean. This ordering is common to all graphed pathways. The regional splitting of these temperature projections has only been compared but not tuned to the CRUTEM5 2021 dataset and not to other ESM data. The northern hemisphere mean temperature for 2015-2020 relative to 1850-1900 for 300 x 2050 pathway in MAGICC was 1.64°C and 0.14°C less than the CRUTEM5 2021 dataset mean (1.78°C) over the same duration. The southern hemisphere mean temperature for 2015-2020 relative to 1850-1900 for 300 x 2050 pathway in MAGICC was 1.21°C and 0.15°C less than the CRUTEM5 2021 dataset mean for 2015-2020 relative to 1857-1900 (1.361°C).

Figure 4 (a-c): Surface Temperature for SSP 1 2.6, SSP 1 1.9, and 300 x 2050 listing the northern, southern, land, and ocean regions, including the world region.



Given the effect of heat exchange and temperature splitting in each region, and even if the temperature evolution for the 300 x 2050 pathway is estimated with high uncertainty and given how long it takes to recover with a high rate of removal, it follows that these next two decades are critical to lowering global temperatures.

Equilibrium Climate Sensitivity, Transient Climate Response, and Transient Response to Cumulative CO₂ Emissions

```
MAGICC ECS setting: 3.257K, MAGICC ΔQ2xC02 setting: 3.71K
Calculating ECS from abrupt-2xC02.
Calculating TCR & TCRE from 1pctC02.
TCR is 2.0882, ECS is 3.2376 kelvin and TCRE is 2.420048 K / 1000 GtC
Calculating ECS from abrupt-0p5xC02.
Calculating TCR & TCRE from 1pctC02-cdr.
TCR is 2.0882, ECS is 3.2555 kelvin and TCRE is 2.420048 K / 1000 GtC
```

Figure 5: MAGICC output listing ECS, TCR, TCRE results, and CMIP6 experiments.

The Equilibrium Climate Sensitivity (ECS) is defined by a doubling or halving of CO₂ concentration over the time it takes for the global temperature to equilibrate. The MAGICC setting for core climate sensitivity (ΔT_{2x}) was raised from the MAGICC 6.801^{2,3} initial defaults of 3K to 3.257K and evolved ECS to 3.24+0.02K over 2500 years, in relation to the heat exchange tunings to allow for a closer proximal distribution to the hemisphere land regions when considering the CRUTEM5 2021 data and HadCRUT5 2020 data analysis. The effect of tuning to the present-day temperature mean also lowered both the Transient Response to Cumulative CO₂ Emissions (TCRE) to 2.4K and the Transient Climate Response (TCR) to 2.1K. Despite present-

day calibration temperature mean trending initially hotter than lower scenarios, SSP 1 2.6 and SSP 1 1.9, ECS didn't trend significantly hotter. This effect is likely given the shorter duration of increased emissions from the least emitting pathway of 300 x 2050. See Section 3.1 of *Introduction of variable climate sensitivities in Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 – Part 1: Model description and calibration, 2011*² for a more in-depth discussion as it pertains to the model emulator MAGICC 6.8x.

MAGICC 6.8 Negative Emissions Calibration


The negative emissions calibration of MAGICC 6.801 was performed by adapting the supplied pymagicc code to utilize the CMIP6 abrupt-0p5xCO2¹² and 1pctCO2-cdr test, then generate ECS, TCR, and TCRE by evaluating the abrupt doubling of emissions and the 1% continual increase in CO₂ concentrations compared to the removal tests abrupt-0p5xCO2 and 1pctCO2-cdr tests. The calibration is valid if the absolute value of the paired tests and subsequent TCR, TCRE, and ECS are found to be equal. The absolute value of the pairs of the TCR and TCRE were equal, and ECS resulted in a slight difference of about 0.0194°C and is still an acceptable demonstration of negative emissions given the tests equilibrate over 2500 years.

When removing all anthropogenic emissions from both fossil fuel emissions and land-use change with the 300 x 2050 pathway, the model was able to achieve 0.07°C over the mean temperature for 1720-1800 or 0.13°C over the mean temperature for 1850-1900. The inclusion of cumulative land-use change for removal was unexpected and unanticipated in the author's previous theory paper. See Supplemental Figures I-K for the tuned MAGICC 6.8 model output running abrupt-0p5xCO2 and comparison to a few abrupt-0p5xCO2 CMIP6 data ^{13,14,15,16,17,18,19,20,21}.

Ocean outgassing is inferred as all fossil fuel emissions had to be removed instead of only the emissions that currently reside in the atmosphere. A removal to roughly realize 302.83 ppm of CO₂ by 2050 (237.85 GtC modeled, compared to (415 ppm - 302.83 ppm) * 2.124 (GtC/ppm) = 238.27 GtC), needed additional removal of 244 GtC to counterbalance the effect of the ocean reestablishing pCO₂ equilibrium (resulted in outgassing), all of which was completed by 2100. The continued removal allowed both temperature and CO₂ to slowly recover and finally converge to match pre-industrial. CO₂ concentration does drop well below 237 ppm; however, CO_{2eq} only drops below 252.99 ppm, all the while, the global temperature mean never goes lower than 0.04°C. Ocean upwelling is predicted to return to baseline by 2100 in the 300 x 2050 pathway, see supplemental figure h. The ocean heat exchange, drop in concentration from emissions forcings, N₂O artifact and GHGs phaseouts introduce enough varying factors to complicate modeling on a reduced complexity model software to merit a more comprehensive understanding of the present-day emissions and temperatures and future evolution of the experiments to be modeled in a full ESM ensemble.

		time	1750-01-01 00:00:00	1817-01-01 00:00:00	1850-01-01 00:00:00	2023-01-01 00:00:00	2050-01-01 00:00:00	2082-01-01 00:00:00	2100-01-01 00:00:00	2550-01-01 00:00:00	2081-2100 mean	2081-2100 1850mn
unit	variable											
ppm	Atmospheric Concentrations CO2		277.0147	284.1080	284.8000	427.3552	302.8265	262.7444	2.3738e+02	2.7882e+02	250.2084	NaN
W / m^2	Radiative Forcing		0.0052	-0.8239	0.1591	2.8734	1.0329	0.3283	-4.0938e-01	6.7958e-03	-0.0567	NaN
K	Surface Temperature		-0.0022	-0.7863	0.0022	1.2993	0.7769	0.4223	6.9181e-02	6.7370e-02	0.2506	0.3235
Gt C / yr	Emissions CO2 MAGICC Fossil and Industrial		0.0000	0.0141	0.0544	3.4667	-4.9806	-4.9806	0.0000e+00	0.0000e+00	-4.7185	NaN
	Emissions CO2 MAGICC AFOLU		0.0000	0.2463	0.4026	0.0000	-1.8515	-1.8515	0.0000e+00	0.0000e+00	-1.7541	NaN
CO2eqppm	Atmospheric Concentrations CO2 Equivalent		277.0147	283.5586	284.7459	453.5959	323.0444	290.8009	2.5299e+02	2.7336e+02	270.6647	NaN
Cumulative Gt C	Emissions CO2 MAGICC Fossil and Industrial Cumulative		0.0000	0.3536	1.2754	481.8968	244.0495	84.6703	9.4418e-05	9.4418e-05	40.1070	NaN
	Emissions CO2 MAGICC AFOLU Cumulative		0.0000	6.5263	17.3104	162.1623	90.7213	31.4733	-2.1904e-03	-2.1904e-03	14.9073	NaN

Table 3: 300 x 2050 CDR pathway data at various time points.






Figure 6 (a-b): Full graphs for Surface Temperature of 300 x 2050 and Radiative Forcing to show 300 x 2050 convergence of about 0.05°C and Radiative Forcing 0.

Accounting of Land Use Change Emissions

With the inclusion of the Global Carbon Budget 2020 emissions data, the land-use change emissions forcings diverged from SSP 1.9 prior to 2020, as shown in figure 7, d. The divergence was unexpected as all emissions data ought to be harmonized by the model to 2005. To work around this modeling behavior, extensive tunings to the heat, hemispheres and soil feedback were applied in addition to removing 42.4 GtC of AFOLU (Land-use change carbon) to match the MAGICC evolved emissions data. The soil carbon feedback was enabled and tuned to allow for positive growth in CO_2 concentration to allow for minimal non-linear evolution to concentration and temperature in the near term mid-2020s, 2100, and beyond through 2550. The combined effect of tunings and workaround additionally increased the Northern/Southern temperature spread.

Given the mixed effects on the land sinks in the ESM results in *Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment from CO_2 , 2020*²⁵, but yet slight decreases with the UVIC ESM in *Asymmetry in the climate-carbon cycle response to positive and negative CO_2 emissions, 2021*²⁴ a very slight rise (0.0674°C) above 0°C relative to 1720-1800 was chosen for the post 2100 temperature calibration. This temperature target remained slight to properly model negative emissions.

The author was unable to find guidance on increasing the land sink or a MAGICC 6.8 setting to allow more below ground mineralization of anthropogenic land-use change CO_2 to remove it from the climate-carbon cycle and ocean or atmosphere sinks. The author doesn't know if there should be a portion of the cumulative land-use emissions that should have turned over to durable storage of CO_2 in a mineralized form. It was suspected and thus written as such in the theory paper²⁹ that land-use change emissions shouldn't need to be removed as they are fully in balance as it were, with the cumulative increases to the land sink since preindustrial. In lack of evidence, this seems in error. However, natural conversion to more permanent storage is a topic open for further investigation.




Figure 7 (a-d): Anthropogenic and Land-Use Change Emissions and cumulative emissions data, and CDR decreasing in green.

Negative Emissions Calibration and Negative Emissions Asymmetry

The 300 x 2050 pathway experiment has a total removal of 644.99+42.2 GtC in negative emissions to match the cumulative total of anthropogenic emissions. If more land-use change emissions were more permanently contained within the land sink whereby

the quantity of negative emissions needed was reduced to about 500GtC, then the removal is within the model ensemble variability of AOGCM model and natural sink uptake studies^{24,25,26}. If all anthropogenic negative emissions since preindustrial 644.99+42.2 GtC or totaling 2520.4 GtCO₂ are needed to reach closer to 0°C over preindustrial, then it's likely that a small quantity more than what was emitted since preindustrial be added this experiment's modeled removal of all of anthropogenic. The magnitude over 500GtC to determine the magnitude of CDR necessary to match pre-industrial temperatures to be completed by 2100 is open for in-depth investigation.

Experimental Temperature Evolution over Model Time Bounds

If negative emissions have a nearly symmetrical climate-carbon cycle response²⁴ extending the experiment 300 x 2050 out to the model limits will evolve an eventual convergence to about 0.06°C. The shading indicates the experiment run with a lower ECS of 2.1°C and a higher ECS of 4.4°C and includes the medium uncertainty region splitting. Given the vast quantity of emissions needed to be removed under such a short time period, it radically limits the amount of near-term emissions allowable and likely is only achievable should we limit emissions to stay well within the carbon budget. A continued dependency on fossil fuels is unable to yield phase-outs in emissions or the deep removals necessary to achieve the scale and scope to match the pre-industrial temperature.




Figure 8: 300 x 2050 pathway with various Equilibrium Climate Sensitivities: 2.01K, 3.3K, and 4.4K from 1720 to 2550 relative to the mean of 1720-1800.

Conclusion

This text outlines starting points for future investigation on pathways that closely match pre-industrial over existing literature. Having a team explore the experiment's thesis of 1.6x preindustrial concentration²⁴ driven from forcings from emissions, and removing upwards of 600GtC (or 2198.4GtCO₂), run with current temperature forcings, and run on a comprehensive model ensemble would best provide more accurate reflections of temperature, holding below 1.5°C, and additional data such as the sea-level rise and AMOC, ENSO, jet-stream turnover²⁷ and evolution over time.

The possibility of being able to eventually match the pre-industrial climate should help to inform the debate to radically accelerate front-loaded, near-term phase-out of anthropogenic emissions sources and scale zero-carbon intensity carbon removals in order to develop a more equal future world.

Data Availability

The experimental setup and data are fully open source, see Supplementary Information for details.

Funding

Anonymous sponsor retired from semiconductor manufacturing, software entrepreneur, Timothee Besset, US economic stimulus.

Competing Interests

None.

Acknowledgments

The pymagicc, MAGICC, and Global Carbon Budget have been instrumental in writing this letter. Shannon thanks the authors of the GCB, HadCRUT5, MAGICC, pymagicc, Scripps, Mauna Loa, NSF, CMIP, and IPCC scientists. Shannon additionally thanks

References

1. Reduced Complexity Model Intercomparison Project Phase 1: introduction and evaluation of global-mean temperature response, Nicholls, Z, R. J. Meinshausen, M., Lewis, J., et al., 2020, DOI: 10.5194/gmd-13-5175-2020
2. Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 – Part 1: Model description and calibration, 2011, Meinshausen, M., Raper, S. C. B., and Wigley, T. M. L., *Atmos. Chem. Phys.*, 11, 1417-1456, DOI: 10.5194/acp-11-1417-2011
3. Emulating atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 – Part 2: Applications, 2011, Meinshausen, M., Wigley, T. M. L., and Raper, S. C. B., *Atmos. Chem. Phys.*, 11, 1457–1471, DOI: 10.5194/acp-11-1457-2011
4. Pymagicc: A Python wrapper for the simple climate model MAGICC, 2018, R. Gieseke, S. N. Willner, M. Mengel, *Journal of Open Source Software*, 3(22), 516, DOI: 10.21105/joss.00516
5. The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview, 2017, Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, et al., *Global Environmental Change*, Volume 42, Pages 153-168, ISSN 0959-3780, DOI: 110.1016/j.gloenvcha.2016.05.009
6. Energy, land-use and greenhouse gas emissions trajectories under a green growth paradigm, SSP1, van Vuuren, D.P., Stehfest, E., Gernaat, D. E.H.J., et al., *Global Environmental Change*, Volume 42, Pages 237-250, ISSN 0959-3780, 2017, DOI: 10.1016/j.gloenvcha.2016.05.008
7. IAMC 1.5°C Scenario Explorer and Data hosted by IIASA, <https://data.ene.iiasa.ac.at/iamc-1.5c-explorer>, <https://data.ene.iiasa.ac.at/iamc-1.5c-explorer/#/license>, Assessment Modeling Consortium & International Institute for Applied Systems Analysis, Huppmann, D., Kriegler, E., Krey, V., et al., (2018) <https://doi.org/10.22022/SR15/08-2018.15429>
8. Scenarios towards limiting global mean temperature increase below 1.5 °C, Rogelj, J., Popp, A., Calvin, K.V. et al., *Nature Clim Change* 8, 325–332, 2018, DOI: 10.1038/s41558-018-0091-3
9. Global Carbon Budget 2021, Friedlingstein, P., Jones, M. W., O’Sullivan, M., et al. (2021), *Earth System Science Data*, DOI: 10.5194/essd-2021-386, GCB Data Sources: <https://doi.org/10.18160/GCP-2021>
10. An updated assessment of near-surface temperature change from 1850: the HadCRUT5 dataset, 2020, *Journal of Geophysical Research (Atmospheres)*, Morice, C.P., Kennedy, J.J., Rayner, N.A., et al., DOI: 10.1029/2019JD032361
11. Global CO₂ concentration for 2020, Dlugokencky, E., Tans, P., NOAA/GML, (2021), https://gml.noaa.gov/webdata/ccgg/trends/CO2/CO2_annmean_gl.txt
12. Land surface air temperature variations across the globe updated to 2019: the CRUTEM5 dataset., 2021, Osborn, T.J., Jones, P.D., Lister, D.H., et al., *Journal of Geophysical Research: Atmospheres*. 126, e2019JD032352, DOI: 10.1029/2019JD032352
13. Nicholls, Z, Lewis, J, Makin, M, et al. Regionally aggregated, stitched and de-drifted CMIP-climate data, processed with netCDF-SCM v2.0.0. *Geosci Data J.* 2020, <https://doi.org/10.5281/zenodo.3903372> and <https://gitlab.com/netcdf-scm/calibration-data>, Using NetCDF CMIP data, <https://gitlab.com/netcdf-scm/calibration-data>, <https://doi.org/10.1002/gdj3.113>, BSD 3-Clause License Copyright (c) 2020
14. CMIP6, Coupled Model Intercomparison Project, CMIP5 terms of use and CMIP6 terms of use, CFMIP, abrupt-0p5xCO₂: <https://view.es-doc.org/?renderMethod=id&project=cmip6&id=8ff7a328-e031-49f1-862c-68be2c5648e8&version=1&client=esdoc-search>
15. MIROC MIROC6 model output prepared for CMIP6 CFMIP abrupt-0p5xCO₂, Version v20190705, Ogura, Tomoo and Watanabe, Masahiro and Hirota, Nagio, 2019, doi.org/10.22033/ESGF/CMIP6.5405, <http://cera-www.dkrz.de/WDCC/meta/CMIP6/CMIP6.CFMIP.MIROC.MIROC6.abrupt-0p5xCO2>, Creative Commons Attribution Share Alike 4.0 International
16. MRI MRI-ESM2.0 model output prepared for CMIP6 CFMIP abrupt-0p5xCO₂. Version v20200107, Yukimoto, S., Koshiro, T., Kawai, H. et al., 2020, doi.org/10.22033/ESGF/CMIP6.6753, <http://cera-www.dkrz.de/WDCC/meta/CMIP6/CMIP6.CFMIP.MRI.MRI-ESM2-0.abrupt-0p5xCO2>, Creative Commons Attribution Share Alike 4.0 International
17. CNRM-CERFACS CNRM-CM6-1 model output prepared for CMIP6 CFMIP abrupt-0p5xCO₂. Version v20190711, Voltaire, Aurore, 2019, doi.org/10.22033/ESGF/CMIP6.3914, <http://cera-www.dkrz.de/WDCC/meta/CMIP6/CMIP6.CFMIP.CNRM-CERFACS.CNRM-CM6-1.abrupt-0p5xCO2>, Creative Commons Attribution Non Commercial Share Alike 4.0 International
18. NASA-GISS GISS-E2.1G model output prepared for CMIP6 CFMIP abrupt-0p5xCO₂. Version v20190524, NASA Goddard Institute For Space Studies (NASA/GISS), 2019, doi.org/10.22033/ESGF/CMIP6.6972, <http://cera-www.dkrz.de/WDCC/meta/CMIP6/CMIP6.CFMIP.NASA-GISS.GISS-E2-1-G.abrupt-0p5xCO2>, Creative Commons Attribution Share Alike 4.0 International
19. IPSL IPSL-CM6A-LR model output prepared for CMIP6 CFMIP abrupt-0p5xCO₂. Version v20180605, Boucher, O., Denvil, S., Levavasseur, G., et al., 2018, doi.org/10.22033/ESGF/CMIP6.5106, <http://cera-www.dkrz.de/WDCC/meta/CMIP6/CMIP6.CFMIP.IPSL.IPSL-CM6A-LR.abrupt-0p5xCO2>, Creative Commons Attribution Non Commercial Share Alike 4.0 Internationals
20. MOHC HadGEM3-GC31-LL model output prepared for CMIP6 CFMIP abrupt-0p5xCO₂. Version v20200829, Webb, M., 2020, doi.org/10.22033/ESGF/CMIP6.5833, <http://cera-www.dkrz.de/WDCC/meta/CMIP6/CMIP6.CFMIP.MOHC.HadGEM3-GC31-LL.abrupt-0p5xCO2>, Creative Commons Attribution Share Alike 4.0 International

21. NCAR CESM2 model output prepared for CMIP6 CFMIP abrupt-0p5xCO₂. Version v20200408, Danabasoglu, G., 2020, doi.org/10.22033/ESGF/CMIP6.7517, <http://cera-www.dkrz.de/WDCC/meta/CMIP6/CMIP6.CFMIP.NCAR.CESM2.abrupt-0p5xCO2>, Creative Commons Attribution Share Alike 4.0 International
22. Estimating Changes in Global Temperature since the Preindustrial Period, 2017, Hawkins, E., Ortega, P., Suckling, E., et al., DOI: 10.1175/BAMS-D-16-0007.1
23. Halocarbon scenarios, ozone depletion potentials, and global warming potentials, 2007, Daniel, J.S., and G.J.M. Velders, A.R. Douglass, et al., Chapter 8 in *Scientific Assessment of Ozone Depletion: 2006*, Global Ozone Research and Monitoring Project—Report No. 50, 572 pp., World Meteorological Organization, Geneva, Switzerland
24. Asymmetry in the climate-carbon cycle response to positive and negative CO₂ emissions, 2021, Zickfeld, K., Azevedo, D., Mathesius, S., et al., *Nat. Clim. Chang.* 11, 613–617, DOI: 10.1038/s41558-021-01061-2
25. Is there warming in the pipeline? A multi-model analysis of the Zero Emissions Commitment from CO₂, 2020, MacDougall, A. H., Frölicher, T. L., Jones, C. D., et al., DOI: 10.5194/bg-17-2987-2020
26. The Carbon Dioxide Removal Model Intercomparison Project (CDR-MIP): Rationale and experimental protocol for CMIP6, 2018, Keller, D. P., Lenton, A., Scott, V., et al., *Geosci. Model Dev.*, DOI: 10.5194/gmd-2017-168
27. Trajectories of the Earth System in the Anthropocene, 2018, Steffen, W., Rockström, J., Richardson, K., et al., *PNAS*, vol. 115 no. 33 8252-8259, DOI: 10.1073/pnas.1810141115
28. Surface air temperature and its variations over the last 150 years, 1999, Jones, P.D., New, M., Parker, D.E., et al., *Reviews of Geophysics* 37, 173-199, DOI: doi.org/10.1029/1999RG900002
29. Alternative Method to Determine a Carbon Dioxide Removal Target, Fiume, S., 2018, ESSOAR Preprint, DOI: 10.1002/essoar.10503117.1