

## **No support for carbon storage of >1000 GtC in northern peatlands**

Comment on the paper by Nichols & Peteet (2019) in *Nature Geoscience* (**12**: 917-921)

[Manuscript Ref. NGS-2019-11-02825]

5 **Zicheng Yu<sup>1,2,\*</sup>, Fortunat Joos<sup>3</sup>, Thomas K. Bauska<sup>4</sup>, Benjamin D. Stocker<sup>5</sup>, Hubertus Fischer<sup>3</sup>, Julie Loisel<sup>6</sup>, Victor Brovkin<sup>7</sup>, Gustaf Hugelius<sup>8</sup>, Christoph Nehrbass-Ahles<sup>9</sup>, Thomas Kleinen<sup>7</sup> and Jochen Schmitt<sup>3</sup>**

<sup>1</sup>Department of Earth and Environmental Sciences, Lehigh University, Bethlehem, PA

10 18015, USA (PeatHunter@gmail.com)

<sup>2</sup>Institute for Peat and Mire Research, School of Geographical Sciences, Northeast Normal University, 130024 Changchun, China

<sup>3</sup>Climate and Environmental Physics, Physics Institute, University of Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland (joos@climate.unibe.ch; hubertus.fischer@climate.unibe.ch; schmitt@climate.unibe.ch)

<sup>4</sup>British Antarctic Survey, High Cross, Madingley Road, Cambridge, CB3 0ET, UK (thausk@bas.ac.uk)

<sup>5</sup>D-USYS, ETH Zürich, LFW C 55.3, Universitätstrasse 2, 8092 Zürich, Switzerland (bestocke@ethz.ch)

20 <sup>6</sup>Department of Geography, Texas A&M University, 3147 TAMU, College Station, TX 77843, USA (julieloisel@tamu.edu)

<sup>7</sup>Max-Planck Institute for Meteorology, Hamburg, Germany (victor.brovkin@mpimet.mpg.de; thomas.kleinen@mpimet.mpg.de)

25 <sup>8</sup>Department of Physical Geography, Stockholm University, 106 91 Stockholm, Sweden (gustaf.hugelius@natgeo.su.se)

<sup>9</sup>Department of Earth Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EQ, UK (cn425@cam.ac.uk)

\*Correspondence author:

30 Emails: PeatHunter@gmail.com; Phones: 484-904-8404

Northern peatlands store large amounts of carbon (C) and have played an important role in the global carbon cycle since the Last Glacial Maximum. Most northern peatlands have established since the end of the deglaciation and accumulated C over the Holocene, leading to a total present-day stock of  $500 \pm 100$  GtC. This is a consolidated estimate, emerging from a diversity of methods<sup>1-5</sup>. Recently, Nichols and Peteet (hereafter N&P)<sup>6</sup> presented an estimate of the northern peat C stock of 1055 GtC—exceeding previous estimates by a factor of two. Here, we argue that this is an overestimate, caused by systematic bias introduced by their inclusion of data that is not representative for the major peatland regions and of records that lack direct measurements of C density. Furthermore, we argue that their estimate cannot be reconciled within the constraints offered by ice-core and marine records of stable C isotopes and estimated contributions from other processes that affected the terrestrial C storage during the Holocene.

45

### Suitability of data and methodology used

As in previous studies<sup>2</sup>, N&P used the *time-history approach* to estimate peatland C stocks and their evolution in time, using the time-varying peatland area and net C accumulation rates. We notice that area-specific net C accumulation rates used by N&P ( $j_c$ ) as shown in their Fig. 2c have a Holocene mean value of 33.4–37.6  $\text{gC m}^{-2} \text{yr}^{-1}$  (median across three methods), and are thus 80-102% higher than reported in previous studies of 18.6  $\text{gC m}^{-2} \text{yr}^{-1}$  (ref. 2) and 46-64% higher than values of 22.9  $\text{gC m}^{-2} \text{yr}^{-1}$  (ref. 3) (Table 1). Why this difference?

**Table 1. Summary of estimates of Holocene peat C accumulation rates and total C storage in northern peatlands**

<b>Number of sites</b>	<b>Number of cores</b>	<b>Accumulation rates (gC m<sup>-2</sup> yr<sup>-1</sup>)</b>	<b>Carbon storage (GtC)</b>	<b>Reference</b>
Unknown	Unknown	33.4-37.6*	1055	Ref. 6 (N&P)
NA**	NA**	NA**	455	Ref. 1
33	33	18.6	546	Ref. 2
127	151	22.9	436	Ref. 3

\*Calculated from the spreadsheets provided in the N&P's Supplementary Information.

\*\*NA: not applicable, as Ref. 1 used peat volume approach based on total peat area, mean peat depth, and C density.

60

N&P calculated C accumulation rates from sedimentation rates (cm yr<sup>-1</sup>) and C density (gC cm<sup>-3</sup>). We argue here that both of these parameters were overestimated by N&P. An

65

important factor that may introduce a high bias in estimates of regional sedimentation rates is the inclusion of additional data from the Neotoma Paleoecology Database (NPD).

The vast majority of these new data used by N&P originate from locations that are not representative for the areas where the vast majority of northern peatland areas are located (their Fig. 1a and Fig. S1). Their use may be motivated as a complement to the relatively

70

limited set of available peat cores with sufficient information to reconstruct accumulation rates. However, these additional data are almost exclusively originating from lower

latitudes than the data underlying previous estimates<sup>2,3,7</sup>, and thus represent peatlands or wetlands located in different climates. Yet, the total peatland area in these regions is small,

if not negligible, compared to the peatland area north of 50° N (ref. 8). A more reasonable

75

approach would have been to treat additional data from outside the boreal and subarctic

regions separately and scale their accumulation rates with the relatively modest peatland area of these respective regions.

80 N&P claim that they calculated C accumulation rates for each of eight peat regions to account for spatial bias. However, as discussed above, their regional delineation matters. Furthermore, using a single average value for C density ( $\text{gC cm}^{-3}$ ) for all sites that are located within these regions and that lack direct measurements is prone to introducing bias. N&P use a median of peat C density measurements from measurements of C content (%) or organic matter content (%) and dry bulk density ( $\text{g cm}^{-3}$ )<sup>3,7</sup>. However, N&P fail to  
85 account for the variability in C density among regions and among different types of peatlands<sup>3</sup>. For example, there is a more than two-fold difference in C density between *Sphagnum* peat ( $0.037 \text{ gC cm}^{-3}$ ;  $n=3332$ ) and humidified peat ( $0.072 \text{ gC cm}^{-3}$ ;  $n= 418$ ) and between western European islands/continental Europe ( $0.028 \text{ gC cm}^{-3}$ ;  $n=449$ ) and western Canada ( $0.076 \text{ gC cm}^{-3}$ ;  $n= 3441$ )<sup>3</sup>. Also, peat likely experiences different degrees  
90 of decomposition and compaction with ages, resulting in highly variable C density often observed along a single peat profile. The propagation estimates of uncertainties of C density in N&P would not resolve the issue about the representativeness of that single median C density value. Previous large-scale syntheses<sup>2,3</sup> used <sup>14</sup>C-dated individual peat profiles to reconstruct their C accumulation history and excluded sites that did not have  
95 direct C density measurements. Those studies thus avoided these biases.

Furthermore, N&P estimated an average sedimentation rate for each region “by dividing the depth of the deepest sample by the most likely calibrated age of the oldest sample in

the region". We argue that the use of only the deepest and oldest peat sample risks  
100 skewing the estimate towards sites with unusually high sedimentation rates. A more  
appropriate approach would be to use mean rates based on multiple observations. In  
addition, sites with the highest sedimentation rates are likely affected by more mineral  
particle input, which inflate mass accumulation rates but is not representative of C  
accumulation rates. In fact, some records from Neotoma appear to show that portions of  
105 sedimentary sequence may represent lake deposits before peat accumulation (see  
Supplementary Information). All these effects were not considered and corrected for by  
N&P.

Unfortunately, we were not able to quantitatively assess the effects of the inclusion of these  
110 non-representative data and the use of mean C density values for filling data gaps. Required  
information was not accessible through the paper by N&P, nor its Supplementary  
Information.

### **Lack of support from global carbon budget constraints**

115 The exceptionally large peat C storage of >1000 GtC in N&P is also not supported by top-  
down constraints from the global C budget reconstructions. To illustrate the effect such  
large perturbations would have on the global carbon cycle we carried out a sensitivity  
analysis using a simple carbon-cycle box model<sup>9</sup>. The model considers the C exchange  
among the atmosphere, land biosphere, oceans and marine sediments. We used the ranges  
120 (median  $\pm$  1 s.d.) from all three scenarios (literature, combined, grid-box) in N&P as model  
inputs. All scenarios essentially yielded the same solutions. Therefore, we only show the

results from the “combined” approach here. We also ran a separate sensitivity experiment by turning off the simple “carbonate compensation” mechanism using just the median scenario. The results show that an increase in peat C storage of >1000 GtC during the  
125 Holocene would induce a decrease in atmospheric CO<sub>2</sub> to below 220 ppm, an increase in atmospheric δ<sup>13</sup>C to a value more than 0.8‰ higher than the observed, and a steady rise in deep ocean δ<sup>13</sup>C -DIC throughout the Holocene (Fig. 1).

Firstly, our box-model calculations demonstrate that the simplified conversion of peat C  
130 uptake into an atmospheric signal of >600 ppm, as shown in their Fig. 2f of N&P, was erroneous due to the neglect of the compensating effect by the ocean that acts to reduce any atmospheric perturbation by up to 80% on the millennial time scale relevant here<sup>10</sup>. We assume that N&P instead converted their estimated terrestrial C stock increase by a division factor of 2.12 GtC per ppm to arrive at a peat C uptake-related decrease in  
135 atmospheric CO<sub>2</sub> of more than 300 ppm over the Holocene. Translating the same peat C uptake into an atmospheric CO<sub>2</sub> signal with our box-model yielded a decrease of about 60 ppm (Fig. 1b)—perfectly consistent with our presumption that the ~80% reduction by ocean uptake was neglected in N&P.

140 Secondly, the experiments suggest that, at face value, exceptionally large peat C storage is difficult to reconcile with the atmospheric and oceanic C budgets. Previously, the observed changes in atmospheric CO<sub>2</sub> concentration and in δ<sup>13</sup>C from ice cores have been used to partition the contributions from the land biosphere and ocean, providing a global constraint on land C budget during the Holocene. The measured increase in CO<sub>2</sub>

145 concentration from 265 ppm at 11 ka to 278 ppm in 1750 CE and the small change in  $\delta^{13}\text{C}$   
(Fig. 1b, c) were used to reconstruct the preindustrial terrestrial net C uptake over the  
Holocene to be about 250 GtC (ref. 11). This total Holocene land C balance reflects a strong  
uptake in the early Holocene through the growth of boreal forests and early peat buildup—  
which is consistent with the observed early-Holocene increase in atmospheric and oceanic  
150  $\delta^{13}\text{C}$  values<sup>12</sup>—and a C release of 50 GtC during the late Holocene<sup>11</sup>. The small decrease in  
land C storage in the last 5 kyr contrasts with the large estimated increase in peat C storage  
of ~400 GtC during the same time period as suggested by N&P (their Fig. 2e). A  
compensating C source of 400-500 GtC with a biogenic  $\delta^{13}\text{C}$  signature would have to be  
invoked to close the budget. A detailed analysis of this budget concluded that CO<sub>2</sub> emissions  
155 from land-use change by early agriculturalists were not sufficient to close the gap between  
peat C uptake and the atmospheric constraint before about 3 ka (ref. 13). The two-fold  
higher estimates of peat C storage by N&P, compared to the record used<sup>13</sup>, make it even  
harder to reconcile the budget. This conflict is not discussed in N&P.

160 N&P speculate that C release from terrestrial cold steppe permafrost that accumulated  
during the glacial time could have compensated the large peat C uptake and thereby satisfy  
the isotopic  $\delta^{13}\text{C}$  mass balance constraint. However, this release occurred mostly during the  
deglacial warming, not during the Holocene, when most of the present extratropical peat C  
storage grows. Rather than balancing the C budget with terrestrial C sources in the  
165 Holocene, N&P suggest that “most important mechanisms for balancing the peatland sink”  
is a continued C release from the deep ocean by the wind-driven upwelling during the  
Holocene. This mechanism requires an even greater loss of C from the deep ocean than

implied by the peatland C sink alone and is not supported by observation and simulation of marine  $\delta^{13}\text{C}$  and carbonate ion changes. For example, an increase in Southern Ocean upwelling would further increase  $\delta^{13}\text{C}$ -DIC in the deep ocean<sup>14</sup> than the already untenable increase  $\delta^{13}\text{C}$ -DIC from peatland regrowth (Fig. 1d), yet  $\delta^{13}\text{C}$  values remained constant after 7 ka, as observed from a stack of benthic  $\delta^{13}\text{C}$  data from 33 deep-ocean (>3000 m) cores around the world oceans<sup>12</sup> (Fig. 1d). Furthermore, the  $\text{CO}_2$  release from the deep ocean would lead to an increase in the carbonate ion concentration and enhanced preservation of carbonates in the deep ocean, but deep ocean cores show the opposite—a reduction in the carbonate ion and an increase in carbonate dissolution during the Holocene<sup>15</sup>. If any oceanic C source contributed to the Holocene  $\text{CO}_2$  rise, it would likely be due to carbonate compensation after deglaciation<sup>12</sup> and surface ocean processes, including shallow water carbonate accumulation such as coral reefs on newly exposed continental shelves<sup>16</sup>. Both processes would cause no significant change in the  $\delta^{13}\text{C}$  value of released  $\text{CO}_2$ , and therefore would not mask the imprint of peat C uptake in the atmospheric  $\delta^{13}\text{C}$  record, but, as demonstrated in our box model experiments, are insufficient to compensate for such a large peat sink.

N&P also relate peat initiation and growth to the atmospheric methane record as archived in polar ice cores. In particular, they relate the strong and rapid increase in  $\text{CH}_4$  at the onset of the Holocene<sup>17</sup> with their peak in peat initiation (Fig. 2b in N&P). While this coincidence is remarkable, we consider that it is problematic to relate peat initiation to the large magnitude and abrupt increase in  $\text{CH}_4$  emissions at that time, as the latter should be related to total area of existing  $\text{CH}_4$  emitting wetlands/peatlands and climate-dependent rates of



CH<sub>4</sub> emissions, not the rates of initiation and peat area increase<sup>18</sup>. It is important to note that the strong increase of CH<sub>4</sub> at that time occurred likely much too quickly to allow for substantial peat area expansion. Therefore the abrupt CH<sub>4</sub> increase is more likely caused by increases in plant productivity, availability of labile C, and suitable CH<sub>4</sub> producing environments in a warm and wet climate<sup>19</sup> at the onset of the Holocene.

In summary, N&P have made an extraordinary claim of doubled C storage in northern peatlands, compared to the estimates available in the literature (500 ±100 GtC). But “Extraordinary claims require extraordinary evidence” (per Carl Sagan), and we conclude that the evidence presented by N&P is not sufficient to support their extraordinary claim.

## References

1. Gorham, E. Northern Peatlands: Role in the carbon cycle and probable responses to climatic warming. *Ecol. Appl.* **1**, 182–195 (1991).
2. Yu, Z., Loisel, J., Brosseau, D. P., Beilman, D. W. & Hunt, S. J. Global peatland dynamics since the Last Glacial Maximum. *Geophys. Res. Lett.* **37**, L13402 (2010).
3. Loisel, J. et al. A database and synthesis of northern peatland soil properties and Holocene carbon and nitrogen accumulation. *Holocene* **24**, 1028–1042 (2014).
4. Jackson, R.B. et al. The ecology of soil carbon: Pools, vulnerabilities, and biotic and abiotic controls. *Annu. Rev. Ecol. Evol. Syst.* **48**, 419–45 (2017).
5. Lindgren, A., Hugelius, G. & Kuhry, P. Extensive loss of past permafrost carbon but a net accumulation into present day soils. *Nature* **560**, 219–222 (2019).
6. Nichols, J.E. & Peteet, D.M.. Rapid expansion of northern peatlands and doubled estimate of carbon storage. *Nature Geoscience* **12**, 917–921 (2019).
7. Treat, C.C. et al. Effects of permafrost aggradation on peat properties as determined from a pan-Arctic synthesis of plant macrofossils. *J. Geophys. Res. Biogeosci.* **121**, 78–94 (2016).
8. Xu, J., Morris, P. J., Liu, J. & Holden, J. PEATMAP: Refining estimates of global peatland distribution based on a meta-analysis. *Catena* **160**, 134–140 (2017).

- 230 9. Bauska, T.K. et al. Carbon isotopes characterize rapid changes in atmospheric carbon dioxide during the last deglaciation. *Proc. Natl. Acad. Sci. USA* **113**, 3465-3470 (2016).
10. Joos, D., et al. Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis. *Atmos. Chem. Phys.* **13**, 2793–2825 (2013).
- 235 11. Elsig, J. et al. Stable isotope constraints on Holocene carbon cycle changes from an Antarctic ice core. *Nature* **461**, 507-510 (2009).
12. Menviel, L. & Joos, F. Toward explaining the Holocene carbon dioxide and carbon isotope records: Results from transient ocean carbon cycle-climate simulations. *Paleoceanography* **27**, PA1207, doi:10.1029/2011PA002224 (2012).
- 240 13. Stocker, B.D., Yu, Z., Massa, C. & Joos, F. Holocene peatland and ice-core data constraints on the timing and magnitude of CO<sub>2</sub> emissions from past land use. *Proc. Natl. Acad. Sci. USA* **114**, 1492-1497 (2017).
- 245 14. Tschumi, T., Joos, F., Gehlen, M., and Heinze, C. Deep ocean ventilation, carbon isotopes, marine sedimentation and the deglacial CO<sub>2</sub> rise. *Clim. Past* **7**, 771–800 (2011).
15. Yu, J., Anderson, R.F. & Rohling, E.J. Deep ocean carbonate chemistry and glacial-interglacial atmospheric CO<sub>2</sub> changes. *Oceanography* **27**, 16–25 (2014).
- 250 16. Brovkin, V. et al. What was the source of the atmospheric CO<sub>2</sub> increase during the Holocene? *Biogeosciences* **16**, 2543–2555 (2019).
17. Chappellaz, J. et al. Synchronous changes in atmospheric CH<sub>4</sub> and Greenland climate between 40 and 8 kyr BP. *Nature* **366**, 443- 445 (1993).
- 255 18. Yu, Z. et al. Evidence for elevated emissions from high-latitude wetlands contributing to high atmospheric CH<sub>4</sub> concentration in the early Holocene. *Glob. Biogeochem. Cycles* **27**,131–140 (2013).
- 260 19. Turetsky, M.R. et al. Short-term response of methane fluxes and methanogens activity to water table and soil warming manipulations in an Alaskan peatland. *J. Geophys. Res.* **113**, G00A10, doi:10.1029/2007JG000496 (2008).
20. Monnin, E. et al. Atmospheric CO<sub>2</sub> concentrations over the last glacial termination. *Science* **291**, 112–114 (2001).
- 265 21. Monnin, E. et al. Evidence for substantial accumulation rate variability in Antarctica during the Holocene, through synchronization of CO<sub>2</sub> in the Taylor Dome, Dome C and DML ice cores. *Earth Planet. Sci. Let.* **224**, 45–54 (2004).

22. Schmitt, J. et al. Carbon isotope constraints on the deglacial CO<sub>2</sub> rise from ice cores. *Science* **336**, 711-714 (2012).

270 23. Peterson, C.D. & Lisiecki, L.E. Deglacial carbon cycle changes observed in a compilation of 127 benthic  $\delta^{13}\text{C}$  time series (20–6 ka). *Clim. Past* **14**, 1229–1252 (2018).

275 24. Lisiecki, L.E., Raymo, M.E. & Curry, W.B. Atlantic overturning responses to Late Pleistocene climate forcings. *Nature* **456**, 85-88 (2008).


280

### **Author contributions**

Z.Y. and F.J. designed the research, T.K.B. carried out the box-model simulation and created the figure, and all authors were involved in writing and revising the manuscript.

285 **Competing interests**

The authors declare no competing interests.



**Fig. 1. Unrealistic consequences of large peat carbon storage.** **A.** Peat C storage change (line) over time with uncertainties (orange band) as reported in N&P<sup>6</sup>. **B.** Observed atmospheric CO<sub>2</sub> concentration from ice core (dots)<sup>20,21</sup> and box-model calculated CO<sub>2</sub> concentration. **C.** Observed atmospheric δ<sup>13</sup>C<sub>CO<sub>2</sub></sub> from ice core (blue shading)<sup>22</sup> and model-calculated value. **D.** Observed deep ocean δ<sup>13</sup>C - DIC from the global ocean (blue shading)<sup>23</sup>, from deep Pacific (dashed blue line)<sup>24</sup> and from a stack of 33 deep-ocean cores<sup>12</sup> and

290

295 model-calculated values. The  $\delta^{13}\text{C}$  values are plotted as anomalies relative to model results  
to highlight divergence in the mid- and late Holocene. In **B**, **C** and **D** solid line and orange  
band show the median values and uncertainties corresponding to peat C storage in **A**.  
Dashed line in **B** represents the outcome without “carbonate compensation” mechanism in  
the model. The box-model calculations show that peat C storage of >1000 GtC would result  
300 unrealistic atmosphere  $\text{CO}_2$  and  $\delta^{13}\text{CO}_2$  values and deep ocean  $\delta^{13}\text{C}$  value, significantly  
diverged from the observations.