

# Exploring coastal climate adaptation through storylines: Insights from Cyclone Idai in Beira, Mozambique

Henrique M.D. Goulart<sup>1,2</sup>, Panagiotis Athanasiou<sup>1</sup>, Kees van Ginkel<sup>1,2</sup>, Karin van der Wiel<sup>3</sup>, Gundula Winter<sup>1</sup>, Izidine Pinto<sup>3</sup>, and Bart van den Hurk<sup>1,2</sup>

<sup>1</sup>Deltares, Delft, The Netherlands

<sup>2</sup>Institute for Environmental Studies, VU University Amsterdam, The Netherlands

<sup>3</sup>Royal Netherlands Meteorological Institute (KNMI), De Bilt, The Netherlands

**Correspondence:** Henrique M.D. Goulart (henrique.goulart@deltares.nl)

## 1 Preprint statement

This is a non-peer reviewed preprint submitted to EarthArXiv.

## 2 Science for Society

Rising seas and stronger storms due to climate change pose significant threats to coastal communities. To effectively plan for adaptation against these events, local climate information is essential. Our study uses storylines — detailed narratives of potential high-impact events — to evaluate adaptation strategies based on local data. We use as case study a powerful tropical cyclone that has caused extensive damage on a coastal city and imagine how these impacts would change under multiple scenarios. Then we calculate the reductions in impacts due to different adaptation strategies. This approach empowers communities to visualise and understand the impacts of future extreme weather events and make informed decisions on how to best adapt against these events.

## 3 Summary

Coastal settlements, facing increasing flood risk from Tropical Cyclones (TCs) under climate change, need local and detailed climate information for effective adaptation. Analysis of historical events and their impacts provides such information. This study uses storylines to evaluate adaptation strategies, focusing on Cyclone Idai's impact on Beira, Mozambique, under different climate conditions and tidal cycles. A storyline of Idai under 3°C warming increases flood impacts by 1.8 times, while aligning Idai with spring tides amplifies these by 21 times. Combining both conditions increases impacts beyond 37 times. An adaptation strategy combining flood protection and accommodation measures reduces impacts by maximum 83%, while a seawall strategy reduces these by 10%. By offering localised, detailed information, storylines can be used to measure the effec-

tiveness of adaptation strategies against extreme events, evaluating their robustness across different scenarios, and quantifying residual impacts, complementing traditional climate risk assessments for informed decision making.

*Keywords:* storylines, coastal adaptation, decision making, coastal flooding, compound events, tropical cyclones, climate change

## 4 Introduction

Human settlements in coastal areas around the world face significant threats from tropical cyclones (TCs) (Dullaart et al., 2021; Ranasinghe et al., 2021). TCs cause extensive floods in coastal regions through heavy precipitation and storm surges (Bevacqua et al., 2019; Gori et al., 2020; Lai et al., 2021), which lead to casualties, property damage, and on longer time scales exacerbate poverty and hinder development in affected areas (Hallegatte et al., 2016a). Climate change is expected to increase the flood hazard from TCs globally (Knutson et al., 2021; Gori et al., 2022), mainly through sea level rise (Woodruff et al., 2013; Knutson et al., 2020) and more extreme precipitation (Patricola and Wehner, 2018; Knutson et al., 2010). This increased risk is particularly severe for low-income and vulnerable regions (Winsemius et al., 2016; Jongman, 2018; Rentschler et al., 2022), where local adaptation capacities are often limited (Fankhauser and McDermott, 2014; Hallegatte et al., 2016b). Consequently, the provision of localised and actionable climate information becomes imperative to support effective coastal adaptation (Nordgren et al., 2016).

There is a gap between traditional climate sciences and decision making (Hazeleger et al., 2015; Sutton, 2019; Rodrigues and Shepherd, 2022). Traditional approaches use probabilities to estimate future climate projections. These projections carry considerable uncertainty, and may impose limitations on exploring the full range of outcomes including the less likely ones (Hazeleger et al., 2015; Shepherd, 2019; Sutton, 2019). This is further exacerbated at the local scale, where uncertainties in the climate system, but also in human and environmental aspects increase (Shepherd and Lloyd, 2021; Sobel, 2021; Lehner and Deser, 2023). Yet, at this scale extreme events, e.g., TCs, generate impacts to society, requiring adaptation strategies to be implemented at a corresponding scale (Sobel, 2021; van den Hurk et al., 2023a). Therefore, probabilistic approaches relying on future climate projections might not fully satisfy the needs for effective adaptation and decision-making at the local scale (Dessai and Hulme, 2004; Shepherd and Sobel, 2020).

An alternative approach to these probabilistic approaches is the use of event storylines (Shepherd et al., 2018). Storylines, in this context, are physically plausible narratives of an event, considering their meteorological and climatic context and societal implications, without assessing prior probabilities of the events or their drivers (Shepherd et al., 2018). The potential to analyse detailed sequential hazard-to-impact chains allows storylines to serve as a bridge between global climate projections and local scale impacts (Shepherd and Lloyd, 2021; Rodrigues and Shepherd, 2022; van den Hurk et al., 2023a). Their contribution to providing decision-oriented information is achieved by expanding a reference event with alternative realisations under explicit assumptions on all – also non-climatic – drivers of the impact, offering a clear and meaningful way to assess and communicate potential impacts under different conditions to decision makers (Shepherd and Sobel, 2020; Sillmann et al., 2020). Previous

studies have adopted storylines to explore the effects of climate change for multiple impact sectors (e.g. Wiel et al., 2020; Goulart et al., 2021; Ciullo et al., 2021).

In flood modelling, risk-based approaches are commonly employed to identify flood risk and obtain cost-effective adaptation measures (de Moel and Aerts, 2011; Ward et al., 2017). However, they suffer from uncertainty in flood event probabilities (de Moel et al., 2015) and underestimate the significance of low-probability high-impact events (Merz et al., 2009). Incorporating climate change projections adds further uncertainty (McInerney et al., 2012; Haasnoot et al., 2013). Scenario-based approaches, which include storylines, offer an alternative to address the need for robust solutions amidst these uncertainties (Hall et al., 2012; Haasnoot et al., 2013; van der Pol et al., 2017). For coastal flooding, storylines were applied to efficiently stress-test flood scenarios (Qiu et al., 2022), to assess impacts from alternative flood events (Goulart et al., 2024), and to explore generic adaptation options across different regions Koks et al. (2023). However, using storylines to inform localised decision making through the integration of local adaptation strategies against specific high impact events is not yet common practice.

This study uses storylines to evaluate the effectiveness of different local adaptation strategies against a high impact event. Specifically, we investigate flood levels and societal impacts from TC Idai (2019) on the city of Beira, Mozambique. Our storylines are built using four different hydrometeorological scenarios (Figure 1), including climate change effects on precipitation and sea level rise, and changes in the timing of the storm relative to tidal cycles. We consider three local coastal adaptation strategies reflecting different approaches to flood protection. Our modelling framework spans the event's meteorological conditions, compound flooding using a hydrodynamic model, and the flood impacts on Beira, specifically population exposure and building damage via an object-based impact model (details for all steps in experimental procedures).


## 5 Results

### 5.1 Flood impacts from Idai in Beira substantially increase with climate change and spring tides


We evaluate the hazards and impacts of TC Idai in Beira for four hydrometeorological scenarios. Idai, one of the strongest storms ever recorded in Southern Africa, caused extensive damage in Beira, Mozambique in 2019 (see Experimental Procedures for more details). In the hydrometeorological Baseline scenario, which reflects the historic event, widespread compound coastal flooding occurs in Beira and specifically along its west coast (Figure 2a). Idai originally made landfall during neap tides, and having the storm coincide with spring tides – the Springtide scenario – leads to a substantially larger inundation extent and depth (Figure 2b). A 3°C climate change by 2100 (3C scenario) leads to an increase in flood extent and depth through precipitation increase and sea level rise (Figure 2c). The 3C scenario shows smaller flood increases compared to the Springtide scenario. This is because sea level rise in 3C scenario leads to lower increase in water levels (0.59 m) than the tidal effect in the Springtide scenario (1.24 m difference between neap and spring tides). Ultimately, the combination of these two scenarios (3C-springtide) leads to the largest flood extent and depth, with most of the study area experiencing flooding (Figure 2d).

We quantify the flood impacts of Idai in Beira in terms of population exposure and building damage. Our results show that approximately 5,000 people are exposed to water depths >15 cm in the Baseline scenario (Figure 3). The Springtide scenario indicates approximately 103,000 people are exposed (20x the Baseline), which is substantially more than the 9,400 exposed


## a) Modelling framework



## b) Hydrometeorological scenarios




## c) Adaptation strategies



**Figure 1. General framework of the study.** a) The modelling framework connecting meteorological conditions, such as wind speed and precipitation, to compound flood and impact. b) The four hydrometeorological scenarios considered: Baseline, 3C, Springtide, and 3C-springtide. c) The three adaptation strategies for the city of Beira: no adaptation, Hold the Line, and Integrated strategies. The red lines indicate walls and dikes; the orange polygon indicates the port region to be elevated, and the yellow polygon indicate the coastal area to retreat from (see Experimental procedures for details).


85 people in the 3C scenario (1.8x the Baseline). The 3C-springtide scenario leads to the highest number of people exposed, around 194,000 (37x the Baseline). In addition to changes in flood extent, a noticeable shift towards higher flood levels for more extreme events is shown (colours in Figure 3). Fraction of people exposed to high flood levels, representing depths above 150 cm, go from 1.6% of the exposed population in the Baseline scenario to 14.5% on the 3C-springtide scenario. Concluding,



**Figure 2. Flood maps of Idai in Beira.** Flood hazard maps of TC Idai under different hydrometeorological scenarios. Different shades of blue indicate flood depth. Note the blue circle in the centre is a small lake within the city bounds.

the more extreme the hydrometeorological scenario of Idai, the more people are directly exposed, and their exposure is to increasingly severe hazards.


90 Economic damage shows similar patterns to population exposure (Figure 3b). The Baseline scenario indicates damages of USD 2.73 million in Beira. The Springtide scenario presents damages of USD 61 million, an increase of 22x the Baseline, and more than the damages in the 3C scenario, USD 4.6 million (1.7x the Baseline). As a consequence of both spring tides and 3C climate scenario, the 3C-springtide peaks at USD 152 million, which is 56x the Baseline, showing non-linear compounding effects. Damage maps at the building level show spatial details on the impacts in Beira (Figure 4) and reveal that informal  
 95 settlements suffer significantly, with numerous cases of total losses in more extreme scenarios.



**Figure 3. Impacts of Idai in Beira.** a) Total population exposure and b) total building damage in the city of Beira from Idai floodings under different hydrometeorological scenarios. Light blue indicates water level between 15 and 50cm, medium blue 50 and 150cm, and dark blue above 150cm.

## 5.2 The Integrated strategy reduces flood impacts more than the Hold the Line strategy


We assess how effectively each adaptation strategy (described in detail in Experimental procedures) performs across the hydrometeorological scenarios (Figure 5). In the Baseline scenario, the Hold the Line strategy, consisting of an extensive seawall along the coast of Beira, reduces population exposure and building damage by approximately 11% and 7.6%, respectively. The Integrated strategy, including dikes around the city centre, port elevation and managed retreat of wetlands, reduces population exposure and damage by 2% and 9.5%, respectively. However, for the counterfactual scenarios, the Integrated strategy consistently outperforms the Hold the Line strategy in reducing population exposure: for the Springtide, 3C, and 3C-springtide scenarios, exposure reductions are 83%, 8%, and 75%, respectively, versus reductions of 9%, 0.3%, and 3.4% with the Hold the Line strategy (Figure 5a). Consequently, we observe a greater decrease in exposure to high water levels with the Integrated strategy compared to the Hold the Line strategy (Figure SI A1). Economic damage follows a similar trend, with the Integrated strategy reducing damages by 84%, 30%, and 81% for the Springtide, 3C and 3C-springtide scenarios, respectively. The Hold the Line strategy shows lower reductions of 10%, 0.7%, and 7.4% (Figure 5b). While none of the included adaptation strategies completely protects Beira from the impacts from Idai across our tested scenarios, the Integrated strategy proves more effective than the Hold the Line in reducing building damage and population exposed, especially for the more extreme hydrometeorological scenarios. These results offer insights on the effectiveness of each adaptation strategy across different scenarios and on the corresponding residual impacts in Beira (Figure 5c,d).



**Figure 4. Damage maps of Beira.** Maps of Beira showing relative economic damage relative to the total value of each building in Beira for each hydrometeorological scenarios. Black contours indicate informal settlements.

## 6 Discussion

Our study develops storylines of Idai to assess the effectiveness of different local coastal adaptation strategies under diverse hydrometeorological scenarios. These include the effects of climate change, sea level rise, and changes in the tidal cycle. We use a comprehensive modelling framework that includes the event's meteorological conditions, compound flood simulation, and the modelling of population exposure and building damage.



**Figure 5. Effectiveness of adaptation strategies.** Reductions in a) population exposure and b) building damage for each adaptation strategy and hydrometeorological scenario. Plain bars represent Hold the Line strategy and hatched bars represent integrated strategy. Maps of relative building damage for c) the Hold the Line and d) Integrated strategies under the 3C-springtide scenario. Black contours show informal settlements.

### 6.1 Insights of using storylines for coastal adaptation in Beira

Our results show that flood impacts in Beira will likely worsen due to climate change, primarily driven by sea level rise. These findings align with existing research showing the threat of rising sea levels to coastal settlements (Hallegatte et al., 2013; Hinkel et al., 2018). We find that a shift in TC Idai's timing to coincide with spring tides could lead to even more severe flooding. This is because the difference in surge height between neap tide and spring tide is more than double the projected SLR by 2100 in a 3°C warming scenario. Our most impactful storyline of Idai is the combination of both climate change and



spring tides, showing again the strong negative effects of climate change for Beira and its inhabitants. In addition, it shows that climate change is one of multiple contributing factors to impacts, as explored here by altering the compound timing of storm landfall and the tidal cycle. This change in timing demonstrates the large range of impacts of local extreme events due to internal variability (Done et al., 2014; Goulart et al., 2023; Lehner and Deser, 2023). Our results highlight the importance of incorporating internal variability next to the role of climate change in adaptation planning (Hinkel et al., 2014; Haasnoot et al., 2020; Herman et al., 2020).

The scope of our study diverges from existing risk-based studies for Beira, such as Eilander et al. (2023b) and van Berchum et al. (2020). Risk-based studies are able to quantify expected annual flood impacts based on probabilities and to determine cost-effective solutions. However, they may not fully account for the complexities of extreme weather events (de Moel et al., 2015; Merz et al., 2009) and climate change uncertainties (McInerney et al., 2012; Haasnoot et al., 2013). While Eilander et al. (2023b) identified managed retreat in highly exposed areas of Beira as the most efficient to reduce expected annual impacts, van Berchum et al. (2020) suggest coastal defences are the most effective long-term measure. Our approach shift focus from estimating the most optimal or cost-effective measures to assessing the performance of specified local adaptation strategies under different scenarios. We find that no strategy fully prevents flooding in Beira from Cyclone Idai, but that the Integrated strategy substantially outperforms the Hold the Line approach in more extreme scenarios. The wide range of impacts across scenarios and strategies in our results enable policymakers and stakeholders to visualise the benefits and limitations of different adaptation strategies when faced with extreme events similar to Cyclone Idai under different scenarios. This includes assessing the robustness of each adaptation strategy against the selected events (Hall et al., 2012; van der Pol et al., 2017) and quantifying residual impacts (Lim et al., 2018; Tanoue et al., 2021), which can inform the planning of complementary measures like evacuation plans or financial aid programs (van den Hurk et al., 2023b).

## 6.2 Validation of simulations, limitations, and recommendations

Our Baseline simulations show more extensive inundation than satellite imagery from the Emergency Management Service (EMS) has shown for Idai landfall in Beira (Copernicus Emergency Management Service, 2019). Satellite imagery has limitations in densely populated areas (Fang Zhang and Liu, 2014). Mester et al. (2023) found this to be the case for Cyclone Idai's impact on Beira, as satellite imagery showed substantially less flooded areas than what has been documented in reports and media (Segerlin et al., 2020). Another study showed that a combined 10-year rainfall and 10-year coastal surge event causes more extensive flooding in Beira than the satellite images (van Berchum et al., 2020), and TC Idai is considered to have a lower probability than once per 10 years (Emerton et al., 2020). Given the uncertainty around the actual flooding extent induced by Idai and our focus on exploring differences between the storylines, we consider the simulated data suitable for this study's purpose.

This study focuses on the physical aspects of climate change, without incorporating socio-economic or land use changes, which greatly influence vulnerability and exposure, and therefore impacts (Koks et al., 2023). Exploring these and other relevant drivers of local change could lead to a more comprehensive impact analysis of future scenarios and adaptation strategies. In this study, we have used "robustness" to refer to how well an adaptation strategy performs across a variety of scenarios (Hall

et al., 2012; Haasnoot et al., 2013; van der Pol et al., 2017). Our focus on Cyclone Idai, despite varying scenarios, implies only one type of event is being examined. Future studies applying storylines for robust adaptation could consider multiple extreme events with different characteristics, e.g. a precipitation-dominated flood event.

160 We adopt three adaptation strategies in our study, but future work could evaluate a broader range of potential adaptation options for the study area, such as drainage systems, nature-based solutions, and evacuation plans. Our adaptation strategy designs are based on recommendations for the city of Beira from local sources (Van Logchem and Queface, 2012). Some adaptation measures, like the managed retreat included in the Integrated strategy, carry considerable social implications (Bongarts Lebbe et al., 2021; Hino et al., 2017; Gussmann and Hinkel, 2021). The findings in this study are mostly exploratory, and actual adaptation  
165 could emerge from collaborative efforts directly involving the communities and stakeholders impacted (Bongarts Lebbe et al., 2021). Our analysis focuses solely on building damage and population exposure. Future work could broaden this scope to include indirect effects, such as interrupted services and economic disruption due to supply shortages (Pant et al., 2018; Tanoue et al., 2020)

### **6.3 Storylines for decision making on adaptation**

170 To effectively support local adaptation and decision making, climate information needs to be usable, relevant and local (Sutton, 2019; Sobel, 2021; Rodrigues and Shepherd, 2022). Our study extends the storyline approach to include adaptation strategies specifically designed for local scale decision making. In this way, they enable stakeholders to visualise the consequences of specific local adaptation measures against high-impact events across relevant possible futures (“what if” scenarios). Consequently, they provide a practical method for quantifying the effectiveness of adaptation strategies under different scenarios, facilitating  
175 the identification of robust adaptation strategies. Finally, they can also be used to quantify residual impacts, which supports the planning of complementary adaptation measures, including early warning systems and evacuation plans (van den Hurk et al., 2023b), and post-disaster recovery mechanisms such as insurance schemes (Kousky, 2019). We find that, by including specific adaptation strategies to local contexts, storylines offer a complementary perspective to traditional probabilistic approaches for informing climate adaptation strategies against high impact events.

## **180 7 Experimental procedures**

### **7.1 Resource availability**

#### **7.1.1 Lead contact**

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Henrique M. D. Goulart (henrique.goulart@deltares.nl).

#### **185 7.1.2 Materials availability**

This study did not generate new unique materials.

### 7.1.3 Data and code availability

The code and data generated for this experiment is available at: [https://github.com/dumontgoulart/storylines\\_for\\_adaptation](https://github.com/dumontgoulart/storylines_for_adaptation). SFINCS is available at <https://sfincs.readthedocs.io> and HydroMT is available at <https://deltares.github.io/hydromt/>. Delft-  
190 FIAT is available at <https://github.com/Deltares/Delft-FIAT>.

## 7.2 Overview

Our storylines explore two components of the event and its impacts: the meteorological hazard and adaptation strategies in the city of Beira (Figure 1). We have developed a modelling framework that captures meteorology, coastal flooding and societal impacts (economic damage to buildings and exposed population) (Figure 1). Based on four distinct meteorological/hydromete-  
195 orological scenarios and three local adaptation strategies, we develop twelve unique storylines. They enable an exploration of the potential impacts of TC Idai on the city of Beira, and the effectiveness of different adaptation strategies in reducing these impacts.

## 7.3 Case study


We explore the impacts of TC Idai on the city of Beira, Mozambique. TC Idai was one of the most impactful TCs to occur in  
200 Southern Africa, affecting mainly Mozambique, Zimbabwe, and Malawi. In Mozambique, 598 casualties were reported, and a further 1600 people were reported injured. Furthermore, it caused damage or complete destruction to nearly 198,000 homes, decimated crop fields, triggered a cholera epidemic, and left an estimated 1.85 million people affected (UN Office for the Coordination of Humanitarian Affairs, 2019).

TC Idai originated off the East coast of Mozambique on March 4, 2019, and it briefly reached Category 4 with peak wind  
205 speeds of 59 m/s (Dube et al., 2021). Idai made landfall twice, with the second one being March 14 near Beira city, the fourth largest city in Mozambique (Figure 6). The region experienced severe impacts mainly from extreme wind speeds and compound coastal flooding, driven by intense precipitation and the storm surge. Though storm surge levels in Beira reached approximately 4 m, the event coincided with a neap tide period which had a limiting effect on water levels (van Berchum et al., 2020; Eilander et al., 2023b).

## 210 7.4 Hydrometeorological scenarios

The Idai storylines are built based on hydrometeorological scenarios designed to explore both the influence of internal variability and climate change around the event. We consider four scenarios:

- the **Baseline** scenario reflects the event as it occurred in 2019. Note that also in this scenario there is some climate change component: global temperatures lie at 1.2 °C above pre-industrial levels, and there is a sea level rise (SLR) of  
215 5 cm relative to the 1995-2014 average.



**Figure 6. Location of Mozambique and Beira.** a) The location of Mozambique (dark gray) and the city of Beira (red square). b) Expanded view of Beira with buildings footprints (black polygons).

- the **Springtide** scenario, a counterfactual scenario that simulates TC Idai occurring in conjunction with spring tides. This introduces an element of internal variability, reflecting natural fluctuations in environmental conditions that can significantly influence local flooding. It involves adjusting the timing of Idai by four days to coincide with the spring tides, while maintaining climatological conditions identical to the baseline.
- 220 – the **3C** scenario, a future counterfactual scenario assuming a global temperature rise of 3°C above pre-industrial levels by 2100. This scenario includes a SLR of 0.59 m and a precipitation increase of 13 % compared to the baseline. This scenario aligns with the global warming projections of 3.2°C by 2100 based on current Nationally Determined Contributions (NDCs) (IPCC, 2023).
- the **3C-springtide** scenario, a compound scenario that combines the springtide event with the future 3°C global warming
- 225 condition.

SLR data was derived from the sixth assessment report (AR6) from the Intergovernmental Panel on Climate Change (IPCC) (IPCC, 2021). We calculate the change in precipitation in a warmer climate using the Clausius-Clapeyron (CC) relation, which establishes a 7 % increase in saturation vapour pressure for each degree of warming. This is in line with findings from recent studies on the increase of precipitation rates of TCs due to climate change in the southern Indian Ocean (Liu et al., 2019; 230 Knutson et al., 2020).

## 7.5 Adaptation strategies

This study assesses the effectiveness of three local coastal adaptation strategies in Beira city. They are based on previous local reports (Van Logchem and Queface, 2012) and designed to provide a clear comparison between distinct approaches to reduce the societal impacts of the Idai event:

- 235 – **No adaptation** strategy, where no further protective measures are adopted.
- **Hold the Line** strategy (Figure 1), focused on protecting the entire land area through the construction of hard infrastructure along the coastline Bongarts Lebbe et al. (2021). Here, it consists of a 2 m wall along the Beira coastline.
- **Integrated** strategy (Figure 1), which combines infrastructure with management and accommodation measures (Bongarts Lebbe et al., 2021; IPCC, 2022). In this study, it consists of a 2 m wall along part of the coast and around the centre  
240 of the city, a managed retreat of settlements from the vulnerable coastal wetlands in the city’s southwest, and raising the port’s elevation by 2 m.

The 2 m height adopted for seawalls and port elevation is based on the 100-year return period surge projections for the region, as identified in Eilander et al. (2023b). All measures, including the managed retreats, are based on recommendations from a local report by the Maputo National Institute for Disaster Management (INGC) (Van Logchem and Queface, 2012).  
245 Managed retreats carry societal implications, which requires careful planning and stakeholder involvement Bongarts Lebbe et al. (2021).

## 7.6 Modelling framework

### 7.6.1 Meteorological data and evaluation

Idai meteorological data for mean sea level pressure (MSLP), wind speed and precipitation is obtained from the high resolution  
250 Integrated Forecast System (IFS) model of the European Centre for Medium-Range Weather Forecasts (ECMWF). It is based on a coupled atmosphere–wave–ocean model (Mogensen et al., 2017), has hourly temporal resolution, and offers the highest spatial resolution among global forecasts, 0.1°, which improves TC simulation (Knutson et al., 2020; Magnusson et al., 2019; Zhang et al., 2023). Previous studies have assessed the capabilities, advancements, and limitations of IFS in simulating TCs (Mogensen et al., 2017; Magnusson et al., 2019; Heming et al., 2019; Becker et al., 2021; Christophersen et al., 2022).

255 There is limited observation data for TC Idai and its impacts in Beira (van Berchum et al., 2020; Mester et al., 2023). We adopt the International Best Track Archive for Climate Stewardship (IBTrACS) (Knapp et al., 2010) to evaluate the IFS-simulated minimum MSLP and maximum wind speeds, and the Integrated Multi-satellitE Retrievals for GPM (IMERG-GPM) (Huffman et al., 2015) for precipitation. Subsequently, we align the values from IFS with observed values, so that floods and impacts in Beira are more accurately simulated. This involves adjusting the model’s mean values around the storm’s centre  
260 during its landfall between March 14th and 15th to match the evaluation data.


## 7.6.2 Compound coastal flooding modelling

We use the Super-Fast INundation of CoastS (SFINCS) model (Leijnse et al., 2021) for both offshore and onshore hydrodynamic simulation. SFINCS is a reduced-physics solver that accurately simulates compound coastal flooding by solving simplified two-dimensional overland flow equations. Its suitability for simulating compound flooding resulting from TCs has been demonstrated in previous studies (Leijnse et al., 2021; Sebastian et al., 2021; Eilander et al., 2023b; Goulart et al., 2024; Nederhoff et al., 2024). A full description of the model is available at Leijnse et al. (2021). The offshore simulation is forced with MSLP and wind speed data from IFS, generating water levels along the coastline of Beira. The onshore simulation is then forced with the generated water levels and precipitation to produce inland flooding levels in Beira. The surface elevation is obtained from a merged dataset that combines several local and global datasets, achieving a 5 m resolution in Beira (Nederhoff et al., 2024). The roughness coefficients are sourced from the Copernicus Global Land Service (Buchhorn et al., 2020) and infiltration rates derived from the GCN250 dataset (Jaafar et al., 2019). For the management and processing of input data, we use the Python package HydroMT (Eilander et al., 2023a).

## 7.6.3 Impact modelling

We use the Delft-FIAT impact model (Slager et al., 2016; Eilander et al., 2023b) to quantify building damages and the population exposed to floods under the different storylines. Delft-FIAT combines flood extent and depths with exposure and vulnerability data, enabling impact modelling at the individual building level. For building exposure, including location and footprint, we use data from OpenStreetMap (Haklay and Weber, 2008). Population data from WorldPop 2020 UN adjusted database at 100 m resolution (Bondarenko et al., 2020) is downscaled to the building level by using the buildings footprints size as weights. Vulnerability curves to estimate the economic damages due to flooding for different types of building are obtained from Huizinga et al. (2017).

## Appendix A: Supplemental information



**Figure A1. Impacts in Beira with different adaptation strategies.** Similar to Figure 3 but showing residual impact for each adaptation strategy.

*Acknowledgements.* This research has been supported by the European Union's Horizon 2020 research and innovation programme under grant agreement No 820712 (project RECEIPT, REremote Climate Effects and their Impact on European sustainability, Policy and Trade). We thank D. Eilander, T. Kelder, K. Nederhoff, R. Hamed, T. Leijnse, T. Tiggeloven and A. Alexander for support and comments on previous

*Author contributions.* HG, KvdW and BvdH contributed to the concept of the study. HG conducted the research and edited the manuscript. HG, IP and KvdW obtained and evaluated the meteorological data. KvG, HG and IP designed and implemented adaptation strategies. HG, PA, GW set up the hydrodynamic and impact models. All authors discussed the analysis and results, and revised the manuscript. BvdH and KvdW supervised the work.

290 *Competing interests.* The authors declare that they have no competing interests.



## References

- Becker, T., Bechtold, P., and Sandu, I.: Characteristics of convective precipitation over tropical Africa in storm-resolving global simulations, *Quarterly Journal of the Royal Meteorological Society*, 147, 4388–4407, <https://doi.org/https://doi.org/10.1002/qj.4185>, 2021.
- 295 Bevacqua, E., Maraun, D., Voudoukas, M. I., Voukouvalas, E., Vrac, M., Mentaschi, L., and Widmann, M.: Higher probability of compound flooding from precipitation and storm surge in Europe under anthropogenic climate change, *Science Advances*, 5, eaaw5531, <https://doi.org/10.1126/sciadv.aaw5531>, 2019.
- Bondarenko, M., Kerr, D., Sorichetta, A., and Tatem, A.: Census/projection-disaggregated gridded population datasets, adjusted to match the corresponding UNPD 2020 estimates, for 51 countries across sub-Saharan Africa using building footprints, <https://doi.org/10.5258/SOTON/WP00683>, available online at <https://www.worldpop.org/>, 2020.
- 300 Bongarts Lebbe, T., Rey-Valette, H., Chaumillon, É., Camus, G., Almar, R., Cazenave, A., Claudet, J., Rocle, N., Meur-Ferec, C., Viard, F., et al.: Designing coastal adaptation strategies to tackle sea level rise, *Frontiers in Marine Science*, 8, 740 602, 2021.
- Buchhorn, M., Smets, B., Bertels, L., Roo, B. D., Lesiv, M., Tsendbazar, N.-E., Herold, M., and Fritz, S.: Copernicus Global Land Service: Land Cover 100m: collection 3: epoch 2015: Globe, <https://doi.org/10.5281/zenodo.3939038>, 2020.
- Christophersen, H., Sippel, J., Aksoy, A., and Baker, N. L.: Recent advancements for tropical cyclone data assimilation, *Annals of the New York Academy of Sciences*, 1517, 25–43, <https://doi.org/https://doi.org/10.1111/nyas.14873>, 2022.
- 305 Ciullo, A., Martius, O., Strobl, E., and Bresch, D. N.: A framework for building climate storylines based on downward counterfactuals: The case of the European Union Solidarity fund, *Climate Risk Management*, 33, 100 349, <https://doi.org/https://doi.org/10.1016/j.crm.2021.100349>, 2021.
- Copernicus Emergency Management Service: [EMSR348] Beira: Delineation Map, Monitoring 2, [https://emergency.copernicus.eu/mapping/ems-product-component/EMSR348\\_03BEIRA\\_01DELINEATION\\_MONIT02/2](https://emergency.copernicus.eu/mapping/ems-product-component/EMSR348_03BEIRA_01DELINEATION_MONIT02/2), product version: v2, 2019.
- 310 de Moel, H. and Aerts, J.: Effect of uncertainty in land use, damage models and inundation depth on flood damage estimates, *Natural Hazards*, 58, 407–425, 2011.
- de Moel, H., Jongman, B., Kreibich, H., Merz, B., Penning-Rowsell, E., and Ward, P. J.: Flood risk assessments at different spatial scales, *Mitigation and Adaptation Strategies for Global Change*, 20, 865–890, 2015.
- 315 Dessai, S. and Hulme, M.: Does climate adaptation policy need probabilities?, *Climate Policy*, 4, 107–128, <https://doi.org/10.1080/14693062.2004.9685515>, 2004.
- Done, J. M., Bruyère, C. L., Ge, M., and Jaye, A.: Internal variability of North Atlantic tropical cyclones, *Journal of Geophysical Research: Atmospheres*, 119, 6506–6519, <https://doi.org/10.1002/2014JD021542>, 2014.
- Dube, K., Chapungu, L., and Fitchett, J. M.: Meteorological and Climatic Aspects of Cyclone Idai and Kenneth, pp. 19–36, Springer International Publishing, Cham, [https://doi.org/10.1007/978-3-030-74262-1\\_2](https://doi.org/10.1007/978-3-030-74262-1_2), 2021.
- 320 Dullaart, J. C., Muis, S., Bloemendaal, N., Chertova, M. V., Couasnon, A., and Aerts, J. C.: Accounting for tropical cyclones more than doubles the global population exposed to low-probability coastal flooding, *Communications Earth & Environment*, 2, 135, 2021.
- Eilander, D., Boisgontier, H., Bouaziz, L. J. E., Buitink, J., Couasnon, A., Dalmijn, B., Hegnauer, M., de Jong, T., Loos, S., Marth, I., and van Verseveld, W.: HydroMT: Automated and reproducible model building and analysis, *Journal of Open Source Software*, 8, 4897, <https://doi.org/10.21105/joss.04897>, 2023a.
- 325

- Eilander, D., Couasnon, A., Sperna Weiland, F. C., Ligtoet, W., Bouwman, A., Winsemius, H. C., and Ward, P. J.: Modeling compound flood risk and risk reduction using a globally applicable framework: a pilot in the Sofala province of Mozambique, *Natural Hazards and Earth System Sciences*, 23, 2251–2272, <https://doi.org/10.5194/nhess-23-2251-2023>, 2023b.
- Emerton, R., Cloke, H., Ficchi, A., Hawker, L., de Wit, S., Speight, L., Prudhomme, C., Rundell, P., West, R., Neal, J., Cuna, J., Harrigan, S.,  
330 Titley, H., Magnusson, L., Pappenberger, F., Klingaman, N., and Stephens, E.: Emergency flood bulletins for Cyclones Idai and Kenneth: A critical evaluation of the use of global flood forecasts for international humanitarian preparedness and response, *International Journal of Disaster Risk Reduction*, 50, 101 811, <https://doi.org/https://doi.org/10.1016/j.ijdr.2020.101811>, 2020.
- Fang Zhang, X. Z. and Liu, D.: Blending MODIS and Landsat images for urban flood mapping, *International Journal of Remote Sensing*, 35, 3237–3253, <https://doi.org/10.1080/01431161.2014.903351>, 2014.
- 335 Fankhauser, S. and McDermott, T. K.: Understanding the adaptation deficit: Why are poor countries more vulnerable to climate events than rich countries?, *Global Environmental Change*, 27, 9–18, <https://doi.org/https://doi.org/10.1016/j.gloenvcha.2014.04.014>, 2014.
- Gori, A., Lin, N., and Xi, D.: Tropical Cyclone Compound Flood Hazard Assessment: From Investigating Drivers to Quantifying Extreme Water Levels, *Earth’s Future*, 8, e2020EF001 660, <https://doi.org/https://doi.org/10.1029/2020EF001660>, e2020EF001660 2020EF001660, 2020.
- 340 Gori, A., Lin, N., Xi, D., and Emanuel, K.: Tropical cyclone climatology change greatly exacerbates US extreme rainfall–surge hazard, *Nature Climate Change*, 12, 171–178, 2022.
- Goulart, H. M. D., van der Wiel, K., Folberth, C., Balkovic, J., and van den Hurk, B.: Storylines of weather-induced crop failure events under climate change, *Earth System Dynamics*, 12, 1503–1527, <https://doi.org/10.5194/esd-12-1503-2021>, 2021.
- Goulart, H. M. D., van der Wiel, K., Folberth, C., Boere, E., and van den Hurk, B.: Increase of Simultaneous Soybean Failures Due To  
345 Climate Change, *Earth’s Future*, 11, e2022EF003 106, <https://doi.org/10.1029/2022EF003106>, e2022EF003106 2022EF003106, 2023.
- Goulart, H. M. D., Benito Lazaro, I., van Garderen, L., van der Wiel, K., Le Bars, D., Koks, E., and van den Hurk, B.: Compound flood impacts from Hurricane Sandy on New York City in climate-driven storylines, *Natural Hazards and Earth System Sciences*, 24, 29–45, <https://doi.org/10.5194/nhess-24-29-2024>, 2024.
- Gussmann, G. and Hinkel, J.: Vested interests, rather than adaptation considerations, explain varying post-tsunami relocation outcomes in  
350 Laamu atoll, Maldives, *One Earth*, 4, 1468–1476, <https://doi.org/https://doi.org/10.1016/j.oneear.2021.09.004>, 2021.
- Haasnoot, M., Kwakkel, J. H., Walker, W. E., and Ter Maat, J.: Dynamic adaptive policy pathways: A method for crafting robust decisions for a deeply uncertain world, *Global environmental change*, 23, 485–498, 2013.
- Haasnoot, M., Kwadijk, J., van Alphen, J., Bars, D. L., van den Hurk, B., Diermanse, F., van der Spek, A., Essink, G. O., Delsman, J., and  
355 Mens, M.: Adaptation to uncertain sea-level rise; how uncertainty in Antarctic mass-loss impacts the coastal adaptation strategy of the Netherlands, *Environmental Research Letters*, 15, 034 007, <https://doi.org/10.1088/1748-9326/ab666c>, 2020.
- Haklay, M. and Weber, P.: OpenStreetMap: User-Generated Street Maps, *IEEE Pervasive Computing*, 7, 12–18, <https://doi.org/10.1109/MPRV.2008.80>, 2008.
- Hall, J. W., Lempert, R. J., Keller, K., Hackbarth, A., Mijere, C., and McInerney, D. J.: Robust Climate Policies Under Uncertainty: A Comparison of Robust Decision Making and Info-Gap Methods, *Risk Analysis*, 32, 1657–1672, <https://doi.org/https://doi.org/10.1111/j.1539-6924.2012.01802.x>, 2012.
- 360 Hallegatte, S., Green, C., Nicholls, R. J., and Corfee-Morlot, J.: Future flood losses in major coastal cities, *Nature climate change*, 3, 802–806, 2013.

- Hallegatte, S., Bangalore, M., Bonzanigo, L., Fay, M., Kane, T., Narloch, U., Rozenberg, J., Treguer, D., and Vogt-Schilb, A.: Shock waves: managing the impacts of climate change on poverty, World Bank Publications, 2016a.
- 365 Hallegatte, S., Vogt-Schilb, A., Bangalore, M., and Rozenberg, J.: Unbreakable: Building the Resilience of the Poor in the Face of Natural Disasters, The World Bank, <https://doi.org/10.1596/978-1-4648-1003-9>, 2016b.
- Hazeleger, W., Hurk, B. J. V. D., Min, E., Oldenborgh, G. J. V., Petersen, A. C., Stainforth, D. A., Vasileiadou, E., and Smith, L. A.: Tales of future weather, *Nature Climate Change*, 5, 107–113, <https://doi.org/10.1038/nclimate2450>, 2015.
- Heming, J. T., Prates, F., Bender, M. A., Bowyer, R., Cangialosi, J., Caroff, P., Coleman, T., Doyle, J. D., Dube, A., Faure, G., Fraser, J., 370 Howell, B. C., Igarashi, Y., McTaggart-Cowan, R., Mohapatra, M., Moskaitis, J. R., Murtha, J., Rivett, R., Sharma, M., Short, C. J., Singh, A. A., Tallapragada, V., Titley, H. A., and Xiao, Y.: Review of Recent Progress in Tropical Cyclone Track Forecasting and Expression of Uncertainties, *Tropical Cyclone Research and Review*, 8, 181–218, <https://doi.org/https://doi.org/10.1016/j.tcr.2020.01.001>, 2019.
- Herman, J. D., Quinn, J. D., Steinschneider, S., Giuliani, M., and Fletcher, S.: Climate Adaptation as a Control Problem: Review and Perspectives on Dynamic Water Resources Planning Under Uncertainty, *Water Resources Research*, 56, e24389, 375 <https://doi.org/https://doi.org/10.1029/2019WR025502>, e24389 2019WR025502, 2020.
- Hinkel, J., Lincke, D., Vafeidis, A. T., Perrette, M., Nicholls, R. J., Tol, R. S. J., Marzeion, B., Fettweis, X., Ionescu, C., and Levermann, A.: Coastal flood damage and adaptation costs under 21st century sea-level rise, *Proceedings of the National Academy of Sciences*, 111, 3292–3297, <https://doi.org/10.1073/pnas.1222469111>, 2014.
- Hinkel, J., Aerts, J. C., Brown, S., Jiménez, J. A., Lincke, D., Nicholls, R. J., Scussolini, P., Sanchez-Arcilla, A., Vafeidis, A., and Addo, 380 K. A.: The ability of societies to adapt to twenty-first-century sea-level rise, *Nature Climate Change*, 8, 570–578, 2018.
- Hino, M., Field, C. B., and Mach, K. J.: Managed retreat as a response to natural hazard risk, *Nature climate change*, 7, 364–370, 2017.
- Huffman, G. J., Bolvin, D. T., Braithwaite, D., Hsu, K., Joyce, R., Xie, P., and Yoo, S.-H.: NASA global precipitation measurement (GPM) integrated multi-satellite retrievals for GPM (IMERG), Algorithm theoretical basis document (ATBD) version, 4, 30, 2015.
- Huizinga, J., De Moel, H., and Szewczyk, W.: Global flood depth-damage functions: Methodology and the database with guidelines, Tech. 385 rep., Joint Research Centre (Seville site), <https://doi.org/https://doi.org/10.2760/16510>, 2017.
- IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, vol. In Press, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896>, 2021.
- IPCC: Climate Change 2022: Impacts, Adaptation and Vulnerability, Summary for Policymakers, Cambridge University Press, Cambridge, 390 UK and New York, USA, 2022.
- IPCC: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Summary for Policymakers, IPCC, Geneva, Switzerland, <https://doi.org/10.59327/IPCC/AR6-9789291691647.001>, 2023.
- Jaafar, H. H., Ahmad, F. A., and El Beyrouthy, N.: GCN250, new global gridded curve numbers for hydrologic modeling and design, 395 *Scientific data*, 6, 145, 2019.
- Jongman, B.: Effective adaptation to rising flood risk, *Nature communications*, 9, 1986, 2018.
- Knapp, K. R., Kruk, M. C., Levinson, D. H., Diamond, H. J., and Neumann, C. J.: The international best track archive for climate stewardship (IBTrACS) unifying tropical cyclone data, *Bulletin of the American Meteorological Society*, 91, 363–376, 2010.

- Knutson, T., Camargo, S. J., Chan, J. C. L., Emanuel, K., Ho, C.-H., Kossin, J., Mohapatra, M., Satoh, M., Sugi, M., Walsh, K., and Wu, L.:  
400 Tropical Cyclones and Climate Change Assessment: Part II: Projected Response to Anthropogenic Warming, *Bulletin of the American Meteorological Society*, 101, E303 – E322, <https://doi.org/https://doi.org/10.1175/BAMS-D-18-0194.1>, 2020.
- Knutson, T. R., McBride, J. L., Chan, J., Emanuel, K., Holland, G., Landsea, C., Held, I., Kossin, J. P., Srivastava, A., and Sugi, M.: Tropical cyclones and climate change, *Nature geoscience*, 3, 157–163, 2010.
- Knutson, T. R., Chung, M. V., Vecchi, G., Sun, J., Hsieh, T.-L., and Smith, A. J.: Climate change is probably increasing the intensity of  
405 tropical cyclones, *Critical Issues in Climate Change Science, Science Brief Review*. <https://doi.org/10.5281/zenodo.4570334>, 2021.
- Koks, E. E., Bars, D. L., Essenfelder, A., Nirandjan, S., and Sayers, P.: The impacts of coastal flooding and sea level rise on critical infrastructure: a novel storyline approach, *Sustainable and Resilient Infrastructure*, 8, 237–261, <https://doi.org/10.1080/23789689.2022.2142741>, 2023.
- Kousky, C.: The role of natural disaster insurance in recovery and risk reduction, *Annual Review of Resource Economics*, 11, 399–418,  
410 2019.
- Lai, Y., Li, J., Gu, X., Liu, C., and Chen, Y. D.: Global Compound Floods from Precipitation and Storm Surge: Hazards and the Roles of Cyclones, *Journal of Climate*, 34, 8319 – 8339, <https://doi.org/10.1175/JCLI-D-21-0050.1>, 2021.
- Lehner, F. and Deser, C.: Origin, importance, and predictive limits of internal climate variability, *Environmental Research: Climate*, 2, 023 001, <https://doi.org/10.1088/2752-5295/acf30>, 2023.
- 415 Leijnse, T., van Ormondt, M., Nederhoff, K., and van Dongeren, A.: Modeling compound flooding in coastal systems using a computationally efficient reduced-physics solver: Including fluvial, pluvial, tidal, wind- and wave-driven processes, *Coastal Engineering*, 163, <https://doi.org/10.1016/j.coastaleng.2020.103796>, 2021.
- Lim, W. H., Yamazaki, D., Koirala, S., Hirabayashi, Y., Kanae, S., Dadson, S. J., Hall, J. W., and Sun, F.: Long-Term Changes in Global Socioeconomic Benefits of Flood Defenses and Residual Risk Based on CMIP5 Climate Models, *Earth's Future*, 6, 938–954,  
420 <https://doi.org/https://doi.org/10.1002/2017EF000671>, 2018.
- Liu, M., Vecchi, G. A., Smith, J. A., and Knutson, T. R.: Causes of large projected increases in hurricane precipitation rates with global warming, *NPJ climate and atmospheric science*, 2, 38, 2019.
- Magnusson, L., Bidlot, J.-R., Bonavita, M., Brown, A. R., Browne, P. A., Chiara, G. D., Dahoui, M., Lang, S. T. K., McNally, T., Mogensen, K. S., Pappenberger, F., Prates, F., Rabier, F., Richardson, D. S., Vitart, F., and Malardel, S.: ECMWF Activities for Improved Hurricane  
425 Forecasts, *Bulletin of the American Meteorological Society*, 100, 445 – 458, <https://doi.org/10.1175/BAMS-D-18-0044.1>, 2019.
- McInerney, D., Lempert, R., and Keller, K.: What are robust strategies in the face of uncertain climate threshold responses? Robust climate strategies, *Climatic change*, 112, 547–568, 2012.
- Merz, B., Elmer, F., and Thielen, A.: Significance of " high probability/low damage" versus " low probability/high damage" flood events, *Natural Hazards and Earth System Sciences*, 9, 1033–1046, 2009.
- 430 Mester, B., Vogt, T., Bryant, S., Otto, C., Frieler, K., and Schewe, J.: Human displacements from Tropical Cyclone Idoi attributable to climate change, *Natural Hazards and Earth System Sciences*, 23, 3467–3485, <https://doi.org/10.5194/nhess-23-3467-2023>, 2023.
- Mogensen, K. S., Magnusson, L., and Bidlot, J.-R.: Tropical cyclone sensitivity to ocean coupling in the ECMWF coupled model, *Journal of Geophysical Research: Oceans*, 122, 4392–4412, <https://doi.org/https://doi.org/10.1002/2017JC012753>, 2017.
- Nederhoff, K., van Ormondt, M., Veeramony, J., van Dongeren, A., Antolínez, J. A. A., Leijnse, T., and Roelvink, D.: Accounting for uncertainties in forecasting tropical-cyclone-induced compound flooding, *Geoscientific Model Development*, 17, 1789–1811,  
435 <https://doi.org/10.5194/gmd-17-1789-2024>, 2024.

- Nordgren, J., Stults, M., and Meerow, S.: Supporting local climate change adaptation: Where we are and where we need to go, *Environmental Science & Policy*, 66, 344–352, <https://doi.org/https://doi.org/10.1016/j.envsci.2016.05.006>, 2016.
- Pant, R., Thacker, S., Hall, J., Alderson, D., and Barr, S.: Critical infrastructure impact assessment due to flood exposure, *Journal of Flood Risk Management*, 11, 22–33, <https://doi.org/https://doi.org/10.1111/jfr3.12288>, 2018.
- 440 Patricola, C. M. and Wehner, M. F.: Anthropogenic influences on major tropical cyclone events, *Nature*, 563, 339–346, <https://doi.org/10.1038/s41586-018-0673-2>, 2018.
- Qiu, J., Liu, B., Yang, F., Wang, X., and He, X.: Quantitative Stress Test of Compound Coastal-Fluvial Floods in China’s Pearl River Delta, *Earth’s Future*, 10, e2021EF002638, <https://doi.org/10.1029/2021EF002638>, e2021EF002638 2021EF002638, 2022.
- 445 Ranasinghe, R., Ruane, A. C., Vautard, R., Arnell, N., Coppola, E., Cruz, F. A., Dessai, S., Saiful Islam, A., Rahimi, M., Carrascal, D. R., et al.: Climate change information for regional impact and for risk assessment, 2021.
- Rentschler, J., Salhab, M., and Jafino, B. A.: Flood exposure and poverty in 188 countries, *Nature communications*, 13, 3527, 2022.
- Rodrigues, R. R. and Shepherd, T. G.: Small is beautiful: climate-change science as if people mattered, *PNAS Nexus*, 1, pgac009, <https://doi.org/10.1093/pnasnexus/pgac009>, 2022.
- 450 Sebastian, A., Bader, D., Nederhoff, C., Leijnse, T., Bricker, J., and Aarninkhof, S.: Hindcast of pluvial, fluvial, and coastal flood damage in Houston, Texas during Hurricane Harvey (2017) using SFINCS, *Natural hazards*, 109, 2343–2362, 2021.
- Segerlin, S., Cannizzaro, N., Gregorio, C., Bokpin, G. A., Ariyan, L., Ngece, N., Bookstaber, M., Rosen, D., Wang, H., Chen, G., Gusah, S., Gyamfi-Yeboah, F., Teye, J. K., Duman, D., Tsige, T., and Gichor, M.: Financing for Resilient and Green Urban Solutions in Beira, Mozambique, United Nations Human Settlements Programme (UN-Habitat), chief Editors and Managers: Xing Quan Zhang, Irina Eichenauer and Wolfgang Ryll, 2020.
- 455 Shepherd, T. G.: Storyline approach to the construction of regional climate change information, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 475, <https://doi.org/10.1098/rspa.2019.0013>, 2019.
- Shepherd, T. G. and Lloyd, E. A.: Meaningful climate science, *Climatic Change*, 169, 1–16, <https://doi.org/10.1007/s10584-021-03246-2>, 2021.
- 460 Shepherd, T. G. and Sobel, A. H.: Localness in Climate Change, *Comparative Studies of South Asia, Africa and the Middle East*, 40, 7–16, <https://doi.org/10.1215/1089201X-8185983>, 2020.
- Shepherd, T. G., Boyd, E., Calel, R. A., Chapman, S. C., Dessai, S., Dima-West, I. M., Fowler, H. J., James, R., Maraun, D., Martius, O., Senior, C. A., Sobel, A. H., Stainforth, D. A., Tett, S. F. B., Trenberth, K. E., van den Hurk, B. J. J. M., Watkins, N. W., Wilby, R. L., and Zenghelis, D. A.: Storylines: an alternative approach to representing uncertainty in physical aspects of climate change, *Climatic Change*, 151, 555–571, <https://doi.org/10.1007/s10584-018-2317-9>, 2018.
- 465 Sillmann, J., Shepherd, T. G., van den Hurk, B., Hazeleger, W., Martius, O., Slingo, J., and Zscheischler, J.: Event-based storylines to address climate risk, *Earth’s Future*, 9, <https://doi.org/10.1029/2020EF001783>, 2020.
- Slager, K., Burzel, A., Bos, E., de Bruijn, K., D., W., Winsemius, H., Bouwer, L., and van der Doef, M.: User Manual Delft-FIAT, Deltares, <https://publicwiki.deltares.nl/display/DFIAT/Delft-FIAT+Home>, last accessed: 22 September 2019, 2016.
- 470 Sobel, A. H.: Usable climate science is adaptation science, *Climatic Change*, 166, 8, 2021.
- Sutton, R. T.: Climate Science Needs to Take Risk Assessment Much More Seriously, *Bulletin of the American Meteorological Society*, 100, 1637 – 1642, <https://doi.org/https://doi.org/10.1175/BAMS-D-18-0280.1>, 2019.

- 475 Tanoue, M., Taguchi, R., Nakata, S., Watanabe, S., Fujimori, S., and Hirabayashi, Y.: Estimation of Direct and Indirect Economic Losses Caused by a Flood With Long-Lasting Inundation: Application to the 2011 Thailand Flood, *Water Resources Research*, 56, e2019WR026092, <https://doi.org/https://doi.org/10.1029/2019WR026092>, e2019WR026092 2019WR026092, 2020.
- Tanoue, M., Taguchi, R., Alifu, H., and Hirabayashi, Y.: Residual flood damage under intensive adaptation, *Nature Climate Change*, 11, 823–826, 2021.
- UN Office for the Coordination of Humanitarian Affairs: Mozambique: Cyclone Idai & Floods Situation Report No. 2 (as of 3 April 2019), <https://reliefweb.int/report/mozambique/mozambique-cyclone-idai-floods-situation-report-no-2-3-april-2019>, accessed: 27-02-2024, 2019.
- 480 van Berchum, E. C., van Ledden, M., Timmermans, J. S., Kwakkel, J. H., and Jonkman, S. N.: Rapid flood risk screening model for compound flood events in Beira, Mozambique, *Natural Hazards and Earth System Sciences*, 20, 2633–2646, <https://doi.org/10.5194/nhess-20-2633-2020>, 2020.
- van den Hurk, B. J., Pacchetti, M. B., Boere, E., Ciullo, A., Coulter, L., Dessai, S., Ercin, E., Goulart, H. M., Hamed, R., Hochrainer-Stigler, S., Koks, E., Kubiczek, P., Levermann, A., Mechler, R., van Meersbergen, M., Mester, B., Middelani, R., Minderhoud, K., Mysiak, J., Nirandjan, S., van den Oord, G., Otto, C., Sayers, P., Schewe, J., Shepherd, T. G., Sillmann, J., Stuparu, D., Vogt, T., and Witpas, K.: Climate impact storylines for assessing socio-economic responses to remote events, *Climate Risk Management*, 40, 100500, <https://doi.org/10.1016/j.crm.2023.100500>, 2023a.
- 485 van den Hurk, B. J., White, C. J., Ramos, A. M., Ward, P. J., Martius, O., Olbert, I., Roscoe, K., Goulart, H. M., and Zscheischler, J.: Consideration of compound drivers and impacts in the disaster risk reduction cycle, *Iscience*, 26, 2023b.
- 490 van der Pol, T. D., van Ierland, E. C., and Gabbert, S.: Economic analysis of adaptive strategies for flood risk management under climate change, *Mitigation and adaptation strategies for global change*, 22, 267–285, 2017.
- Van Logchem, B. and Queface, A.: Responding to Climate Change in Mozambique: Synthesis Report, Synthesis report, Maputo INGC, 2012.
- Ward, P. J., Jongman, B., Aerts, J. C., Bates, P. D., Botzen, W. J., Diaz Loaiza, A., Hallegatte, S., Kind, J. M., Kwadijk, J., Scussolini, P., 495 et al.: A global framework for future costs and benefits of river-flood protection in urban areas, *Nature climate change*, 7, 642–646, 2017.
- Wiel, K. V. D., Selten, F. M., Bintanja, R., Blackport, R., and Screen, J. A.: Ensemble climate-impact modelling: extreme impacts from moderate meteorological conditions, *Environmental Research Letters*, 15, <https://doi.org/10.1088/1748-9326/ab7668>, 2020.
- Winsemius, H. C., Aerts, J. C., Van Beek, L. P., Bierkens, M. F., Bouwman, A., Jongman, B., Kwadijk, J. C., Ligtvoet, W., Lucas, P. L., Van Vuuren, D. P., et al.: Global drivers of future river flood risk, *Nature Climate Change*, 6, 381–385, 2016.
- 500 Woodruff, J. D., Irish, J. L., and Camargo, S. J.: Coastal flooding by tropical cyclones and sea-level rise, *Nature*, 504, 44–52, 2013.
- Zhang, Z., Wang, W., Doyle, J. D., Moskaitis, J., Komaromi, W. A., Heming, J., Magnusson, L., Cangialosi, J. P., Cowan, L., Brennan, M., Ma, S., Das, A. K., Takuya, H., Clegg, P., Birchard, T., Knaff, J. A., Kaplan, J., Mohapatra, M., Sharma, M., Masaaki, I., Wu, L., and Blake, E.: A review of recent advances (2018–2021) on tropical cyclone intensity change from operational perspectives, part 1: Dynamical model guidance, *Tropical Cyclone Research and Review*, 12, 30–49, <https://doi.org/https://doi.org/10.1016/j.tcr.2023.05.004>, 2023.