
Trends of hydroclimatic intensity in Colombia

Oscar Mesa^{1,*}, Viviana Urrea¹ and Andrés Ochoa¹

¹*Departamento de Geociencias y Medio Ambiente, Universidad Nacional de Colombia, Medellín, Colombia*

Correspondence*:

Oscar Mesa, Departamento de Geociencias y Medio Ambiente, Universidad Nacional de Colombia, Carrera 80 # 64-223, 050041 Medellín, Colombia
ojmesa@unal.edu.co

2 ABSTRACT

Prediction of changes in precipitation in upcoming years and decades caused by global climate change associated with the greenhouse effect, deforestation and other anthropic perturbations is a practical and scientific problem of high complexity and huge consequences. To advance toward this challenge we look at the daily historical record of all available rain gauges in Colombia to estimate an index of the intensity of the hydrologic cycle (Giorgi et al., 2011). The index is the product of precipitation intensity and dry spell length. Theoretical reasons indicate that global warming should lead to increasing trends in either one of the factors or both. Our results indicate that there is no clear picture, there are gauges with positive and negative significant trends, and most of the gauges do not show a significant trend. We present the geographic distribution of results within regions and concerning the elevation in the Andes Cordillera. Results seem to agree with previous reports of total annual precipitation trends.

Keywords: Precipitation trends, Climate change, Colombia, Hydrologic cycle, dry spell length

1 INTRODUCTION

Predicting the effect of climate change on Colombia's hydrology, more specifically on precipitation, is not a small matter. To illustrate, only in the electricity sector recent studies for the Mining and Energy Planning Unit Macías and Andrade (2014) estimate that the impacts of the decrease in precipitation imply an increase in annual investment of US\$ 290 million per year for the period 2013-2015.

But not only the impact is of magnitude, but the scientific problem of such prediction is also very complex, such models are not necessarily accurate. The focus of this work is the impact on precipitation, although global climate change impacts many more aspects such as temperature, sea level, coastal erosion, páramo ecosystem loss, vector-borne diseases, biodiversity, agriculture, and others.

Understanding the temporal-spatial variability of precipitation is a challenge of great importance due to the environmental, social, economic and cultural implications of the distribution of water resources in any country. On the other hand, the analysis of hydrologic processes under a climate change context, in addition to incorporating greater complexity to these processes, leads to the need to join efforts towards the understanding of national hydrology to manage the resource better. The broad objective of this work is to contribute to that challenging question, focusing on the use of precipitation information and mathematical and statistical techniques to identify and interpret evidence of climate change in Colombia.

30 Given the complexity of rainfall fields and the tropical climate in the rough topography derived by the
31 Andes cordillera, rainfall records in Colombia are generally scarce, both because of their quality, missing
32 data, length of the records and spatial coverage. Therefore, it is a unique challenge for Hydrology to predict
33 the impact of climate change over Colombia rainfall at spatial and temporal scales suitable for critical
34 applications. Among those applications, one can mention the planning for the sustainable development of
35 the territory and its hydraulic resources, disaster prevention. Some of the theoretical questions are a better
36 knowledge of the influence of macro-climatic phenomena like El Niño-Southern Oscillation (ENSO). In
37 turn, a better understanding may allow better predictions.

2 PREVIOUS WORK

38 This short review has two parts. First, we present the main results of previous studies about climate change
39 impact on Colombian rainfall trends. Then we briefly show the general context of how global warming
40 impact precipitation.

41 Various works describe the climatology of the precipitation in Colombia (Snow, 1976; Oster, 1979;
42 Eslava, 1993; Mesa et al., 1997; Mejía et al., 1999). The central control is the passage, twice a year, of the
43 Inter-Tropical Convergence Zone that marks the rainy seasons of April-May and September-November in
44 the Andes, and the seasons with the lowest rainfall in December-February and June. -August. The spatial
45 distribution is marked by the sources of humidity in the Caribbean, the Pacific, and the Amazon, by the
46 topography and prevailing winds. The inter-annual variability is controlled mainly by the phenomenon of
47 ENSO in the tropical Pacific (Poveda and Mesa, 1997; Poveda et al., 2011).

48 Several studies have found evidence of climate change in Colombia using various statistical techniques
49 with different record lengths (e.g. Smith et al., 1996; Mesa et al., 1997; Quintana-Gomez, 1999; Vuille
50 et al., 2003; Ochoa and Poveda, 2008; Pabón, 2009; Cantor and Ochoa, 2011; Cantor, 2011; Carmona and
51 Poveda, 2014; Hurtado and Mesa, 2015). In summary, these studies identify increasing trends in mean
52 and minimum temperature records in a significant number of stations. Besides, they find mixed trends in
53 precipitation, with a similar percentage of stations for each trend and 20% without a statistically significant
54 trend for the set of considered series of up to 40 years of records. For precipitation stations with longer
55 records, the majority (63%) shows an increasing trend and only 16% decreasing trend. A clear geographical
56 pattern is not identified to locate areas with a particular trend, except in the Pacific plain that has the highest
57 definite upward trend, explained by an increasing trend of the influence of moisture in the Pacific and the
58 Chocó Jet.

59 These conclusions coincide with the IDEAM report, Mayorga et al. (2011), which analyzed 310 rainfall
60 stations with monthly records in the period 1970-2010 using RCLIMDEX. This is a program developed
61 by the CCI / CLIVAR working group (Climate and Ocean: Variability, Predictability and Change) part
62 of the World Climate Research Program of the World Meteorological Organization and Unesco for the
63 detection of climate change as a coordinated international effort to have indexes calculated with the same
64 methodology that can be integrated internationally (Peterson, 2005). Of the 310 precipitation stations, 71 %
65 show increasing trend, seven % without trend, 22 % decreasing trend. The conclusion related to the rising
66 temperature is also confirmed, with an increase of 0.17 K per decade. It is essential to quote the report:
67 “... the generalized increase in precipitation in the country is noticeable, highlighting the north-western
68 zone (Antioquia and Chocó), Vichada, the Piemonte de Putumayo and the island of Providencia, where
69 the most significant increase. The opposite occurs on the island of San Andrés, on the eastern slope of the

70 Eastern Cordillera (Arauca and Casanare) and in large areas of the Upper Cauca, where there is a decrease
71 in precipitation. The largest decrease occurs in the south-west of the territory. ”

72 It is clear that the observed trends may be due to other causes besides increasing greenhouse gas global
73 warming; deforestation and urbanization among others, not to mention observational issues.

74 Concerning the impact of deforestation, [Salazar \(2011\)](#) estimates through a numerical experiment that a
75 possible drastic future change in coverage in the Amazon area would bring about a reduction in precipitation
76 in Colombia of an order of magnitude of 300 mm/year.

77 The warming of the Colombian Andes has led to the complete extinction of eight tropical glaciers, and
78 the six remaining snow-caps lose ice at accelerated rates ([Rabatel et al., 2013](#)). The páramos, unique and
79 strategic ecosystems to supply water to several cities, including Bogotá and Medellín are also in danger by
80 warming and other anthropogenic activities ([Ruiz et al., 2008](#)).

81 The heating means an increase of the air saturation vapor pressure according to the Clausius-Clapeyron
82 equation. Overall, this increase is has been accompanied by an increase in evaporation and, therefore, in
83 absolute humidity, to keep the relative humidity relatively constant ([Stevens et al., 2013](#)). A few series of
84 tank evaporation and relative humidity confirm this observation for Colombia ([Mesa et al., 1997](#)). The
85 consequences of this are essential to understanding the physical mechanisms associated with the impact of
86 warming on precipitation and river flow.

87 [Mesa et al. \(1997\)](#) reports that a good part of the flow series in the Magdalena-Cauca basins have a
88 decreasing trend. Recently, [Carmona and Poveda \(2014\)](#) confirm and reinforce this conclusion about the
89 trends from their analysis of the same series with new years of observations and other series not analyzed
90 in the initial work. The number of stations with a decreasing trend is higher than the number showing an
91 increasing trend. The percentage that decreases varies from 61% to 100% depending on the length of the
92 record, the first value for records of 25 years and the last for records over 50 years. Whereas, 0 to 34% of
93 the analyzed streamflow gauges show an increase. The positive regional trend for the Atrato and San Juan
94 flows coincide with areas of large increasing trends in precipitation.

95 [Hurtado and Mesa \(2015\)](#) developed a reanalysis of the precipitation field in Colombia, comprising 384
96 fields of monthly precipitation in the period 1975-2006 at a spatial resolution of 5 minutes of arc. For
97 the reanalysis, they used records of 2270 rain gauges and various satellite-derived products for the most
98 recent period. Then using Empirical Orthogonal Functions, Principal Component Analysis and statistical
99 tests they looked for changes or trends. According to their results, both the Mann - Whitney mean change
100 test and the simple t trend test indicate increasing precipitation trends mainly in the Pacific, Orinoco, and
101 Amazon basin regions. In most of the Andean region, there are no changes or trends.

102 Another line of work has been the use of global climate models (GCM): [Pabón \(2005\)](#); [Ruiz \(2007, 2010\)](#);
103 [IDEAM-Colombia \(2010\)](#). These models adequately represent the core of the physical processes, although
104 not the totality, particularly rainfall, tropical convection, and topography. Besides, spatial resolution is not
105 generally adequate for regional studies. In general, there is the possibility of correcting these deficiencies
106 through the use of mesoscale models that have finer spatial resolution and better represent convective
107 processes. This methodology, known as downscaling, has advanced, but it is not free of difficulties.

108 [Ruiz \(2010\)](#) and [Pabón \(2005\)](#) analyze the results of the low-resolution global models to conclude that
109 “annual precipitation would be reduced in some regions and would increase in others. In the regions where
110 there is a certain degree of coincidence in most of the models and current trends about an increase are: the
111 North and Central Pacific, Middle Magdalena, Sabana de Bogotá, Sogamoso River Basins, Catatumbo,

112 Arauca, Piedemonte Llanero, Central Orinoquia, Central Amazon and Amazonian piedmont. In those
113 places the increase in rainfall compared with the typical period 1961-1990 could be between 10 and 15
114 % by 2050 and between 15 and 25 % for the year 2080. For the other regions, the study says that it is
115 difficult to arrive at a concrete result since the different models present contradictory results”. The results
116 of the different IPCC models are available in <http://www.climatewizard.org/>. From there one can conclude
117 that predicted change in rainfall for Colombia in scenario A1B for the average of the models indicate an
118 increase of around ten % for all of Colombia, except the northernmost zone. If the models or scenarios are
119 examined individually the general trend is the same, although the magnitudes vary. In general, for most of
120 Colombia, they predict positive trends, except for the most northern part where they predict a decreasing
121 trend.

122 [Pabón \(2005\)](#); [Ruiz \(2007, 2010\)](#); [IDEAM-Colombia \(2010\)](#) went beyond low-resolution models into
123 higher resolution using the regional model PRECIS (Providing Regional Climates for Impacts Studies)
124 of the United Kingdom and the global model GSM-MRI (Global Spectral Model) of Japan. The results
125 obtained can be summarized by quoting the Colombian 2nd Communication: “The areas that on average,
126 by the end of the 21st century, would have the greatest reductions in precipitation would be the departments
127 of Huila, Putumayo, Nariño, Cauca, Tolima, Córdoba, Bolivar and Risaralda where the rains would be
128 reduced close to 15% with respect to the 1971-2000 climatology. Likewise, it is possible that rainfall
129 increases near 10%, in large areas of Chocó. From the point of view of the most pessimistic scenarios
130 (A2, for example), the most significant reductions in rainfall throughout the 21st century would occur in
131 Córdoba, Cauca, Bolívar, Caldas, Sucre, Valle, Antioquia, Nariño and Risaralda , where it would rain
132 between 70 and 80% of the rainfall recorded during the period 1971-2000 (that is, reductions between 20%
133 and 30%).”

134 However, some contradictions are striking. The minimum consistency required is that during the historical
135 period the results of the models correspond to the observations. The trends in precipitation deduced from
136 low-resolution global models do seem to be consistent with the trends observed in historical records.
137 However, there is no consistency between the predictions of high-resolution models and low-resolution
138 models. This single observation disqualifies their results of the high-resolution models.

139 Supported by these IDEAM results, UPME has analyzed the vulnerability of the Colombian electricity
140 sector, which on average supplies 70% through hydro-electricity ([ACON-OPTIM, 2013](#)). The monthly
141 flows that feed each reservoir were projected using the decreasing trends of precipitation. As expected,
142 they found a generalized decrease for the three climate change scenarios analyzed (A2, B2, and A1B).
143 It is worth mentioning the analysis they make of these results: “Although the results obtained coincide
144 with reductions in the water supply of the reservoirs, in many cases the percentage changes in the monthly
145 flows are exaggerated, reaching reductions greater than 50% of the current condition. Those values are
146 considered unlikely considering that the recorded flow series do not show sufficient statistical evidence to
147 demonstrate that there is any linear decreasing trend”. Such studies give rise to policies and actions that
148 have clear, practical consequences.

149 [Urán \(2015\)](#) carried out an analysis of the scaling between precipitation and temperature limited by the
150 Clausius-Clapeyron using 86 stations of precipitation and 9 temperature stations over the Antioquia region
151 of Colombia, with 15 minutes resolution. He also used rain derived from TRMM data (Tropical Rainfall
152 Measure Mission) with rainfall intensities every 3 hours. He found that for temporal scales greater than 12
153 hours the trends are no longer significant trends. For the finer temporal scales trends become significant
154 for extreme deciles of the distribution. He reports a close scaling due to the Clausius-Clapeyron relation
155 limiting the intensification of precipitation following the ideas of [O’gorman and Schneider \(2009\)](#).

156 In response to global warming, the hydrological cycle also changes. A warmer atmosphere means more
157 radiative cooling of the troposphere, which is a growing function of temperature. The highest infra-red
158 radiation emission corresponds to the balance required to compensate for the greater radiation absorbed.
159 To the extent that changes in cloudiness or the absorption of radiation by water vapor offset the necessary
160 radiative cooling, changes in precipitation may occur. Regionally, the winds determine where there is an
161 increase or a decrease. If the winds change little, compared to the humidity they transport, the wet regions
162 import more water, and there could be more rain, while the dry ones could be drier (Mitchell et al., 1987;
163 Soden and Held, 2006; Wentz et al., 2007).

164 Giorgi et al. (2011) introduce a new measure of hydroclimatic intensity (HY-INT), which integrates
165 metrics of precipitation intensity and dry spell length. The responses of these two metrics to global warming
166 are deeply interconnected. They found clear increasing trends of HY-INT in global and regional climate
167 models. Depending on the region, the increase in HY-INT is due to an increase in precipitation intensity,
168 dry spell length, or both. They also examined late twentieth-century observations and concluded that
169 they also exhibit dominant positive HY-INT trends, providing a hydroclimatic signature of late-twentieth-
170 century warming. Precipitation intensity increases because of increased atmospheric water holding capacity.
171 However, increases in mean precipitation need increases in surface evaporation rates, which are lower than
172 for atmospheric moisture. Global warming increase potential evapotranspiration which may probably result
173 in an increase in actual evaporation, or evapotranspiration in plants, only if adequate moisture is available.
174 So potentially there is more drying, but in drought situations part of any extra energy goes into raising
175 temperatures, thereby amplifying warming over dry land. This feedback leads to a reduction in the number
176 of wet days and an increase in dry spell length.

3 STUDY AREA AND DATA

177 We analyzed data from 1706 sites in the whole territory of Colombia, 1062 in the Andes region. The
178 other sites are in the Amazon (77), the Caribbean 398, the Orinoco (91) and the Pacific (78) regions. Data
179 comprise daily time series of rainfall amount. Since the method requires no missing data (Section 4) we
180 trim the series to the common period between 1970 and 2014. We also fill the missing data with data from
181 the CHIRPS (Funk et al., 2015) data-set. A condition for this filling was that the percentage of missing
182 days did not exceed 30%. Otherwise, the whole year is missing. The authors tested the performance of
183 CHIRPS in Colombia in previous studies (Urrea et al., 2019; Urrea, 2017; Urrea et al., 2016). Of the
184 original IDEAM dataset, we dropped for our analysis all the stations with more than 50% missing data
185 in the common period and others that do not pass a minimum quality check. The final data-set has 1629
186 stations.

187 Figure 1 shows the IDEAM the rain gauge network. The network covers a range of elevations from sea
188 level in the Caribbean and Pacific coasts to 4150 meters above sea level in the Andes. Notice also the low
189 density of the gauge network in the Amazon and Orinoco regions.

190 Colombian climate is tropical, mean annual temperature is high, above 25°C at sea level, the diurnal
191 range of temperature exceeds the annual range, and the annual range is minimal, less than 5°C (Snow,
192 1976). Precipitation is abundant in comparison with any other place in the world. There are places of the
193 Pacific coast with perennial rain with mean annual totaling 12200 mm. Over the Orinoco and Amazon
194 basins in Colombia, the mean annual precipitation varies from 2000 to 7000 mm per year. In the Caribbean
195 coast the average is of the order of 1500 mm, but to the north, there are places with near 300 mm/year. In
196 the Andean region mean annual precipitation ranges from 1000 to 3000 mm/year.




Figure 1. Location of the IDEAM rain gauge network that was used in this work. The bottom left graph shows the vertical distribution of the rain stations. The map also shows the five natural regions of Colombia (IGAC, 1997).

197 Atmospheric moisture is transported toward Colombia by the trade winds from the Caribbean sea, the
 198 Atlantic ocean through both Amazon and Orinoco basins that themselves contribute with recirculating
 199 moisture. Also from the Pacific ocean, westerly winds contribute to the massive convergence of moisture
 200 over Colombia. The migration of the inter-tropical convergence zone and three low-level jet streams (Chocó,
 201 Caribbean, and South America) are part of the complex circulation given rise to that high precipitation
 202 (Poveda et al., 2014).

4 METHODS

203 Hydroclimatic intensity (Giorgi et al., 2011) is evaluated using the HY-INT indicator

$$\text{HY-INT} = \text{INT} \times \text{DSL}, \quad (1)$$

204 where INT and DSL are mean intensity during wet days and mean dry spell duration for each year in the
 205 record. In both cases, one works with scaled variables using the respective inter-annual mean as scale factor
 206 (Giorgi et al., 2011). Therefore the long-term average of both INT and DSL are 1.

207 We also evaluate the trends of P, the total annual precipitation for each year; and LW, the average length
 208 of a wet spell in each year. We also counted the number of wet and dry days in each year as well as the
 209 number of dry and wet spells in each year.

210 Also, we constructed an extreme indicator of the hydrologic cycle generalizing (Giorgi et al., 2011) ideas.
 211 For that, we computed the maximum daily intensity for each year (INTX) and the maximum dry spell
 212 length for each year (DSLX). In other words, for this proposal instead for the average of the corresponding

213 variable for each year, we take the maximum. Their product gives the HY-INTX indicator of the strength of
 214 the hydrologic cycle.

215 For each year in the record we computed each one of the variables mentioned above. Therefore for each
 216 gauge and each variable, we have a time series. We then proceed to evaluate the existence of trends in those
 217 time series for each gauge.

218 4.1 Trend analysis

219 We use the Mann-Kendall test (Mann, 1945; Kendall, 1955) for autocorrelated data (Hamed and
 220 Ramachandra-Rao, 1998) to evaluate the existence of trends in the time series, and the Sen's slope
 221 estimator (Sen, 1968) for calculating the magnitude of the trend. A summary of these techniques follows.
 222 More details in the cited references.

223 The null hypothesis of the Mann-Kendall test is that the data come from independent and identically
 224 distributed random variables (iid) and hence no long-term trend exists. When the data are iid, the statistic
 225 S ,

$$S = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \text{sgn}(x_j - x_i), \quad (2)$$

226 has asymptotic normality with mean zero and variance $\text{Var}[S]$

$$\text{Var}[S] = \frac{n(n-1)(2n+5)}{18} - \frac{1}{18} \sum_{j=1}^m t_j(t_j-1)(2t_j+5). \quad (3)$$

227 In Eq. 2 n is the sample size, x_t is the value of the time series at time t , and $\text{sgn}(x_j - x_i)$ is defined by

$$\text{sgn}(x_j - x_i) = \begin{cases} 1 & \text{if } x_j - x_i > 0, \\ 0 & \text{if } x_j - x_i = 0, \\ -1 & \text{if } x_j - x_i < 0. \end{cases} \quad (4)$$

228 The sum in the last term of Eq. 3 accounts for the reduction in variance due to the existence of tied ranks
 229 (Hamed, 2008). In Eq. 3 m is the number of groups of tied ranks and t_j is the number of ranks in group j .

230 The standardized test statistic Z is calculated by

$$Z = \begin{cases} \frac{S-1}{\sqrt{\text{Var}[S]}} & \text{if } S > 0, \\ 0 & \text{if } S = 0, \\ \frac{S+1}{\sqrt{\text{Var}[S]}} & \text{if } S < 0. \end{cases} \quad (5)$$

231 The null hypothesis of no trend is rejected if $|Z|$ exceeds the value $|Z_{1-\alpha/2}|$ of the standard normal
 232 distribution for a given significance level α .

233 The result of Mann-Kendall test is sensitive to autocorrelation in the data. Positive autocorrelation
 234 increases the probability of rejecting the null hypothesis when it is true, and negative autocorrelation
 235 increases the probability of not rejecting the null hypothesis when it is false. This effect occurs because of
 236 a bias in the estimation of $\text{Var}[S]$. Hamed and Ramachandra-Rao (1998) suggested the empirical formula

237 in Eq. 6 for calculating $\text{Var}[S]$ in the presence of autocorrelation.

$$\text{Var}[S_{ac}] = \text{Var}[S] \times \left[1 + \frac{2}{n(n-1)(n-2)} \sum_{i=1}^{n-1} (n-i)(n-i-1)(n-i-2)\rho_s(i) \right] \quad (6)$$

238 where $\rho_s(i)$ is the auto-correlation function of the ranks of the observations.

239 The Sen's non-parametric method (Sen, 1968) estimates the long-term linear trend slope of a time series
240 as the median value of the slopes between all pairs of points in the series. For $N = n \cdot (n-1)/2$ pairs of
241 data in the series, the N slopes, Q , are calculated as shown in Eq. 7. The median of Q is the Sen's slope
242 estimator.

$$Q_i = \frac{x_j - x_k}{j - k}, \quad i = 1, 2, \dots, N; \quad 1 \leq j \leq n-1; \quad j \leq k \leq n. \quad (7)$$

243 4.2 The HY-INT trend

244 Even though HY-INT is not a linear function of INT and DSL, the long-term trend slope of HY-INT is
245 a function of the trend slopes of INT and DSL. Equivalently, one can estimate it from the time series of
246 HY-INT. Taking the time derivative of Eq. 1 one gets

$$\frac{d\text{HY-INT}}{dt} = \text{INT} \frac{d\text{DSL}}{dt} + \text{DSL} \frac{d\text{INT}}{dt}. \quad (8)$$

247 And because all the variables are scaled, what one needs is the logarithmic derivative

$$\frac{1}{\text{HY-INT}} \frac{d\text{HY-INT}}{dt} = \frac{1}{\text{DSL}} \frac{d\text{DSL}}{dt} + \frac{1}{\text{INT}} \frac{d\text{INT}}{dt}. \quad (9)$$

248 Therefore the temporal trend slopes satisfy

$$m_{\text{HY-INT}} = m_{\text{INT}} + m_{\text{DSL}}. \quad (10)$$

5 RESULTS

249 Neglecting data autocorrelation in trend analysis increases the probability of error in the Mann-Kendall test
250 result (Kulkarni and von Storch, 1992; von Storch, 1995). We compared the results of the classic MK test
251 and the MK test for autocorrelated data proposed by Hamed and Ramachandra-Rao (1998) in our 1629
252 series data set, and conclude that ignoring the auto-correlation may lead to false trends of the order of 20%
253 of the gauges, and false no trends in of the order of 10% of the gauges.

254 Table 1 presents in each row the four elements of the confusion matrix for the Mann-Kendall test that does
255 not take into account auto-correlation in comparison with the one that does. We take this last one as the
256 correct method for the comparison. The most significant error (from 18 to 25%) comes from false trends.
257 However, there are also errors due to false no trends (from 11 to 15%). As a result, the accuracy (total
258 number of hits) is between 60 and 71%. It is always recommended to take into account the auto-correlation
259 of the series to evaluate the significance of trends.

260 We also considered the possible implication of the definition of the year. Besides the calendar year, we
261 considered a hydrology year starting on April 1st. The idea was that the end of the calendar year might

Table 1. Confusion matrices for the evaluation method of the significance of the trends for each of the indicated variables without taking into account auto-correlation. For the purpose of the illustration the Mann-Kendall test with auto-correlation is considered the true one.

Variable	true trend R-R	false trend R-NR	false no trend NR-R	true no trend NR-NR
Number of gauges (Percentage of the total number of gauges)				
P	45 (3%)	293 (18%)	180 (11%)	1111 (68%)
INT	116 (7%)	400 (25%)	245 (15%)	868 (53%)
DSL	62 (4%)	316 (19%)	184 (11%)	1067 (66%)
HY-INT	108 (7%)	393 (24%)	212 (13%)	916 (56%)




Figure 2. Two examples of trend analysis for (top to bottom) P, DSL, INT and HY-INT for two representative stations. Left: Susacón in Boyacá, at 2550 masl. Right: La Línea El Porvenir in Risaralda, at 1955 masl.

split the longest dry spell. Because the dry season usually starts in mid-December and ends in March in Colombia. However, as we show below, the dominant fact for the significant trends is an increase in the number of wet days (70%), not the length of the dry season.

Figure 2 illustrates two of the 1629 cases of the trend analysis. Notice the treatment of the missing years that may come for any missing day. For the Susacón gauge in the left part of the Figure, the trends in P, INT, and HY-INT are decreasing and statistically significant. However, the trend in DSL is not. For La Línea El Porvenir station in the right side of the Figure DSL, INT, and HY-INT have significant positive trends, whereas P has an increasing non statistically significant trend.




Figure 3. Histograms of the HYINT (left), INT (center) and DSL (right) trend slopes of the 1629 rain gauges used. Non significant trends (H₀: Stationary hypothesis not rejected, NR) in grey and significant trends in black (H₀: Stationary hypothesis rejected, R).




Figure 4. Same as Figure 3 for the trend slope of the total annual precipitation (left), the number of wet days in the year(center) and number of wet spells in the year(right).

270 Table 2 summarizes the results of the trend analysis for the 1629 stations and the more relevant variables.
 271 The first observation is that only a minority of the stations show significant trends. Among the variables,
 272 INT shows the largest percentage of significant trends, but only reaching a 22.2% of the stations. The least
 273 percentage is for the variable DSLX with only 9.3%. A second observation is that the extreme variables do
 274 not show a larger percentage of significant trends in comparison with the corresponding average variables:
 275 For INTX the percentage is 9.3 in comparison with 22.2 for INT. Similarly, for DSLX is 9.3 in comparison
 276 with 15.1 for DSL; and for HYINTX is 9.4, whereas for HY-INT it is 19.6. Among the stations with
 277 statistically significant trends, the analysis of positive and negative trends is interesting. There is a clear
 278 majority among the significant ones for positive trends for P, the number of wet days, LW, Number of both
 279 dry and wet runs, INTX, and INT. Similarly, there is a majority of a significant negative trend for DSLX,
 280 HYINTX, number of dry days, DSL, and HY-INT.

281 Only 40% of the significant HY-INT stations have a positive trend, even though one of its factors, INT,
 282 has a majority of positive trends (57%) among the significant ones. However, that percentage reduces to
 283 43% among the ones with significant HY-INT trend. The other reason is the sign of the trends in DSL,
 284 69.1% negative among significant DSL trends. This percentage only is reduced to 61% when one considers
 285 stations with significant HY-INT trend. Besides, the histograms of the slopes of the trends in Figure 3
 286 illustrate well this point. Therefore for Colombia's humid climate HY-INT, the indicator proposed by
 287 Giorgi et al. (2011) to measure the strength of the hydrologic cycle, only makes sense partially concerning
 288 INT, but not for dry spells. Because, they are not increasing in length significantly, not even among the
 289 small number of stations with significant trends that on the contrary are getting shorter for the average dry
 290 run and more so for the extreme ones.




Figure 5. Left: Dispersion diagram of the DSL trend slope vs. INT trend slope for all stations with significant HY-INT trend slope. Notice that because of Eq 10 the trend slope of HY-INT is the sum of the trend slopes of INT and DSL. This equation explains the slanted iso-lines for the HY-INT trend slope. Right: Dispersion diagram of the HY-INT trend slope vs. precipitation trend slope for all stations with significant HY-INT trend slope




Figure 6. Precipitation trend slope vs. mean annual precipitation (A) and HY-INT trend slope vs. mean annual precipitation (B) for the stations with significant trend. n_+ and n_- are number of stations with positive (negative) slope. S_+ and S_- are the corresponding slopes.

291 Figure 4 shows the histograms of the trend slopes for P, the number of wet days and the number of wet
 292 spells. Again, the majority of the stations do not have significant trends. However, among the significant
 293 ones, there is a clear majority of positive trends.

294 Figure 5 allows further analysis of the result about HY-INT discussed above. Notice that almost all
 295 stations with positive trend slope for HY-INT have positive trend slopes for both INT and DSL. Conversely,
 296 almost all with negative trends for HY-INT have negative trends for INT and DSL. Therefore, most of the
 297 points in the figure are in either the first or the third quadrant (89%). Of the 320 stations with a significant
 298 trend in HT-INT, 124 and 170 do not have significant trends for INT and DSL respectively.

299 Figure 5 also shows the relation between the trend slopes for P and HY-INT. There are two groups of
 300 stations also, depending on the sign of the HY-INT trend. However, among both, there are positive and
 301 negative trends in P.

302 For the significant trends, the magnitude of the trends seems to be related linearly with P, see Figures 6A
 303 and 6B. This tendency is more clear for the P trend than for the HY-INT trend. The magnitude of the P

Table 2. Basic statistical analysis of the significant trends of the different variables using the calendar year and taking into account autocorrelation: P: Total Annual Precipitation, INT: averaged scaled intensity on wet days, DSL: averaged scaled dry run length; HYINT=INT×DSL; N stands for number; LW: averaged scaled length of wet runs. In addition extreme variables are computed for the hydrologic year (April 1st to March 31): INTX: the maximum daily intensity; DSLX: the maximum dry run length; HY-INTX=INTX×DSLX; and LWX: The maximum wet run length. Column symbols: N Rej is the number of stations for which the null hypothesis of no trend was rejected; % Rej/T is the percentage of total stations that rejected the test; N Positive is the number of stations with significant positive trend; and finally % Pos/Rej is the percentage of those with positive trend among the ones with significant trend.

Variable	N Rej	% Rej/T	N Positive	% Pos/Rej
P	225	13.8	165	73.3
INT	361	22.2	206	57.1
DSL	246	15.1	76	30.9
HY-INT	320	19.6	128	40.0
N Wet Days	281	17.2	200	71.2
N Dry Days	282	17.3	80	28.4
LW	286	17.6	183	64.0
N Wet Runs	229	14.1	134	58.5
N Dry Runs	235	14.4	144	61.3
INTX	148	9.3	85	57.4
DSLX	148	9.3	29	19.6
HYINTX	149	9.4	39	26.2
LWX	228	14.4	146	64.0

304 trend slope increases with precipitation irrespective of the sign. Whereas for the HY-INT case, positive
305 trend slope decreases with P and the negative slope increases.

306 As expected from the small number of stations with significant trends, the space distribution does not
307 seem to show any pattern. Maps in figure 7 show in the location of each station the sign of the trend in
308 HY-INT, INT, DSL, and P. Nevertheless, Table 3 shows that though the percentage of stations with a
309 significant trend is small, there is a majority of positive trends for total annual precipitation overall (73%)
310 and an overwhelming majority in the Pacific (94%) and Caribbean (91%) regions. These results accord with
311 previous studies for the Pacific region. For the Caribbean region, the results show a trend in the opposite
312 direction to the predictions of GCM's (see section 2).

313 Bearing in mind that only a fraction (10 to 20%) of the stations show statistically significant trends,
314 among the significant ones the Pacific region exhibits differences with the overall behavior. For the HY-INT
315 variable, the percentage of significant trends is 32% whereas overall is only 20% and the percentage of
316 increasing trends goes from 40% (overall) to 50%. For the INT variable, the corresponding figures go
317 from 57% to 74%. Similarly, for the number of wet spells, the percentage of increasing trends among
318 the significant ones goes from 59% overall to 67% for the Pacific region. For the number of wet days in
319 the year, the percentage of significant trends is 27% whereas overall is only 17%. Moreover, for DSLX
320 the significant negative trends go from 80% to 93%. Together with the observation about total annual
321 precipitation, these changes indicate that the region tends to a more humid condition.

322 For the rest of the regions, the significant trends in HY-INT are not notably different from the total.
323 Except for the Orinoco region that we analyze below.

324 Besides the changes in P mentioned above, for the Caribbean region, the percentage of increasing trends
325 among the significant trends differs from the overall in the number of wet spells (59% overall vs. 76%) and




Figure 7. Maps of the trend sign for HY-INT (A) and INT (B), DSL (C) and P (D). Significant positive (increasing) trends in blue, negative (decreasing) trends in red and no significant trends in gray.

326 the INT variable (57% overall vs. 63%). With the caveat of the low number of stations with statistically
 327 significant trends, these results also point in the direction of a wetter region.

328 For the Amazon region, the variables with different behavior than the overall are in the decreasing trends
 329 of DSL (69% overall vs. 92%) and DSLX (80% overall vs. 94%). That means that the lengths of dry spells
 330 are getting shorter.

331 The Orinoco region has few stations and much fewer significant trends. Among those the variables
 332 HY-INT, number of wet days, and the number of wet spells show notorious differences in the percentage of
 333 increasing trends with the countrywide numbers: 40 vs. 55%, 71 vs. 47%, and 59 vs. 64% respectively (Off
 334 each pair the first figure is for the whole country and the second one is for the region).

335 The Andes region has a substantial number of all the stations in the country, and therefore for all the
 336 variables, the behavior of each trend is close to the one of the whole country, except for the number of wet
 337 spells. For that variable, the percentage of positive trends among the significant ones is only 46% whereas
 338 the country average is 59%. This reduction is possible because for the other four regions the percentage
 339 increases.

Table 3. Regional summary of the trends of rain gauges for Total Annual Precipitation and HY-INT. Column symbols as in Table 2

Region	N Rej	% Rej/T	N Positive	% Pos/Rej
Total Annual Precipitation				
Amazon	8	11	6	75
Andes	133	13	85	64
Caribbean	57	15	52	91
Orinoco	9	10	5	56
Pacific	18	24	17	94
Total	225	14	165	73
HY-INT				
Amazon	16	21	5	31
Andes	188	19	74	39
Caribbean	81	21	31	38
Orinoco	11	13	6	55
Pacific	24	32	12	50
Total	320	20	128	40

340 Figure 8 shows the distribution of the record length of all the stations coding with gray the ones
 341 corresponding to a significant trend in HY-INT. The percentage of stations with significant trend out of
 342 the total number of stations in each record length bin does not change significantly except at the tails. In
 343 other words, the probability that the HY-INT trend of a given station is statistically significant is pretty
 344 much uniform as a function of the record length. This observation indicates that for the set of rain gauges
 345 analyzed the record length does not explain the lack of significant trends. On the contrary, the reason for
 346 the limited number of significant stations with significant trend needs other explanation.

347 We also looked into the possible dependence of trend slopes on latitude, longitude, elevation, and
 348 seasonality of the annual regime of precipitation. However, there were not any pattern worth mentioning.

6 DISCUSSION

349 Our results about the trends in annual precipitation agree in some way with the previous studies reported
 350 in Section 2 that considered rain gauges: Increasing trends prevail over decreasing trends among the




Figure 8. Histogram of the the record length (number of complete years) of the 1629 rain gauges used in this study. Stations with significant trend for HY-INT in gray.

351 statistically significant ones. However, we found a large number of stations with no significant trends.
 352 Probably the number of stations we considered explain these differences.

353 The principal result of this work is that the existing records of precipitation in the rain gauge network
 354 of Colombia do not show a clear signal of statistically significant trends. Only approximately 20% of the
 355 gauges present significant trends. This observation is valid for all the variables we studied, even with a
 356 lower percentage of significant trends for some of the variables. Given the overwhelming evidence of
 357 global climate change, this result claims for an explanation.

358 Also, the lack of a space pattern for the trends in the Andes is another evidence that is striking. The few
 359 stations with significant trends are mixed in space with nearby gauges with no significant trend or even
 360 significant trend of the opposite sign.

361 One possibility is to blame short records. Although one wishes to have longer records, this does not seem
 362 to be a good explanation. The percentage of significant trends out of the number of gauges in a given bin of
 363 record length remains approximately constant.

364 One factor that may play a role is the irregularity of the precipitation field.

365 The humid tropical climate of Colombia is probably a factor for the explanation. However, (Stevens et al.,
 366 2013) has argued that changes in precipitation need to be coupled with changes in the global circulation.
 367 The sole increase in absolute moisture is not sufficient to change the amount of precipitation, at least in an
 368 average sense. In that sense, there are no reports of changes in the trade winds or the low-level jets that
 369 bring moisture to Colombia, except for the Chocó jet.

370 The issue of changes in ENSO due to climate change is a big area of debate, see for instance Kohyama
 371 and Hartmann (2017). Some argue that with global warming ENSO could become more frequent and
 372 intense, but others that it could become weaker or more located on central rather than eastern Pacific.
 373 The effect of ENSO over Colombia is to produce dryer weather. Therefore the issue is very relevant to

374 elucidate the effects of climate change over Colombian precipitation. In that sense, our results seem to
375 suggest that ENSO is not becoming more intense or frequent. Also, the fact that majority of significant
376 trends for total annual precipitation is positive for the Pacific region accords with the nonlinear ENSO
377 warming suppression and a possible strengthening of the Chocó jet.

378 Other result is about the HY-INT index of [Giorgi et al. \(2011\)](#) to quantify the intensity of the hydrologic
379 cycle. For many parts of the globe, it may be true that rainfall intensity and dry spell length are deeply
380 interconnected. However, our results suggest that it is not the case for a humid tropical climate like
381 Colombia's. At least for the few gauges with significant trends the two factors in the definition of HY-INT,
382 rainfall intensity and dry spell length do not necessarily go together. For instance, of the 57% (205 out of
383 361) of the station with significant INT trend have a positive trend; but of those, 45% have positive DSL
384 trend and 55% negative one (92 and 113 respectively).

385 We wanted to complement HY-INT, the indicator of the intensity of the hydrologic cycle, by defining
386 an extreme version, HYINTX, the product of the maximum daily rainfall times the maximum dry spell
387 length. For Colombia, this indicator did not give any good results. Even they were weaker than the original
388 HY-INT indicator. One reason for this failure seems to be that dry spell length tends to decrease even
389 though the trend in maximum intensity is positive.

AUTHOR CONTRIBUTIONS

390 All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it
391 for publication.

CONFLICT OF INTEREST STATEMENT

392 The authors declare that the research was conducted in the absence of any commercial or financial
393 relationships that could be construed as a potential conflict of interest.

FUNDING

394 Universidad Nacional de Colombia provided support for this work.

ACKNOWLEDGMENTS

395 We acknowledge Colombian meteorological service, IDEAM, for providing the rainfall data.

REFERENCES

- 396 ACON-OPTIM (2013). *Estudio para determinar la vulnerabilidad y las opciones de adaptación del sector*
397 *energético Colombiano frente al cambio climático. Informe final*. Tech. rep., Unidad de Planeación
398 Minero Energética, UPME-0223-290-dic/2013-V1. 416 págs., Bogotá
- 399 Cantor, D. (2011). *Evaluación y análisis espaciotemporal de tendencias de largo plazo en la*
400 *hidroclimatología colombiana. Master Thesis in Water Resources Engineering*. Master Thesis in
401 Water Resources Engineering, Universidad Nacional de Colombia, Medellín
- 402 Cantor, D. and Ochoa, A. (2011). Señales de cambio climático en series de lluvia en Antioquia. In *IX*
403 *Congreso Colombiano de Meteorología* (Bogotá: IDEAM - Universidad Nacional de Colombia), 11 p.
404 doi:10.13140/RG.2.1.1573.0326

- 405 Carmona, A. M. and Poveda, G. (2014). Detection of long-term trends in monthly hydro-climatic series
406 of Colombia through Empirical Mode Decomposition. *Climatic Change* 123, 301–313. doi:10.1007/
407 s10584-013-1046-3
- 408 Eslava, J. A. (1993). Climatología y diversidad climática de Colombia. *Revista de la Academia Colombiana*
409 *de Ciencias Exactas, Físicas y Naturales* 18, 507–538
- 410 Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., et al. (2015). The climate hazards
411 infrared precipitation with stations—a new environmental record for monitoring extremes. *Scientific*
412 *Data* 2. doi:10.1038/sdata.2015.66
- 413 Giorgi, F., Im, E.-S., Coppola, E., Diffenbaugh, N. S., Gao, X. J., Mariotti, L., et al. (2011). Higher
414 Hydroclimatic Intensity with Global Warming. *Journal of Climate* 24, 5309–5324. doi:10.1175/
415 2011JCLI3979.1
- 416 Hamed, K. H. (2008). Trend detection in hydrologic data: The Mann–Kendall trend test under the scaling
417 hypothesis. *Journal of Hydrology* 349, 350–363. doi:10.1016/j.jhydrol.2007.11.009
- 418 Hamed, K. H. and Ramachandra-Rao, A. (1998). A modified Mann-Kendall trend test for autocorrelated
419 data. *Journal of Hydrology* 204, 182–196. doi:10.1016/S0022-1694(97)00125-X
- 420 Hurtado, A. F. and Mesa, O. J. (2015). Cambio climático y variabilidad espacio–temporal de la precipitación
421 en Colombia. *Revista EIA* 12, 131–150
- 422 IDEAM-Colombia (2010). *2a Comunicación Nacional ante la Convención Marco de las Naciones Unidas*
423 *sobre cambio climático* (Bogotá)
- 424 IGAC (1997). *Mapa de Regiones Naturales de Colombia* (Bogotá, Colombia: Instituto Geográfico Agustín
425 Codazzi)
- 426 Kendall, M. G. (1955). *Rank Correlation Methods* (London, UK: Griffin), 2 edn.
- 427 Kohyama, T. and Hartmann, D. L. (2017). Nonlinear ENSO warming suppression (news). *Journal of Climate*
428 30, 4227–4251
- 429 Kulkarni, A. and von Storch, H. (1992). Monte Carlo experiments on the effect of serial correlation on the
430 Mann-Kendall test of trend. *Meteorologische Zeitschrift* 4, 82–85. doi:10.1127/metz/4/1992/82
- 431 Macías, A. M. and Andrade, J. (2014). *Estudio de generación eléctrica bajo escenario de cambio climático*.
432 Tech. rep., Unidad de Planeación Minero Energética, UPME, Bogotá
- 433 Mann, H. B. (1945). Nonparametric Tests Against Trend. *Econometrica* 13, 245–259. doi:
434 0012-9682(194507)13:3<245:NTAT>2.0.CO;2-U
- 435 Mayorga, R., Hurtado, G., and Benavides, H. (2011). *Evidencias de cambio climático en Colombia con*
436 *base en información estadística*. Tech. rep., IDEAM–METEO/001-2011 Nota técnica del IDEAM
- 437 Mejía, J. F., Mesa, O. J., Poveda, G., Vélez, J. I., Hoyos, C. D., Mantilla, R. I., et al. (1999). Distribución
438 espacial y ciclos anual y semianual de la precipitación en Colombia. *Dyna* 127, 7–26
- 439 Mesa, O., Poveda, G., and Carvajal, L. F. (1997). *Introducción al Clima de Colombia* (Santa Fe de Bogotá,
440 D.C., Colombia: Universidad Nacional de Colombia)
- 441 Mitchell, J. F., Wilson, C., and Cunningham, W. (1987). On CO₂ climate sensitivity and model dependence
442 of results. *Quarterly Journal of the Royal Meteorological Society* 113, 293–322
- 443 Ochoa, A. and Poveda, G. (2008). Distribución espacial de señales de cambio climático en Colombia. In
444 *Proc. XXIII Latin American Hydraulics Meeting, IAHS, Cartagena, Colombia, September*
- 445 Oster, R. (1979). Las precipitaciones en Colombia. *Colombia Geográfica* 6, 5–147
- 446 O’gorman, P. A. and Schneider, T. (2009). Scaling of precipitation extremes over a wide range of climates
447 simulated with an idealized GCM. *Journal of Climate* 22, 5676–5685
- 448 Pabón, J. D. (2005). Escenarios de cambio climático para territorio Colombiano. *Documento INAPPDF-B*

- 449 Pabón, J. D. (2009). El cambio climático global y su manifestación en Colombia. *Cuadernos de Geografía: Revista Colombiana de Geografía* 12, 111–119
- 451 Peterson, T. (2005). Climate change indices. *WMO Bulletin* 54, 83–86
- 452 Poveda, G., Álvarez, D. M., and Rueda, Ó. A. (2011). Hydro-climatic variability over the Andes of
453 Colombia associated with ENSO: a review of climatic processes and their impact on one of the earth's
454 most important biodiversity hotspots. *Climate Dynamics* 36, 2233–2249
- 455 Poveda, G., Jaramillo, L., and Vallejo, L. F. (2014). Seasonal precipitation patterns along pathways of
456 South American low-level jets and aerial rivers. *Water Resources Research* 50, 98–118. doi:10.1002/
457 2013WR014087
- 458 Poveda, G. and Mesa, O. J. (1997). Feedbacks between hydrological processes in tropical South America
459 and large-scale ocean-atmospheric phenomena. *Journal of Climate* 10, 2690–2702
- 460 Quintana-Gomez, R. A. (1999). Trends of maximum and minimum temperatures in northern South
461 America. *Journal of Climate* 12, 2104–2112
- 462 Rabatel, A., Francou, B., Soruco, A., Gomez, J., Cáceres, B., Ceballos, J., et al. (2013). Current state of
463 glaciers in the tropical andes: a multi-century perspective on glacier evolution and climate change. *The*
464 *Cryosphere* 7, 81–102
- 465 Ruiz, D., Moreno, H. A., Gutiérrez, M. E., and Zapata, P. A. (2008). Changing climate and endangered
466 high mountain ecosystems in Colombia. *Science of The Total Environment* 398, 122–132. doi:10.1016/j.
467 scitotenv.2008.02.038
- 468 Ruiz, J. F. (2007). *Escenarios de cambio climático, algunos modelos y resultados de lluvia para Colombia*
469 *bajo el escenario A1B. Nota Técnica.* Tech. rep., IDEAM – METEO/003-2007
- 470 Ruiz, J. F. (2010). *Cambio climático en temperatura, precipitación y humedad relativa para Colombia*
471 *usando modelos meteorológicos de alta resolución. Panorama 2011 – 2100.* Tech. rep., IDEAM Nota
472 técnica 005/2010
- 473 Salazar, J. F. (2011). *Regulación biótica del ciclo hidrológico en múltiples escalas.* Ph.D. thesis, PhD
474 thesis. Universidad Nacional de Colombia. Medellín
- 475 Sen, P. K. (1968). Estimates of the Regression Coefficient Based on Kendall's Tau. *Journal of the American*
476 *Statistical Association* 63, 1379. doi:10.2307/2285891
- 477 Smith, R., Poveda, G., Mesa, O., Pérez, C., and Ruiz, C. (1996). En búsqueda de señales del cambio
478 climático en Colombia. In *IV Congreso Colombiano de Meteorología, Sociedad Colombiana de*
479 *Meteorología, Bogotá*
- 480 Snow, J. (1976). The Climate of Northern South America. In *World survey of climatology Volume 12:*
481 *Climates of Central and South America*, ed. W. Schwerdtfeger (Amsterdam, Netherlands: Elsevier),
482 chap. 6. 295–403
- 483 Soden, B. J. and Held, I. M. (2006). An assessment of climate feedbacks in coupled ocean-atmosphere
484 models. *Journal of Climate* 19, 3354–3360
- 485 Stevens, B., Bony, S., et al. (2013). What are climate models missing. *Science* 340, 1053–1054
- 486 Urán, J. D. (2015). *Cambios en los valores extremos de precipitación en regiones tropicales asociados a*
487 *cambio climático.* Master's thesis, Posgrado en Aprovechamiento de Recursos Hidráulicos. Universidad
488 Nacional de Colombia Sede Medellín
- 489 Urrea, V. (2017). *Variabilidad espacial y temporal del ciclo anual de lluvia en Colombia.* Master thesis in
490 water resources engineering, Universidad Nacional de Colombia, Medellín
- 491 Urrea, V., Ochoa, A., and Mesa, O. (2016). Validación de la base de datos de precipitación CHIRPS para
492 Colombia a escala diaria, mensual y anual en el período 1981-2014. In *XXVII Congreso Latinoamericano*
493 *de Hidráulica* (Lima, Perú: IAHS), 11

- 494 Urrea, V., Ochoa, A., and Mesa, O. (2019). Seasonality of Rainfall in Colombia. *Water Resources Research*
495 doi:10.1029/2018WR023316
- 496 von Storch, H. (1995). Misuses of Statistical Analysis in Climate Research. In *Analysis of Climate*
497 *Variability: Applications of Statistical Techniques*, eds. H. von Storch and A. Navarra (Berlin: Springer-
498 Verlag), chap. 2. 11–26
- 499 Vuille, M., Bradley, R. S., Werner, M., and Keimig, F. (2003). 20th century climate change in the tropical
500 andes: observations and model results. In *Climate variability and change in high elevation regions: past,*
501 *present & future* (Springer). 75–99
- 502 Wentz, F. J., Ricciardulli, L., Hilburn, K., and Mears, C. (2007). How much more rain will global warming
503 bring? *Science* 317, 233–235