# Challenges in using modern pollen analogs for Cenozoic paleoecology: examples from the European Neogene

Valentí Rull<sup>1,2</sup>

<sup>1</sup>Spanish National Research Council (CSIC), Botanic Institute of Barcelona, Pg. del Migdia s/n, 08038 Barcelona, Spain. Email: <u>vrull@csic.es</u>

<sup>2</sup>Institut Català de Paleontologia Miquel Crusafont (ICP-CERCA), Universitat Autònoma de Barcelona, c/Columnes s/n, 08193 Cerdanyola del Vallès, Barcelona, Spain. Email: <u>valenti.rull@icp.cat</u>

## Abstract

The use of modern pollen analogs in paleoecology is well established in Quaternary studies; however, the reliability of this approach decreases with increasing geological age due to evolutionary changes. Establishing a definitive chronological boundary beyond which modern pollen analogs remain reliable is currently unfeasible. This limitation affects not only paleoenvironmental reconstructions based on the niche conservatism hypothesis for individual extant species, but also community- and biome-level reconstructions that assume compositional and environmental constancy over time. This paper discusses three case studies from the European Neogene—focusing on individual species, communities, and biomes—where the use of modern pollen analogs under the untested assumption of niche conservatism is common. These studies include: (1) quantitative paleoclimatic estimations using the climatic envelope of extant species, (2) comparisons between Middle Miocene European mangroves and presentday Middle-East mangrove communities, and (3) the use of Artemisia pollen as an indicator of dry steppe biomes. The main conclusion is that assuming species' niche conservatism and extrapolating this assumption to communities and biomes is unwarranted and may lead to unsupported conclusions. Therefore, empirical testing of species' niche conservatism across the relevant time spans is essential. Alternative and complementary methods—such as the fossil community approach and the use of pollen-independent proxies (e.g., stable isotopes, biomarkers)—are strongly recommended.

## Keywords

Modern analogs, niche conservatism, paleoclimatology, communities, biomes, Europe, Neogene

# 1. Introduction

The use of modern pollen analogs is common in Quaternary paleoecology and paleoclimatology, based on the assumption that the ecological and environmental requirements of extant species have remained constant over the last ~2.6 million years (e.g., Birks & Birks, 1980; Delcourt & Delcourt, 1991; Cronin, 2009; Bradley, 2015; Rull, 2020). This assumption has led to detailed quantitative estimates of climatic and other environmental parameters using transfer functions, which apply the environmental envelope of extant species to fossil pollen taxa (Birks et al., 2012). However, this assumed niche constancy—also known as niche conservatism (Wiens et al., 2010)—declines over time due to evolutionary processes, progressively reducing the likelihood of finding reliable modern analogs for fossil pollen taxa. Therefore, niche-conservatism uncertainty (NCU) increases with time, and establishing a chronological boundary for the reliability of modern analogs beyond the Quaternary is currently unfeasible (Fauquette, 2017; Matthaeus et al., 2023).

In spite of this, the use of modern analog approaches in Cenozoic paleoecology and paleoclimatology remains common. These methods range from detailed paleoclimatic estimates to reconstructions of past community and biome dynamics, based on the assumption that extant communities and biomes have also remained compositionally and environmentally stable over time. The concept of compositional stability also emerged from Quaternary studies (Prentice et al., 1996; Ni et al., 2010) and was later adopted for the Neogene (e.g., Mahler et al., 2022; Altoaguirre et al., 2023). Some macroecological surveys support this perspective. For example, phylogenetic niche conservatism—the tendency of lineages to retain niche-related traits through speciation events (Crisp & Cook, 2012), or more succinctly, the retention of ecological traits over time among related species (Wiens et al., 2010)—has been considered a major factor structuring vegetation types and biomes globally (Crisp et al., 2009). Similar results have been reported for the European Cenozoic (Lososová et al., 2020).

However, these studies rely on phylogenetic inference and modelling, rather than empirical paleontological evidence, and do not resolve the NCU conundrum, neither for species' niche traits nor for community or biome composition. Therefore, assuming niche conservatism across taxonomic, geographical, and temporal scales is unwarranted and may lead to questionable past ecological and biogeographical reconstructions. As a consequence, modern analog approaches that uncritically adopt niche conservatism as a premise should be approached with caution.

Among the many examples in the literature that rely on these assumptions, we have selected three from the European Mediterranean Neogene including inferences involving individual species, communities and biomes. The first case involves the assumption of climatic niche conservatism of extant species to quantitatively estimate Neogene climates, particularly paleotemperature and paleoprecipitation, which has been widely used in Europe (Utescher et al., 2014; Suc et al., 2018). The second example is the use of modern mangrove communities from the Middle East as modern analogs for those that existed along the northern Mediterranean coasts during the Middle Miocene Climatic Optimum (MMCO) (Plaziat et al., 2001; Popescu et al., 2021). This case was chosen in light of a recent study that provides a detailed description of Middle East mangroves and their environmental context (Waleed, 2024). The third example concerns the widespread use of *Artemisia* fossil pollen as a straightforward indicator of temperate, mostly cold and arid, steppe biomes (Zagwijn, 1960; Suc, 1984). This case is revisited in response to a recent thorough study on *Artemisia* pollen morphology and the broad range of biomes inhabited by species of this genus (Lu et al., 2022). While these cases have been analyzed in detail in other publications, this paper provides a succinct summary.

The main aim of this discussion is to highlight the need to avoid using unsupported general modern-analog assumptions as premises for interpreting past ecosystems and their environmental features. The intention is not to dismiss the use of modern-analog approaches, but rather to improve their reliability by taking into account the specific characteristics of each taxon and community under study. A comprehensive review of the subject is beyond the scope of this paper, which focuses on specific case studies from the currently defined European continent.

# 2. Species level: paleoclimatic estimates

Pollen-based Cenozoic paleoclimatic estimates using modern analog approaches—such as the widely used Coexistence Approach (Mosbrugger & Utescher, 1997) or the Climate Amplitude Method (Fauquette et al., 1998), among others—assume climatic niche conservatism in individual species and community constancy over evolutionary timescales. In this framework, precise quantitative estimates of past climatic variables are obtained from fossil taxa that share pollen morphology with extant species—known as nearest living relatives (NLRs)—by

transferring the climatic envelope of these NLRs to their assumed fossil representatives. These methods have been challenged from both theoretical and practical perspectives.

Theoretically, the assumptions of niche conservatism and community constancy have been considered unrealistic, as they remain empirically untested in most reconstructions. Testing climatic niche conservatism requires independent paleoclimatic and palynological evidence. Non-empirical methods—such as molecular phylogenetic inference and species distribution modeling—suggest that the presence or absence of niche conservatism in plants throughout the Cenozoic depends on both the geological timescale and the taxa under study, with no general rule being universally applicable (e.g., Svenning, 2003; Abbink et la., 2004; Byrstriakova et al., 2011; Loera et al., 2012; Martín-Hernanz et al., 2014; Weeks et al., 2014; Benítez-Benítez et al., 2017; Zhang et al., 2021; Vieira et al., 2023; Qian et al., 2024; Quirk et al., 2024; Fernández-Mazuecos & Glover, 2025). Therefore, niche conservatism in the target taxa over the time interval of interest cannot be assumed a priori.

Concerns have also been raised about the establishment of reliable botanical affinities, which largely depends on the taxonomic resolution achievable in the fossil pollen record—typically at the genus or family level—compared to NLRs. This is especially critical in genera and families with homogeneous pollen morphology. Another drawback is that modern-analog-based methods implicitly overlook the possibility that extinct species, as well as species originated after the time interval under study, with the same pollen morphology that extant taxa may be present in fossil assemblages. Additionally, the potential for modern-analog approaches to result in circular reasoning when studying vegetation responses to climate shifts has been highlighted. Reviews and discussions addressing these issues can be found in Grimm & Potts (2016), Lu et al. (2022); Matthaeus et al. (2023), Quirk et al. (2024) and Rull et al., 2024a).

Another untested assumption is that pollen morphology and ecophysiological traits linked to climatic requirements evolve in a similar manner. However, Krassilov (2000) showed that "sporophytic and pollen characters show only weak correlation over their evolutionary history." According to the author, in some cases, pollen morphology is a conservative trait compared to other plant characteristics, while in other cases, the opposite is true. There is no general rule that can be universally applied; each case requires specific analysis.

The chronological uncertainty associated to niche conservatism (NCU) would also apply to community composition. In this case, empirical palynological data show that the taxonomic composition of many communities has actually varied over time, even during the most recent Quaternary period, largely due to range shifts of extant species (Jackson & Williams, 2004; Williams & Jackson, 2007; Rull, 2020). These compositional changes have been even more pronounced throughout the Paleogene and the Neogene, due to evolutionary factors such as speciation and extinction, in addition to large-scale dispersal (e.g., Rull, 2024). Therefore, extant communities cannot be assumed a priori to be modern analogs for past Cenozoic communities.

From a practical standpoint, pollen-based paleoclimatic estimates derived from modern-analog methods have consistently shown significant discrepancies when compared with pollenindependent and modeling approaches (e.g., You et al., 2009; Herold et al., 2011; Goldner et al., 2014; Zhang et al., 2019; Salocchi et al., 2021; Botsyun et al., 2022; Gilson et al., 2022). According to Popescu et al., (2021), this could be due to the excessive weight that pollen-based methods assign to megathermal plants, particularly those from mangroves (see next section), in pollen-based paleoclimatic reconstructions.

In summary, given the current state of knowledge, pollen-based paleoclimatic reconstructions using modern-analog approaches should be interpreted with caution and await further

developments that may improve their reliability, with emphasis on empirical paleoecological testing of previous assumptions. The use of pollen-independent methods of paleoclimatic estimation, especially those based on isotopic and biomarker proxies, is highly recommended (Rull et al., 2024a).

## 3. Communitiy level: Mediterranean mangroves

Mangroves are intertidal forest communities currently restricted to tropical and subtropical coasts between approximately 30°N and 30°S (Spalding et al., 2010). However, during the Miocene—particularly the Middle Miocene Climate Optimum (MMCO; 17–15 Ma), when global temperatures were 4–8 °C higher than today (Westerhold et al., 2020)—pollen records indicate that the northern limit of mangrove distribution extended to the northern Mediterranean coasts, reaching up to approximately 45°N (Plaziat et al., 2001; Popescu et al., 2021). These MMCO Mediterranean mangroves have been referred as *Avicennia*-only mangroves, as they were dominated by *Avicennia* (Acanthaceae) – one of the two globally distributed genera of mangrove-forming trees (Tomlinson, 2016; Duke, 2017) – whereas *Rhizophora* (Rhizophoraceae), the other conspicuous mangrove genus, along with other mangrove-forming tree genera, was absent (Popescu et al., 2021).

The MMCO *Avicennia*-only Mediterranean mangroves have been considered taxonomically and ecologically comparable to those currently found in the arid and semi-arid regions of the Middle East, particularly along the coasts surrounding the Arabian Peninsula (Jiménez-Moreno and Suc, 2007; Suc et al., 2018; Popescu et al., 2021). These regions represent the northernmost boundary of modern mangrove distribution, where *Avicennia* is the dominant mangrove-forming tree. If this comparison holds, modern Middle-East mangroves could serve as modern analogues for reconstructing the ecological characteristics of MMCO Mediterranean mangroves. However, this comparison has yet to be fully explored from biogeographical and paleoenvironmental perspectives (Rull et al., 2024b).

The Miocene Avicennia-only Mediterranean mangroves were considered impoverished mangroves located at the northernmost edge of mangrove distribution, likely reflecting a temperature-driven latitudinal pattern (Popescu et al., 2021). These authors estimated a mean annual temperature (MAT) of 18–20 °C and mean annual precipitation (MAP) exceeding 1000 mm/year, except in some localities of the Iberian Peninsula—particularly in the southwestern sector—where climates were drier (see also Postigo-Mijarra et al., 2009; Barrón et al., 2010). As mentioned above, these pollen-based estimates were higher than others based on pollen-independent proxies and paleoclimate modeling, which yielded MAT values of 11–17 °C and sea surface temperatures (SST) of around 20 °C (You et al., 2009; Goldner et al., 2011).

Regarding current Middle Eastern mangroves, a recent review covering most African and Asian coasts surrounding the Arabian Peninsula has revealed highly diverse environmental conditions (Waleed, 2025). MAT vary significantly across the region, ranging from 20 °C in the NE to 30 °C in the SE. SST can reach up to 28 °C in summer, while MAP is generally below 100 mm/year, with evaporation rates typically ten times higher. Additionally, the absence of large permanent freshwater sources limits sediment and nutrient supply essential for mangrove growth and results in hypersaline coastal conditions exceeding 40 ppt. These extremely hot, arid and hypersaline environments are generally unfavorable for mangrove development (Quisthoudt et al., 2012).

Only two mangrove-forming tree species—Avicennia marina and Rhizophora mucronata—are capable of surviving the harsh environmental conditions of the Middle East. Avicennia marina typically dominates the mangrove canopy, whereas Rhizophora mucronata is rare or entirely

absent, found only in a few locations with some freshwater input (Waleed, 2025). Due to the intense environmental stress, both species exhibit stunted, dwarf growth forms when compared to populations thriving in more favorable climates. Overall, the sparse, low-stature, and species-poor *Avicennia*-dominated mangroves of the Middle East reflect a unique regional adaptation to aridity and related environmental pressures (Osland et al., 2017), rather than a pattern driven by latitude-dependent temperature gradients. In contrast, other Asian mangroves occurring at similar latitudes (approximately 10°–30° N), but under more favorable environmental regimes, rank among the most diverse and healthy mangrove ecosystems in the pantropical zone (Tomlinson, 2016; Duke, 2017).

These observations suggest that the present-day Middle East mangroves represent a latitudinal anomaly within the northern tropical/subtropical mangrove belt—one that is independent of the global temperature gradient and instead more closely linked to regional aridity and hypersalinity. Consequently, these extant mangroves are unlikely to serve as appropriate modern analogs for the Miocene Mediterranean mangroves, considered as a temperature-driven latitudinal feature. This view is further supported by the contrasting climatic conditions estimated for Miocene mangroves, particularly the absence of extreme heat and aridity. Additionally, some studies from the NE Iberian Peninsula do not support the presence of hypersaline conditions (Bitzer, 2004). Combined with the absence of a present-day northern latitudinal zone dominated solely by *Avicennia* mangroves, the above observations make it difficult to identify suitable modern analogs for the MCO Mediterranean mangroves. In the Southern Hemisphere, *Avicennia* mangroves occur in the Mediterranean antipodes (New Zealand and southeastern Australia), where minimum temperatures—a latitude-dependent factor—are the primary drivers of mangrove distribution and diversity (Osland et al., 2017). Further comparisons with these systems could offer valuable insights.

The available evidence suggests that potential modern analogs for MCO Avicennia-only mangroves would have thrived in mild, humid climates. Today, Avicennia-only mangroves occur sporadically, either as pioneering communities on recently formed mudflats or in hypersaline areas within broader mangrove ecosystems across tropical and subtropical regions (Thatoi et al., 2016). It is plausible that MMCO mangroves developed under similar conditions, though on a much broader regional scale. However, further research is needed to determine whether meaningful parallels exist between ancient and modern Avicennia stands. If no clear analogs can be identified, MMCO Mediterranean mangroves may represent an extinct mangrove type—a phenomenon not uncommon in mangrove history, particularly throughout the Cenozoic (Rull, 2024). In paleoecology, communities with no modern counterparts are relatively frequent, even during the Quaternary, when the species involved were already extant (Jackson and Williams, 2004; Williams and Jackson, 2007).

## 4. Biome level: Artemisia dry steppes

The presence of significant amounts of *Artemisia* (Asteraceae) pollen in fossil records has traditionally been considered an indicator of arid and semi-arid biomes, particularly steppes, in temperate regions (Wang, 2004). This interpretation applies to pre-Holocene records, as *Artemisia* species are common weeds and ruderal plants associated with human activities such as grazing and other intensive or extensive agricultural practices, which began to play a significant role during the Early Holocene Neolithization (Roberts, 2014). In the Neogene, steppes dominated by *Artemisia*, with *Ephedra* present, began to be important at the end of the Miocene (6.2-5.6 Ma) in the eastern Mediterranean area (Anatolian), coinciding with the Antarctic glaciation and the desiccation of the Mediterranean Sea. At the beginning of the Pliocene, these steppes expanded across the entire Mediterranean region, reaching the westernmost part during the Middle Pleistocene (Suc et al., 2018).

According to the most widely accepted view, the European Pleistocene was characterized by the alternation of cold steppes—dominated by *Artemisia*, *Ephedra*, and other Asteraceae—during glacial phases, and deciduous forests with thermophilous elements during interglacials (Zagwijn, 1960; Suc, 1984). However, Subally et al. (1999) noted that this pattern was disrupted, or even reversed, in some Mediterranean regions, with *Artemisia* reaching its highest values during interglacials, as indicated by independent isotopic proxies. The same authors analyzed the ecological niches of Mediterranean *Artemisia* species and found that, although steppe representatives were dominant, other species were distributed along environmental gradients of elevation—and thus temperature—and hydroclimatic conditions. Subally & Quézel (2002) suggested that *Artemisia* peaks during glacial and interglacial periods may correspond to different species, and recommended identifying *Artemisia* pollen at the species level and incorporating independent paleoclimatic proxies.

The main limitation is that *Artemisia* pollen is highly homogeneous, making identification at the species level challenging—particularly under light microscopy (LM), which is the standard method used in pollen analysis. Some attempts have been made in this direction, but they primarily aimed to resolve phylogenetic questions rather than support environmental reconstruction (e.g., Jiang et al., 2005; Ghahraman et al., 2007; Hayat et al., 2010; Hussain et al., 2019). Suc et al. (2004) were the first to attempt linking *Artemisia* pollen morphology with environmental preferences. Using scanning electron microscopy (SEM), they found that the key distinguishing feature was the morphology and size of spinules, and they preliminarily associated these traits with various Mediterranean species differing in thermal and edaphic requirements. Unfortunately, this classification was not further developed.

Recently, Lu et al. (2022) conducted a comprehensive pollen-morphological study that included species from all known *Artemisia* lineages (Sanz et al., 2008; Malik et al., 2017), along with the biomes and vegetation types in which they occur globally, accompanied by corresponding environmental characterizations. The study confirmed the high level of intra-generic pollen homogeneity and reinforced the value of spinule morphological traits as key diagnostic features. Lu et al. (2022) were able to distinguish only three main pollen types within a genus comprising over 500 species (World Flora Online; <u>https://wfoplantlist.org/</u>). These pollen types were designated as SWS (short and wide spinules), LNS (long and narrow spinules), and SG (sparse granules).

Lu et al. (2022) highlighted the wide range of biomes and climatic conditions in which *Artemisia* species occur—from forests to deserts, including grasslands and coastal saline environments—across various climatic zones, with a particular emphasis on temperate regions. The three defined pollen-morphological types showed little correlation with bioclimatic zones. Species with the SWS pollen type have a global distribution and occupy a wide variety of habitats, with a slight preference for humid forests and grasslands. The LNS pollen type was characteristic of generalist species found in forests, grasslands, deserts, and coastal areas, typically in regions with higher temperatures. Species with the SG pollen type were associated with low-elevation open biomes, such as dry grasslands, deserts, and saline environments. In summary, only species with the SG pollen type—and some with LNS—can be considered indicators of arid biomes; the others lack clear indicator value.

It can be concluded that *Artemisia*, when treated as a bulk pollen category, is not a reliable indicator of any specific biome or vegetation type. The difficulty in identifying pollen at the species level, along with the lack of correlation between pollen morphology and environmental preferences, complicates bioclimatic interpretation. Moreover, the need for scanning electron microscopy (SEM) to distinguish pollen types limits the applicability of the defined pollen-

morphological groups in routine pollen analysis. As a result, the interpretation of *Artemisia* fossil pollen records still depends on supporting methods, such as the fossil assemblage approach or the use of independent paleoenvironmental proxies (Rull, 2025). It would be worthwhile to investigate whether recent advances—such as Fourier transform infrared spectroscopy (FTIR) (Kendel & Zimmermann, 2020) and automated pollen counting using SEM (Smyth et al., 2015)—could offer useful support.

# 5. Discussion

The main drawbacks of using modern pollen analogs for the European Neogene stem from uncertainties surrounding niche conservatism (NCU) and its extrapolation to community- and biome-level taxonomic composition and environmental requirements. The most common approach is to assume niche conservatism and constancy in community or biome composition a priori, often without critical evaluation—an approach that can lead to unsupported conclusions. In the case of pollen, these assumptions are especially problematic due to additional factors, including limitations in pollen morphology and taxonomic resolution, difficulties in establishing accurate botanical affinities, and the possible presence of extinct species or taxa that evolved after the period under study. As such, finding empirical evidence for niche conservatism in the taxa examined over the relevant time span is essential. Identifying reliable modern analogs and clearly defining the spatiotemporal contexts in which they are applicable should therefore be a key objective.

Alternative and complementary approaches based on direct observation of the paleontological record—without reliance on potential modern analogs—are strongly recommended. One such method is the fossil community approach, which statistically defines fossil pollen assemblages based solely on paleontological evidence and incorporates pollen-independent paleoclimatic and paleoenvironmental proxies such as stable isotopes, biomarkers, and similar indicators (Rull et al., 2024a). This approach can be applied to individual species, communities, and biomes, and it provides empirical evidence for testing niche conservatism. As a result, it can help reduce the NCU for the taxa and time interval under study. Paleoclimatic modeling can also contribute to testing niche conservatism, though the underlying assumptions and premises of the models must be carefully considered.

# Acknowledgements

This contribution did not receive funding specifically for its development. The CERCA Programme, Generalitat de Catalunya, provided logistic support.

# References

- Abbink, O.A., Van Konijnenburg-Van Cittert, J.H.A., Visscher, H. 2004. A sporomorph ecogroup model for the Northwest European Jurassic Lower Cretaceous: concepts and framework. Netherlands Journal of Geosciences/Geologie en Mijnbouw 83, 17–31.
- Altoaguirre, Y., Postigo-Mijarra, J.M., Casas-Gallego, M., Moreno-Domínguez, R., Barrón, E. 2023. Mapping the Late Miocene Pyrenean forests of the La Cerdanya Basin, Spain. Forests 14, 1471.
- Barrón, E., Rivas-Carvalho, R., Postigo-Mijarra, J.M., Alcalde-Olivares, C., Vieira, M., Castro, L., Pais, J., Valle-Hernández, M. 2010. The Cenozoic vegetation of the Iberian Peninsula: A synthesis. Review of Palaeobotany and Palynology, 162, 382-402.
- Benítez-Benítez, C., Escudero, M., Rodríguez-Sánchez, F., Martín-Bravo, S., Jeménez-Mejías, P.
  2017. Pliocene-Plesitocene ecological niche evolution shapes the phylogeography of a Mediterranean plant group. Molecular Ecology 27, 1696-1713.

Birks, H.J.B., Birks, H.H. 1980. Quaternary Palaeoecology. Edward Arnold, London.

- Birks, H.J.B., Lotter, A.F., Juggins, S., Smol, J.P. 2012. Tracking Environmental Change Using Lake Sediments. Vol. 5: Data Handling and Numerical Techniques. Kluwer, Dordrecht.
- Bitzer, K. 2004. Estimating paleogeographic, hydrological and climatic conditions in the upper Burdigalian Vallès-Penedès basin (Catalunya, Spain). Geologica Acta, 2, 321-331.
- Botsyun, S., Ehlers, T.A., Koptev, A., Böhme, M., Methner, K., Risi, C., Stepanek, C., Mutz, S.G., Werner, M., Boateng, D., Mulch, A. 2022. Middle Miocene climate and stable oxygen isotopes in Europe based on numerical modelling. Paleoceanography and Paleoclimatology 37, e2022PA004442.
- Bradley, R.S. 2015. Paleoclimatology. Reconstructing Climates of the Quaternary. Elsevier-Academic Press, Oxford.
- Bystriakova, N., Schnedier, H., Coomes, D. 2011. Evolution of the climatic niche in scaly tree ferns (Cyatheaceae, Polypodiopsida). Botanical Journal of the Linnean Society 165, 1-19.
- Crisp, M.D., Arroyo, M.T.K., Cook, L.G., Gandolfo, M.A., Jordan, G.J., McGlone, M.S., Weston, P.H., Westoby, M., Wilf, P., Linder, H.P. 2009. Phylogenetic biome conservatism on a global scale. Nature 485, 754-756.
- Crisp, D., Cook, L.G. 2012. Phylogenetic niche conservatism: what are the underlying evolutionary and ecological causes? New Phytologist 196, 681-694.
- Cronin, T.H. 2009. Paleoclimates. Understanding Climate Change Past and Present. Columbia University Press, New York.
- Delcourt, H.R. Delcourt, P.A., 1991. Quaternary Ecology. Chapman and Hall, London.
- Duke, N.C. (2017). Mangrove floristics and biogeography revisited: further deductions from biodiversity hot spots, ancestral discotinuities, and common evolutionary processes. In Rivera-Monroy, V.H., Lee, S.Y., Kristensen, E., Twilley, R.R. (Eds.), Mangrove Ecosystems: A Global Biogeographic Perspective. Springer, pp. 17-53.
- Fauquette, S. 2017. Paléoenviroments en Europe et Región Méditerranéene au Cenozoic. Universtité Montpellier, Montpellier.
- Fauquette, S., Guiot, J., Suc, J.-P. 1998. A method for climatic reconstruction of the Mediterranean Pliocene using pollen data. Palaeogeography, Palaeoclimatology, Palaeoecology 144, 183–201.
- Fernández-Mazuecos, M., Glover, B.J. 2025. Climatic and edaphic niche shifts during plant radiation in the Mediterranean biodiversity hotspot. Annals of Botany 135, 717-734.
- Ghahraman, A., Nourbakhsh, N., Mehdigoli, K., Attar, F. 2007. Pollen morphology of *Artemisia* L. (Asteraceae) in Iran. Iran Journal of Botany 13, 21-29.
- Gilson, M.E., McCoy, J., O'Keefe, J.N.K., Nuñez Otaño N.B., Pound, M.J. 2022. Resconstructing terrestrial paleoclimates: a comparison of the co-existence approach, Bayesian and probability reconstruction techniques using the UK Neogene. Paleoceanography and Paleoclimatology 37, e2022PA004358.
- Goldner, A., Herold, N., & Huber, M. (2014). The challenge of simulating the warmth of the mid-Miocene climatic optimum in CESM1. Climate of the Past, 10, 523–536.
- Grimm, G.W., Potts, A.J. 2016. Fallacies and fantasies: the theoretical underpinnings of the Coexistence Approach for palaeoclimate reconstruction. Climate of the Past 12, 611-622.
- Hayat, M.Q., Ashraf, M., Khan, M.A., Yasmin, G., Shaheen, N., Jabeen, S. 2010. Palynological study of the genus Artemisia (Asteraceae) and its systematic implications. Pakistan Journal of Botany 42, 751-763.
- Herold, N., Huber, M., Müller, R. D. 2011. Modeling the Miocene climate optimum. Part I: Land and atmosphere. Journal of Climate, 24, 6353–6372.
- Hussain, A., Potter, D., Hayat, M.Q., Sahreen, S., Bokhari, S.A.I. 2019. Pollen morphology and its systematic implication on some species of *Artemisia* L. from Gilgit-Baltisan Pakistan. Bangladesh Journal of Plant Taxonomy 26, 157-168.

- Jackson, S.T., Williams, J.W. 2004. Modern analogs in Quaternary paleoecology: here today, gone yesterday, gone tomorrow? Annual Reviews of Earth and Planetary Sciences 32, 496-537.
- Jiang, L., Qin,W., Ye, L.-Z., et al. 2005. Pollen morphology of *Artemisia* L. and its systematic significance. Wuhan University Journal of Natural Sciences 10, 448-454.
- Jiménez-Moreno, G., & Suc, J.-P. 2007. Middle Miocene latitudinal climatic gradient in Western Europe: evidence from pollen reccords. Palaeogeography, Palaeoclimatology, Palaeoecology, 253, 208–225.
- Kendel, A., Zimmermann, B. 2020. Chemical analysis of pollen by FT-Raman and FTIR secpteroscopies. Frotiers in Plant Science 11, 352.
- Krassilov, V.A. 2000. Factors of palynomorphological evolution. Paleontological Journal 34, S2-S13.
- Loera, I., Sosa, V., Ickert-Bond, S.M. 2012. Diversification in North American arid lands: Niche conservatism, divergence and expansion of habitat explain speciation in the genus *Ephedra*. Molecular Phylogenetics and Evolution 65, 437-450.
- Lososová, Z., Divíšek, J., Chytrý, M., Götzenberger, L., Těšitel, J., Mucina, L. 2020. Macroevolutionary patterns in European vegetation. Journal of Vegetation Science 32, e12942.
- Lu, L.-L., Jiao, B.-H., Qin, F., Xie, G., Lu, K.-Q., Li, J.-F., Sun, B., Li, M., Ferguson, D.K., Gao, T.-G., Yao, Y.-F., Wang, Y.-F. 2022. *Artemisia* pollen dataset for exploring the potential ecological indicators in deep time. Earth System Science Data 14, 3961-3995.
- Mahler, S., Shatilova, I., Bruch, A.A. 2022. Neogene long-term trends in climate of the Colchic vegetation refuge in Western Georgia uplift versus global cooling. Review of Palaeobotany and Palynology 296, 104546.
- Malik, S., Vitales, D., Hayat, M.Q., Korovkov, A.A., Garnatje, T., Vallès, J. 2017. Phylogeny and biogeography of *Artemisia* subgenus *Seriphidium* (Asteraceae: Anthemideae). Taxon 66, 934-952.
- Martín-Hernanz, S., Albaladejo, R.G., Lavergne, S., Rubio, E., Grall, A., Aparicio, A. 2012. Biogeographic history and environmental niche evolution in the palearctic genus *Helianthemum* (Cistaceae). Molecular Phylogenetics and Evolution 163, 107238.
- Matthaeus, W.J., Macarewich, S.I., Richey, J., Montañez, I.P., McElwain, J.C., White, J.D., Wilson, J.P., Poulsen, C.J. 2023. A systems approach to understanding how plants transformed Earth's environment in deep time. Annual Review of Earth and Planetary Sciences 51, 551-580.
- Mosbrugger, V., Utescher, T. 1997. The coexistence approach: a method for quantitative reconstructions of tertiary terrestrial palaeoclimate data using plant fossils. Palaeogeography, Palaeoclimatology, Palaeoecology 134, 61–86.
- Ni, J., Yu, G., Harrison, S.P., Prentice, I.C. 2010. Palaeovegetation in Chinaduring the Late Quaternary: biome reconstructions based on a global scheme of plant functional types. Palaeogeography, Palaeoclimatology, Palaeoecology 289, 44-61.
- Osland, M.J., Feher, L.C., Griffith, K.T., Cavanaugh, K.C., Enwright, N.M., Day, R. H., Stagg, C.L., Krauss, K.W., Howard, R.J., Grace, J.B., Rogers, K. 2017. Climatic controls on the global distribution, abundance, and species richness of mangrove forests. Ecological Monographs, 87, 341-359.
- Plaziat, J.-C., Cavagnetto, C., Koeninger, J.-C., Baltzer, F. 2001. History and biogeography of the mangrove ecosystem, based on a critical reassessment of the paleontological record. Wetlands Ecology and Management, 9, 161-179.
- Popescu, S.-M., Suc, J.-P.; Fauquette, S., Bessedik, M., Jiménez-Moreno, G., Robin, C., Labrousses, L. 2021. Mangrove distribution and diversity during three Cenozoic thermal maxima in the Northern Hemisphere (pollen records from the Arctic-North Atlantic-Mediterranean regions). Journal of Biogeography 48, 2771-2784.

- Postigo-Mijarra, J.M., Barrón, E., Gómez Manzaneque, F., Morla, C. 2009. Floristic changes in the Iberian Peninsula and Balearic Islands (south-west Europe) during the Cenozoic. Journal of Biogeography, 36, 2025-2043.
- Prentice, I.C., Guiot, J., Huntley, B., Jolly, D., Cheddadi, R. 1996. Reconstructing biomes from palaeoecological data: a general method and its application to European pollen data at 0 and 6 ka. Climate Dynamics 12, 185-194.
- Qian, H., Qian, S., Zhang, J., Kessler, M. 2024. Effects of climate and environmental heterogeneity on the phylogenetic structure of regional angiosperm floras worldwide. Nature Communications 15, 1079.
- Quirk, Z.J., Smith, S., Acosta, R.P., Poulsen, C.J. 2024. Where did they come from, where did they go? Niche conservatism in woody and herbaceous plants and implications for plant-based paleoclimatic reconstructions. American Journal of Botany 111, e16426.
- Quisthoudt, K., Schmitz, N., Randin, C.F., Dahdouh-Guebas, F., Robert, E.M.R., Koedam, N. 2012. Temperature variation among mangrove latitudinal range limits worldwide. *Trees*, 26, 1919–1931.
- Roberts, N. 2014. The Holocene: An Environmental History. Wiley-Blackwell, Hoboken.
- Rull, V. 2020. Quaternary Ecology, Evolution and Biogeography. Elsevier/Academic Press, London.
- Rull, V. 2024. Origin and Evolution of Caribbean Mangroves. Springer Nature, Cham.
- Rull, V. 2025. Insights on the indicator capacity of *Artemisia* pollen in pre-Holocene paeoecology. *Palynology*.
- Rull, V., Alba, D., Casanovas-Vilar, I. 2024a. Middle Miocene vegetation of the Vallès-Penedès Basin (NE Iberian Peninsula), as inferred from fossils pollen records: State of the art and future prospects. Review of Palaeobotany and Palynology 321, 105042.
- Rull, V., Vicente, A., Casanovas-Vilar. I. 2004b. Middle Miocene Mediterranean mangoves and potential modern analogs. EartArXiv, doi 10.31223/X52Q6j.
- Salocchi, A.C., Krawielicki, J., Eglinton, T.I., Fioroni, C., Fontana, D., Conti, S., Picotti, V. 2021. Biomarker constraints on Mediterranean climate and ecosystem transitions during the Early-Middle Miocene. Palaeogeography, Palaeoclimatology, Palaeoecology, 562, 110092.
- Sanz, M., Vilatersana, R., Hidalgo, O., Garcia-Jacas, N., Susanna, A., Schneeweiss, G.M., Vallès, J. 2008. Molecular phylogeny and evolution of floral characters of *Artemisia* and allies (Anthemideae, Asteraceae): evidence from nrDNA ETS and ITS sequences. Taxon 57, 66-78.
- Smyth, A.P., Seales, B., Bradley, P.M. 2015. A pollen profile by scanning electron microscopy bracketing the mid-Holocene *Tsuga canadensis* declines at Pouywater Pond Bog, Holden, Massachussetts. Grana 54, 263-270.
- Spalding, M., Kainuma, M., Collins, L. 2010. World Atlas of Mangroves. Routledge, London.
- Subally, D., Quézel, P. 2002. Glacial or interglacial: *Artemisia*, a plant indicator with dual responses. Review of Palaeobotany and Palynology 120, 123-130.
- Subally, D., Bilodeau, G., Tamrat, E., Ferry, S., Debard, E. 1999. Cyclic climatic records during the Olduvai subchron (uppermost Pliocene) on Zakynthos Island (Ionia Sea). Geobios 32, 793-803.
- Suc J.-P. 1984. Origin and evolution of the Mediterranean vegetation and climate in Europe. Nature, 307: 429-432.
- Suc, J.-P., Fauquette, S., Popescu, S.-M. 2004. L'investigation palynologique du Cénozoïque passe par les herbiers. In Pierrel, R., Reduron, J.-P. (Eds.), Actes du Colloque "Les Herbiers : un Outil d'Avenir, Tradition et Modernité". Villeurbanne, France, pp. 67-87.
- Suc, J.-P., Popescu, S.-M., Fauquette, S., Bessedik, M., Jiménez-Moreno, G., Bachiri Taoufiq, N., Zheng, Z., Médail, F, Klotz, S. 2018. Reconstruction of Mediterranean flora, vegetation and climate for the last 23 million years based on an extensive pollen dataset. Ecologia Mediterranea, 44, 53-85.
- Svenning, J.-C. 2003. Deterministic Plio-Pleistocene extinctions in the European cool-temperate flora. Ecology Letters 6, 646-653.

- Thatoi, H., Samantaray, D., Das, S.K. 2016. The genus *Avicennia*, a pioneer group of dominant mangrove plant species with potential medicinal values: a review. Frontiers in Life Science, 9, 267–291.
- Tomlinson, P.B. 2016. The Botany of Mangroves. Cambridge University Press.
- Utescher, T., Bruch, A.A., Erdei, B., François, L., Ivanov, D., Jacques, F.M.B., Kern, A.K., Liu, Y.-S.C., Mosbrugger, V., Spicer, R.A. 2014. The coexistence approach—theoretical background and practical considerations of using plant fossils for climate quantification. Palaeogeography, Palaeoclimatology, Palaeoecology 410, 58–73.
- Vieira, M., Zetter, R., Grímsson, F., Denk, T. 2023. Niche evolution versus niche conservation and habitat loss determine persistense and extirpation in the Late Neogene European Fagaceae. Quaternary Science Reviews 300, 107896.
- Waleed, W.A., Abdel-Maksoud, Y.K., Kanwar, R.S., Sewilam, H. 2025. Mangroves in Egypt and the Middle east: current status, threats, and opprotunities. International Journal of Environmental Science and Technology, 22, 1225-1262.
- Wang, W. 2004. On the origin and development of *Artemisia* (Astearaceae) in the geological past. Botanical Journal of the Linnean Society 145, 331-336.
- Weeks, A., Zapata, F., Pell, S.K., Daly, D.C., Mitchell, J.D., Fine, P.V.A. 2014. To move or to evolve: contrasting patterns of intercontinental connectivity and climatic niche evolution in "Terebinthaceae" (Anacardiaceae and Burseraceae). Frontiers in Genetics 5, 409.
- Westerhold, T., Marwan, N. Drury, A.J., Liebrand, D., Agnini, G., Anagostou, E., Barnet, J.S.K., Bohaty, S.M., De Vleesschouwer, D., Florindo, F., Frederichs, T., Hodell, D.A., Holbourn, A.E., Kroon, D., Lauretand, V., Littler, K., Lourens, L.J., Lyle, M., Pälike, H., Röhl, U., Tian, J., Wilkens, R.H., Wilson, P.A., Zachos, J.C. 2020. An astronomically dated record of Earth's climate and its predictability over the last 66 million years. Science, 369, 1383–1387.
- Wiens, J.J., Ackerly, D.D., Allen, A.P., Anacker, B.L., Buckley, L.B., Cornell, H.V., Damschen, E.I., Davies, J.T., Grytnes, J.-A., Harrison, S.P., Hawkins, B.A., Holt, R.D., McCain, C.M., Stephens, P.R. 2010. Niche conservatism as an emerging principle in ecology and conservation biology. Ecology Letters 13, 1310-1324.
- Williams, J.W., Jackson, S.T. 2007. Novel climates, no-analog communities, and ecological surprises. Frontiers in Ecology and the Environment 5, 475-482.
- You, Y., Hubern, M., Müller, R. D., Poulsen, C. J., Ribbe, J. 2009. Simulation of the Middle Miocene climate optimum. Geophysical Research Letters, 36, L04702.
- Zagwijn, W.H. 1960. Aspects of the Pliocene and early Pleistocene vegetation in The Netherlands. Mededelingen van de Geologische Stichting C 3, 1-78.
- Zhang, J., Lenz, O.K., Wang, P., Hornung, J. 2021. The Eco-Plant model and its implication on Mesozoic dispersed sporomorphs for Bryophytes, Pteridophytes, and Gymnosperms. Review of Palaobotany and Panlynology 293, 104503.
- Zhang, Z.-Y., Cheng, D.-M., Li, C.-S., Hu, W., Zhan, X.-H., Ji, H.-L. 2019. The complexity of climate reconstructions using the coexistence approach on Qinghai-Tibetan Plateau. Journal of Palaeogeography 8, 5.