



THIS MANUSCRIPT HAS BEEN SUBMITTED TO THE JOURNAL OF GLACIOLOGY AND HAS NOT BEEN PEER-REVIEWED.

### **Ice dynamic and hydrological response to ice-dammed lake drainages at Isunnguata Sermia, West Greenland**

Journal:	<i>Journal of Glaciology</i>
Manuscript ID	Draft
Manuscript Type:	Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Livingstone, Stephen; The University of Sheffield, School of Geography and Planning</p> <p>Storror, Robert; Sheffield Hallam University, Department of the Natural and Built Environment</p> <p>Doyle, Samuel; The University of Sheffield; Aberystwyth University</p> <p>Thorpe, Sian; The University of Sheffield, School of Geography and Planning</p> <p>Moffatt, Angus; The University of Sheffield, School of Geography and Planning</p> <p>Sole, Andrew; The University of Sheffield, School of Geography and Planning</p> <p>Chudley, Thomas ; Durham University</p> <p>Gimbert, Florent; University of Grenoble Alpes, CNRS, IRD, Institut des Géosciences de l'Environnement (IGE)</p> <p>Graly, Joseph; Northumbria University,</p> <p>Licht, Kathy; Indiana University Indianapolis</p> <p>Winter, Kate; Northumbria University, Geography</p> <p>Jayarapu, Asini; Indiana University Indianapolis</p> <p>Bagshaw, Elizabeth; University of Bristol</p> <p>Barruol, Guilhem; University of Grenoble Alpes, CNRS, IRD, Institut des Géosciences de l'Environnement (IGE)</p> <p>Bauer, Kayla; Indiana University Indianapolis</p> <p>Bianchi, Gianluca; Cardiff University, School of Earth and Environmental Sciences</p> <p>Buzzard, Sammie; Northumbria University</p>

	<p>Clason, Caroline; Durham University, Department of Geography</p> <p>Craw, Lisa; Cardiff University, School of Earth and Environmental Sciences</p> <p>Davison, Benjamin; The University of Sheffield, School of Geography and Planning</p> <p>Edwards, Laura; Liverpool John Moores University</p> <p>Gilhooly, Bill; Indiana University Indianapolis</p> <p>Hamilton, Trinity; University of Minnesota College of Biological Sciences</p> <p>Hansen, Chris; University of Minnesota College of Biological Sciences</p> <p>Hawkins, Jonathan; Cardiff University, School of Earth and Environmental Sciences</p> <p>Ing, Ryan; university of edinburgh, School of Geosciences</p> <p>Jatta, Moses; Indiana University Indianapolis</p> <p>Jones, Andrew; Sheffield Hallam University, Department of the Natural and Built Environment</p> <p>Kennedy, Tori; Indiana University Indianapolis</p> <p>Killingbeck, Siobhan; Swansea University</p> <p>Le Bris, Tifenn; University of Grenoble Alpes, CNRS, IRD, Institut des Géosciences de l'Environnement (IGE)</p> <p>McCerery, Rebecca; Northumbria University, Geography and Environmental Sciences</p> <p>Messerli, Alex; Asiaq Greenland Survey</p> <p>Michel, Alexandre; University of Grenoble Alpes, CNRS, IRD, Institut des Géosciences de l'Environnement (IGE)</p> <p>Napoleoni, Felipe; university of edinburgh, School of Geosciences</p> <p>Peacey, Matthew; Aberystwyth University</p> <p>Prior-Jones, Michael; Cardiff University, School of Earth and Ocean Science</p> <p>Ross, Neil; Newcastle University, School of GPS</p> <p>Veness, Remy; Sheffield Hallam University, Institute of Social Sciences</p> <p>Woodie, Kayla; Indiana University Indianapolis</p> <p>Young, Tun Jan; University of St Andrews</p> <p>Hepburn, Adam; Aberystwyth University, Department of Geography and Earth Sciences</p> <p>Booth, Adam; University of Leeds, School of Earth and Environment</p>
Keywords:	Glacier hydrology, Jökulhlaups (GLOFs), Subglacial processes
Abstract:	<p>Ice-marginal lakes are increasingly common around Greenland and are important for modulating glacier runoff and dynamics. This study investigates the evolution of a ~3 km<sup>2</sup> and up to ~100 m deep ice-dammed lake at Isunnguata Sermia, West Greenland. Satellite observations between 1987 and 2024, and field observations of a 2023 drainage using passive seismics, GNSS and time-lapse imagery reveal that the lake drains subglacially and has undergone 12 fill-drain cycles since 1987, a drainage periodicity of 1-3 years. Peak lake volume has decreased since 2010, associated with glacier thinning. Lake drainage can perturb the wider subglacial hydrology system, including triggering the release of stored subglacial water along the flood path in 2019. During the extreme melt year of 2012, the lake drained but did not refill, suggesting that subglacial leakage under the ice dam was sustained by record runoff. Transient ice flow acceleration was observed during the late season drainage in 2023 when the subglacial hydrological system was less efficient and therefore more easily overwhelmed. Our results indicate that ice-dammed lake fill-drain cycles, and the downstream impact on subglacial hydrology and ice dynamics, are modulated by ice dam thickness, melt supply and the antecedent subglacial hydraulic capacity.</p>

SCHOLARONE™  
Manuscripts

**Ice dynamic and hydrological response to ice-dammed lake drainages at Isunnguata Sermia, West Greenland**

Stephen J. Livingstone<sup>1</sup>, Robert D. Storrar<sup>2</sup>, Samuel H. Doyle<sup>1,3</sup>, Sian Thorpe<sup>1</sup>, Angus Moffatt<sup>1</sup>, Andrew J. Sole<sup>1</sup>, Thomas R. Chudley<sup>4</sup>, Florent Gimbert<sup>5</sup>, Joseph A. Graly<sup>6</sup>, Kathy Licht<sup>7</sup>, Kate Winter<sup>6</sup>, Asini Jayarapu<sup>7</sup>, Elizabeth A. Bagshaw<sup>8</sup>, Guilhem Barruol<sup>5</sup>, Kayla Bauer<sup>7</sup>, Gianluca Bianchi<sup>9</sup>, Sammie Buzzard<sup>6</sup>, Caroline C. Clason<sup>4</sup>, Lisa Craw<sup>9</sup>, Benjamin Davison<sup>1</sup>, Laura A. Edwards<sup>10</sup>, Bill Gilhooly<sup>7</sup>, Trinity Hamilton<sup>11</sup>, Chris Hansen<sup>11</sup>, Jonathan Hawkins<sup>9</sup>, Ryan Ing<sup>12</sup>, Moses Jatta<sup>7</sup>, Andrew H. Jones<sup>2</sup>, Tori Kennedy<sup>7</sup>, Siobhan Killingbeck<sup>13</sup>, Tifenn Le Bris<sup>5</sup>, Rebecca McCerery<sup>6</sup>, Alexandra Messerli<sup>14</sup>, Alexandre Michel<sup>5</sup>, Felipe Napoleoni<sup>12</sup>, Matthew W. Peacey<sup>3</sup>, Michael R. Prior-Jones<sup>9</sup>, Neil Ross<sup>15</sup>, Remy Veness<sup>2</sup>, Kayla Woodie<sup>7</sup>, Tun Jan Young<sup>16</sup>, Adam Hepburn<sup>3</sup>, Adam Booth<sup>17</sup>

Corresponding author: [s.j.livingstone@sheffield.ac.uk](mailto:s.j.livingstone@sheffield.ac.uk)

- 1School of Geography and Planning, University of Sheffield, Sheffield, United Kingdom
- 2Geography, Environment and Planning, Sheffield Hallam University, Sheffield, United Kingdom
- 3Centre for Glaciology, Department of Geography and Earth Sciences, Aberystwyth University, Aberystwyth, United Kingdom
- 4Department of Geography, Durham University, Durham, United Kingdom
- 5Univ. Grenoble Alpes, CNRS, INRAE, IRD, Grenoble INP, IGE, 38000 Grenoble, France
- 6Department of Geography and Environmental Sciences, Northumbria University, Newcastle upon Tyne, United Kingdom
- 7Earth and Environmental Sciences, Indiana University Indianapolis, Indianapolis, United States
- 8School of Geographical Sciences, University of Bristol, Bristol, United Kingdom
- 9School of Earth and Environmental Sciences, Cardiff University, Cardiff, United Kingdom
- 10School of Biological and Environmental Sciences, Liverpool John Moores University, Liverpool, United Kingdom
- 11College of Biological Sciences, University of Minnesota, Minneapolis, United States
- 12School of Geosciences, Edinburgh University, Edinburgh, United Kingdom
- 13Department of Geography, Faculty of Science and Engineering, Swansea University, United Kingdom
- 14Asiaq Greenland Survey, Nuuk, Greenland
- 15School of Geography, Politics and Sociology, Newcastle University, Newcastle upon Tyne, United Kingdom
- 16School of Geography & Sustainable Development, University of St Andrews, St Andrews, United Kingdom
- 17School of Earth and Environment, University of Leeds, Leeds, United Kingdom

**ABSTRACT**

Ice-marginal lakes are increasingly common around Greenland and are important for modulating glacier runoff and dynamics. This study investigates the evolution of a ~3 km<sup>2</sup> and up to ~100 m deep ice-dammed lake at Isunnguata Sermia, West Greenland. Satellite observations between 1987 and 2024, and field observations of a 2023 drainage using passive seismics, GNSS and time-lapse imagery reveal that the lake drains subglacially and has undergone 12 fill-drain cycles since 1987, a drainage periodicity of 1-3 years. Peak lake volume has decreased since 2010, associated with glacier thinning. Lake drainage can perturb the wider subglacial hydrology system, including triggering the release of stored subglacial water along the flood path in 2019. During the extreme melt year of 2012, the lake drained but did not refill, suggesting that subglacial leakage under

the ice dam was sustained by record runoff. Transient ice flow acceleration was observed during the late season drainage in 2023 when the subglacial hydrological system was less efficient and therefore more easily overwhelmed. Our results indicate that ice-dammed lake fill-drain cycles, and the downstream impact on subglacial hydrology and ice dynamics, are modulated by ice dam thickness, melt supply and the antecedent subglacial hydraulic capacity.

## INTRODUCTION

Ice-marginal lakes form as water becomes dammed behind ice or moraines, or trapped in overdeepened basins. Over 3300 ice-marginal lakes exist around the fringe of the Greenland Ice Sheet (GrIS) (How and others, 2021), constituting ~10% of the ice margin (Carrivick and others, 2022). The number of ice-marginal lakes in Greenland has increased over the last three decades, likely in response to enhanced meltwater runoff and glacier recession (Carrivick and Quincey, 2014; How and others, 2021). This is significant as lakes trap sediment and nutrients, delay the transfer of water to the ocean (Carrivick and Tweed, 2013), and can accelerate glacier recession and mass loss (Kirkbride, 1993; King and others, 2019; Sutherland and others, 2020; Mallalieu and others, 2021).


Ice-dammed lakes can drain catastrophically, producing high-magnitude floods, known as glacial lake outburst floods (GLOFs) or jökulhlaups (e.g., Thórarinnsson, 1939; Liestøl, 1956). GLOFs have been well studied around the world (see Lützow and others, 2023 for a global database) due to the hazard they pose to communities (e.g., Post and Mayo, 1971; Haeberli, 1983; Carrivick and Tweed, 2016), their downstream ecological and geomorphological impact (e.g. Russell and others, 2006, 2011; Tomczyk and others, 2020) and their potential to modify ice dynamics (e.g., Einarsson and others, 2016). Ice-dammed lakes slowly (months to years) accumulate meltwater and rainfall before rapidly (days to weeks) draining due to unstable failure of the ice dam. Potential dam failure mechanisms include dam flotation (Thórarinnsson, 1939; Glen, 1954), syphoning by the subglacial drainage system (Whalley, 1971), breaches between the ice dam and bedrock (Haeberli, 1983; Walder and Costa, 1996) and Darcian flow through basal sediments (Fowler and Ng, 1996).

During the rising flood stage of a subglacial GLOF, hydrographs are typically characterised by exponentially- ('slow') or linearly- ('fast') rising discharge. Slow drainage is thought to be controlled by the opening of a subglacial channel due to a positive feedback between the rate of frictional melting of its ice walls and frictional heat generated by the turbulence of the water flowing through it (Nye, 1976). Fast lake drainage is caused by the initial hydraulic uplift of the ice by a sheet flood rather than drainage through a discrete conduit(s) (Björnsson, 1992; Flowers and others, 2004; Einarsson and others,

2017). The hydraulic gradient driving water flow is controlled by the lake level and therefore decreases as the lake drains, eventually resulting in rapid termination as channel closure by ice deformation exceeds melt opening (Nye, 1976). This resealing of the ice dam allows water to begin accumulating again, consequently leading to cyclic fill-drain behaviour. Temporal trends suggest that over time the magnitude of GLOFs from ice-dammed lakes globally have decreased (Rick and others, 2023; Veh and others, 2023), consistent with glacier mass loss reducing the thickness of the ice dam and thus the maximum lake level (Thórarinnsson, 1939; Matthews and Clague, 1993; Evans and Clague, 1994).

GLOFs are relatively under-studied in Greenland compared to other glaciated regions of the world, such as Iceland and the Himalayas (Lützow and others, 2023) and there are limited multi-decadal records, which are important for determining controls on flood timing and size, as well as drainage mechanisms (Ng and Liu, 2009). However, they are likely to be common events, with a recent study identifying 326 lakes in Greenland experiencing a total of 541 GLOFs between 2008 and 2022 (Dømgaard and others, 2024). This includes rapid (days) drainage of large ( $>1\text{ km}^3$ ) volumes of water (e.g., Kjeldsen and others, 2017; Grinsted and others, 2017; Carrivick and Tweed, 2019). In this paper, we investigate the evolution of a  $\sim 3\text{ km}^2$  ice-dammed lake adjacent to the northern margin of Isunnguata Sermia, West Greenland (Fig. 1). This lake was included in the GLOF inventory of Dømgaard and others, (2024), with drainages detected in satellite imagery in 2010, 2020 and 2022, but based on only 18 observations spanning 2008-2022. We build on this work, using optical imagery and time-stamped Digital Surface Models (DSMs) to comprehensively reconstruct lake-level evolution since 1987. This analysis allows us to determine controls on flood frequency and magnitude, and its downstream impacts. Field measurements are used to analyse the hydraulic and glaciodynamic processes of the most recent GLOF in 2023.





**Figure 1.** Location of Isunnguata Sermia, West Greenland (blue star in inset map) and ice-dammed lake at the northern margin of the ice tongue. Region of Interest (ROI) 1 (black circle) is used to calculate median ice thickness change of the glacier downstream of the ice-dammed lake from ArcticDEM strips. Cross profile x-x' is used to calculate ice-surface elevation profiles for Fig. 4a. ROIs D1-3 (downstream of the lake) and U1 (upstream of the lake) are used to calculate velocities from feature tracking (see Methods). Blue dashed polygons represent ice-surface elevation anomalies. SGL1-3 are anomalies (expected to represent subglacial lakes) previously reported in Livingstone et al., (2019). Anomalies 1-3 are reported herein. Scientific instruments were installed in summer 2023 and captured the GLOF that occurred in the autumn of the same year. Time-lapse cameras are labelled 1-6. Background is Sentinel-2 optical imagery (2024-08-13). Note river inflow (blue arrow) at the eastern end of the lake, originating from under Isunnguata Sermia.

## DATA AND METHODS

### Ice-dammed lake evolution - Remote Sensing (1987-2024)

To evaluate the long-term evolution of the ice-dammed lake, shorelines were manually digitized from 313 optical satellite images and time-stamped ArcticDEM DSM strips between 1987 and June 2024. USGS/NASA Landsat 5-8 (30 m resolution), ASTER (15 m resolution) and Sentinel-2 (10 m resolution) imagery was digitized using the Google Earth Engine Digitisation Tool (GEEDiT) v. 2.03 (Lea, 2018). This was supplemented by images from Planet Labs that were acquired using the 3 m resolution PlanetScope instrument. Time-stamped ArcticDEM DSM strips at 2 m resolution (Porter et al., 2023) were extracted and co-registered using the *pDEMtools* python package (Chudley and Howat, 2024). Co-registration was performed against the ArcticDEM mosaic following Nuth and Kääb (2011), with stable ground identified using the BedMachine v5 (Morlighem

and others, 2022) land mask. Shorelines for both ArcticDEM strips and Planet imagery were manually digitized in QGIS v3.34. Shoreline digitization involved mapping the boundary between water and land for regions where the edge of the lake could be clearly discriminated. Images obscured by clouds or where the presence of icebergs made it difficult to identify the shoreline were excluded from the dataset. Shoreline elevations were extracted from the bathymetry determined using the co-registered 2019-08-11 ArcticDEM DSM strip when the lake was fully drained.

Median elevations were calculated for each time-stamped lake-shoreline. The uncertainty in shoreline-derived lake levels was calculated from the standard deviation ( $1\sigma$ ) of elevation values, with the lower image resolutions typically resulting in greater uncertainty (mean uncertainty:  $\pm 9.4$  m for Landsat;  $\pm 5.4$  Planet;  $\pm 4.7$  for ArcticDEM). Lake shoreline elevations were converted to volume using the *Surface Water Storage* plugin in QGIS, based on the 2019-08-11 ArcticDEM DSM strip, producing a scaling relationship between lake shoreline elevation and lake volume (Fig. S1). Volume uncertainties were calculated by applying the elevation-volume scaling relationship to the standard deviation of elevation values. We do not account for errors associated with variations in glacier front position and stranded icebergs in the ArcticDEM DSM strip. Freeboard was calculated by manual picking of the ice surface elevation at the front of the calving face, along a transect orientated normal to the ice margin and aligned with the centre of the lake. The minimum value along this transect was used to identify the lake surface height, which was subtracted from the ice surface height.

The downstream impacts and footprint of each GLOF were investigated using optical satellite imagery in GEEDiT. This included identifying large calving events linked to, or preceding, the GLOF, coincident drainages of smaller downstream ice-dammed lakes, and flood limits of the proglacial braided river system.

To investigate the coupling between lake fill-drain cycles and ice-surface elevation change, median and median absolute difference (MAD) ice surface elevation was calculated. This was done within a 500 m diameter region of interest ~500 m upstream of the main lake calving front for 51 co-registered ArcticDEM DSM strips between 2009 and 2023 (Fig. 1, ROI1). Ice surface elevation profiles across the main calving front (Cross-profile x-x' in Fig. 1) were used to extract changes in ice surface, lake height and freeboard of the ice-front from the ArcticDEM.

Ice-surface elevation anomalies (Fig. 1, anomalies 1-3) occurred at the same time as some lake drainage events. Following Livingstone and others (2019), relative elevation changes for each anomaly were calculated by subtracting the mean ice-surface elevation of the anomaly from the mean elevation of a 500 m buffer around it. This approach



182 allowed the dynamic effect to be isolated from longer term thinning signals, and to remove  
183 the influence of any vertical and horizontal offsets between DSMs.

#### 185 **Ice surface motion (2016-2024)**

186 Ice-surface velocity was estimated from feature and speckle tracking of Sentinel-1a  
187 (2014/4 - present) and Sentinel-1b (2016/4 - 2021/12) Interferometric Wide swath mode  
188 Single-Look Complex Synthetic Aperture Radar (SAR) amplitude images. We used  
189 approximately 2000 Sentinel-1a/b 12-day repeat-pass image pairs between January  
190 2016 and May 2025. Prior to tracking, Sentinel image pairs were focused and co-located  
191 using the Generic Mapping Toolbox for SAR imagery (GMTSAR; Sandwell et al., 2011a,  
192 2011b). Each image was split into many overlapping 'image patches', with tracking of  
193 each image patch undertaken in MATLAB, within a version of PIVsuite (Thielicke and  
194 Stamhuis, 2014; <https://uk.mathworks.com/matlabcentral/fileexchange/45028-pivsute>)  
195 adapted for ice flow (Tuckett and others, 2019; Davison and others, 2020). Ice velocity  
196 data from all coincident image pairs were filtered for anomalous data, mosaicked, co-  
197 located and stacked. The stacked data were smoothed using a 3x3 pixel moving median  
198 kernel and resampled to 200 m. Additional filtering of the stacked data based on temporal  
199 variations in velocity magnitude and flow direction removed remaining spurious estimates.

200 Time series of ice motion were created by taking the median value from within regions of  
201 interest (Figure 1); one upstream of the lake drainage (U1) and three downstream (D1-  
202 3). All 12-day velocity estimates from within each ROI were smoothed using a second  
203 degree 24-day Savitsky-Golay filter. Analysis of apparent motion over non-moving  
204 bedrock areas indicates average uncertainties of 20 - 50 m/yr.

#### 205 **Field measurements of 2023 lake drainage**

206 During late September and early October 2023 a range of sensors placed on and in-front  
207 of Isunnguata Sermia recorded a GLOF associated with subglacial drainage of the ice-  
208 dammed lake.

#### 210 ***Time-lapse camera imagery and proglacial river stage***

211 Six SLTL-6210MC Plus 12 Megapixel Scouting time-lapse cameras were installed near  
212 the margin of Isunnguata Sermia (Fig. 1) to monitor subglacial upwelling and proglacial  
213 accreted ice between September 2022 and April 2024. Each camera took 4 repeat photos  
214 per day from 11:00 – 14:00; each photo is time- and temperature-stamped.

215 Time lapse imagery from camera 3 (Fig.1) was used to find the relative stage of a section  
216 of the braided proglacial river during the ice-dammed lake drainage. The Segment  
217 Anything Model (SAM; Kirillov and others, 2023) was used to create a mask of the river  
218 for each time lapse image. From this, a mean average height of the wetted shoreline was  
219 calculated in the region where the shoreline is against the quasi-vertical bank, thus

providing the most representative estimate of river surface height (Fig. S2). The shoreline position was then converted from pixels to metres with respect to a reference measurement taken when the camera was deployed. Stage is displayed relative to the lowest stage during this period.

### ***Ice surface motion***

Time series of horizontal and vertical ice surface motion were determined from dual frequency (L1 + L2) Global Navigation Satellite System (GNSS) data recorded by a Leica GS10 receiver at 0.1 Hz. GNSS station C was located at 67.186N, -50.203W, 2.0 km south of the ice-dammed lake and 5.8 km east of the glacier terminus (Fig. 1). The GNSS antenna was mounted on a 6-m-long aluminium pole drilled 5 m into the ice surface. Rapid re-freezing of the hole secured the antenna pole within the ice. Data were post-processed kinematically (King, 2004) relative to a bedrock-mounted reference station using the differential carrier-phase positioning software Track v1.53 (Chen, 1998). The baseline length was 4.0 km and precise ephemerides from the International GNSS Service were used (Dow and others, 2009). Positioning uncertainties were estimated at ~0.01 m in the horizontal and ~0.02 m in the vertical by calculating the standard deviation of the residuals from linear regression applied to the position time series during a month-long period of steady ice motion in November 2023. Small (< 5 min) gaps in the position record were linearly interpolated before a second-order 6 h low-pass Butterworth filter was applied. The position record was then resampled to 10 minute medians and differentiated to calculate velocity, which was further filtered using a 6 h moving average. Daily mean velocities were calculated from the filtered 10-min interval velocity record. To prevent phase shifts, phase preserving filters, differentiation, and resampling methods were used throughout.

### ***Passive seismic***

A Digos DATA-CUBE<sup>3</sup> Type 2 data logger with a 4.5 Hz 3-component geophone sampling at 100 Hz was installed adjacent to the proglacial river (Fig. 1) to measure seismic tremor generated by river discharge. This approach follows previous studies which found a correlation between seismic noise in the 2 – 10 Hz frequency range and water discharge, attributed to turbulent flow in the river channel (Schmandt and others, 2013; Díaz and others, 2014; Gimbert and others, 2014). Seismic power was calculated using Welch's averaging method (Welch, 1967) applied to the vertical component of ground motion, on 3 s time windows with 50% overlap. Power was averaged over 60 s time windows.

### ***Surface melt estimates***


To estimate daily surface runoff (melt plus rain) at GNSS-C we used COSIPY - a COupled Snowpack and Ice surface energy and mass-balance model in PYthon (see Sauter and others, 2020). We forced this model with data from the PROMICE KAN\_L

automatic weather station situated at ~630 m a.s.l. approximately 14 km SE of the ice-dammed lake (Fausto and others, 2021) and precipitation data from the nearest ERA5 Land reanalysis model grid cell (Muñoz Sabater and others, 2019). Air temperatures were interpolated to the elevation of GNSS-C using a constant lapse rate of  $-5.3\text{ }^{\circ}\text{C km}^{-1}$  and air pressure was interpolated using the barometric equation.

## RESULTS

### Ice-dammed lake evolution (1987-2024)


Shoreline mapping of the ice-dammed lake revealed 12 fill-drain cycles between 1987 and 2024 (Fig. 2). Prior to the launch of the Landsat 7 satellite in 1999, data are sparser with at least one GLOF likely missed between 1987, when the lake was relatively full ( $0.164 \pm 0.044\text{ km}^3$ ), and 1993 when the lake was filling from a lower level ( $0.077 \pm 0.035\text{ km}^3$ ). Lake evolution is characterised by slow filling over 1-3 years (mean: 2.8 years), with satellite imagery indicating recharge from water emanating from under ice flowing into the eastern side of the lake (Fig. 1). Filling is followed by rapid subglacial drainage beneath the southwest ice-contact boundary of the lake. There is a clear seasonal pattern of filling; lake levels typically rise by 40-50 m during the summer melt season (June-August), with slower filling towards the start and end, while the winter (October-March) is characterised by no significant (i.e. within error) or slow increases in water level (Fig. 2).



**Figure 2.** Ice-dammed lake volume evolution (1987-2024) based on shorelines mapped using optical imagery and the time-stamped ArcticDEM strips, and application of an elevation-volume scaling relation (see methods and Figure S1). Note that 1987 to 1993 is poorly constrained (black dashed line). See Methods section for details of error bars.


Lake drainages occur throughout the melt season, ranging from May-June (e.g., 2012) through to late September / early-October (e.g. 2023), but with the majority (at least 6) occurring between July and August when melt tends to be greatest. The lake volume at which drainage occurs varies, but with a statistically significant negative relationship (Pearson  $r = -0.71$ ,  $p < 0.05$ ) through time (Fig. 3a, Fig. S3). Up to 2010, lake volumes reached  $>0.176 \text{ km}^3$  ( $>365 \text{ m a.s.l.}$ ) before drainage occurred, with a peak lake volume of  $0.218 \pm 0.04 \text{ km}^3$  (218 million cubic metres) recorded during the 2010 drainage. Since 2010, 50% of the drainages have occurred at lake volumes  $<0.176 \text{ km}^3$ , with the 2019 and 2023 levels peaking at just  $0.151 \pm 0.019 \text{ km}^3$  and  $0.111 \pm 0.01 \text{ km}^3$  respectively. This  $\sim 0.05 \text{ km}^3$  reduction in volume and 20 m drop in maximum lake level since 2010 corresponds with a  $\sim 10 \text{ m}$  reduction in median ice-surface elevation within ROI1, dropping from  $\sim 413 \text{ m a.s.l.}$  in 2010 to  $\sim 403 \pm 4.9 \text{ m a.s.l.}$  in 2023 (Figs. 3a and 4a). Within  $\sim 200 \text{ m}$  of the calving-front, Isunnguata Sermia responds to changes in lake level (Fig. 4a), lifting up to 37 m as the lake becomes full and the ice likely begins to float (Fig. 4b). The freeboard height during periods when the lake is full and the ice-front is floating is 12-16 m, which corresponds to a maximum ice thickness of 145-193 m assuming hydrostatic equilibrium, an ice density of  $917 \text{ kg m}^{-3}$  and water density of  $1000 \text{ kg m}^{-3}$ . The lake basin is completely or nearly emptied during GLOFs, with the lake level dropping around 100 m and the freeboard height increasing (Fig. 4b). The denser temporal record of satellite observations from 2009 onwards has enabled 6 GLOF events to be constrained to  $<10$  days, with the 2022 event occurring in just 4 days or less (Table 1), with a minimum mean discharge of  $480 \text{ m}^3 \text{ s}^{-1}$ .

The downstream ice velocity response to drainage of the ice dammed lake is variable (Fig. 5). Of the four drainage events within the window of the Sentinel-1 ice velocity data (2016-2025), there was no discernable impact on ice flow during the 2017 and 2019 events, but a relative increase in ice flow at downstream sites during the 2022 and 2023 events. The 2022 drainage event was associated with a 20-60% speed-up at downstream ROIs compared to the upstream ROI, whilst the 2023 event coincided with a 50-110% relative acceleration, which was the largest GLOF and non-GLOF-related velocity anomaly between 2016-2025.




**Figure 3.** (a) Time series of ice-dammed lake volume (black line; symbols are the same as Figure 2) and median ice-surface elevation (orange stars, ROI1 in Fig 1) from 2009 to 2024. See Methods Section for details of error bars. (b-d) Time series of mean relative ice surface elevation from 2009 to 2024 for three ice-surface elevation anomalies on Isunnguata Sermia (see Fig. 1) (b-d). (b) Anomaly 1; (c) Anomaly 2; (d) Anomaly 3. Vertical dotted lines mark drainage events of the ice-dammed lake visible in (a).





**Figure 4.** (a) Cross-profile showing ice-lake interactions at the calving front from 2009-2022 based on ArcticDEM DSM strips along profile x-x' in Fig. 1. Note the overall pattern of declining elevation (ice thinning) over the 13-year period, with ~10 m thinning furthest upglacier and nearly 40 m thinning within 100 m of the lake. Within 200 m of the lake, the ice surface is uplifted when the lake is filled. (b) Lake height, ice-height and freeboard (difference between the two) calculated from the ArcticDEM DSM (2009-2024). Note that not all drainage events are captured by the ArcticDEM alone.





**Figure 5.** (a) Ice velocity temporal anomaly compared to the whole time period mean for three ROIs (see Figure 1 for locations). U1 is upglacier and D1-3 are downglacier from the ice dammed lake. Vertical blue lines and labels mark the first satellite record of each ice-dammed lake drainage (see Table 1). Subplot (b) shows the difference between the ice-velocity anomalies for D1-3 and U1. Note that both the 2022 and 2023 lake drainage events coincide with a large positive anomaly indicating a relative speed up of the downstream ROIs. The raw time series for each ROI are shown in Figure S4.

**Table 1.** Dates of ice-dammed lake drainage based on satellite imagery (yyyy-mm-dd). This includes the dates immediately pre-, during and post-drainage, and associated lake volumes and water levels. The drainage duration is the number of days between the pre- and post-drainage imagery. Drainage durations were only noted where they can be reasonably constrained by satellite imagery and the ArcticDEM. The satellite imagery used to identify the lake shorelines is noted in brackets. L = Landsat and S = Sentinel.

Drainage dates and source	Lake volume (km <sup>3</sup> ) and [lake level (m a.s.l.)]
---------------------------	--

Pre-drainage	Drainage & post-drainage	Pre-drainage	Post-drainage	Drainage duration
1994-07-12 (L5)	1995-05-05 (L5)	$0.201 \pm 0.065$ [373 $\pm$ 9.3]	$0.015 \pm 0.008$ [291 $\pm$ 4.7]	
2000-09-17 (L7)	2001-03-17 (L7)	$0.191 \pm 0.072$ [370 $\pm$ 10.6]	$0.005 \pm 0.013$ [275 $\pm$ 12.9]	
2003-08-05 (ASTER)	2003-09-24 (L7)	$0.205 \pm 0.014$ [374 $\pm$ 2.0]	$0.011 \pm 0.008$ [285 $\pm$ 5.8]	
2006-05-11 (L7)	2006-07-05 (ASTER) 2006-07-21 (ASTER)	$0.176 \pm 0.047$ [365 $\pm$ 7.1]	$0.015 \pm 0.005$ [291 $\pm$ 2.7] $0.001 \pm 0.005$ [262 $\pm$ 10.3]	
2007-08-27 (L7)	2007-09-12 (L7)	$0.167 \pm 0.045$ [363 $\pm$ 6.9]	$0.010 \pm 0.010$ [284 $\pm$ 7.1]	
2010-07-02 (L7)	2010-07-10 (ArcticDEM) 2010-07-18 (L7)	$0.218 \pm 0.040$ [377 $\pm$ 5.6]	$0.021 \pm 0.008$ [297 $\pm$ 4.1] $0.007 \pm 0.013$ [280 $\pm$ 11.0]	8 days
2012-05-04 (L7)	2012-06-03 (L7)	$0.165 \pm 0.034$ [362 $\pm$ 5.2]	$0.014 \pm 0.012$ [289 $\pm$ 7.1]	
2015-07-24 (L8)	2015-07-31 (L8)	$0.181 \pm 0.044$ [366 $\pm$ 6.5]	$0.002 \pm 0.007$ [266 $\pm$ 10.3]	7 days
2017-08-15 (S2)	2017-08-16 (Planet) 2017-08-21 (S2) 2017-08-24 (S2)	$0.160 \pm 0.026$ [360 $\pm$ 4.1]	$0.096 \pm 0.022$ [338 $\pm$ 4.3] $0.008 \pm 0.006$ [282 $\pm$ 4.7] $0.002 \pm 0.003$ [267 $\pm$ 5.6]	9 days
2019-08-02 (S2)	2019-08-07 (S2) 2019-08-09 (S2) 2019-08-11 (ArcticDEM)	$0.151 \pm 0.019$ [355 $\pm$ 4.7]	$0.031 \pm 0.013$ [306 $\pm$ 5] $0.0003 \pm 0.002$ [259 $\pm$ 4.8] $0.001 \pm 0.001$ [261 $\pm$ 1.6]	5 days
2022-06-30 (S2)	2022-07-02 (S2) 2022-07-03 (L9) 2022-07-04 (S2)	$0.168 \pm 0.009$ [362 $\pm$ 1.4]	$0.068 \pm 0.031$ [327 $\pm$ 7.0] $0.013 \pm 0.012$ [289 $\pm$ 7.3] $0.002 \pm 0.002$ [267 $\pm$ 3.2]	4 days
2023-09-30 (S2)	2023-10-01 (Planet) 2023-10-02 (Planet) 2023-10-05 (Planet)	$0.111 \pm 0.010$ [344 $\pm$ 1.9]	$0.072 \pm 0.017$ [329 $\pm$ 3.7] $0.058 \pm 0.012$ [322 $\pm$ 3] $0.001 \pm 0.005$ [264 $\pm$ 8.9]	5 days


**2019 GLOF - Cascade of downstream subglacial events**

The 2019 GLOF drained 0.12 km<sup>3</sup> over ~5 days, with the lake level dropping 98 m between the 4th and 9th August (Table 1, Fig. 6). This gives a mean discharge of 278 m<sup>3</sup> s<sup>-1</sup>, which is likely to underestimate peak discharge. The foreland showed limited evidence of a change in river stage during and following the ice-dammed lake drainage (Fig. 6). However, the GLOF generated a cascade of downstream subglacial events detected from repeat ArcticDEM DSM data and optical imagery. Downstream of the lake three quasi-circular regions of the ice surface subsided during the same period

(Anomalies 1-3 in Figure 7a). Time series of relative elevation change for each anomaly (Fig. 3b-d) suggest similar behaviours, with subsidence starting between 16th June and 11th August, which was largely complete by 25th August and at its lowest level on 22nd September. Given the subsidence persisted after 11th August when the ice-dammed lake was fully drained, we suggest these events were most likely triggered by the GLOF, although we lack the temporal resolution to confirm this.

Anomaly 1 is located 2.6 km downstream of the ice-dammed lake at the western end of a large overdeepened trough where the ice is ~230 m thick (Fig. 7b). A 0.36 km<sup>2</sup> surface depression with a mean drop of 15 m and maximum drop of 38 m formed between 16th June and 22nd September 2019 (Fig. 3b). Anomalies 2 and 3 are both located within 1.5 km of the glacier terminus, close to the main meltwater outlet and above relatively thin ice (120-130 m). They formed 0.11 and 0.10 km<sup>2</sup> depressions respectively, with mean surface lowering of 5.2 and 5.7 m and maximum surface lowering of 14 and 16 m across the same period (Fig. 3c-d). Across the full time-series, Anomalies 1 and 3 are characterised by periods of slow uplift followed by rapid subsidence, similar to the three ice-surface elevation patterns previously detected further to the east (SGL 1-3 in Figure 1). These were previously interpreted as recording subglacial lake filling and drainage (Livingstone et al., 2019). Anomaly 1 previously underwent a 4.3 m mean elevation drop between 3rd August and 31st October 2011, with crescentic crevasses apparent on the ice surface in 2012. Anomaly 3 also subsided between 31st October 2011 and 14th February 2013, and might therefore be associated with the 2012 GLOF, although the drainage event is poorly constrained. Anomaly 2 is characterised by a long-term trend of subsidence relative to the surrounding ice surface. There is a more subtle (few metres) pattern of subsidence, evident between 6th June 2019 and 22nd September 2019 (Fig. 7a). This subsidence, which follows the W-E axis of the basal overdeepening across Anomaly 1 (Fig. 7b) southwards towards Anomaly 2 and then west to the outflow at the terminus, suggests the wider pattern of subglacial drainage is associated with the anomalies and represents a potential drainage flowpath during GLOFs (arrows on Figure 7a).






**Figure 6.** Cloud-free true colour optical satellite imagery (PlanetScope) of the terminus (left column: note the absence of significant changes to the proglacial river and floodplain) and ice-dammed lake (right column) before (4th August), towards the end of (8th August) and after (9th August) the 2019 drainage event. Images © 2019 Planet Labs PBC.



394



**Figure 7.** (a) Anomaly plot of ice-surface elevation change from 2019-06-16 to 2019-09-22 from differencing ArcticDEM DSMs. Dashed arrows represent the interpreted subglacial drainage pathways based on the pattern of ice-surface subsidence. The long-dashed line emanating from the ice-dammed lake represents the inferred lake drainage pathway based on both the pattern of subsidence and changes in proglacial river stage. Note during the period the ice-dammed lake drained, three areas of the ice surface subsided (anomalies 1-3). (b) Ice thickness based on the Bedmachine v5 DEM (Morlighem et al., 2017).

395

396

397

398

399

400

401

402

403

## 2023 GLOF - remote sensing and field observations

### *Remote sensing observations*


Satellite imagery of the ice-dammed lake shows that it drained  $0.11 \text{ km}^3$  over 5 days between 30th September and 5th October 2023, with the lake level dropping 80 m (Table 1, Fig. 8). This drainage corresponds to a mean discharge of  $255 \text{ m}^3 \text{ s}^{-1}$  across the 5 days, with a mean of  $\sim 471 \text{ m}^3 \text{ s}^{-1}$  across a 23 hour period between 30th September and 1st October. During drainage between the 1st and 2nd October, there was a  $89,669 \text{ m}^2$  calving event from the glacier terminus into the ice-dammed lake. Between the 1st and 2nd October water ponded on the ice surface towards the northern terminus (Fig. 8). The lack of ponding off-ice, the origin at crevasses and the drop in elevation towards the margin, suggest this water rose to the ice surface from the bed through the crevasses during the GLOF.

Increased proglacial river flow is observed in both satellite and time-lapse imagery from 24th September, despite low rainfall and melt, suggesting that lake drainage could have begun earlier (Figs. 8-10). This includes an increase in water levels from 25th September recorded by time-lapse cameras (e.g., Fig. 9), increased activity of the northern proglacial river system and an increase in stage in the southern proglacial river system between the 24th and 30th September (Fig. 8).

Time-lapse photography at  $67.181^\circ\text{N}$   $-50.334^\circ\text{W}$  (camera 4 in Fig. 1) began on 21st September 2023. The camera pointed westward, across a stream between the Isunnguata Sermia terminus and a subglacial upwelling (Fig. 9 and supplementary [video](#)). Here, turbulent subglacial waters emerged into an area of ponded water surrounded by accreted ice, which overlay the outwash plain. The water level recorded by the time-lapse camera slowly fell from 11:00 (local Western Greenland Time) on 21st September until 13:00 on 25th September. One hour later, the water began rising. By 11:00 on 26th September, the water had risen and at 11:00 on 27th September, another increase in water level was visible. During this period, no turbulent upwelling was evident. At 11:00 on 28th September, water overflowed its basin and the point of emerging water was apparent in the same position as 21st September. By 29th September, the emerging waters began erupting higher than the surrounding accreted ice. Additionally, a mound of dislodged accreted ice or frozen outwash appeared in the pond overnight. These mounds disappeared on 30th September and were replaced by other mounds, including a prominent one in the foreground of the image. The eruption height reached a maximum on 1st October and continued through 2nd October. By 11:00 on 3rd October, the water level fell while vigorous eruptions continued. This is consistent with satellite imagery indicating that between the 3rd and 5th October the northern proglacial river had returned to its previous state and the southern river had decreased in stage. In the 11:00




443 photograph on 5th October, the eruption had ceased and ponded water drained, leaving  
444 a landscape of accreted ice.  
445



446




**Figure 8.** Cloud-free true colour optical satellite imagery (PlanetScope) of the terminus (left column) and ice-dammed lake (right column) before (24th September), during (2nd October) and after (5th October) the 2023 drainage event. Images © 2023 Planet Labs PBC.



**Figure 9.** Time-lapse of the upwelling at the terminus of the glacier in 2023. Camera location: 67.181N -50.334W (Fig. 1 - Camera 4). Black circle highlights the point of turbulent upwelling. Pink dashed line is a water level reference line. Arrows on 29th September images highlight the formation and then movement of a mound of dislodged accreted ice or frozen outwash. See link for time-lapse video of the upwelling - <https://vimeo.com/1103612163>.

**Passive seismic observations**

During the first three weeks of September 2023, seismic energy in the frequency range 2-10 Hz gradually decreased at the foreland passive seismic station (Fig. 10a). This frequency range is characteristic of turbulent water flow (Gimbert and others, 2014), and the energy decrease was consistent with reduced runoff and proglacial river discharge in response to colder air temperatures during the late melt season. However, from 26th September, seismic energy began to increase rapidly, peaking around 2-4th October and then rapidly decreasing, with a pronounced period of low seismic energy on 5-6th October that fell below the longer term trend (Fig. 10b). The increase in seismic power was contemporaneous with an increase in river stage (Fig. 10b) suggesting that the record of seismic power can be used as a proxy for river discharge. During the subsequent melt event on 6-7th October, there was no correlation between high seismic energy and river stage (Fig. 10c), potentially reflecting a shift in where water is discharging from under the ice.




**Fig. 10.** (a) The passive seismic record from the forefield of Isunnguata Sermia from 29th August 2023 to 24th October 2023. Location of the terminus passive seismic station is marked in Figure 1. (b) Average seismic power within the 2-10 Hz frequency range (marked by the red lines in panel a). Note, the GLOF is picked out by a rapid increase in seismic energy (red line) starting on 26th September and peaking around 2-4th October (green shading), before rapidly decreasing. From 5-6th October there is a distinct low in seismic energy (orange shading). The change in seismic power is consistent with changes in relative proglacial river stage during the GLOF (blue line). (c). COSIPY daily total melt and rainfall at the GNSS site.

### ***Ice dynamic observations***


At the glacier scale, there was a 50-100 m yr<sup>-1</sup> speed-up of ice flow into the southwestern ice-contact boundary of the ice-dammed lake and across the northern part of the terminus during the 2023 GLOF (Fig. 11, Fig. S5). The latter coincided with the emergence of turbid meltwater at the ice surface (Fig. 8). More generally, there was a slight ~20 m yr<sup>-1</sup> speed-up of the tongue of Isunnguata Sermia downstream of the ice-dammed lake during the period when the lake drained (Fig. 11c). This acceleration was related to an increase in westerly flow, particularly over the central part of the tongue, and greater spreading of the near-terminus (Fig. S5). However, this magnitude of speed-up was similar to areas upstream of the lake, although they tended to have a patchier spatial signal.

The GNSS station, located 1-2 km up-glacier from the inferred drainage route of the ice-dammed lake (Fig. 7a), also recorded evidence of a change in ice dynamics during the 2023 GLOF (Fig. 12). There was a subtle increase in daily average horizontal velocity during the GLOF (from ~75 m yr<sup>-1</sup> to ~80 m yr<sup>-1</sup>) between 23rd September and 3rd October (Fig. 12a), although the initial speed up coincided with a small rainfall event on 24th September (Fig. 12c). This velocity increase was accompanied by a switch from ice-surface uplift to ice-surface subsidence of several centimetres over about 7 days (Fig. 12b). On 4-5th October horizontal velocity increased sharply, peaking at 93 m yr<sup>-1</sup> on 5th October (Fig. 12a,d), and there was a steepening of the subsidence signal (~1 cm/day), which continued until 7th October (Fig. 12b,e). The velocity spike and increased subsidence lagged maximum lake discharge by 1-2 days and coincided with shutdown of the proglacial drainage system (Figs. 8-10). The trajectory of the GNSS also shifted slightly during the GLOF, temporarily trending further northwards towards the ice-dammed lake (Fig. 12d,e) until 7th October. A velocity spike of >100 m yr<sup>-1</sup>, and a switch to ice-surface uplift, on 7-8th October was attributed to a melt and rainfall event (Fig. 12c).





**Fig. 11.** Isunnguata Sermia ice velocity: a. 2016 - 2025 average ice velocity.; b. Average ice velocity for image pairs that include the 2023 ice-dammed lake drainage. Arrows in (b) and (c) are flow direction vectors and magnitudes; c. Ice velocity anomaly for the 2023 ice-dammed lake drainage period compared to the long-term average. Positive values (blue) represent pixels for which ice velocity was greater during the lake drainage period.



**Fig. 12.** Ice surface motion recorded by GNSS C in 2023: a. Daily mean horizontal surface velocity; b. Ice-surface elevation change; c. COSIPY daily total melt and rainfall at GNSS-C; d. The track of the GNSS coloured by 6-hr average horizontal velocity anomaly. e. As for (d) but coloured by elevation deviation relative to 26th September 00:00. The green shade highlights the period of lake drainage and corresponds to the time period of subplots (d) and (e). The dotted lines in (d) and (e) represent an extrapolation of the linear fit from 26-27th September



## DISCUSSION

We have reconstructed the evolution of a large ( $0.218 \pm 0.040 \text{ km}^3$ ) ice-dammed lake at the northern margin of Isunnguata Sermia, West Greenland from remote sensing and field measurements. Our results reveal multiple fill-drain cycles with gradual filling of the lake followed by rapid subglacial drainage over multiple days with a periodicity of 1-3 years. These GLOF events impact downstream and upstream ice dynamics, subglacial hydrology and runoff.


### GLOF magnitude and frequency

Reconstruction of lake-level evolution since 1987 reveals no obvious change in the frequency or annual timing of GLOFs over the 37-year study period, but a  $\sim 0.05 \text{ km}^3$  reduction in maximum lake volume and a  $\sim 20 \text{ m}$  decrease in the maximum lake level since 2010. The general reduction in lake volume since 2010 appears to coincide with glacier thinning, which could have reduced the water pressure in the lake needed to approach or exceed the overburden pressure of the ice dam (Thorarinsson, 1939; Tweed and Russell, 1999). However, this relationship breaks down in some years; for example, the lake reached a greater volume in 2022 ( $0.168 \pm 0.009 \text{ km}^3$ ) than in 2019 ( $0.151 \pm 0.019 \text{ km}^3$ ). This variation might reflect error in the mapping process or the temporal availability of satellite imagery - missing the maximum lake level. Alternatively, changes in geometry might not be the sole cause of the variations in water level, with drainage not always occurring when the ice dam reaches flotation (e.g., Björnsson, 1992; Huss and others, 2007; Veh and others, 2023).

The reduction in lake volume does not lead to a reduction in GLOF frequency (Fig. 3a). This is consistent with recent large-scale studies (see also Veh and others, 2023; Rick and others, 2023). In Figure 13 we explore potential runoff controls on GLOF frequency by summing modelled daily discharge from the main outlet of Isunnguata Sermia (Mankoff et al., 2020) over the recharge period between GLOFs. Although used as a proxy of runoff for lake recharge, we note that the catchment for the ice-dammed lake is likely much smaller than the main outlet. Overall, there is no clear relationship between catchment runoff and lake recharge volume. However, six of the eight recharge periods exhibited similar peak lake volumes ( $0.149\text{-}0.167 \text{ km}^3$ ) and total catchment runoff ( $9.2\text{-}11.6 \text{ km}^3$ ), suggesting a similar rate of filling (Fig. 13).

There are two clear outliers in the fill-drainage cycles. The first is the 2007 to 2010 recharge period, which had the lowest total catchment runoff but an anomalously large ( $0.208 \text{ km}^3$ ) lake recharge volume (Fig. 13). One possible explanation is that our crude proxy of using total runoff from the main Isunnguata Sermia catchment misses changes in area of the lake's catchment, driven by changing ice thickness and water pressure (e.g., Chu et al., 2016).

The other outlier is the 2012 to 2015 recharge period, which coincided with approximately double the catchment runoff ( $23.46 \text{ km}^3$ ) (Fig. 13). This discrepancy can be explained by changes in the stability of the ice dam. The preceding 2012 lake drainage, which occurred relatively early in the melt season (May-June), was not followed by substantial lake filling until 2013 (Fig. 3), despite the record melt that year (Nghiem and others, 2012). This behaviour indicates that during this period the lake experienced a net loss of water, with the most obvious route for increased water loss via the subglacial channel(s) created during the 2012 GLOF. This is consistent with elevated runoff in 2012 (Nghiem and others, 2012) causing melt-enlargement to exceed creep closure of subglacial channels (Röthlisberger, 1972), with the early season GLOF enabling the channel(s) to remain open as surface melt increased thereafter. Further evidence that the ice dam can allow water to escape is provided by slight decreases in lake volume in winter, when outputs must have exceeded inputs. This behaviour is more evident when the lake is fuller and water pressure therefore higher prior to 2015 (Figs. 2-3a) (Gilbert, 1971; Matthews and Clague, 1993).



**Figure 13.** Summed discharge from the main outlet of Isunnguata Sermia (Mankoff and others, 2020) between drainage events of the ice-dammed lake against the change in lake volume over the same period where lake drainages are well constrained. The dates represent the period of lake filling (recharge).

**Lake drainage mechanism: inferences from the 2023 GLOF**

The rapid rising limb and sharp falling limb of the discharge hydrograph, inferred by the seismic and stage records at the terminus in 2023 (Fig. 10), is consistent with channelised drainage and the classical slow outburst flood model (Nye, 1976). However, at Isunnguata Sermia water emerges at multiple points along the terminus, suggesting a more complex multi-path drainage configuration. The spatial and temporal evolution of proglacial drainage and increase in downstream ice motion observed during the 2023 GLOF also suggests the channelised subglacial system was at least temporarily overwhelmed. Similar observations of other slow-rising GLOFs (Huss and others, 2007; Magnússon and others, 2007; Sugiyama and others, 2008) have been linked to water pressure exceeding ice overburden, driving water out of the channel, reducing basal friction and enabling other flowpaths to form (Clarke, 2003). Such short-term reconfiguration of subglacial systems highlights the importance of understanding the role of glacier lake drainage in local hydrologic and ice-dynamic perturbations.

## **Perturbation of the subglacial hydrological system**

### ***Subglacial water release during 2019 GLOF***

The 2019 GLOF coincided with ice-surface subsidence in three downstream regions (Fig. 7a). Anomalies 1 and 3 are interpreted as distinct subglacial lakes based on their surface elevation time series of slow uplift (subglacial lake filling) followed by rapid subsidence (subglacial lake drainage) (e.g., Livingstone and others, 2019). The coincidence of the rapid drainage of these lakes with the 2019 ice-dammed GLOF suggests that GLOFs can trigger secondary releases of stored subglacial water along their flood path, which then contribute to output flood waters (e.g., Huss and others, 2007; Shangguan and others, 2017). Mejia and others (2021) suggest that subglacial flood waves from rapid supraglacial lake drainages can dewater isolated cavities by opening connections to them. Thus, we might expect release of stored water to be linked to GLOFs that have overwhelmed the subglacial drainage system causing vertical uplift and increased sliding, in turn leading to growth of cavities initiating new connections (Mejia and others, 2021). However, the 2019 GLOF occurred during peak melt season when we might expect the additional meltwater discharge from the GLOF to be largely accommodated by an efficient subglacial channelised system (Bartholomew and others, 2010); an interpretation supported by the absence of an observable ice-dynamic response (Fig. 5). An alternative mechanism is that the 2019 GLOF caused an increase in the local hydraulic gradient, possibly due to a pressure wave, or via expansion of low pressure channels, which was sufficient to trigger drainage of water stored subglacially (e.g., Jóhannesson, 2002; Kingslake and Ng, 2013; Dow and others, 2018). More generally, there is no obvious reason why the 2019 GLOF triggered the release of subglacially stored water compared to other years, suggesting a complex response that could also be related to the drainage route, antecedent conditions and stage of the subglacial lakes. This makes the estimation of proglacial discharges challenging.

The lack of a subsequent uplift pattern at Anomaly 2 following subsidence in 2019 suggests that either it was a subglacial lake that didn't refill once drained and therefore a transient feature, or given its position close to the meltwater outlet at the terminus, it could be recording collapse of ice above a subglacial channel (Egli and others, 2021; Hösli and others, 2025). The latter would be consistent with the GLOF causing rapid enlargement of the subglacial channel, thinning the ice roof and making it more prone to collapse. Although Anomaly 3 is even closer to the terminus, tunnel collapse might be favoured at Anomaly 2, for example, if it was over thinner ice or at a location where the hydraulic gradient is steeper resulting in greater turbulent melting of the subglacial channel roof.

### ***Subglacial water storage after the 2023 GLOF***

From 5-6th October, at the end of the 2023 GLOF, there is a distinct reduction in the observed seismic power to below the pre-event background signal, suggesting proglacial water flow is beneath the detectable threshold. This is supported by time-lapse imagery close to the glacier terminus, revealing an abrupt drop in water flow over the same period. This could reflect low melt inputs into the subglacial drainage system during this period, or a wider perturbation of the subglacial hydrology by the GLOF, temporarily inhibiting or changing the configuration of subsequent water flow from the wider catchment to the terminus. This contrasts with other studies that have observed a tail of above-average discharge thought to be associated with the release of stored water (e.g., Anderson and others, 2003; Huss and others, 2007). We postulate that the creation of large low-pressure channels (and potential evacuation of stored water similar to 2019) during the 2023 GLOF reduced the hydraulic gradient, initially inhibiting drainage (Dow and others, 2022) out of the overdeepened trough at Isunnguata Sermia (Lindbäck and others, 2014). Drainage up a reverse slope is supported by active ice accretion in the immediate foreland (Fig. 9), consistent with supercooled water (Cook and others, 2006; Cook and Swift, 2012). We suggest the relatively quiet seismic period from 5-6th October was characterised by re-pressurisation and re-filling of subglacial water stores until there was sufficient water to re-escape the overdeepening.

### ***Ice dynamic response to GLOFs***

At the western ice-water boundary of the ice-dammed lake, the latter stages of lake filling are typically characterised by ice-surface uplift (Fig. 4), indicating that water is penetrating up to ~200 m under the ice dam (either due to flexure or movement along fractures) causing flotation (e.g. Nye, 1976; Anderson and others, 2003; Walder and others, 2005; Bigelow and others, 2020). By neglecting subglacial storage, our lake volume estimates from satellite data therefore underestimate the volume of water. During GLOFs there is further evidence of mechanical damage, with calving of the marginal ice-front recorded by satellite data in multiple years (e.g., Figs. 6,8). This calving is likely driven by a rapid

reduction in the hydrostatic pressure as the lake drains and the ice-front freeboard height increases.

The impact of ice-dammed lake drainage on ice velocity is variable (Fig. 5). Both the 2017 and 2019 GLOFs produced no detectable ice velocity signal in our Sentinel-1 data. These events occurred in August during peak melt season, and therefore we suggest the limited impact on ice motion is because the antecedent hydraulic capacity of the subglacial drainage system was sufficient to accommodate the additional water without spikes in water pressure (e.g., Livingstone and others, 2022). Indeed, there is also no observable change in the proglacial river and flood plain during the 2019 GLOF (Fig. 6) indicating that drainage was largely accommodated by the existing river system. In contrast, both the 2022 and 2023 GLOFs were associated with increased ice-flow downstream of the ice-dammed lake (Figs. 5, 11-12). Compared to 2019 the 2022 GLOF was of greater magnitude (mean discharge of  $480 \text{ m}^3 \text{ s}^{-1}$ ) and occurred earlier in the melt season (beginning of July) when the drainage system would have been more inefficient. It therefore follows that the 2022 GLOF was able to overwhelm the antecedent drainage system and elicit a detectable velocity response (e.g., Iken & Bindenschadler, 1986; Kingslake and Ng, 2013).

The 2023 GLOF produced the largest observed downstream velocity anomaly (Fig. 5). This is consistent with the subglacial drainage system being overwhelmed due to its low antecedent hydraulic capacity after the end of the melt season (October), and following a period of declining melt and proglacial river flow. The larger acceleration of the northern part of the terminus (Fig. 11b-c) suggests the drainage system here was particularly sensitive to melt inputs. This is supported by the outflow of subglacial water to the ice surface through crevasses in this region (Fig. 8), suggesting high subglacial water pressures (Roberts, 2005; Roberts and others, 2010; Bowling and others, 2021). GNSS C, upstream of the ice-dammed lake, records an initial ~6% speed-up of ice flow and a switch from ice-surface uplift to subsidence between 24th September and 3rd October. This signal is consistent with longitudinal extension causing vertical compression of the ice at C, driven by downstream acceleration from increased basal sliding of the lower glacier tongue during the GLOF. The GNSS subsequently shows a sharp increase in ice-velocity and ice-surface subsidence from 4th October (Fig. 12), which lags peak discharge by ~1-2 days and is associated with ice flow deflection towards the ice-dammed lake. This is interpreted to record increased ice flow into the ice-dammed lake (Fig. 11b-c) as it re-adjusts following the rapid lowering of hydrostatic pressure (lake water-level drop) (Riesen and others, 2010), and re-grounding of floating ice causing an increase in driving stress.



## CONCLUSIONS

We studied the evolution of a  $\sim 3 \text{ km}^2$  ice-dammed lake adjacent to the northern margin of Isunnguata Sermia, West Greenland that drains subglacially every 1-3 years. We used remote sensing observations to monitor lake volume changes between 1987 and 2024, and to quantify the downstream subglacial hydrological and ice dynamic impact. In 2023 a GLOF was instrumented with a number of sensors, including GNSS, a passive seismic station and time-lapse cameras deployed on and at the terminus of Isunnguata Sermia.

We demonstrate no clear change in the frequency or annual timing of GLOFs at Isunnguata Sermia, but a reduction in peak lake volume since 2010, likely due to glacier thinning in response to climate change. Predicting future changes in lake volume and drainage timing is complicated by the penetration of a wedge of water under the ice and leakage through the ice dam. In particular, net loss of water through the ice dam during the extreme melt year of 2012 suggests that some ice dams might become more leaky as the climate warms, reducing their potential to fill.

Our results demonstrate the potential for GLOFs to impact downstream ice dynamics and hydrology, and to trigger the release of water stored along the subglacial flood-path. The concomitant drainage of subglacial lakes during the 2019 GLOF demonstrates the potential for ice-dammed lake drainages to perturb the wider subglacial hydrological system. But the absence of similar releases of stored water during other GLOFs suggests antecedent subglacial hydrological conditions, drainage route and the impact of drainage on subglacial water pressure also impact this process.

Two ice velocity mechanisms were identified. The first velocity response, characterised by increased flow into the lake basin, is caused by the rapid removal of hydrostatic pressure at the lake-ice boundary and an increase in driving stress (ice-surface slope) following lake lowering. The second is increased basal sliding caused by the GLOF pressurising the subglacial drainage system. The impact of this is related to whether the discharge exceeds the hydrological capacity of the existing subglacial drainage system. In particular, GLOFs that occurred in the early- or late-melt season when subglacial drainage is inefficient elicited a greater ice velocity response.

## Author contribution statement

The research was conceptualised by SL, RS and AS. SL wrote the manuscript with review and edits from RS, SD, RI, ST, AM, AS, TC, JG, CC, EB, KL, KW, BD, AHJ, SK, AB, AH and NR. Fieldwork to deploy and maintain the sensors was carried out by everyone other than AJ, AH and BD. BD automated the ice velocity processing and AS carried out the remote sensing of velocity. SD and AM processed the GNSS data. ST and FG processed the passive seismic data. ST also calculated river stage from the time-lapse imagery. SL



and RS carried out the shoreline mapping and reconstruction of freeboard height and water volumes. TC did the DEM production/ co-registration and RI the COSIPY modelling. Figures were produced by SL, RS, AM, AS, ST and KL. AJ produced the supplementary video of the upwelling. Funding for the project was acquired by SL, RS, SB, AS, KL, KW, JG, EB, NR, LE, BG, TH, MPJ and AB.

## Acknowledgements

The SLIDE team was supported by NERC Grant NE/X000257/1. The NICE team was supported by NSF/NERC grants OPP-2039854 and NE/W004860/1, and is grateful for the support of Asiaq Greenland Survey. Both teams acknowledge the support of Kangerlussuaq International Science Support. TRC was supported by a Leverhulme Early Career Fellowship (ECF-2022-589). MPJ, JH and LC were supported by a UKRI Future Leaders Fellowship (MR/V022237/1). The ice-dammed lake shorelines, passive seismic data, time-lapse imagery and GNSS are available at the Polar Data Centre at XXXX. The authors thank the National Aeronautics and Space Administration [NASA]/USGS for the acquisition and free distribution of Landsat and ASTER images. The European Space Agency (ESA) is acknowledged for providing the Sentinel-1 data from the Copernicus Programme used in this study. We thank Planet Labs for providing access to their daily imagery through the education and research program. ArcticDEM is provided by the Polar Geospatial Center under NSF-OPP awards 1043681, 1559691, 1542736, 1810976, and 2129685. The PROMICE weather station data sets are available from How and others, (2022). Maps were created using QGIS.

## References

- Anderson SP and 6 others (2003) Integrated hydrologic and hydrochemical observations of Hidden Creek Lake jökulhlaups, Kennicott glacier, Alaska. *Journal of Geophysical Research: Earth Surface*, 108(F1).
- Bartholomew I, Nienow P, Mair D, Hubbard A, King MA, and Sole A (2010) Seasonal evolution of subglacial drainage and acceleration in a Greenland outlet glacier. *Nature Geoscience*, 3(6), pp.408-411.
- Bigelow DG, Flowers GE, Schoof CG, Mingo LD, Young EM and Connal BG (2020) The role of englacial hydrology in the filling and drainage of an ice-dammed lake, Kaskawulsh Glacier, Yukon, Canada. *Journal of Geophysical Research: Earth Surface*, 125(2), p.e2019JF005110.
- Björnsson H (1992) Jökulhlaups in Iceland: prediction, characteristics and simulation. *Annals of Glaciology*, 16, pp.95-106.
- Bowling J and 10 others (2021) Surface outburst of a subglacial flood from the Greenland Ice Sheet. <https://doi.org/10.21203/rs.3.rs-569793/v1>.

- 789 Carrivick JL, Tweed FS (2013) Proglacial lakes: character, behaviour and geological importance.  
790 *Quaternary Science Reviews* 78 pp.34-52.
- 791 Carrivick JL and Quincey DJ (2014) Progressive increase in number and volume of ice-marginal  
792 lakes on the western margin of the Greenland Ice Sheet. *Global and Planetary Change*, 116,  
793 pp.156-163.
- 794 Carrivick JL and Tweed FS (2016) A global assessment of the societal impacts of glacier outburst  
795 floods. *Global and Planetary Change*, 144, pp.1-16.
- 796 Carrivick JL and Tweed FS (2019) A review of glacier outburst floods in Iceland and Greenland  
797 with a megafloods perspective. *Earth-Science Reviews*, 196, p.102876.
- 798 Carrivick JL and 9 others, (2022) Ice-marginal proglacial lakes across Greenland: Present status  
799 and a possible future. *Geophysical Research Letters*, 49(12), p.e2022GL099276.
- 800 Chen G (1998). GPS Kinematic Positioning for the Airborne Laser Altimetry at Long Valley,  
801 California (Massachusetts Institute of Technology).
- 802 Chu W, Creyts TT and Bell RE (2016) Rerouting of subglacial water flow between neighboring  
803 glaciers in West Greenland. *Journal of Geophysical Research: Earth Surface*, 121(5), pp.925-  
804 938.
- 805 Chudley TR and Howat IM (2024) pDEMtools: conveniently search, download, and process  
806 ArcticDEM and REMA products. *Journal of Open Source Software*, 9(102), p.7149.
- 807 Clarke GK (2003) Hydraulics of subglacial outburst floods: new insights from the Spring–Hutter  
808 formulation. *Journal of Glaciology*, 49(165), pp.299-313.
- 809 Cook SJ, Waller RI and Knight PG (2006) Glaciohydraulic supercooling: the process and its  
810 significance. *Progress in Physical Geography*, 30(5), pp.577-588.
- 811 Cook SJ and Swift DA (2012) Subglacial basins: Their origin and importance in glacial systems  
812 and landscapes. *Earth-Science Reviews*, 115(4), pp.332-372.
- 813 Davison BJ and 6 others, (2020) Subglacial drainage evolution modulates seasonal ice flow  
814 variability of three tidewater glaciers in southwest Greenland. *Journal of Geophysical Research:*  
815 *Earth Surface*, 125(9). doi:10.1029/2019jf005492
- 816 Díaz J, Ruíz M, Crescentini L, Amoruso A and Gallart J (2014). Seismic monitoring of an Alpine  
817 mountain river. *Journal of Geophysical Research: Solid Earth*, 119(4), pp.3276-3289.
- 818 Dømgaard M, Kjeldsen K, How P and Bjørk A (2024) Altimetry-based ice-marginal lake water  
819 level changes in Greenland. *Communications Earth & Environment*, 5(1), p.365.
- 820 Dow JM, Neilan RE and Rizos C (2009) The International GNSS Service in a changing landscape  
821 of Global Navigation Satellite Systems. *J. Geod.* 83, 191198.

- 822 Dow CF, and 6 others (2018) Dynamics of active subglacial lakes in Recovery Ice Stream. *Journal*  
823 *of Geophysical Research: Earth Surface*, 123(4), pp.837-850.
- 824 Einarsson B, Magnússon E, Roberts MJ, Pálsson F, Thorsteinsson T and Jóhannesson T (2016)  
825 A spectrum of jökulhlaup dynamics revealed by GPS measurements of glacier surface motion.  
826 *Annals of Glaciology*, 57(72), pp.47-61.
- 827 Einarsson B, Jóhannesson T, Thorsteinsson T, Gaidos E and Zwinger T (2017) Subglacial flood  
828 path development during a rapidly rising jökulhlaup from the western Skaftá cauldron, Vatnajökull,  
829 Iceland. *Journal of Glaciology*, 63(240), pp.670-682.
- 830 Evans SG and Clague JJ (1994) Recent climatic change and catastrophic geomorphic processes  
831 in mountain environments. In *Geomorphology and Natural Hazards* (pp. 107-128). Elsevier.
- 832 Flowers GE, Björnsson H, Pálsson F and Clarke GK (2004) A coupled sheet-conduit mechanism  
833 for jökulhlaup propagation. *Geophysical Research Letters*, 31(5).
- 834 Fowler AC and Ng FSL (1996) The role of sediment transport in the mechanics of jökulhlaups.  
835 *Annals of Glaciology*, 22, pp.255-259.
- 836 Gilbert R (1971) Observations on ice-dammed summit lake, British Columbia, Canada. *Journal of*  
837 *Glaciology*, 10(60), pp.351-356.
- 838 Gimbert F, Tsai VC and Lamb MP (2014) A physical model for seismic noise generation by  
839 turbulent flow in rivers. *Journal of Geophysical Research: Earth Surface*, 119(10), pp.2209-2238.
- 840 Glen JW (1954) The stability of ice-dammed lakes and other water-filled holes in glaciers. *Journal*  
841 *of Glaciology*, 2(15), pp.316-318.
- 842 Grinsted A, Hvidberg CS, Campos N and Dahl-Jensen D (2017). Periodic outburst floods from an  
843 ice-dammed lake in East Greenland. *Scientific Reports*, 7(1), p.9966.
- 844 Guizar-Sicairos M, Thurman ST, and Fienup JR (2008) Efficient subpixel image registration  
845 algorithms. *Optics Letters*, 33(2), 156. <https://doi.org/10.1364/OL.33.000156>
- 846 Haeberli W (1983) Frequency and characteristics of glacier floods in the Swiss Alps. *Annals of*  
847 *Glaciology*, 4, pp.85-90.
- 848 Hösli L and 10 others (2025) Subglacial cavity collapses on Swiss glaciers: spatio-temporal  
849 distribution and mass loss contribution. *Journal of Glaciology*, pp.1-35.
- 850 How, P and 10 others (2021). Greenland-wide inventory of ice marginal lakes using a multi-  
851 method approach. *Scientific reports*, 11(1), p.4481.
- 852 How P and others (2025) PROMICE and GC-Net automated weather station data in Greenland.  
853 doi:10.22008/FK2/IW73UU.

- 854 Iken A and Bindshadler RA (1986) Combined measurements of subglacial water pressure and  
855 surface velocity of Findelengletscher, Switzerland: conclusions about drainage system and sliding  
856 mechanism. *Journal of Glaciology*, 32(110), pp.101-119.
- 857 Jóhannesson T (2002) Propagation of a subglacial flood wave during the initiation of a jökulhlaup,  
858 *Hydrol. Sci. J.*, 47, 417–434.
- 859 Huss M, Bauder A, Werder M, Funk M and Hock R (2007) Glacier-dammed lake outburst events  
860 of Gornersee, Switzerland. *Journal of Glaciology*, 53(181), pp.189-200.
- 861 King O, Bhattacharya A, Bhambri R and Bolch T (2019) Glacial lakes exacerbate Himalayan  
862 glacier mass loss, *Sci. Rep.*, 9, 18145.
- 863 King M (2004). Rigorous GPS data-processing strategies for glaciological applications. *J. Glaciol.*  
864 50, 601607.
- 865 Kingslake J and Ng F (2013) Modelling the coupling of flood discharge with glacier flow during  
866 jökulhlaups. *Annals of Glaciology*, 54(63), pp.25-31.
- 867 Kirillov A and 10 others (2023) Segment anything. In *Proceedings of the IEEE/CVF International*  
868 *Conference on Computer Vision* (pp. 4015-4026).
- 869 Kirkbride MP (1993) The temporal significance of transitions from melting to calving termini at  
870 glaciers in the central Southern Alps of New Zealand. *The Holocene*, 3(3), pp.232-240.
- 871 Kjeldsen KK, Khan SA, Bjørk AA, Nielsen K and Mouginit J (2017) Ice-dammed lake drainage in  
872 west Greenland: Drainage pattern and implications on ice flow and bedrock motion. *Geophysical*  
873 *Research Letters*, 44(14), pp.7320-7327.
- 874 Lea JM (2018) The Google Earth Engine Digitisation Tool (GEEDiT) and the Margin change  
875 Quantification Tool (MaQiT)—simple tools for the rapid mapping and quantification of changing  
876 Earth surface margins. *Earth Surface Dynamics*, 6(3), pp.551-561.
- 877 Liestøl O (1956) Glacier dammed lakes in Norway. *Norwegian Journal of Geography*, 15:3-4, 122-  
878 149. <https://doi.org/10.1080/00291955608542772>
- 879 Lindbäck K and 8 others (2014) High-resolution ice thickness and bed topography of a land-  
880 terminating section of the Greenland Ice Sheet. *Earth System Science Data*, 6(2), pp.331-338.
- 881 Livingstone SJ, Sole AJ, Storrar RD, Harrison D, Ross N and Bowling J (2019). Brief  
882 communication: Subglacial lake drainage beneath Isunguata Sermia, West Greenland:  
883 geomorphic and ice dynamic effects. *The Cryosphere*, 13(10), pp.2789-2796.
- 884 Livingstone SJ and others (2022) Subglacial lakes and their changing role in a warming climate.  
885 *Nature Reviews Earth & Environment*, 3(2), pp.106-124.
- 886 Lützow N, Veh G and Korup O (2023) A global database of historic glacier lake outburst floods.  
887 *Earth System Science Data Discussions*, 2023, pp.1-27.




- 888 Magnússon E, Rott H, Björnsson H and Pálsson F (2007). The impact of jökulhlaups on basal  
889 sliding observed by SAR interferometry on Vatnajökull, Iceland. *Journal of Glaciology*, 53(181),  
890 pp.232-240.
- 891 Mallalieu J, Carrivick JL, Quincey DJ and Raby CL (2021) Ice-marginal lakes associated with  
892 enhanced recession of the Greenland Ice Sheet. *Global and Planetary Change*, 202, p.103503.
- 893 Mankoff KD and 9 others (2020) Greenland liquid water discharge from 1958 through 2019. *Earth*  
894 *System Science Data*, 12(4), pp.2811-2841.
- 895 Mathews WH and Clague JJ (1993) The record of jökulhlaups from Summit Lake, northwestern  
896 British Columbia. *Canadian Journal of Earth Sciences*, 30(3), pp.499-508.
- 897 Morlighem M and 10 others (2017) BedMachine v3: Complete bed topography and ocean  
898 bathymetry mapping of Greenland from multibeam echo sounding combined with mass  
899 conservation. *Geophysical Research Letters*, 44(21), pp.11-051.
- 900 Morlighem M, and others (2022) IceBridge BedMachine Greenland (IDBMG4, Version 5) [Data  
901 set]. Boulder, Colorado USA. *NASA National Snow and Ice Data Center Distributed Active*  
902 *Archive Center*. <https://doi.org/10.5067/GMEVBWFLWA7X>
- 903 Muñoz-Sabater J and others (2021) ERA5-Land: a state-of-the-art global reanalysis dataset for  
904 land applications. *Earth System Science Data* 13(9), 4349–4383. doi:10.5194/essd-13-4349-  
905 2021.
- 906 Ng F and Liu S (2009) Temporal dynamics of a jökulhlaup system. *Journal of Glaciology*, 55(192),  
907 pp.651-665.
- 908 Nghiem SV and 8 others (2012) The extreme melt across the Greenland ice sheet in 2012.  
909 *Geophysical Research Letters*, 39(20).
- 910 Nuth C and Kääb A (2011) Co-registration and bias corrections of satellite elevation data sets for  
911 quantifying glacier thickness change. *The Cryosphere*, 5(1), 271-290. [https://doi.org/10.5194/tc-](https://doi.org/10.5194/tc-5-271-2011)  
912 [5-271-2011](https://doi.org/10.5194/tc-5-271-2011).
- 913 Nye JF (1976) Water flow in glaciers: jökulhlaups, tunnels and veins. *Journal of Glaciology*,  
914 17(76), pp.181-207.
- 915 Post A and Mayo LR (1971) Glacier dammed lakes and outburst floods in Alaska (No. 455). *US*  
916 *Geological Survey*.
- 917 Rick B, McGrath D, McCoy SW and Armstrong WH (2023) Unchanged frequency and decreasing  
918 magnitude of outbursts from ice-dammed lakes in Alaska. *Nature Communications*, 14(1), p.6138.
- 919 Riesen P, Sugiyama S and Funk M (2010) The influence of the presence and drainage of an ice-  
920 marginal lake on the flow of Gornergletscher, Switzerland. *Journal of Glaciology*, 56(196), pp.278-  
921 286.

- 922 Roberts MJ (2005) Jökulhlaups: a reassessment of floodwater flow through glaciers. *Reviews of*  
923 *Geophysics*, 43(1).
- 924 Roberts MJ, Russell AJ, Tweed FS and Knudsen Ó (2000) Ice fracturing during jökulhlaups:  
925 implications for englacial floodwater routing and outlet development. *Earth Surface Processes*  
926 *and Landforms*, 25(13), 1429-1446.
- 927 Röthlisberger H (1972) Water pressure in intra-and subglacial channels. *Journal of Glaciology*,  
928 11(62), pp.177-203.
- 929 Russell AJ, Roberts MJ, Fay H, Marren PM, Cassidy NJ, Tweed FS and Harris T (2006) Icelandic  
930 jökulhlaup impacts: implications for ice-sheet hydrology, sediment transfer and geomorphology.  
931 *Geomorphology*, 75(1-2), pp.33-64.
- 932 Russell AJ, Carrivick JL, Ingeman-Nielsen T, Yde JC and Williams M (2011) A new cycle of  
933 jökulhlaups at Russell Glacier, Kangerlussuaq, West Greenland. *Journal of Glaciology*, 57(202),  
934 pp.238-246.
- 935 Sandwell D, Mellors R, Tong X, Wei M and Wessel P (2011a) An InSAR processing system based  
936 on Generic Mapping Tools. UC San Diego: Scripps Institution of Oceanography.
- 937 Sandwell D, Mellors R, Tong X, Wei M and Wessel P (2011b) Open radar interferometry software  
938 for mapping surface deformation. *Eos, Transactions American Geophysical Union*, 92(28), 234.  
939 <https://doi.org/10.1029/2011EO280002>
- 940 Sauter T, Arndt A and Schneider C (2020) COSIPY v1.3 – an open-source coupled snowpack  
941 and ice surface energy and mass balance model. *Geoscientific Model Development* 13(11),  
942 5645–5662. doi:10.5194/gmd-13-5645-2020.
- 943 Schmandt B, Aster RC, Scherler D, Tsai VC and Karlstrom K (2013) Multiple fluvial processes  
944 detected by riverside seismic and infrasound monitoring of a controlled flood in the Grand Canyon.  
945 *Geophysical Research Letters*, 40(18), pp.4858-4863.
- 946 Shangguan D, Ding Y, Liu S, Xie Z, Pieczonka T, Xu J and Moldobekov B (2017). Quick release  
947 of internal water storage in a glacier leads to underestimation of the hazard potential of glacial  
948 lake outburst floods from Lake Merzbacher in Central Tian Shan Mountains. *Geophysical*  
949 *Research Letters*, 44(19), pp.9786-9795.
- 950 Sugiyama S, Bauder A, Huss M, Riesen P and Funk M (2008) Triggering and drainage  
951 mechanisms of the 2004 glacier-dammed lake outburst in Gornergletscher, Switzerland. *Journal*  
952 *of Geophysical Research: Earth Surface*, 113(F4).
- 953 Sutherland JL, Carrivick JL, Gandy N, Shulmeister J, Quincey DJ and Cornford S (2020)  
954 Proglacial lakes control glacier geometry and behavior during recession. *Geophysical Research*  
955 *Letters*, 47(19), p.e2020GL088865.


- 956 Thorarinsson S (1939) Chapter IX. The ice dammed lakes of Iceland with particular reference to  
957 their values as indicators of glacier oscillations. *Geografiska Annaler*, 21(3–4), 216–242.  
958 <https://doi.org/10.1080/20014422.1939.11880679>.
- 959 Tomczyk, AM, Ewertowski MW and Carrivick JL (2020) Geomorphological impacts of a glacier  
960 lake outburst flood in the high arctic Zackenberg River, NE Greenland. *Journal of Hydrology*, 591,  
961 p.125300.
- 962 Tweed FS and Russell AJ (1999) Controls on the formation and sudden drainage of glacier-  
963 impounded lakes: implications for jökulhlaup characteristics. *Progress in Physical Geography*,  
964 23(1), pp.79-110.
- 965 Tuckett PA and 6 others (2019) Rapid accelerations of Antarctic Peninsula outlet glaciers driven  
966 by surface melt. *Nature Communications*, 10. doi:10.1038/s41467-019-12039-2
- 967 Veh G and 8 others (2023) Less extreme and earlier outbursts of ice-dammed lakes since 1900.  
968 *Nature*, 614(7949), pp.701-707.
- 969 Walder JS and 6 others (2005) Fault-dominated deformation in an ice dam during annual filling  
970 and drainage of a marginal lake. *Annals of Glaciology*, 40, pp.174-178.
- 971 Welch PD (1967) The use of fast Fourier transform for the estimation of power spectra: A method  
972 based on time averaging over short, modified periodograms. *IEEE Transactions on Audio and*  
973 *Electroacoustics* 15(2): 70–73.
- 974 Whalley WB (1971) Observations of the drainage of an ice-dammed lake – Strupvatnet, Troms,  
975 Norway. *Norsk Geografisk Tidsskrift* 25, 165–74.
- 976  
977  
978

979 SUPPLEMENTARY FIGURES






981 **Figure S1.** Elevation-Volume scaling relationship derived from the 11/08/2019 ArcticDEM DSM  
 982 strip when the lake was empty.




983 **Figure S2.** Examples of the Time-lapse imagery used to estimate the river stage at the terminus  
 984 of the glacier in 2023. Camera location: -50.334W 67.181N -50.335W (Fig. 1 - Camera 3). Red  
 985 and green stars show the background and foreground points used in the Segment Anything Model  
 986 (SAM). Blue line is the output of the SAM showing the water mask. The height of the water mask  
 987 averaged in the area delineated by the white dotted line is taken as the relative river stage.


988  
 989  
 990



**Figure S3.** Robust linear regression of maximum pre-drainage lake volume through time. A statistically significant negative linear relationship was observed (Pearson  $r = -0.71$ ,  $p = <0.05$ ).



**Figure S4.** Ice velocity time-series for ROIs D1-3 and U1 (see Fig. 1 for locations). (a) is the U1 time-series, and (b-d) are the D1 to D3 times-series, respectively.



**Figure S5.** Isunnguata Sermia ice velocity: a-b. 2014 - 2025 average ice velocity of the U and V components, respectively; c-d. Average ice velocity of the U and V components for image pairs that include the 2023 ice-dammed lake drainage; e-f. Ice velocity anomaly in the U and V direction for the 2023 ice-dammed lake drainage period compared to the long-term average. Positive values (blue) represent pixels for which ice velocity was greater during the lake drainage period.

**Supplementary Video.** Isunnguata Sermia outburst (2023) - <https://vimeo.com/1103612163>