

# **22 Years Later: Evaluating Glover & Smith's 2003 Predictions of the 2025 Deep-Sea Ecosystem**

Authors: Kalyan Rao\*<sup>1</sup>

<sup>1</sup>The Westminster Schools, Marine Biology, Atlanta, GA, USA

\*[kalyanrao@westminster.net](mailto:kalyanrao@westminster.net)

## **Peer Review Status:**

*This is a non-peer reviewed preprint submitted to EarthArXiv*

This paper will be submitted to *Frontiers in Marine Science* for peer review.

## **Abstract**

Earth's deep-sea ecosystem remains one of the least explored and understood ecosystems on the planet. Human interest in the resources available in the deep-sea must be balanced with potential harm to the deep-sea ecosystem. Accurately predicting the effect on human impact on the deep-sea floor can help guide policies and actions today. This research evaluates prior predictions of human impact on the deep-sea floor ecosystem to aid in more accurate predictions in the future. In 2003, Glover & Smith published the frequently cited "The deep-sea floor ecosystem: current status and prospects of anthropogenic change by the year 2025" that made 15 predictions about the state of the deep-sea in 2025. In this paper we now examine their predictions for 2025 and assess the accuracy of each prediction. We classify each of the predictions as either accurate, partially accurate, or inaccurate. Using current studies and evaluations of the deep-sea, alongside news sources detailing deep-sea progress, we compare the state of the ocean today with Glover & Smith's predictions to test their accuracy. 7 of their predictions proved to be accurate, 4 were partially accurate, and 4 more were inaccurate. We identified common themes of both accurate and inaccurate predictions to guide future predictions. Predictions regarding technological development were largely inaccurate while predictions concerning sources of damage to ecosystems were largely accurate.

## **Introduction**

The ability to forecast future ecological and industrial developments in the deep-sea is critical for shaping policy decisions and conservation processes. As deep-sea data is comparatively sparse, and there are limited points of comparison, making such predictions can be challenging. It is important to critically assess previous predictions so that future predictions do not contain similar errors when prior predictions proved inaccurate. Glover & Smith's influential 2003 article made 15 predictions about the state of human impacts in the deep-sea by 2025. Glover and Smith categorized their predictions into 5 broader categories: 1) deep-sea waste disposal, 2) deep-sea fisheries, 3) deep-sea oil and gas drilling, 4) deep-sea mineral and methane fuel extraction, and 5) climate change impacts on the deep sea. Now in 2025, it is possible to review and reflect on their predictions. Understanding which predictions were accurate, and which ones were not, yields valuable information about the drivers of change in deep-sea ecosystems and the limitations of long-range forecasting in a data-poor environment. Here we assessed the accuracy of Glover & Smith's predictions by comparing them to peer-reviewed literature, policy documents, industry reports, and news reports that reveal the current state of the deep-sea.

## **Evaluation**

Glover & Smith's article contains information and predictions regarding the deep-sea in 2025, and they then summarize their predictions with 15 succinct bullet points. We use these 15 points to evaluate their predictions. To find current references on the deep-sea to compare to Glover & Smith's article we conducted a literature search dated subsequent to 2003. As the deep-sea is an environment with the majority of human innovation relevant to the predictions coming in the 2020's, peer reviewed literature was not sufficient to get up to date information on new developments. We also included industry reports and policy documents that revealed new progress to broaden our assessment and understanding of the current state of the deep sea. To determine the accuracy of each prediction, we evaluated if it was fully recognized in the current state of the deep-sea. If all aspects of a prediction matched the current state of the deep-sea, it

was deemed accurate. A prediction was evaluated as partially accurate if only some of its facets were accurate. We also evaluated predictions as partially accurate if it is only applicable to a few areas of the deep sea. A prediction that hadn't proved true in any facet was designated as inaccurate.

## **Results**

### **Deep Sea Waste in 2025**

1. Ships will continue to sink, but will not pose any greater environmental hazard than today.

Assessment: Accurate

Ships have continued to sink from 2003 to 2025. Over 340 ships have sunk in the past decade alone, and there has been a downward trend in the number of ships sinking yearly (Allianz, 2025). Bulk carriers, ships that frequently cross the deep-sea and have the potential to greatly affect the deep-sea due to their contents, sank 71 times between 2005 and 2014 but only sunk 20 times between 2015 and 2024 (Intercargo, 2016, 2025). Additionally, sunken ships do not pose any greater environmental hazard than in 2003. The introduction of doubled hulls to oil tankers, which are mandated in US waters, reduced oil spills by 20-62% on average (Yip et al., 2011). Sinking ships do continue to sink and pose less environmental hazard than in 2003.

2. Sewage sludge and dredge spoil disposal will have begun in the deep sea, with significant localized impacts.

Assessment: Partially Accurate

Sewage sludge disposal does not occur at the deep sea, but dredge spoils are disposed of at deep sea. At the 2022 London Convention, the International Maritime Organization (IMO) removed sewage sludge from the list of permitted waste for dumping. The IMO cited that sewage dumping had been declining for the past decades and that many regional and domestic policies banned sewage sludge dumping. Additionally, there exist many safer alternatives to removing sewage waste than dumping it at sea (IMO, 2022). While sewage sludge dumping was in practice before its ban, it did have significant localized impacts, causing measurable changes to benthic ecology near the dumpsite (US Department of Commerce, 2014). Dredge spoil dumping does occur at sea, but it is often focused around established sites (OSPAR, 2017). Deep-sea disposal of dredge spoils is preferred to other methods of disposal since it has the least environmental impact (Svensson et al., 2022). The sites of dredge spoil dumping do often see some localized significant environmental impacts including metal pollution, faunal smothering, and fluxes in turbidity (Stronkhorst et al., 2003; Chen et al., 2018; Palanques et al., 2022). The effects are mostly localized and don't affect areas beyond the dump site (Blake et al., 2009).

3. Large quantities of CO<sub>2</sub> (up to 1 Gt C yr<sup>-1</sup>) will be sequestered in the deep ocean; major environmental effects will initially be localized to within 10–100 km of injection points, but traceable far-field effects will begin to be observed.

Assessment: Inaccurate

We could find no data to suggest that large quantities of CO<sub>2</sub> are being sequestered in 2025. The technology remains experimental and simulation heavy (Aman et al., 2025). CO<sub>2</sub> Sequestration mainly does not occur because of the dangers it poses to deep-sea ecosystems. Sequestered CO<sub>2</sub> increases the pH of surrounding waters, harming organisms, especially those with calcium carbonate skeletons. Many deepwater species can not adapt to instant drastic changes in pH so are extremely sensitive. Ocean acidification leads to a high mortality rate of meiofauna and can jeopardize entire food webs (Barry et al., 2004).

### **Deep-Sea Fisheries in 2025**

4. All existing (2003) deep-sea fisheries will be commercially extinct.

Assessment: Inaccurate

Not all 2003 deep-sea fisheries are commercially extinct. While many are commercially extinct, some fisheries from 2003 still exist today such as New Zealand's Orange Roughy fishery which is still active today (Moor et al., 2025). The fishery is still alive due to auditing and regulation that prevent overfishing. The species saw a rapid decline towards the end of the 20th century but has since then seen a recovery in population (MSC, 2016).

5. New fisheries will have been discovered and 'mined', but each fishery will have 20–30 year lifespans or less.

Assessment: Accurate

Most deep-sea fisheries often see a boom-bust cycle in which they are rapidly farmed over a short period of time until the area is commercially depleted. Due to this fact, it is difficult for deep-sea fisheries to be commercially active for more than 20-30 years. Deep-sea fisheries suffer from the reproductive habits of many deep-sea fish which often produce a small number of offspring. Additionally, their mating can be erratic and episodic, leading to slow recovery (Maribus gGmbH, 2013).

6. Significant environmental impacts will have occurred on seamounts and deep-water coral beds; extinctions may have been prevented through the setting up of marine protected areas.

Assessment: Accurate

There have been significant and irreversible impacts on seamounts and deep-water coral beds. Approximately 70% of fishing ships deploy demersal trawling nets that can reach depths up to 2000m (Maribus gGmbH, 2013). Trawling causes major irreversible damage to deep-sea ecosystems, and there are no legally binding regulations on them currently (Maribus gGmbH, 2013; Goode et al., 2020). In reef habitats, heavy fishing results in 90% of the reef becoming bare rock, and heavy fishing on seamounts can lead to 97% of coral being destroyed (Koslow and Glowlett-Holmes, 1998; Clark and O'Driscoll, 2003). Marine protected areas (MPAs) have been established to protect species in pristine areas from being destroyed (Goode et al., 2020). Converting previously fished areas into MPAs does not greatly affect the ecosystem. There is little regrowth or recolonization of corals, demonstrating the low-resilience and slow recovery time of deep-sea communities (Huvenne et al., 2016).

## **Deep-Sea Oil and Gas Drilling in 2025**

7. Deep (500 m) drilling will be commonplace, and may dominate the offshore market by 2025. Ultra-deep (1000 m) drilling rigs will be coming on line at depths up to 4000 m.

Assessment: Accurate

To evaluate Glover & Smith's prediction, we must quantify the term "commonplace". The authors decide that a greater 10% market share in offshore drilling is commonplace. Deep drilling is now commonplace, and ultra-deep drilling rigs have come on line at depths up to 4000m. There is limited information on the exact number of oil rigs globally; the majority of reports are regional (e.g. Gulf of Mexico or Brazil). Data differs between regions because of differences in size of the continental shelf. Additionally, the definition of "deep" differs between studies. To assess the prevalence of deep-drilling within the market we only considered reports that consider "deep" between 400-600m of water depth. While it is difficult to exactly judge how much deep-water drilling comprises the market, most sources place the figure between 20% and 55%, establishing deep drilling as commonplace (Rystad, 2024; Precedence, 2025; Research Nester, 2025). Ultra-deep drilling rigs are now implemented. By 2006 there were already more than 500 ultra-deep drilling rigs globally, and in the United States alone there are 38 active platforms at least 1000 meters deep (Shaughnessy et al., 2007; BOEM, 2025). Glover & Smith were very accurate when predicting that ultra-deep water drilling will not pass 4000m. The deepest drill currently being constructed is the Komodo-1 drill in the Colombia basin which will reach a depth of 3950m (Carvajal-Arenas et al., 2024).

8. Significant, but localized, impacts will be occurring at the deep-sea floor surrounding deep oil drilling structures, but oil companies will be mandated to minimize the impacts and acquire baseline survey data before commencing drilling operations.

Assessment: Partially accurate

The effects of deepwater drilling are localized, rarely spreading further than 2 km from the rigs. Within 100 m of the drilling sees the greatest impacts. This damage can come from anchors which rip up biogenic habitats on the ocean floor causing scars as they are dragged along the seabed (Hall–Spencer et al., 2002; Watling, 2014). Direct impacts of infrastructure installation such as sediment resuspension and burial also occur within a 100m radius. Within 200-300m sees the most common ecological impacts on the population and community levels (Cordes et al., 2016). Organisms from a few 100 m to 1km from the drilling are susceptible to sound induced behavior disruption (Southall et al., 2008). Discharges of water-based and low-toxicity oil-based drilling muds and produced water can extend over 2 km from the source, and evidence suggests that benthic organisms within 2 km of the source will be affected by produced water (Bakke et al., 2013; Cordes et al., 2016). There are mandates around the world for oil companies to minimize impacts, but varied regulations exist in different maritime jurisdictions and for areas beyond national jurisdiction (Mazor et al., 2014; Katsanevakis et al., 2015). Some mandates do not require acquisition of survey data (Cordes et al., 2016). The United Nations Convention on the Law of the Sea allows for individual countries to decide the regulations within their exclusive economic zone (EEZ), and there has been no significant effort to standardize EEZ regulations across countries (Cordes et al., 2016). Many of the world's leading nations in offshore drilling, including the United States, the UK, and Norway, all require risk assessments from organizations prior to starting drilling (Larsson and Purser, 2011). Not every oil company is required to acquire survey baseline data, though many of them do (Cordes et al., 2016).

9. Large-scale accidents will be an occasional hazard to marine life, but are unlikely to cause more than temporary ecological incidents.

Assessment: Partially Accurate

Large scale accidents do occasionally happen, but the effects are much longer than temporary incidents, especially for larger accidents such as the Deepwater Horizon oil spill. While small oil spills are unlikely to have persistent effects, large scale accidents can have long-term ecosystem perturbations. The oil affects all ranges of organisms including benthic fauna, deep-water corals, coastal wetlands, marine mammals, and seabirds. It is virtually impossible to clean up all of the oil lost from drilling accidents, and it can last for decades once spilled (Barron et al., 2020). Additionally, the rig accidents expel tons of harmful gases that can never be recaptured (Hu, 2025). In oil spread to saltmarshes, contaminated sediments lead to extreme death, and it can take up to decades for meiofauna to recover. In heavily oiled-areas, oil-induced erosion can cause permanent regional declines in abundance and diversity of meiofauna (Pant et al., 2025). Deep-water oil spills lead to immediate death of many deep-sea species that are less resilient and take longer to reproduce, often taking years to come close to pre-accident abundance and diversity. Most corals take multiple decades to recover after an oil spill, and some sites that are heavily affected show little evidence of possible recovery (Schwing et al., 2020). In benthic species, polycyclic aromatic hydrocarbon levels can remain elevated for up to 6+ years (Gohlke et al., 2011). Many fish exposed to oils during drilling accidents have cardiotoxic effects that restrict fitness (Pasparakis et al., 2019). Oil spills also impact the maternal health of dolphins

which has effects on their reproductive success, affecting the population in the long-term (Murawski et al., 2021). Large-scale accidents are an occasional hazard to marine life, but the effects are not temporary and can harm ecosystems for generations.

## **Deep-sea mineral and methane fuel extraction in 2025**

10. The first deep-sea marine mineral extraction of gold-rich hydrothermal sulphide deposits will have proved commercially successful.

Assessment: Inaccurate

The first deep-sea mineral extractions have not proved commercially successful. Japan conducted the only large-scale extraction of hydrothermal sulfide deposits in 2017 in the Okinawa Trench. The project wasn't commercial, instead run by Japan Organization for Metals and Energy Security, a government organization. They successfully excavated around 650 kilograms of cobalt, nickel, and gold (Giseburt, 2024). The project was overall a success, but was unable to prove commercial viability. The only other project to mine deep-sea hydrothermal deposits was the Solwara 1 project. Working with the Papua New Guinea (PNG) government, the Canadian based company, Nautilus mineral cooperation, attempted to excavate the Manus Basin. In 2011, PNG granted a 20 year mining license in the Bismarck Sea. Over the next few years the project received criticism from many climate activists and many financial setbacks (Hutt, 2018). In 2019, due to major pushback against the project and other environmental concerns, PNG and other Melanesian countries banned deep-sea mining for 10 years in the Udaune Declaration on Climate Change (Cannon, 2023).

11. Developing countries will begin to encourage high-risk mining ventures on offshore deposits.

Assessment: Accurate

Many island nations in Oceania view deep-sea mining as a potential catalyst for development. The Cook Islands, Nauru, Tonga and Kiribati are among the region's most active proponents (Abe et al., 2025). The Clarion-Clipperton Zone in the Eastern Pacific Ocean is the region with the most activity in deep-sea mining with many nations aiming to claim parts of it. The International Seabed Authority (ISA) has granted exploration licenses to explore mining in the Clarion-Clipperton zone to 31 nations including developing countries such as Cook Islands, Nauru, Tonga, and Kiribati (Keating-Bitonti, 2025). While many developing countries support deep-sea mining, other developing nations such as Palau and Costa Rica oppose deep-sea mining ventures. Nauru is one of the main proponents, sponsoring The Metals Company. In 2021 Nauru gave a two year notice to the ISA to draft their regulations on deep-sea mining, but they failed to develop a mining code by the 2023 deadline (Alberts, 2023). Instead the ISA promised to have rules set in place during 2025 which have not been released yet, but are currently being developed in the July 2025 ISA conference (Letra, 2025).

12. Developed countries will continue to rely on land-based mineral extraction, but will maintain strategic interests in offshore deposits.

Assessment: Accurate

Many developed nations are in the process of exploring offshore deposits in addition to relying on land-based mining. Of the 31 countries who were granted exploration licenses in the Clarion-Clipperton zone, many are developed and already have land-based mines such as France, China, Germany, and Russia (Keating-Bitonti, 2025).

13. Offshore oil industries, faced with declining oil and gas reserves, will start to extract methane hydrate at commercially viable quantities.

Assessment: Inaccurate

There have been successful methane hydrate extractions and tests done by governments, but it has not been commercially realized (Yin and Linga, 2019). In 2013 Japan became the first country to excavate methane hydrate from deep-waters, but the project was temporary and didn't extend into the commercial market (Yiallourides, 2019). The methane hydrate deposit in the South China Sea was first discovered in 2007, and recently China announced that they have successfully extracted it with 16 thousand cubic meters of gas being extracted daily (Zhengzheng, 2017; Akhmetkaliyeva, 2020). Due to its high cost, methane hydrate extraction commercialization has yet to have been achieved and is only done by larger governments (Akhmetkaliyeva, 2020).

14. The vast bulk of the bathyal and abyssal deep sea will remain untouched by mineral and methane extraction and there will be no convincing evidence of species extinctions.

Assessment: Accurate

The vast majority of the bathyal and abyssal deep sea has been untouched by mineral and methane extraction. Outside of Exclusive Economic Zones, the ISA has granted a few exploration licenses, leaving the majority of the deep-sea untouched. Over 80% of the ocean still remains unobserved (McVeigh, 2021). There is no convincing evidence of species extinctions in the deep sea. There are very few studies that investigate the impacts of mining activities in the deep-sea, so there is little evidence of extinctions or how large the other environmental impacts are (Cuvelier et al., 2018). Even in the Clarion-Clipperton Zone, the most studied region of mining activity, around 90% of species are estimated to be undescribed, with thousands undiscovered (Jackson, 2023). We lack the technology to accurately monitor the species of the deep-sea, and the new technology isn't being created fast enough (Thomas et al., 2022). If there



were species extinctions, there would not be convincing evidence since the majority of species are not tracked. Around seamounts there has been a little tracking of species, but there have not been any known extinctions yet, though some species have become endangered (McVeigh, 2021).

### Climate change impacts on the deep sea in 2025

15. Generally speaking, it will be too early for major effects of climate change (such as changes in thermohaline circulation) to be felt in the deep ocean, although subtle shifts in benthic standing crop and biogeochemical recycling in sea floor sediments may be detectable

Assessment: Partially Accurate

Data show that it is not too early to notice major effects due to climate change. Major changes have been felt in the deep-ocean, many of them due to global warming. The abyssal ocean has warmed up significantly and contributes a large amount to global heat content increases (Johnson and Purkey, 2024). Thermohaline circulation has been affected by climate change too. The Atlantic Meridional Ocean Circulation (AMOC) has slowed down recently. Anthropogenic warming is the most likely cause of the slowdown, and climate change models that simulate a weakened AMOC match real world data (Baker et al., 2025; Li and Liu, 2025). Subtle shifts in benthic standing crop and biogeochemical recycling have been detectable. There has been an increase in bottom water temperature and sediment total organic carbon and a decrease in meiofaunal biomass which indicates long-term shifts in benthic ecosystems (Zhou et al., 2023). There have been observable significant changes to benthic populations which are correlated with an increase of particulate organic carbon flux (Ruhl and Smith, 2004; Ruhl, 2007). There has also been a northward shift of species linked to rising temperatures and retreating ice cover (Goethel et al., 2019). There have been noticeable major effects due to climate change in the deep ocean and shifts in benthic standing crop and biogeochemical recycling have been detected.

Prediction	Assessment
1. Ships will continue to sink with no greater environmental risk.	Accurate
2. Sewage sludge and dredge spoil disposal will occur in the deep sea with significant localized impacts.	Partially accurate
3. Large quantities of CO <sub>2</sub> will be sequestered and have localized major environmental effects.	Inaccurate
4. All existing (2003) deep-sea fisheries will be commercially extinct.	Inaccurate
5. New fisheries will have been discovered and ‘mined’, but each fishery will have 20–30 year lifespans or less.	Accurate

6. Significant environmental impacts will have occurred on seamounts and deep-water coral beds; extinctions may have been prevented through the setting up of marine protected areas.	Accurate
7. Deep drilling will be commonplace and ultra-deep drilling rigs will be coming on line.	Accurate
8. Significant localized impacts will be occurring at the deep-sea floor surrounding deep oil drilling structures, but oil companies will be mandated to minimize the impacts and acquire baseline data.	Partially accurate
9. Large-scale accidents will be an occasional hazard to marine life, but are unlikely to cause more than temporary ecological incidents.	Partially accurate
10. The first deep-sea marine mineral extraction will have proved commercially successful.	Inaccurate
11. Developing countries will encourage high-risk mining ventures.	Accurate
12. Developed countries will rely on land-based mineral extraction, but will maintain strategic interests in offshore deposits.	Accurate
13. Offshore oil industries will start to extract methane hydrate.	Inaccurate
14. The vast bulk of the bathyal and abyssal deep sea will remain untouched and there will be no convincing evidence of species extinctions.	Accurate
15. Generally speaking, it will be too early for major effects of climate change to be felt in the deep ocean, although subtle shifts in benthic standing crop and biogeochemical recycling in sea floor sediments may be detectable.	Partially accurate

## Discussion

Glover & Smith accurately predicted 7 aspects of the deep-sea in 2025. 4 predictions were partially accurate, and 4 more predictions were inaccurate. Within the 5 broad categories described in Glover & Smith's paper (deep-sea waste disposal, deep-sea fisheries, deep-sea oil and gas drilling, deep-sea mineral and methane fuel extraction, and climate change impacts on the deep sea), there were similar levels of accuracy. Each one contained both accurate and inaccurate predictions. As assessment of past deep-sea predictions is an essential guide to improving future predictions, we should seek to unearth common themes linking either accurate and inaccurate predictions. To further assess the predictions, we created common themes in which to group predictions and analyze correlations between accuracy and types of predictions. We categorized the predictions into 4 themes: technological progress (predictions: 3, 7, 10, 13), sources of damage to ecosystems (1, 2, 8, 9, 14), effects of damage on ecosystems (4, 5, 6, 9, 15), and policy/regulatory (2, 8, 11, 12).

## **Technological predictions**

The most striking pattern in the accuracy of various themes was the failure of predictions about technology. Of the 4 technological predictions, only drilling depth proved to be accurate. This reflects an over-optimism in technological development. These predictions may have failed because they did not accurately consider technological, economic, and regulatory barriers. Predictions 3, 10, and 13 illustrate the three different reasons explaining why the technological predictions failed. CO<sub>2</sub> sequestration (prediction 3) does not exist because of environmental concerns that were not anticipated in 2003. This highlights the possibility for ethical and policy decisions to regulate deployment even when a technology exists. Commercial marine mineral mining (prediction 10) failed due to economic reasons. The unanticipated gap between technological viability and economic feasibility was too large for the industry to become successful. Methane hydrate extraction (prediction 13) floundered and remains in the pilot-project phase. Only government-funded research has proven successful at extracting methane hydrate, and the technology has not reached the private sector. Glover & Smith's prediction on deep-sea drilling depths (prediction 7) was accurate because it extrapolated from 2003 data. Instead of predicting innovation and utilization of new technologies, it instead predicted continued progress in drilling technology which was already being commercially used in 2003.

## **Sources of damage predictions**

Predictions concerning the sources of damage to deep-sea ecosystems were all accurate except for the prediction that sewage sludge would be dumped at sea (prediction 2). All of the sources of damage that Glover & Smith predicted were already damaging ecosystems in 2003. These predictions succeeded by recognizing that humans would continue their activities that harmed the deep-sea. Unlike new technologies which may face sudden regulatory changes and economic barriers, current industrial activities may be more resistant to change. Instead of predicting change in the deep-sea environment, the predictions about sources of damage anticipated a lack of change and were accurate. However, our analysis does reveal important nuances. The prediction that sewage sludge would be disposed of at deep-sea, a seemingly predictable trend, was inaccurate due to policy making at the 2002 London Convention. This highlights how policy can alter otherwise predictable and stable trajectories.

## **Effects of damage to ecosystems predictions**

The accuracy of predictions about ecosystem responses to human activities were mixed. The predictions about the declines of fisheries (predictions 5 and 6) succeeded because they were based on decades of fishing records and the known vulnerability of deep-sea ecosystems. The fisheries extinction prediction (prediction 4) presents an interesting case where the prediction would have been accurate if not for human intervention. Glover & Smith specifically mentioned the Orange Roughy in their article as a species that was rapidly decreasing in catch size. They acknowledged that "current management practices" were leading to its extinction but didn't predict that the management process would change (Glover & Smith, 2003). This prediction would have been accurate if humans did not intervene and save the Orange Roughy fishery. The predictions about the effects of drilling accidents and climate change (predictions 9 and 15) on

deep-sea ecosystems underestimated the magnitude of their damage. Prediction 15 most likely failed due to the unanticipated rapid acceleration of climate change in the past two decades.

### **Policy/Regulatory Predictions**

The accuracy of predictions about ecosystem responses to human activities were mixed. Glover & Smith's simpler predictions that countries will be interested in offshore mining ventures (predictions 11 and 12) were both accurate. They most likely were accurate because they merely predicted an expressed interest instead of systemic changes in the deep-sea. The prediction that oil companies would be mandated to acquire baseline survey data (prediction 8) was partially accurate since it does not apply to every company around the world. This reflects that policy decisions not made by an international organization will instead fall to regional managements that differ between locations. We cannot expect regulations to be the same around the world unless it is made by a global governing body. The prediction about dumping sewage sludge at deep-sea (prediction 2) proved to be inaccurate because of the IMO, a global governing body. Sewage dumping was banned to protect the deep-sea, reflecting that on top of leading innovation, policy making will also reverse old practices.

### **Study Limitations**

Our analysis faced several limitations. The sparse data environment of the deep-sea is an inherent challenge when determining accuracy. Some assumptions might be premature as more data is collected about the deep sea. Additionally, the ternary evaluation of predictions may oversimplify complex predictions and limit directionality of the interpretations. Future studies could enhance this analysis by developing more nuanced evaluatory metrics that account for partial success.

### **Conclusion**

This current evaluation of Glover & Smith's (2003) predictions about the 2025 deep-sea ecosystem reveals insights on the ability of long-term forecasting in a data-poor environment. Of the 15 predictions made, 7 were accurate, 4 were inaccurate, and 4 more were partially accurate. Overall, there was a mix of accurate, partially accurate, and inaccurate predictions with few correlations between result categories. Within the 5 broad categories of predictions that Glover and Smith used, there were a mix of accurate and inaccurate predictions. We created different categories of themes to find correlations between accuracy and type of prediction. Notably, predictions concerning technological innovation were largely inaccurate and current development lagged the predictions. Predictions on sources of damage to the deep-sea were mostly accurate. This analysis reveals the difficulty in making predictions about a region with sparse amounts of data. These findings underscore the need for enhanced data collection and improved methods for better forecasting the deep-sea ecosystem. This study also highlights the importance of integrating socioeconomic and regulatory factors in future predictions. By identifying the limitations and strengths of past predictions we can improve future predictions that will shape policy decisions and conservation processes.

## Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Author Contributions

The author confirms being the sole contributor to this work and has approved it for publication.

## Funding

Details of all funding sources should be provided, including grant numbers if applicable. Please ensure to add all necessary funding information, as after publication this is no longer possible.

## Acknowledgments

The author thanks all the editors involved in this issue and the reviewers for their insightful comments and feedback on this paper.

## References

- Abe, M., Maharroof, F., and Eldacrory, F. (2025). "Policy Implications in Deep-Sea Mining: Opportunities and Challenges for Kiribati. Kolonia: United Nations Multi-Country Office for Micronesia.
- Akhmetkaliyeva, S. (2020). Successful Mining of "Flammable Ice". *Eurasian Res. Inst.* Available at: <https://www.eurasian-research.org/publication/successful-mining-of-flammable-ice/> (Accessed July 23, 2025).
- Alberts, E. C. (2023). Deep-sea mining rules delayed two more years; mining start remains unclear. *Mongabay*. Available at: <https://news.mongabay.com/2023/07/deep-sea-mining-rules-delayed-two-more-years-mining-start-remains-unclear/>
- Allianz (2025). Safety and Shipping Review 2025. Allianz Commercial. Available at: <https://commercial.allianz.com/content/dam/onemarketing/commercial/commercial/reports/commercial-safety-shipping-review-2025.pdf>
- Aman, P., Sharma, G., and Sarika, K. (2025). Methane production from gas hydrates through CO<sub>2</sub> sequestration: A review. *Gas Sci. Eng.* 134, 205532. doi: 10.1016/j.jgsce.2024.205532
- Baker, J. A., Bell, M. J., Jackson, L. C., Vallis, G. K., Watson, A. J., and Wood, R. A. (2025). Continued Atlantic overturning circulation even under climate extremes. *Nature* 638, 987–994. doi: 10.1038/s41586-024-08544-0

- Bakke, T., Klungsøyr, J., and Sanni, S. (2013). Environmental impacts of produced water and drilling waste discharges from the Norwegian offshore petroleum industry. *Mar. Environ. Res.* 92, 154–169. doi: 10.1016/j.marenvres.2013.09.012
- Barron, M. G., Vivian, D. N., Heintz, R. A., and Yim, U. H. (2020). Long-Term Ecological Impacts from Oil Spills: Comparison of *Exxon Valdez*, *Hebei Spirit*, and Deepwater Horizon. *Environ. Sci. Technol.* 54, 6456–6467. doi: 10.1021/acs.est.9b05020
- Barry, J. P., Buck, K. R., Lovera, C. F., Kuhn, L., Whaling, P. J., Peltzer, E. T., et al. (2004). Effects of direct ocean CO<sub>2</sub> injection on deep-sea meiofauna. *J. Oceanogr.* 60, 759–766.
- Blake, J. A., Maciolek, N. J., Ota, A. Y., and Williams, I. P. (2009). Long-term benthic infaunal monitoring at a deep-ocean dredged material disposal site off Northern California. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 56, 1775–1803. doi: 10.1016/j.dsr2.2009.05.021
- BOEM (2025). Offshore Statistics by Water Depth. Available at: <https://www.data.boem.gov/Leasing/OffshoreStatsbyWD/Default.aspx> (Accessed July 23, 2025).
- Cannon, J. (2023). Pacific alliance adopts moratorium on deep-sea mining, halting resurgent PNG project. *Mongabay*. Available at: <https://news.mongabay.com/2023/09/pacific-alliance-adopts-moratorium-on-deep-sea-mining-halting-resurgent-png-project/>
- Carvajal-Arenas, L. C., Torrado, L., Ramos, J. P., and Mann, P. (2024). Is the Colombian Basin on its way to become the next big frontier? *GEO ExPro Mag.*
- Chen, C.-F., Chen, C.-W., Ju, Y.-R., Kao, C.-M., and Dong, C.-D. (2018). Impact of disposal of dredged material on sediment quality in the Kaohsiung Ocean Dredged Material Disposal Site, Taiwan. *Chemosphere* 191, 555–565. doi: 10.1016/j.chemosphere.2017.10.091
- Clark, M., and O'Driscoll, R. (2003). Deepwater fisheries and aspects of their impact on seamount habitat in New Zealand. *J. Northwest Atl. Fish. Sci.* 31, 441–458.
- Cordes, E. E., Jones, D. O. B., Schlacher, T. A., Amon, D. J., Bernardino, A. F., Brooke, S., et al. (2016). Environmental Impacts of the Deep-Water Oil and Gas Industry: A Review to Guide Management Strategies. *Front. Environ. Sci.* 4. doi: 10.3389/fenvs.2016.00058
- Cuvelier, D., Gollner, S., Jones, D. O. B., Kaiser, S., Arbizu, P. M., Menzel, L., et al. (2018). Potential Mitigation and Restoration Actions in Ecosystems Impacted by Seabed Mining. *Front. Mar. Sci.* 5. doi: 10.3389/fmars.2018.00467
- Giseburt, A. (2024). Japan prepares to mine its deep seabed by decade's end. *Mongabay*. Available at: <https://news.mongabay.com/2024/03/japan-prepares-to-mine-its-deep-seabed-by-decades-end/>
- Goethel, C. L., Grebmeier, J. M., and Cooper, L. W. (2019). Changes in abundance and biomass of the bivalve *Macoma calcaria* in the northern Bering Sea and the southeastern Chukchi Sea from 1998 to 2014, tracked through dynamic factor analysis models. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 162, 127–136. doi: 10.1016/j.dsr2.2018.10.007

- Gohlke, J. M., Doke, D., Tipre, M., Leader, M., and Fitzgerald, T. (2011). A Review of Seafood Safety after the Deepwater Horizon Blowout. *Environ. Health Perspect.* 119, 1062–1069. doi: 10.1289/ehp.1103507
- Goode, S. L., Rowden, A. A., Bowden, D. A., and Clark, M. R. (2020). Resilience of seamount benthic communities to trawling disturbance. *Mar. Environ. Res.* 161, 105086. doi: 10.1016/j.marenvres.2020.105086
- Hall–Spencer, J., Allain, V., and Fosså, J. H. (2002). Trawling damage to Northeast Atlantic ancient coral reefs. *Proc. R. Soc. Lond. B Biol. Sci.* 269, 507–511. doi: 10.1098/rspb.2001.1910
- Hu, S. (2025). Stop Offshore Drilling | NRDC. *NRDC*. Available at: <https://www.nrdc.org/stories/offshore-drilling-101> (Accessed July 25, 2025).
- Hutt, D. (2018). After the loss of a ship, deep sea mining plans for PNG founder. *Mongabay*. Available at: <https://news.mongabay.com/2018/12/after-the-loss-of-a-ship-deep-sea-mining-plans-for-png-founder/>
- Huvenne, V. A. I., Bett, B. J., Masson, D. G., Le Bas, T. P., and Wheeler, A. J. (2016). Effectiveness of a deep-sea cold-water coral Marine Protected Area, following eight years of fisheries closure. *Biol. Conserv.* 200, 60–69. doi: 10.1016/j.biocon.2016.05.030
- IMO (2022). Dumping of sewage sludge at sea to be prohibited worldwide. Available at: <https://www.imo.org/en/mediacentre/pressbriefings/pages/dumping-of-sewage-sludge-.aspx> (Accessed July 23, 2025).
- Intercargo (2016). Years 2005 to 2015 and the Trends. International Association of Dry Cargo Shipowners. Available at: <https://www.intercargo.org/wp-content/uploads/2016/03/casualty-report-2015-printing-version-v4.pdf>
- Intercargo (2025). Years 2015 to 2024 and Trends. International Association of Dry Cargo Shipowners. Available at: <https://www.intercargo.org/wp-content/casualty-report/2025/>
- Jackson, J. (2023). Deep-Sea Mining May Harm Thousands of Species—Before They Are Even Discovered | The Pew Charitable Trusts. Pew Charitable Trust. Available at: <https://www.pew.org/en/research-and-analysis/data-visualizations/2023/deep-sea-mining-may-harm-thousands-of-species-before-they-are-even-discovered> (Accessed July 23, 2025).
- Johnson, G. C., and Purkey, S. G. (2024). Refined Estimates of Global Ocean Deep and Abyssal Decadal Warming Trends. *Geophys. Res. Lett.* 51. doi: 10.1029/2024gl111229
- Katsanevakis, S., Levin, N., Coll, M., Giakoumi, S., Shkedi, D., Mackelworth, P., et al. (2015). Marine conservation challenges in an era of economic crisis and geopolitical instability: The case of the Mediterranean Sea. *Mar. Policy* 51, 31–39. doi: 10.1016/j.marpol.2014.07.013
- Keating-Bitonti, C. (2025). Seabed Mining in Areas Beyond National Jurisdiction: Issues for Congress. Available at: <https://www.congress.gov/crs-product/R47324> (Accessed July 23, 2025).

- Koslow, J. A., and Glowlett-Holmes, K. (1998). The seamount fauna off southern Tasmania: Benthic communities, their conservation and impacts of trawling. Environment Australia & The Fisheries Research and Development Corporation.
- Larsson, A. I., and Purser, A. (2011). Sedimentation on the cold-water coral *Lophelia pertusa*: Cleaning efficiency from natural sediments and drill cuttings. *Mar. Pollut. Bull.* 62, 1159–1168. doi: 10.1016/j.marpolbul.2011.03.041
- Letra, L. (2025). Statement at the thirtieth session of International Seabed Authority Council, Part II - International Seabed Authority., (Kingston, Jamaica: International Seabed Authority). Available at: <https://www.isa.org.jm/news/statement-at-the-thirtieth-session-of-international-seabed-authority-council-part-ii/> (Accessed July 23, 2025).
- Li, K.-Y., and Liu, W. (2025). Weakened Atlantic Meridional Overturning Circulation causes the historical North Atlantic Warming Hole. *Commun. Earth Environ.* 6. doi: 10.1038/s43247-025-02403-0
- Maribus gGmbH (2013). World Ocean Review 2: The Future of Fish – The Fisheries of the Future. Maribus gGmbH. Available at: <https://worldoceanreview.com/en/wor-2/fisheries/deep-sea-fishing/> (Accessed July 23, 2025).
- Mazor, T., Possingham, H. P., Edelist, D., Brokovich, E., and Kark, S. (2014). The Crowded Sea: Incorporating Multiple Marine Activities in Conservation Plans Can Significantly Alter Spatial Priorities. *PLoS ONE* 9, e104489. doi: 10.1371/journal.pone.0104489
- McVeigh, K. (2021). Deep-sea mining may push hundreds of species to extinction, researchers warn. *The Guardian*. Available at: <https://www.theguardian.com/environment/2021/dec/09/mining-may-push-hundreds-of-deep-sea-mollusc-species-to-extinction> (Accessed July 23, 2025).
- Moor, A., Jones, M., Wilson, I., Zhang, J., Lilkendey, J., D.L. Trip, E., et al. (2025). Age-based population dynamics of the four-rayed rattail *Coryphaenoides subserrulatus* from the Chatham Rise, New Zealand. *N. Z. J. Mar. Freshw. Res.*, 1–10. doi: 10.1080/00288330.2025.2488385
- MSC (2016). Orange Roughy: The extraordinary turnaround | Marine Stewardship Council. *Mar. Steward. Counc.* Available at: <https://www.msc.org/en-au/what-we-are-doing/msc-theory-of-change/sustainable-fishing/sustainable-fishery-examples/orange-roughy-the-extraordinary-turnaround> (Accessed July 23, 2025).
- Murawski, S. A., Grosell, M., Smith, C., Sutton, T., and Halanych, K. M. (2021). Impacts of Petroleum, Petroleum Components, and Dispersants on Organisms and Populations | Oceanography. doi: 10.5670/oceanog.2021.122
- OSPAR (2017). Dumping and Placement of Dredged Material. *OSPAR Comm.* Available at: <https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/pressures-human-activities/dumping-and-placement-dredged-material/> (Accessed July 23, 2025).



- Palanques, A., Guillén, J., Puig, P., and Durán, R. (2022). Effects of long-lasting massive dumping of dredged material on bottom sediment and water turbidity during port expansion works. *Ocean Coast. Manag.* 223, 106113. doi: 10.1016/j.ocecoaman.2022.106113
- Pant, M., Fleeger, J. W., Johnson, D. S., Riggio, R., Hou, A., Deis, D. R., et al. (2025). Recovery of Saltmarsh Macroinfauna After the Deepwater Horizon Oil Spill. *Estuaries Coasts* 48, 87. doi: 10.1007/s12237-025-01520-5
- Pasparakis, C., Esbaugh, A. J., Burggren, W., and Grosell, M. (2019). Impacts of deepwater horizon oil on fish. *Comp. Biochem. Physiol. Toxicol. Pharmacol. CBP* 224, 108558. doi: 10.1016/j.cbpc.2019.06.002
- Precedence (2025). Offshore Drilling Market Navigates Growth with Energy Demand and Deepwater Innovations. Precedence Research. Available at: <https://www.precedenceresearch.com/offshore-drilling-market> (Accessed July 25, 2025).
- Research Nester (2025). Offshore Drilling Rigs Market size to surpass \$183.9 billion by 2037 | 6.8% CAGR Forecast. Research Nester. Available at: <https://www.researchnester.com/reports/offshore-drilling-market/4895> (Accessed July 25, 2025).
- Ruhl, H. A. (2007). Abundance and size distribution dynamics of abyssal epibenthic megafauna in the northeast Pacific. *Ecology* 88, 1250–1262. doi: 10.1890/06-0890
- Ruhl, H. A., and Smith, K. L. (2004). Shifts in Deep-Sea Community Structure Linked to Climate and Food Supply. *Science* 305, 513–515. doi: 10.1126/science.1099759
- Rystad (2024). Upstream majors dive into deeper waters as exploration budgets stay flat in 2024. Rystad Energy. Available at: <https://www.rystadenergy.com/news/upstream-majors-drilling-production-oil-gas-exploration> (Accessed July 25, 2025).
- Schwing, P. T., Montagna, P. A., Joye, S. B., Paris, C. B., Cordes, E. E., McClain, C. R., et al. (2020). A Synthesis of Deep Benthic Faunal Impacts and Resilience Following the Deepwater Horizon Oil Spill. *Front. Mar. Sci.* 7. doi: 10.3389/fmars.2020.560012
- Shaughnessy, J., Daugherty, W., Graff, R., and Durkee, T. (2007). More Ultradeepwater Drilling Problems., in *SPE/IADC Drilling Conference*, (Amsterdam, The Netherlands: SPE). doi: 10.2118/105792-ms
- Southall, B. L., Bowles, A. E., Ellison, W. T., Finneran, J. J., Gentry, R. L., Greene, C. R., et al. (2008). Marine mammal noise-exposure criteria: initial scientific recommendations. *Bioacoustics* 17, 273–275. doi: 10.1080/09524622.2008.9753846
- Stronkhorst, J., Ariese, F., van Hattum, B., Postma, J. F., de Kluijver, M., Den Besten, P. J., et al. (2003). Environmental impact and recovery at two dumping sites for dredged material in the North Sea. *Environ. Pollut.* 124, 17–31. doi: 10.1016/S0269-7491(02)00430-X

- Svensson, N., Norén, A., Modin, O., Karlfeldt Fedje, K., Rauch, S., Strömvall, A.-M., et al. (2022). Integrated cost and environmental impact assessment of management options for dredged sediment. *Waste Manag.* 138, 30–40. doi: 10.1016/j.wasman.2021.11.031
- Thomas, E. A., Böhm, M., Pollock, C., Chen, C., Seddon, M., and Sigwart, J. D. (2022). Assessing the extinction risk of insular, understudied marine species. *Conserv. Biol.* 36, e13854. doi: 10.1111/cobi.13854
- US Department of Commerce, N. O. and A. A. (2014). NOAA Ocean Explorer: Deep East 2001. Available at: <https://oceanexplorer.noaa.gov/explorations/deepeast01/background/dumping/dumping.html> (Accessed July 23, 2025).
- Watling, L. (2014). Trawling exerts big impacts on small beasts. *Proc. Natl. Acad. Sci.* 111, 8704–8705. doi: 10.1073/pnas.1407305111
- Yiallourides, C. (2019). Offshore Methane Hydrates in Japan: Prospects, Challenges and the Law. *Br. Inst. Int. Comp. Law*. Available at: <https://www.biicl.org/projects/offshore-methane-hydrates-in-japan-prospects-challenges-and-the-law> (Accessed July 23, 2025).
- Yin, Z., and Linga, P. (2019). Methane hydrates: A future clean energy resource. *Chin. J. Chem. Eng.* 27, 2026–2036. doi: 10.1016/j.cjche.2019.01.005
- Yip, T. L., Talley, W. K., and Jin, D. (2011). The effectiveness of double hulls in reducing vessel-accident oil spillage. *Mar. Pollut. Bull.* 62, 2427–2432. doi: 10.1016/j.marpolbul.2011.08.026
- Zhengzheng, H. (2017). China's first gas hydrate extraction successful. *CGTN*. Available at: [https://news.cgtn.com/news/3d67544f786b7a4d/share\\_p.html](https://news.cgtn.com/news/3d67544f786b7a4d/share_p.html) (Accessed July 23, 2025).
- Zhou, H., Abdul, K. M., Yu, Q., Liu, X., Hua, E., Lou, Y., et al. (2023). Detection of global change with meiofauna and its benthic environment in a shelf sea cold water mass ecosystem. *Reg. Stud. Mar. Sci.* 62, 102918. doi: 10.1016/j.rsma.2023.102918